

Research

Plasma oxidized low density lipoprotein cholesterol correlates inversely with testosterone in young adult male smokers

Maria Onomhaguan Ebesunun^{1,&}, Olurakinwo Lanre Bankole¹, Olayiwola Oduwole²

¹Chemical Pathology& Immunology, Obafemi Awolowo College of Health Sciences Olabisi Onabanjo,University, Sagamu Campus, Nigeria, ²Institute of Reproductive and Developmental Biology, Imperial College London, Faculty of Medicine, Hammersmith Hospital Campus, London W12 0NN

[&]Corresponding author: Maria Onomhaguan Ebesunun, Department of Chemical Pathology & Immunology, Obafemi Awolowo College of Health Sciences, Olabisi Onabanjo University,Sagamu Campus, Nigeria

Key words: Cigarette, CVD, Ox-LDL, testosterone, cholesterol

Received: 08/09/2013 - Accepted: 07/05/2014 - Published: 04/11/2014

Abstract

Introduction: there are indications that oxidized low density lipoprotein cholesterol (Ox-LDL) may play an important role in cardiovascular disease (CVD) events. In most developing countries, the interplay between the different lipid fractions and cigarette smoking has not been studied. This study assessed the effect of cigarette smoking on the alterations in plasma lipid fractions and their associations with the gonadal hormone, testosterone (T). **Methods:** one hundred and sixty male participants, consisting of eighty smokers and eighty apparently healthy non-smokers were recruited. Anthropometric indices and biochemical parameters were determined using standard procedures. **Results:** significant increases were obtained in plasma total cholesterol (TC), triglyceride (TG), oxidized low density lipoprotein (Ox-LDL) and Ox-LDL/TT ratio (p. **Conclusion:** this study showed an inverse correlation between Ox-LDL and testosterone as well as strong association between the number of tobacco and cigarettes usage per day. These changes in part, could be major causes of premature CVD and decreased fertility in young adults.

Pan African Medical Journal. 2014; 19:241 doi:10.11604/pamj.2014.19.241.3354

This article is available online at: <http://www.panafrican-med-journal.com/content/article/19/241/full/>

© Maria Onomhaguan Ebesunun et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

The health consequences of smoking cigarettes and other tobacco related products are well known. They are important major health hazard in the development of atherosclerosis and cardiovascular disease (CVD) [1]. The connections between cigarette smoking and CVD have been addressed by several investigators [2-4]. In the INTERHEART study, Frick et al [5] indicated an overall population attributable risk of about 38% between smoking and myocardial infarction. Tobacco smoke in general contains numerous compounds majorly among which are nicotine, carbon monoxide and various carcinogens [3]. These compounds do have effects on the metabolic and biological processes of the body including the modification of lipoproteins, especially the LDLC [6], the oxidized form of which has been shown to accumulate excessively in arterial walls, leading to increased peroxidation both in vitro and in vivo [7]. Recent experimental and clinical data also suggest that exposure to cigarette and other tobacco related smoke increases oxidative stress, thereby compounding the potential mechanism(s) for initiating cardiovascular dysfunction through the activities of nicotine in cigarettes which enhances fat oxidation [8]. Experimental and clinical data [9] indicates that nicotine in cigarette smoke changes lipoproteins, particularly LDLC into the oxidized form(ox-LDL) thus quickening its uptake by macrophages, thereby leading to the progression of atherosclerosis [3]. There is also a growing interest following several studies on the link between cigarette smoking and low testosterone levels in men in relation to the development of coronary heart disease[10,11]. Overall, males have been shown to have a higher risk of CVD in comparison to premenopausal women, with the incidence in women increasing after menopause and a concomitant decrease in the gender difference in mortality and morbidity [12, 13]. In a developing country like Nigeria, where there is not much enlightenment on the implications and adverse effects of smoking cigarette and other tobacco related products are not known, a sizeable number of the male population are involved in this habit. The aim of this study therefore, was to ascertain the relationship between smoking, the different fractions of lipids and gonadal hormone. The results we believe will highlight the relationship between these parameters in our targeted subjects.

Methods

One hundred and sixty apparently healthy male participants consisting of eighty smokers (44.4±1.7 years) and eighty non-smokers (35.1±1.0 years) were recruited for this study. The smokers were recruited from motor parks and military barracks; areas where the habit is a common occurrence. The smokers consumed at least 4 sticks of cigarette daily. Demographic data such as lifestyle pattern, smoking habit, alcohol use, socioeconomic status and medical history were obtained from each participant based on questionnaire adapted from the work of Ebesun et al, [14]. The controls were men who had never been involved in cigarette smoking at any stage of their life. Informed consent was also obtained from participants before commencement of study.

Anthropometric measurements

The weight of each participant in kilogram was taken using a Seca adult weighing scale. The height in centimeter was taken using a fixed meter rule without foot wear. The body mass index (BMI) in kg/m² of each subject was calculated using weight/height²

Sample collection

Blood samples were collected from all subjects after an overnight fast (10-12 hours) into EDTA bottles. The blood samples were immediately placed on ice bag and plasma was separated within short time of collection. The plasma samples were thereafter stored in at -20° C until analyzed for biochemical parameters.

Biochemical analysis

Total cholesterol was determined using the method of Alain et al, [15] while the triglyceride was estimated by the enzymatic method of David and Buccolo [16]. LDLC and VLDLC were precipitated out by the addition of phosphotungstic acid in the presence of magnesium ions. After centrifugation, the cholesterol concentration in the supernatant representing the HDLC fraction was determined with the method used for total cholesterol estimation. The LDLC was calculated using the formula of Friedwald et al, [17]. Ox-LDLC was analyzed using an Enzyme Immunoabsorbent Assay (EIA) method based on the principle of competitive binding between Ox-LDLC in the test sample and oxidized LDL-HRP conjugate for a constant amount of rabbit anti-oxidized LDLC. Testosterone analysis was based on the principle of competitive binding between T in the test sample and testosterone-HRP conjugate for a constant amount of rabbit anti-testosterone. Accuracy and precision of biochemical tests were monitored by the inclusion of commercial quality control samples with each batch of assay.

Statistical analysis

All results were subjected to statistical analysis using SPSS version 14 software (SPSS Inc. Chicago, Illinois) for windows. Student t-test was used for comparison of variables. Post Hoc analysis was also carried out. Pearson correlation coefficient was used to assess the relationship between variables. The results were expressed in mean ± S.E and the level of significance was accepted at p < 0.05.

Results

Results are presented in tables 1-3 below: **Table 1** shows biophysical and biochemical parameters in smokers and non-smokers. Smokers were older than non-smokers (p

Table 2 shows the biophysical and biochemical parameters in smokers based on the duration (years) of cigarettes smoked. These were made up of two groups; group 1 consisted of smokers that have been smoking between 5 to 13 years, while group 2 consisted of those that have smoked between 14 to 22 years. Smokers in group 1 were young than smokers in group2 (p0.001) were not significantly different in the two groups. Comparison of parameters between military men and motor park vendors did not show any significant changes between the groups.

Table 3 shows Pearson correlation (r) coefficient, this indicated that plasma TC was significantly correlated with HDLC (r=0.958, p \leq 0.001). **Figure 1** shows a graph of Ox-LDLC in smokers and non-smokers. The smokers showed significantly higher value of plasma Ox-LDLC. **Figure 2** shows a graph of testosterone in smokers and non-smokers. The smoker had a lower mean plasma testosterone.

Discussion

A significant and clinically relevant finding in this study was that the plasma level of Ox-LDL was significantly increased in smokers and the level of significance was associated with increasing number of cigarettes and tobacco usage per day. Furthermore, this study showed a low level of mean total testosterone in smokers although this increase was not statistically significant, but was inversely correlated with Ox-LDL. Thus suggesting that as the Ox-LDL increases plasma T decreases. The possible reason for this is that after receptor-mediated endocytosis of LDL, cholesterol removed from LDL is used for diverse metabolic processes required for healthy functioning cells including the synthesis of testosterone. However, modified/oxidized LDL is not recognised by the receptor and instead is taken up by scavenger receptor macrophages. The uptake of Ox-LDL does not lead to its degradation. Instead it accumulates in the macrophages as cholesteryl esters, therefore foam cells and these amass in the sub-endothelial space, contributing to atherosclerotic plaques [18]. We could not demonstrate whether the low mean plasma testosterone proceeds or were a direct effect of smoking. More pronounced responses regarding hormones are found in most cases with heavy smokers. Since most of the subjects are light smokers, this could be the cause of the variance found between this study and those of others [19-21].

An epidemiology study in the North Eastern Nigeria [22] indicated that smoking is common among the low income earners. This was also revealed from the questionnaire on socioeconomic status used in this study. The reason for this is not known, but we speculate that since the majority of our subjects are motor park vendors/military men who had little or no education, and therefore do not have any idea on the pathophysiology of smoking. The possible mechanism linking low testosterone and elevated Ox-LDL in smokers is plausible. Smoking is known to be a modifier of the male reproductive hormones, T [23,24]. Its association with impaired lipid profile could be a major factor that promotes diseases like atherosclerosis and CVD especially in heavy smokers. Ox-LDL is increasingly formed in smokers possibly, as a result of the reaction between the lipid fractions in LDL particles and the reactive oxygen species generated by the toxicants in cigarette and tobacco smoke [9, 25]. Cigarette and tobacco smoke are established source of free radicals generation [2] which increases oxidative stress and other associated complications like endothelium damage and lipid oxidation. A high level of Ox-LDL in the body therefore, is deleterious. According to previous studies [26,27], Ox-LDL is also a major risk factor associated with atherosclerotic plaque formation. Significant and higher levels of plasma total cholesterol, triglyceride, and LDL were obtained in this study. Several suggestions have been put forward; among which are that the effect of nicotine in the cigarette and tobacco smoke causes lipolysis through stimulation of catecholamine, and therefore increasing cholesterol, TG, free fatty acids and LDL [20, 2]. Changes in lipid profile in smokers and the risk of increased tobacco related diseases have also been indicated to be dose dependent [28] and may as well be related to the duration of the habit. The role of dietary differences between smokers and non-smokers as well as free radical-mediated oxidative stress as a pivotal step for the development of atherosclerosis could also be relevant.

This study also demonstrated that smokers have significantly lower levels of plasma HDL than the controls subjects. The reason for this low level is not clear; it may possibly be a pointer to a significant cardiovascular risk. Available evidence indicates that for every milligram decrease in plasma HDL due to cigarette smoking

or any other factors, there is a 2.5% chance of increased premature CVD development [29].

Conclusion

The evidence from this study suggests a strong relationship between increased plasma Ox-LDL and number of tobacco/cigarettes usage per day. The inverse correlation between testosterone and Ox-LDL obtained in this study indicates that increases in Ox-LDL can lead to reduce plasma testosterone in smokers. These changes in part, could lead to decreased fertility and premature CVD events in young adults.

Competing interests

Authors declared no competing interests.

Authors' contributions

MO was involved in the conception, design, supervision of sample collection, acquisition of data and drafted the manuscript for publication. OLB was involved in the collection, analysis and acquisition of data as well as writing up of the manuscript for academic purpose. LO was involved in the original concept of the study and drafting of the manuscript for publication. All authors read and approved the final version of the manuscript.

Acknowledgments

We wish to acknowledge Margaret Solomon who was kind enough to assist in the purchase of the oxidized LDL kit. Mrs Doreen Okeke for the excellent technical assistance. We are also grateful to all the volunteers who willingly participated in this study

Tables

Table 1: biophysical and biochemical parameters of smokers and non-smokers (Mean + S.E.M)

Table 2: biophysical and biochemical parameters of smokers based on the duration (in years) of cigarette smoked. (Mean + S.E.M)

Table 3: Pearson correlation coefficient of biochemical parameters in smokers

Figure 1: oxidised-low density lipoprotein cholesterol in smokers and non smokers

Figure 2: total testosterone in smokers and non smokers

References

1. WHO. Why is tobacco a public health priority? Tobacco free Initiative. Retrieved 2010. <http://www.who.int/tobacco/about/en/index.html>. Accessed on September 23. **Google Scholar**

2. Price F, Mowbray PI, Lee AJ, Rumley R, Lowe GD, Fowkes FG. Relationship between smoking and cardiovascular risk factors in the development of peripheral arterial disease and coronary artery disease: Edinburgh Artery study. *European Heart Journal*.1999; 20(5): 344-353. **PubMed | Google Scholar**
3. Ebesunun MO, Adetunji RA, Umahion KO. Plasma vitamin C; Bétacarotene, lipids and lipoproteins in young adult smokers and smokers. *British Journal of Pharmacology and Toxicology*. 2011; 2(2): 71-74. **PubMed | Google Scholar**
4. Teo KK, Ounpuu S, Hawken S, Pandey MR. Tobacco use and risk of myocardial infarction in 52 countries in the INTERHEART study: a case-control study. *Lancet*. 19 august 2006; 368 (9536):647-58. **PubMed | Google Scholar**
5. Frick, EB, Bittar RE, Carvalho MH, Zugaib M. Impact of plasma oxidized low- density ipoprotein removal on atherosclerosis. *Circulation*. 2008; 118(1):75-6 83. **PubMed | Google Scholar**
6. Black HR, Laragh BG, Brenner BM. Smoking and cardiovascular disease; Hypertension: Pathophysiology, Diagnosis and Management. 2nd edition. New York, NY: Raven Press Ltd; 1995;2621-2647. **PubMed | Google Scholar**
7. Tang J, Morris JK, Wald NJ, Hole D, Shipley M, Tunstall Pedoe H. Mortality in relation to tar yield of cigarettes: a prospective study of four cohorts. *BMJ*. 1995;311(7019): 1530-1533. **PubMed | Google Scholar**
8. Department of Health and Human Services. The Health consequences of smoking and nicotine addiction: A report of the Surgeon General. DHHS. 1998; Pub no:(CDC) 88-8406. **Google Scholar**
9. Andersson, K, Arner P. Systemic nicotine stimulates human adiposetissuelipolysis through local cholinergic and catecholaminergic receptors. *International Journal of Obesity Related Metabolic Disorder*. 2000; 25(8):1225-32. **PubMed | Google Scholar**
10. John A, Ambrose, Rajat S, Barua. The pathophysiology of cigarette smoking and cardiovascular disease. *Journal of American College of Cardiology*. 2004; 43(10):1731. **PubMed | Google Scholar**
11. Brovarri JP, Lauwerys RS, Roels HE. Association between testosterone level sand smoking in men. *JAMA*. 1999; 336(10):699-702. **PubMed | Google Scholar**
12. Juul EM, Carlson SG, KerniceMJ. Associations between low total testosterone levels and oxidized-LDL. *Advances in Nephrology Necker Hospital*. 2008; 19(2):9-38. **PubMed | Google Scholar**
13. Hamaya BH, Coustegi FT. Mehanisms of lipid peroxidation ; a textbook of Cell Metabolism. 2000; 5th ed. Argy II publishing , U.K. : 2207-33. **Google Scholar**
14. Ebesunun MO, Agbedana EO, Taylor GOL, Oladapo OO. Plasma lipids lipoprotein (a),homocysteine and other cardiovascular disease risk factors in Nigerians with CVD. *App Physiol Nutri Metab*. 2008; 33(2):282-289. **PubMed | Google Scholar**
15. Allain CC, Poon LS, Chan CSG, Richmond W, Fu PC. Enzymatic determination of total serum cholesterol. *Clinical Chemistry*. 1974; 20(4): 470-3. **PubMed | Google Scholar**
16. Buccolo G, David H. Quantitative determination of serum triglycerides by the use of enzymes. *Clinical Chemistry*. 1973; 19(5): 476-82. **PubMed | Google Scholar**
17. Friedwald WT, Levy IR, Fredrickson DS. A method for estimating LDL-C without centrifugation. *Clinical Chemistry*. 1972;18(3):499 - 502. **PubMed | Google Scholar**
18. Asem N, Asem BB. The Many Faces of Cholesterol: How Modifications in LDL and HDL Alter Their Potential to Promote or Prevent Atherosclerosis. *The UCLA USJ*. 2009; 22: 11-17. **PubMed | Google Scholar**
19. Barud W, Palusinski R, Beltowski J, Wójcicka G. Inverse relationship between total testosterone and anti-oxidized low density lipoprotein antibody levels in ageing males. *Atherosclerosis*. 2002; 164:(2)283-288. **PubMed | Google Scholar**
20. BriggsMS, Goldstem JC. A regular pathway for testosterone. *International Journal of Obesity Related Metabolic Disorders*. 2003; 25(8):1225-32. **PubMed | Google Scholar**
21. Willet S, Beckley. Association between smoking and testosterone levelsin men. *American Journal Obstectic and Gynecology*. 1994; 117(5): 1047-1051. **PubMed | Google Scholar**
22. Ayankogbe OO, Inem OA, Bamgbola OA. Attitudes and determinant of cigarette smoking among rural dwellers in South West Nigeria. *Nigeria Medical Practitioner*. 2003; 44(1): 70-74. **PubMed | Google Scholar**
23. Shoji T, Nishizawa Y, Fukumoto M, Shimamura K, Kimura J. Inverse relationship between circulating oxidized low density lipoprotein (ox-LDL) and anti-oxLDL antibody levels in healthy subjects. *Atherosclerosis*. 2000;148 (1): 171-177. **PubMed | Google Scholar**
24. Briggs MS, Goldstem JC. A regular pathway for testosterone. *International Journal of Obesity Related Metabolic Disorders*. 2003;25(8): 1225-32. **PubMed | Google Scholar**
25. Odey FA, Okokon IB, Ogeeche JO, JT, Ekanem EE. Prevalence of cigarette smoking among adolescents in Calabar city, south-eastern Nigeria. *Journal of Medicine and Medical Sciences*. 2012; 3(4):237-242. **PubMed | Google Scholar**
26. Kapoor D, Jone TH. Smoking and hormones in health and endocrine disorders. *European Journal of Endocrinology*. 2005; 152(4):491-499. **PubMed | Google Scholar**
27. Venturi S, Donati FM, Gareeboo H, Roussety F . Ethnic differences inischamic heart disease and stroke mortality in Mauritius between 1989 and 1994. *Ethn Health*. 1998;3 (1-2):45-54. **PubMed | Google Scholar**
28. Craig WY, Palomaki GE, Haddow JE. Cigarette smoking and serum lipid and lipoprotein concentrations: an analysis of published data. *British Medical Journal*. 1989; 298(1):784-788. **PubMed | Google Scholar**

29. Goldbourt U, Yaani SM, Mansley J H. Isolated low HDL cholesterol as a risk factor for coronary heart disease mortality: a 21 year follows up of 8000 men. *Atherosclerosis Thromb Vascular Biology*.1997; 17(1):107-13. **PubMed | Google Scholar**

Table 1: biophysical and biochemical parameters of smokers and non-smokers(Mean+ S.E.M)

Variables	Smokers n=80	Non-smokers n=80	t-value	p-value
BMI (kg/m ²)	21.3±3.6	22.4±2.8	.86	NS
TC(mg/dl)	211.7±1.3	174.4±2.6	-12.7	p<0.001
TG (mg/d)	129.9±1.2	115.2±2.4	-5.6	p<0.001
HDLC(mg/dl)	40.2±0.8	47.1±0.7	6.8	p<0.001
LDLC(mg/dl)	143.7±1.8	102.6±2.8	-12.3	p<0.001
Ox-LDL (ng/dl)	65.4±0.7	44.4±0.4	-19.8	p<0.001
TT (ng/dl)	4.7±0.1	4.9±0.1	.73	NS
LDLC/Ox-LDL	13.9±0.1	9.1±0.1	.106	NS
Ox-LDL/TT	13.9±0.1	9.1±0.1	27.1	p<0.001
Age (years)	44.4±1.7	35.1±1.0	-11.2	p<0.001

BMI=Body Mass Index. TC=Total Cholesterol. TG=Triglyceride.
 TT =TotalTestosterone HDLC=High Density Lipoprotein Cholesterol.
 LDLC= Low DensityLipoprotein Cholesterol. Ox-LDL=Oxidized LDLC.
 p= Level of significance. NS= not significant.

Table 2: biophysical and biochemical parameters of smokers based on the duration (in years) of cigarette smoked. (Mean + S.E.M)


Variables	GP1=5-13years(n=35)	GP2=14-22 years (n=45)	t- value	p-value
BMI (kg/m ²)	22.3±3.6	21.7±2.8	10.2	NS
TC (mg/dl)	206.4±1.5	214.3±2.6	-9.7	p<0.001
TG (mg/dl)	126.1±1.2	132.8±2.4	-8.6	p<0.001
HDLC (mg/dl)	39.8±0.8	36.4±0.7	11.8	p<0.001
LDLC (mg/dl)	132.7±1.8	157.2±2.8	-18.3	p<0.001
Ox-LDL (ng/dl)	55.6±0.7	67.4±0.4	-19.8	p<0.001
TT (ng/dl)	4.5±0.1	4.7±0.1	5.3	NS
LDLC/Ox-LDL	2.4±0.1	2.3±0.1	9.6	NS
Ox-LDL/TT	12.9±0.1	14.3±0.1	-24.1	p<0.001
Age (years)	32.8±1.7	39.1±1.0	-16.8	p<0.001

BMI=Body Mass Index. TC=Total Cholesterol. TG=Triglyceride.
 TT =TotalTestosterone HDLC=High Density Lipoprotein Cholesterol.
 LDLC= Low DensityLipoprotein Cholesterol. Ox-LDL=Oxidized LDLC.
 GP= group.p = Level of significance. NS= not significant.

Table 3: Pearson correlation coefficient of biochemical parameters in smokers


Variables	TC (mg/dl)	HDLC (mg/dl)	LDLC(mg/dl)	Ox-LDLC(ng/dl)	TT (ng/dl)
TC (mg/dl)		.958**	.989**	.974**	.632*
HDLC (mg/dl)	.958**		-.975**	.955**	.577*
LDLC (mg/dl)	.989**	-.957**		.981**	.615*
Ox-LDLC (ng/dl)	.974	.955**	.981**		-.649*
TT (ng/dl)	.632*	.577*	.615*	-.649*	

Pearson Correlation coefficient significant at **p<0.01
* P<0.05
TT =Total Testosterone HDLC=High Density Lipoprotein Cholesterol.
LDLC= Low Density Lipoprotein Cholesterol. Ox-LDLC=Oxidized LDLC



CI= Confidence Interval at 95%.

Figure 1: oxidised-low density lipoprotein cholesterol in smokers and non smokers



CI= Confidence Interval at 95%.

Figure 2: total testosterone in smokers and non smokers