



ARTICLE

The Effect of a Chitosan/TiO₂-Nanoparticle/ Rosmarinic Acid-Based Nanocomposite Coating on the Preservation of Refrigerated Rainbow Trout Fillets (*Oncorhynchus mykiss*)

Pınar Kizilkaya^{1,*} and Mükerrerem Kaya²

¹Department of Food Technology, Ardahan Vocational School of Technical Sciences, Ardahan University, Ardahan 75002, Turkey

²Department of Food Engineering, Faculty of Agriculture, Atatürk University, Erzurum 25240, Turkey

OPEN ACCESS

Received June 1, 2023

Revised August 9, 2023

Accepted August 10, 2023

*Corresponding author : Pınar Kizilkaya
Department of Food Technology, Ardahan
Vocational School of Technical Sciences,
Ardahan University, Ardahan 75002, Turkey
Tel: +90-5077893903
E-mail: pinarkizilkaya@ardahan.edu.tr

*ORCID
Pınar Kizilkaya
<https://orcid.org/0000-0002-8420-7222>
Mükerrerem Kaya
<https://orcid.org/0000-0001-6340-828X>

Abstract The aim of this study was to determine the effect of chitosan (CH)-based nanocomposite coating applications [chitosan+TiO₂ (CHT) and chitosan+TiO₂+rosmarinic acid (CHTRA)] on changes in quality attributes of rainbow trout fillets during cold storage (4°C). Fish fillets were randomly divided into four groups and subjected to treatments (CH, CHT, CHTRA, and control). After treatments, the groups were packaged under a modified atmosphere (40% CO₂+30% O₂+30% N₂) and stored at 4°C for 18 days. During cold storage, the samples were subjected to physico-chemical and microbiological analyses. During storage, CH, CHT, and CHTRA treatments showed lower aerobic mesophilic and psychrotrophic bacteria counts than the control. However, the differences between coating treatments were not significant. The highest mean pH value was determined in the control group. As the storage time increased, the thiobarbituric acid reactive substances value increased. At the end of the storage period, no significant differences were observed between the treatments, including in the control group. The total volatile basic nitrogen (TVB-N) level in the control group was above 25 mg/100 g on day 15 of storage. However, the TVB-N level in the treatment groups was below 20 mg/100 g on day 18. It was also determined that coating application×storage period interaction had a significant effect on all color parameters (p<0.01). At the end of storage, the highest CIE L* was observed in CHTRA treatment. However, the value of this treatment did not differ from that of the CH treatment.

Keywords chitosan, nanocomposite coating, rosmarinic acid, TiO₂, trout fillet

Introduction

The use of biomaterials in food packaging has recently attracted attention because of health and environmental concerns. The rising cost of petroleum products and the increasing environmental damage caused by their use as packaging materials has led to

a growing interest in alternative materials that can be used instead of petroleum-derived materials (Salimiraad et al., 2022). Biopolymers are one of the popular alternative materials (Rahman et al., 2021). These biodegradable packaging materials can include proteins, lipids, polysaccharides and their combinations (Zabihollahi et al., 2020). Chitosan (CH) is a commonly used carbohydrate-derived biodegradable polymer (Wang et al., 2021). CH, obtained from deacetylated chitin, is a polysaccharide composed of glucosamine and N-acetylglucosamine copolymer (Jiang et al., 2022). The fact that CH is non-toxic, biodegradable, biocompatible (Ambaye et al., 2022; Bento et al., 2020), low-cost, sustainable, and renewable makes it increasingly researched (Silva et al., 2021). Also, CH shows broad-spectrum antimicrobial activity (bacteria, yeast, and molds; Yu et al., 2021). Moreover, CH is commonly used in food preservation and packaging owing to its good film-forming properties. An important way to develop many features (mechanical, barrier, antimicrobial, and antioxidant, etc.) of CH-based films and coatings is to combine CH with different organic or inorganic materials (Qu and Luo, 2021).

Despite the development of new food processing techniques that improve food quality, microbial contamination remains a major safety concern for all foods (Nwabor et al., 2020). With modern food processing techniques, a targeted reduction in germs can be achieved during production. However, post-production contamination is still the main factor in microbial food deterioration. Therefore, edible antimicrobial packaging (films or coatings) is of great importance for preventing microbial spoilage (Kumar et al., 2020).

Edible packaging is a biopolymer that can be produced and developed from renewable materials (such as polysaccharides and proteins; Hoque et al., 2021). However, biopolymeric films show poor mechanical and barrier properties. Nanocomposites are a new material class and possess at least one nanoscale size. They have become important in the development of the physico-mechanical and thermal features of these films (Hosseini et al., 2022; Nath et al., 2022). Titanium dioxide (TiO₂) nanoparticles are interesting substances utilized in the production of nanocomposite films and coatings (He et al., 2016). TiO₂ is a semiconductor metal oxide that is considered a promising material because of its chemical stability, low toxicity, and low cost (Jovanović et al., 2015; Lin et al., 2015). The addition of phenolic substances to these environmentally friendly materials can provide new properties for packaging materials (Heydari-Majd et al., 2019; Nath et al., 2022).

One of the most efficient, naturally water-soluble phenolic co-pigments, rosmarinic acid (Zhao et al., 2021), is an ester of caffeic acid and 3,4-dihydroxy phenyl lactic acid (Petersen and Simmonds, 2003) and lipids. It is of interest to the food industry as a natural antioxidant and antibacterial agent (Marchev et al., 2021). It has been reported that rosmarinic acid can be used mainly in the manufacture of nanocomposite packaging (Sani et al., 2017).

Rainbow trout (*Oncorhynchus mykiss*) plays an important role in human nutrition because of its high protein and omega-3 fatty acid contents. Therefore, it is highly valued in the market and often sold as fresh fillets (Hosseini et al., 2022). However, rainbow trout and other seafood products are also susceptible to microbiological and chemical deterioration due to their high water activity and pH values, free amino acids, and polyunsaturated fatty acids (Volpe et al., 2015).

The aim of this study was to determine the effects of adding nanoparticles and phenolic substances to edible coatings on the quality of cold-stored rainbow trout under a modified atmosphere. For this purpose, rainbow trout fillets were applied five treatments [control, CH, chitosan+TiO₂ (CHT), and chitosan+TiO₂+rosmarinic acid (CHTRA)]. After these treatments, rainbow trout fillets were packaged under a modified (40% CO₂+30% O₂+30% N₂) atmosphere. During cold storage (4°C) for 18 days, the samples were subjected to physico-chemical [(pH, thiobarbituric acid reactive substances (TBARS), total volatile basic nitrogen (TVB-N), instrumental color parameters (CIE L*, CIE a*, and CIE b*)], and microbiological analyses [total aerobic mesophilic bacteria (TMAB) and psychrotrophic bacteria].

Materials and Methods

Materials

CH with a deacetylation degree of 75%–85% and medium molecular weight (Sigma-Aldrich, St. Louis, MO, USA), glycerol (85%, Merck, Darmstadt, Germany), acetic acid (100%, Merck), and rosmarinic acid (HPLC-grade, purity \geq 98%; Sigma-Aldrich) were utilized for the preparation of edible coatings. Titanium dioxide nanoparticles (TiO₂) with a 30–50 nm particle size and high purity (99%) were acquired from a nanotechnological products company (Nanografi, Ankara, Turkey). The packaging material employed was Polyamide/Polyethylene (PA/PE) bags (15×25 cm, 3- seal bags GB 70) obtained from Südpack Verpackungen (Ochsenhausen, Germany) company, with an oxygen permeability of 40 cm³/m²/day.atm. at 23°C, nitrogen permeability of 24 cm³/m²/day.atm. at 23°C, carbon dioxide permeability of 145 cm³/m²/day.atm. at 23°C, and water vapor permeability of <3 g/m²/day.atm. at 23°C.

Preparation of CH coating and nanocomposite coatings to apply the composite and nanocomposite coating materials, 168 skinned fillets were obtained from 84 rainbow trout (*O. mykiss*), weighing an average of 350–400 g, sourced from the Atatürk University Faculty of Fisheries Application Center (Erzurum, Türkiye).

Preparation of chitosan (CH) coating and nanocomposite coatings

In this study, three different coating materials were prepared: CH (2% CH suspension; CH), CH (2% CH suspension) with TiO₂ nanoparticles (1.5%; CHT), and CH (2% CH suspension) with TiO₂ nanoparticles (1.5%) and 5 ppm rosmarinic acid (CHTRA).

The CH solution was prepared using a modified solvent-casting method described by Kanmani and Rhim (2014) and Nowzari et al. (2013). A 2% CH solution was prepared by dissolving CH in 1% (v/v) acetic acid. To ensure complete dissolution, the solution was stirred for 24 h at 50°C using a magnetic stirrer (MSH-20, Daihan Scientific, Wonju, Korea). After 24 h of mixing, 1% (v/v) glycerol was added as a plasticizer and the solution was stirred for an additional 6 h. TiO₂ nanoparticles (10 mg/L) and/or rosmarinic acid (0.005 mg/mL) previously prepared using ultrasound (BANDELIN electronic, Berlin, Germany) were slowly added to the coating solutions. The mixture was stirred at 24,000 rpm with a homogenizer (Ultra-Turrax T25, IKA T25, IKA-Werke, Staufen, Germany) during the addition process, followed by an additional 20 min of stirring. The nanocomposite coating solutions, containing titanium dioxide and rosmarinic acid underwent 20 min of ultrasound treatment and 10 min of UV irradiation (Lin et al., 2015).

Application of coating solutions to fillets

The coating process for trout fillets was conducted using the immersion method. Trout fillets were immersed in the prepared coating solutions for 1 min to facilitate the coating process. Subsequently, the coated trout fillets were dried at 4°C for 12 h. The dried samples were then packaged using a packaging machine (Multivac A 300/16, Multivac, Wolfertschwenden, Germany) under modified atmospheric conditions (40% CO₂+30% O₂+30% N₂). The packaged samples were stored at 4±1°C for 18 days. Trout fillets without any coating process but with direct modified atmosphere packaging (MAP) were designated as the control group.

Microbiological analysis

For microbiological analysis, 25 g of sample was homogenized with 225 mL of sterile physiological saline solution (0.85%

NaCl) in a stomacher (Lab Stomacher Blander 400-BA 7021, Seward Medical, Newport Gwent, UK) for 1 min. Serial dilutions were prepared from this homogenate and microbiological analyzes were carried out on days 0, 3, 6, 9, 12, 15, and 18 of the storage period. For enumeration of aerobic for TMAB, Plate Count Agar (PCA, Merck) was used. The plates were incubated at 30°C for 48 h (Baumgart et al., 1993); For psychrotrophic bacteria, PCA (Plate Count Agar, Merck) was used and the plates were incubated at 10°C for 7 days (Vanderzant and Splittstoesser, 1992); Enterobacteriaceae were determined on Violet Red Bile Dextrose (VRBD, Merck), the plates were incubated at 30°C for 2 days under anaerobic condition using (Anaerocult, Merck); colonies larger than 1 mm were counted (Baumgart et al., 1993). All microbiological analyses were carried out using the surface spread plate method. The results were expressed as Log CFU/g.

Physical and chemical analyses

To determine the pH value, 10 g sample was homogenized in 100 mL of distilled water. The mixture was homogenized using an Ultra-Turrax (IKA T25, IKA-Werke, Staufen, Germany) for 1 min. The pH values were calibrated using appropriate buffer solutions (pH 4.00 and pH 7.00) and measured using a pH meter (Schott, Lab Star pH, Thermo Fisher Scientific, Mainz, Germany).

The color intensities of the cross-sectional surface of the samples were determined using a colorimeter device (CR-400, Konika Minolta, Osaka, Japan). The CIE L*, CIE a*, and CIE b* were determined based on criteria established by the International Commission on Illumination for three-dimensional color measurement (Commission Internationale de l'Éclairage).

The TBARS analysis was conducted according to the method described by Lemon (1975), and the TBARS values were expressed as μmol malondialdehyde (MDA)/kg. A steam distillation method was used to determine the TVB-N level of the samples. The results obtained were given in mg TVB-N/100 g (European Commission, 2005).

Statistical analysis

In the study, coating application [control: uncoated (C), CH, CHT, and CHTRA)] and storage time (at $4\pm 0.5^\circ\text{C}$, 0, 3, 6, 9, 12, 15, and 18 days) were considered as the factors. The experiments were set up in two replications using a random complete blocks trial plan in a 4×7 factorial order. The data obtained were subjected to analysis of variance, and significant mean values of the main sources of variation were compared using the Duncan multiple comparison test. The SPSS analyses were performed using the SPSS 22 software package (SPSS 22.0, 2013).

Results and Discussion

Microbiological properties, pH and thiobarbituric acid reactive substances (TBARS) values, and total volatile basic nitrogen (TVB-N) levels

The overall effects of coating application and storage period on the microbiological and physico-chemical properties of rainbow trout fillets were given in Table 1 (mean \pm SD). The coating application had a very significant effect TMAB and total psychrotrophic bacteria ($p<0.01$). The control group showed the highest mean TMAB count. Similar results were observed for psychrotrophic bacteria. The mean lowest TMAB count was determined in CHTRA treatment. However, no significant difference was observed between CHTRA and CHT treatments with regard to psychrotrophic bacteria ($p>0.05$).

Table 1. The overall effect of coating application and storage period on the microbiological and physico-chemical properties of rainbow trout fillets

Factor	n	TMAB (Log CFU/g)	Psychrotrophic bacteria (Log CFU/g)	pH	TBARS ($\mu\text{mol MDA/kg}$)	TVB-N (mg/100 g)
Coating application (CA)						
Control	28	6.08 \pm 1.64 ^c	5.98 \pm 1.73 ^c	6.56 \pm 0.07 ^c	27.75 \pm 25.65 ^a	21.69 \pm 8.25 ^b
CH	28	4.12 \pm 1.04 ^b	4.02 \pm 1.14 ^a	6.41 \pm 0.08 ^a	26.48 \pm 23.22 ^a	17.49 \pm 1.24 ^a
CHT	28	4.18 \pm 1.09 ^b	4.28 \pm 0.86 ^b	6.45 \pm 0.08 ^b	26.66 \pm 23.90 ^a	17.05 \pm 1.25 ^a
CHTRA	28	3.91 \pm 1.11 ^a	4.14 \pm 0.93 ^{ab}	6.43 \pm 0.09 ^{ab}	31.26 \pm 29.68 ^b	17.34 \pm 1.30 ^a
Significance		**	**	**	*	**
Storage period (SP)						
0 d	16	3.22 \pm 0.70 ^a	3.39 \pm 0.58 ^a	6.50 \pm 0.11 ^b	3.13 \pm 2.15 ^a	15.16 \pm 0.76 ^a
3 d	16	3.44 \pm 0.74 ^a	3.47 \pm 0.60 ^a	6.48 \pm 0.10 ^b	4.77 \pm 2.68 ^a	16.62 \pm 0.87 ^b
6 d	16	3.81 \pm 0.72 ^b	3.96 \pm 0.62 ^b	6.42 \pm 0.09 ^a	9.79 \pm 1.93 ^b	17.12 \pm 0.58 ^b
9 d	16	4.30 \pm 1.08 ^c	4.33 \pm 1.08 ^c	6.47 \pm 0.06 ^b	19.79 \pm 4.73 ^c	17.32 \pm 0.53 ^b
12 d	16	5.12 \pm 1.15 ^d	4.96 \pm 1.15 ^d	6.41 \pm 0.09 ^a	33.06 \pm 6.70 ^d	18.27 \pm 1.81 ^c
15 d	16	5.68 \pm 1.32 ^e	5.68 \pm 1.27 ^e	6.48 \pm 0.09 ^b	53.19 \pm 9.51 ^e	20.45 \pm 4.21 ^d
18 d	16	6.44 \pm 1.32 ^f	6.44 \pm 1.33 ^f	6.48 \pm 0.11 ^b	72.34 \pm 12.91 ^f	23.80 \pm 9.07 ^e
Significance		**	**	**	**	**
CA \times SP		**	**	**	**	**

Mean \pm SD.^{a-f} Different letters indicate statistical difference ($p < 0.05$) in each column. * $p < 0.05$; ** $p < 0.01$.TMAB, total aerobic mesophilic bacteria; TBARS, thiobarbituric acid reactive substances; MDA, malondialdehyde; TVB-N, total volatile basic nitrogen; CH, chitosan; CHT, chitosan+TiO₂ nanoparticles; CHTRA, chitosan+TiO₂ nanoparticles+rosmarinic acid.

On the other hand, it was determined that the coating application \times storage period interaction had a very significant effect on both bacterial groups ($p < 0.01$; Table 1). As shown in Fig. 1, the TMAB in the control group increased more rapidly after the 6th day compared to coating treatments. While the mean TMAB count of the control group was 10^8 CFU/g at the end of storage, the number did not exceed 10^6 CFU/g in the coated treatments. Moreover, the differences between coated treatments were not significant (Fig. 1). Psychrotrophic bacteria are the main group of microorganisms responsible for the spoilage of fresh fish stored at low temperatures (4°C). Therefore, the count of these bacteria is a reliable indicator of the quality of cold-stored fish meat (Shokri et al., 2020). The interaction of coating application and storage period also had a very significant effect on the number of psychrotrophic bacteria ($p < 0.01$; Table 1). The number of psychrotrophic bacteria showed a similar trend to the number of TMAB (Fig. 2). All results indicated that coating treatments led to a significant reduction in bacterial counts. TiO₂ or rosmarinic acid had no additional effect on the reduction of the psychrotrophic bacteria. As shown in Fig. 2, the coating treatment resulted in lower psychrotrophic bacterial counts than the control on all days of analysis. The differences between the coating treatments were not significant (Fig. 1). Ojagh et al. (2010) reported comparable increases in TMAB count in rainbow trout coated with CH enriched with cinnamon oil during cold storage, suggesting a prolonged storage period. Likewise, Echeverría et al. (2018) found a decrease of 2 logarithmic units in the TMAB counts on 15 day of the storage in nanocomposite-coated samples of tuna fish compared to the control group. On the other hand, in a study




Fig. 1. Effect of interaction of coating application×storage period on total aerobic mesophilic bacteria (A) and psychrotrophic bacteria (B) of rainbow trout fillets during cold storage. ^{A-F} Different capital letters significant differences between storage periods for coating application. ^{a,b} Different small letters indicate significant differences between coating applications for the storage periods. C, uncoated; CH, chitosan; CHT, chitosan+TiO₂; CHTRA, chitosan+TiO₂+rosmarinic acid.

examining the effect of quince seed gum containing thyme or thyme essential oil on the shelf-life of rainbow trout fillets, it was reported that the number of psychrotrophic bacteria count in control group reached 10^8 CFU/g on day 18. In comparison, it remained at levels of 10^5 – 10^6 CFU/g in fillet samples with quince seed mucilage films containing thyme essential oil (Jouki et al., 2014). In a study investigating the effect of an edible active coating based on CH-sage essential oil nanoemulsion on the shelf life of rainbow trout fillets, psychrotrophic bacterial counts in the control group samples exceeded 10^8 CFU/g on day 16 of storage, while the coating treatment resulted in a count of 10^4 – 10^5 CFU/g (Shokri et al., 2020).

The antimicrobial properties of the CH coatings have already been reported in previous studies (Priyadarshi and Rhim, 2020). López-Caballero et al. (2005) found that a coating consisting of CH dissolved in acetic acid and gelatin exhibited an inhibitory effect on the Gram-negative flora of fish cakes. In our study, at the end of storage, the counts of Enterobacteriaceae were below the detectable level ($<10^2$ CFU/g; data not shown). Comparable results were also observed in another study on fish fillets (Volpe et al., 2015). Furthermore, Hisar et al. (2004) reported that MAP significantly decreased the count of Enterobacteriaceae in fillets.

The overall effects of coating application and storage period on pH value of rainbow trout fillets were given in Table 1 (mean±SD). The lowest mean pH value was observed in the control group. Changes in the average pH values were also




Fig. 2. Effect of interaction of coating application×storage period on pH value of rainbow trout fillets during cold storage. ^{A,B} Different capital letters significant differences between storage period for coating application. ^{a-c} Different small letters indicate significant differences between coating applications for the storage periods. C, uncoated; CH, chitosan; CHT, chitosan+TiO₂; CHTRA, chitosan+TiO₂+rosmarinic acid.

observed during storage. As shown in Fig. 2, on the 15th day and 18th day of storage, higher pH values were observed in the control group than in the treatment groups. According to these results, the pH value in the coating groups was 6.50 or below at the end of storage, whereas the pH value of the control group was above 6.50. In other words, the pH change in the coating applications was limited. Berizi et al. (2018) also reported that trout fillets coated with CH and permanganate extract showed a lower pH value at the end of frozen storage than the control. An increase in pH during storage can adversely affect the quality of the product, especially with regard to sensory properties such as color, odor and texture (Alak et al., 2010).

The lipids of fresh fish are very susceptible to oxidation, which leads to changes in the quality characteristics of fish. The coating application had a significant effect on the TBARS value of rainbow trout fillets ($p < 0.05$). The lowest mean TBARS value was observed in the CHTRA treatment. The differences among the other groups were not significant. In contrast, TBARS values increased with increasing storage time (Table 1). As shown in Fig. 3, the control group had a lower value than the other groups on the 9th day of storage, and the TBARS value for control was 13.52 $\mu\text{mol MDA/kg}$ ($< 1 \text{ mg MDA/kg}$). On




Fig. 3. Effect of the interaction of coating application×storage period on thiobarbituric acid reactive substances (TBARS) value of rainbow trout fillets during cold storage. ^{A-F} Different capital letters significant differences between storage period for coating application. ^{a-c} Different small letters indicate significant differences between coating applications for the storage periods. MDA, malondialdehyde; C, uncoated; CH, chitosan; CHT, chitosan+TiO₂; CHTRA, chitosan+TiO₂+rosmarinic acid.

the other hand, the TBARS value for coating treatments were under 2 mg MDA/kg. Karki et al. (2023) reported that the tolerable TBARS value of fish products is 1 mg MDA/kg (100 μ mol MDA/kg is equivalent to 7.2 mg/kg MDA). On the other hand, Xiong et al. (2021) reported that the threshold of TBARS value for oxidative rancidity and sensory acceptability ranged from 1–2 mg MDA/kg. On the following days of storage, significant increases in TBARS values were observed in all groups. The TBARS value on the 12th day of storage was below 2 only in CHT treatment. On the 15th day of storage, the TBARS value increased significantly in all groups (>3.0 mg MDA/kg). The highest TBARS value during cold storage was found in the CHTRA group at the end of storage. However, with regard to TBARS, no significant differences were observed among all treatments, including the control (Fig. 3). On the other hand, another study reported that CH coating on salmon fillets provided better results against lipid oxidation than gelatin coating (Xiong et al., 2021).

The determination of volatile nitrogenous compounds such as trimethylamine, dimethylamine, and ammonia, collectively referred to as the TVB-N (Shokri et al., 2020). The control group exhibited the highest mean TVB-N levels during storage. There was no significant difference between the coating groups. As the storage time increased, the mean TVB-N level increased (Table 1). The interaction of the coating application and storage period had a very significant effect ($p < 0.01$) on the TVB-N level. These results were consistent with the microbiological results. It was observed that CH and TiO₂ nanoparticles contributed to a significant decrease in the TVB-N values and significantly impacted the shelf-life of trout fillets. However, rosmarinic acid had no effect on TVB-N level (Fig. 4). As shown in Fig. 4, the highest TVB-N value was observed in the control group on the 12th, 15th and 18th days of storage. However, the differences between the coating treatments were not significant. It was reported that maximum acceptable TVB-N level for rainbow trout is 25 mg/100 g (Giménez et al., 2002). In our study, the TVB-N levels in the coated samples remained below this acceptable limit during storage period. In contrast, the TVB-N level in the control group exceeded the acceptable limit of 25 mg/100 g after 15 days (Fig. 4). Similarly, it has been reported that after 16 days of refrigerated storage, rainbow trout fillet samples coated with CH and CH combined with other substances remained below the acceptable limit, while the TVB-N level of the control group samples increased to 40 mg/100 g (Ojagh et al., 2010). In addition, López-Caballero et al. (2005) demonstrated that a protective CH-gelatin coating applied to fish balls significantly reduced TVB-N level. On the other hand, Korkmaz (2016) studied the effect of quinoa edible film on rainbow trout fillets and reported a TVB-N value of 20.35 ± 0.49 mg/100 g in the control group and 18.65 ± 0.21




Fig. 4. Effect of the interaction of coating application×storage period on total volatile basic nitrogen (TVB-N) level of rainbow trout fillets during cold storage. ^{A–D} Different capital letters significant differences between storage period for coating application. ^{a,b} Different small letters indicate significant differences between coating applications for the storage periods. C, uncoated; CH, chitosan; CHT, chitosan+TiO₂; CHTRA, chitosan+TiO₂+rosmarinic acid.

mg/100 g in the experimental group after 12 days of storage.

CIE L*, CIE a*, and CIE b*

Physico-chemical changes that occur during storage can affect the appearance and texture of the fish (Zarandona et al., 2021). The coating application had a very significant effect the CIE L*, CIE a*, and CIE b* of rainbow trout fillets ($p<0.01$). The storage period was found to be very effective for the CIE L* and CIE b* ($p<0.01$). This factor also affected CIE a* ($p<0.05$; Table 2). Control and CH treatment had lower CIE L* than nanocomposite groups. In another study on sea bream, the highest CIE L* was found in the coating groups. As shown in Fig. 5, the lowest initial CIE L* was observed in the CH treatment. At the end of storage, the highest CIE L* was obtained with CHTRA treatment. However, the value of this treatment did not differ from that of the CH treatment. Considering these results, it can be concluded that the coating process (composite or nanocomposite coating) has a positive effect on the CIE L*. However, no remarkable changes were observed in the CIE a* and CIE b* in this study. Duan et al. (2010) also reported no changes in the CIE L*, CIE a* and CIE b* of fish samples stored at 2°C for three weeks.

Conclusion

The results demonstrated that the coating application (CH, CHT, or CHTRA) effectively inhibited microbial growth in

Table 2. The overall effect of coating application and storage period on the CIE L*, CIE a*, and CIE b* of rainbow trout fillets

Factor	n	CIE L*	CIE a*	CIE b*
Coating application (CA)				
Control	28	47.90±2.23 ^a	0.84±1.41 ^b	11.22±3.46 ^a
CH	28	48.29±3.97 ^a	-0.32±1.41 ^a	12.36±3.92 ^{ab}
CHT	28	50.51±2.54 ^b	0.20±1.69 ^{ab}	13.79±2.82 ^b
CHTRA	28	51.02±3.45 ^b	0.68±2.39 ^b	13.76±3.95 ^b
Significance		**	**	**
Storage period (SP)				
0 d	16	45.72±2.37 ^a	0.44±1.52 ^{abc}	10.56±3.83 ^a
3 d	16	48.76±2.22 ^b	0.18±1.10 ^{abc}	12.94±2.96 ^{bc}
6 d	1	49.95±3.84 ^{bc}	0.30±1.66 ^{abc}	12.97±3.92 ^{bc}
9 d	16	49.36±2.78 ^{bc}	-0.42±1.42 ^a	11.67±2.87 ^{ab}
12 d	16	50.65±3.06 ^c	0.06±1.26 ^{ab}	14.88±3.64 ^c
15 d	16	50.95±2.70 ^c	1.07±2.82 ^c	14.62±3.36 ^c
18 d	16	50.63±3.41 ^c	0.83±2.16 ^{bc}	11.85±3.55 ^{ab}
Significance		**	*	**
Coating×Storage				
		**	**	**

Mean±SD.

^{a-c} Different letters indicate statistical difference ($p<0.05$) in each column. * $p<0.05$; ** $p<0.01$.

CH, chitosan; CHT, chitosan+TiO₂ nanoparticles; CHTRA, chitosan+TiO₂ nanoparticles+rosmarinic acid.




Fig. 5. Effect of the interaction of coating application×storage period on CIE L* of rainbow trout fillets during cold storage. ^{A-D} Different capital letters significant differences between storage period for coating application. ^{a-c} Different small letters indicate significant differences between coating applications for the storage periods. C, uncoated; CH, chitosan; CHT, chitosan+TiO₂; CHTRA, chitosan+TiO₂+rosmarinic acid.

rainbow trout fillets during cold storage. In addition coating application showed the lower TVB-N level than control group. The TBARS value increased as the storage time increased. At the end of storage, no significant differences were observed between treatments in terms of TBARS (Fig. 3). It was also determined that the pH change in the coating applications was limited. On the other hand, the coating process has a positive effect on the CIE L*.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

We would like to thank Atatürk University, Eastern Anatolia High Technology Application and Research Center (DAYTAM) for giving us the opportunity to perform our study.

Author Contributions

Conceptualization: Kizilkaya P, Kaya M. Data curation: Kizilkaya P. Formal analysis: Kizilkaya P. Methodology: Kizilkaya P, Kaya M. Software: Kizilkaya P. Validation: Kizilkaya P. Investigation: Kizilkaya P. Writing - original draft: Kizilkaya P. Writing - review & editing: Kizilkaya P, Kaya M.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

Alak G, Aras Hisar S, Hisar O, Kaban G, Kaya M. 2010. Microbiological and chemical properties of bonito fish (*Sarda sarda*)

- fillets packaged with chitosan film, modified atmosphere and vacuum. *Kafkas Univ Vet Fak Derg* 16:S73-S80.
- Ambaye TG, Vaccari M, Prasad S, van Hullebusch ED, Rtimi S. 2022. Preparation and applications of chitosan and cellulose composite materials. *J Environ Manag* 301:113850.
- Baumgart J, Eigener V, Firnhaber J, Hildebrandt G, Reenen Hoekstra ES, Samson RA, Spicher G, Timm F, Yarrow D, Zschaler R. 1993. *Mikrobiologische untersuchung von lebensmitteln*. 3rd ed. Behr's, Hamburg, Germany.
- Bento R, Pagán E, Berdejo D, de Carvalho RJ, García-Embid S, Maggi F, Magnani M, de Souza EL, García-Gonzalo D, Pagán R. 2020. Chitosan nanoemulsions of cold-pressed orange essential oil to preserve fruit juices. *Int J Food Microbiol* 331:108786.
- Berizi E, Hosseinzadeh S, Shekarforoush SS, Barbieri G. 2018. Microbial, chemical, textural and sensory properties of coated rainbow trout by chitosan combined with pomegranate peel extract during frozen storage. *Int J Biol Macromol* 106:1004-1013.
- Duan J, Cherian G, Zhao Y. 2010. Quality enhancement in fresh and frozen lingcod (*Ophiodon elongates*) fillets by employment of fish oil incorporated chitosan coatings. *Food Chem* 119:524-532.
- Echeverría I, López-Caballero ME, Gómez-Guillén MC, Mauri AN, Montero MP. 2018. Active nanocomposite films based on soy proteins-montmorillonite- clove essential oil for the preservation of refrigerated bluefin tuna (*Thunnus thynnus*) fillets. *Int J Food Microbiol* 266:142-149.
- European Commission [EC]. 2005. Commission regulation (EC) No 2074/2005: Total volatile basic nitrogen (TVB-N) limit values for certain categories of fishery products and analysis methods to be used. European Commission, Brussels, Belgium. pp 36-39.
- Giménez B, Roncalés P, Beltrán JA. 2002. Modified atmosphere packaging of filleted rainbow trout. *J Sci Food Agric* 82:1154-1159.
- He Q, Zhang Y, Cai X, Wang S. 2016. Fabrication of gelatin-TiO₂ nanocomposite film and its structural, antibacterial and physical properties. *Int J Biol Macromol* 84:153-160.
- Heydari-Majd M, Ghanbarzadeh B, Shahidi-Noghabi M, Najafi MA, Hosseini M. 2019. A new active nanocomposite film based on PLA/ZnO nanoparticle/essential oils for the preservation of refrigerated *Otolithes ruber* fillets. *Food Packag Shelf Life* 19:94-103.
- Hisar ŞA, Hisar O, Yanik T. 2004. Balıklarda mikrobiyolojik, enzimatik ve kimyasal bozulmalar. *Atatürk Üniv Ziraat Fak Derg* 35:261-265.
- Hoque M, Gupta S, Santhosh R, Syed I, Sarkar P. 2021. Biopolymer-based edible films and coatings for food applications. In *Food, medical, and environmental applications of polysaccharides*. Pal K, Banerjee I, Sarkar P, Bit A, Kim D, Anis A, Maji S (ed). Elsevier, Amsterdam, Netherlands. pp 81-107.
- Hosseini SF, Ghaderi J, Gómez-Guillén MC. 2022. Tailoring physico-mechanical and antimicrobial/antioxidant properties of biopolymeric films by cinnamaldehyde-loaded chitosan nanoparticles and their application in packaging of fresh rainbow trout fillets. *Food Hydrocoll* 124:107249.
- Jiang J, Chen X, Zhang GL, Hao H, Hou HM, Bi J. 2022. Preparation of chitosan-cellulose-benzyl isothiocyanate nanocomposite film for food packaging applications. *Carbohydr Polym* 285:119234.
- Jouki M, Yazdi FT, Mortazavi SA, Koocheki A, Khazaei N. 2014. Effect of quince seed mucilage edible films incorporated with oregano or thyme essential oil on shelf life extension of refrigerated rainbow trout fillets. *Int J Food Microbiol* 174:88-97.

- Jovanović B, Whitley EM, Kimura K, Crumpton A, Palić D. 2015. Titanium dioxide nanoparticles enhance mortality of fish exposed to bacterial pathogens. *Environ Pollut* 203:153-164.
- Kanmani P, Rhim JW. 2014. Physical, mechanical and antimicrobial properties of gelatin based active nanocomposite films containing AgNPs and nanoclay. *Food Hydrocoll* 35:644-652.
- Karki S, Chowdhury S, Dora KC, Murmu P, Nath S. 2023. Influence of a traditional herb (*Bergenia ciliata*) from darjeeling himalayas on lipid oxidation and sensorial quality of sous-vide cooked tilapia (*Oreochromis niloticus*). *J Aquat Food Prod Technol* 32:256-268.
- Korkmaz F. 2016. Yenilebilir biyofilm olarak kinoa (*Chenopodium quinoa*)' nın gökkuşağı alabalığı (*Onchorynchus mykiss*) filetolarının raf ömrü üzerine etkisinin araştırılması. M.S. thesis, Atatürk Univ., Erzurum, Turkey.
- Kumar S, Mukherjee A, Dutta J. 2020. Chitosan based nanocomposite films and coatings: Emerging antimicrobial food packaging alternatives. *Trends Food Sci Technol* 97:196-209.
- Lemon DW. 1975. An improved TBA test for rancidity. Fisheries and Marine Service, Halifax, NS, Canada.
- Lin B, Luo Y, Ten Z, Zhang B, Zhou B, Wang Q. 2015. Development of silver/titanium dioxide/chitosan adipate nanocomposite as an antibacterial coating for fruit storage. *LWT-Food Sci Technol* 63:1206-1213.
- López-Caballero ME, Gómez-Guillén MC, Pérez-Mateos M, Montero P. 2005. A chitosan-gelatin blend as a coating for fish patties. *Food Hydrocoll* 19:303-311.
- Marchev AS, Vasileva LV, Amirova KM, Savova MS, Koycheva IK, Balcheva-Sivenova ZP, Vasileva SM, Georgiev MI. 2021. Rosmarinic acid: From bench to valuable applications in food industry. *Trends Food Sci Technol* 117:182-193.
- Nath D, Chetri R, Santhosh R, Sarkar P. 2022. Nanocomposites in food packaging. In *Food, medical, and environmental applications of nanomaterials*. Pal K, Sarkar A, Sarkar P, Bandara N, Jegatheesan V (ed). Elsevier, Amsterdam, Netherlands. pp 167-203.
- Nowzari F, Shábanpour B, Ojagh SM. 2013. Comparison of chitosan-gelatin composite and bilayer coating and film effect on the quality of refrigerated rainbow trout. *Food Chem* 141:1667-1672.
- Nwabor OF, Singh S, Paosen S, Vongkamjan K, Voravuthikunchai SP. 2020. Enhancement of food shelf life with polyvinyl alcohol-chitosan nanocomposite films from bioactive *Eucalyptus* leaf extracts. *Food Biosci* 36:100609.
- Ojagh SM, Rezaei M, Razavi SH, Hosseini SMH. 2010. Effect of chitosan coatings enriched with cinnamon oil on the quality of refrigerated rainbow trout. *Food Chem* 120:193-198.
- Petersen M, Simmonds MSJ. 2003. Rosmarinic acid. *Phytochemistry* 62:121-125.
- Priyadarshi R, Rhim JW. 2020. Chitosan-based biodegradable functional films for food packaging applications. *Innov Food Sci Emerg Technol* 62:102346.
- Qu B, Luo Y. 2021. A review on the preparation and characterization of chitosan-clay nanocomposite films and coatings for food packaging applications. *Carbohydr Polym Technol Appl* 2:100102.
- Rahman S, Konwar A, Majumdar G, Chowdhury D. 2021. Guar gum-chitosan composite film as excellent material for packaging application. *Carbohydr Polym Technol Appl* 2:100158.
- Salimiraad S, Safaeian S, Basti AA, Khanjari A, Nadoushan RM. 2022. Characterization of novel probiotic nanocomposite films based on nano chitosan/ nano cellulose/ gelatin for the preservation of fresh chicken fillets. *LWT-Food Sci Technol* 162:113429.
- Sani MA, Ehsani A, Hashemi M. 2017. Whey protein isolate/cellulose nanofibre/TiO₂ nanoparticle/rosemary essential oil nanocomposite film: Its effect on microbial and sensory quality of lamb meat and growth of common foodborne

- pathogenic bacteria during refrigeration. *Int J Food Microbiol* 251:8-14.
- Shokri S, Parastouei K, Taghdir M, Abbaszadeh S. 2020. Application an edible active coating based on chitosan-*Ferulago angulata* essential oil nanoemulsion to shelf life extension of rainbow trout fillets stored at 4°C. *Int J Biol Macromol* 153:846-854.
- Silva AO, Cunha RS, Hotza D, Machado RAF. 2021. Chitosan as a matrix of nanocomposites: A review on nanostructures, processes, properties, and applications. *Carbohydr Polym* 272:118472.
- Vanderzant C, Splittstoesser DF. 1992. Compendium of methods for the microbiological examination of foods. American Public Health Association, Washington, DC, USA.
- Volpe MG, Siano F, Paolucci M, Sacco A, Sorrentino A, Malinconico M, Varricchio E. 2015. Active edible coating effectiveness in shelf-life enhancement of trout (*Oncorhynchus mykiss*) fillets. *LWT-Food Sci Technol* 60:615-622.
- Wang H, Ding F, Ma L, Zhang Y. 2021. Edible films from chitosan-gelatin: Physical properties and food packaging application. *Food Biosci* 40:100871.
- Xiong Y, Kamboj M, Ajlouni S, Fang Z. 2021. Incorporation of salmon bone gelatine with chitosan, gallic acid and clove oil as edible coating for the cold storage of fresh salmon fillet. *Food Control* 125:107994.
- Yu J, Wang D, Geetha N, Khawar KM, Jogaiah S, Mujtaba M. 2021. Current trends and challenges in the synthesis and applications of chitosan-based nanocomposites for plants: A review. *Carbohydr Polym* 261:117904.
- Zabihollahi N, Alizadeh A, Almasi H, Hanifian S, Hamishekar H. 2020. Development and characterization of carboxymethyl cellulose based probiotic nanocomposite film containing cellulose nanofiber and inulin for chicken fillet shelf life extension. *Int J Biol Macromol* 160:409-417.
- Zarandona I, López-Caballero ME, Montero MP, Guerrero P, de la Caba K, Gómez-Guillén MC. 2021. Horse mackerel (*Trachurus trachurus*) fillets biopreservation by using gallic acid and chitosan coatings. *Food Control* 120:107511.
- Zhao L, Pan F, Mehmood A, Zhang H, Ur Rehman A, Li J, Hao S, Wang C. 2021. Improved color stability of anthocyanins in the presence of ascorbic acid with the combination of rosmarinic acid and xanthan gum. *Food Chem* 351:129317.