



MicroRNA-21 mediates bone marrow mesenchymal stem cells protection of radiation-induced lung injury during the acute phase by regulating polarization of alveolar macrophages

Pengtao Bao, Weiguo Zhao, Mi Mou, Xiaofei Liu

Department of Respiration, the Eighth Medical Center of PLA General Hospital, Beijing 100091, China

Contributions: (I) Conception and design: W Zhao; (II) Administrative support: W Zhao, P Bao; (III) Provision of study materials or patients: P Bao, M Mou; (IV) Collection and assembly of data: P Bao, M Mou, X Liu; (V) Data analysis and interpretation: P Bao, M Mou, X Liu; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Weiguo Zhao. Department of Respiration, the Eighth Medical Center of PLA General Hospital, Beijing 100091, China. Email: wgzhao309@126.com.

Background: Radiation-induced lung injury (RILI) often occurs in patients with non-small cell lung cancer (NSCLC) after radiotherapy, and the prognosis of patients with RILI is usually poor. This work plan to investigate the expression patterns of microRNA-21(miR-21) in NSCLC patients with RILI and the protective effects of miR-21 over-expressed bone marrow mesenchymal stem cells (BMSCs) against RILI in rat model.

Methods: MiR-21 expressions were determined in both serum samples and bronchoalveolar lavage fluid (BALF) samples from NSCLC patients after radiation therapy. The correlation between miR-21 expression and the follow-up clinical characterizations were determined. Further, miR-21 over-expressed BMSCs were transplanted into RILI rats and the protective effects were evaluated. BMSCs and alveolar macrophages (AMs) were co-cultured *in vitro* and the macrophage M1 polarization markers were determined by ELISA and qRT-PCR assays.

Results: Expression of miR-21 was significantly increased in NSCLC patients with RILI compared with control group, especially before or at 4 weeks after radiation therapy commenced. The miR-21 levels were highly correlated with IL-12, TNF- α , and IL-6 expressions and the severity of RILI. Animal based experiments demonstrated that BMSCs treatment had a remarkable effect on alleviating alveolitis in RILI rats, and miR-21 over-expression could enhance this effect significantly. Cell based experiments demonstrated that BMSCs notably inhibited M1 polarization of AMs and this inhibition is in a miR-21 dependent manner.

Conclusions: These results indicated that BMSCs could blocked the proinflammatory pathway of macrophage through miR-21 over-expression, thus could be a potential therapeutic strategy for RILI.

Keywords: Non-small cell lung cancer (NSCLC); radiation-induced lung injury (RILI); polarization; bone marrow mesenchymal stem cells (BMSCs); microRNA (miRNAs)

Submitted Aug 27, 2019. Accepted for publication Nov 26, 2019.

doi: 10.21037/tcr.2019.12.77

View this article at: <http://dx.doi.org/10.21037/tcr.2019.12.77>

Introduction

Radiation-induced lung injury (RILI) often occurs after radiotherapy in patients with non-small cell lung cancer (NSCLC). The prognosis of patients with RILI is usually

poor (1). Reports have indicated that lung injury, with symptoms only during radiotherapy for lung cancer, has an incidence rate reaching 30% (2). Currently, RILI treatment options are rather limited, and many of which lead to higher recurrence rates (3). Therefore, finding markers that

can indicate the occurrence of RILI, and finding effective therapeutic method to prevent or treat RILI, are all very important goals on the way to improved prognosis of patients receiving radiotherapy.

The mechanisms through which microRNAs (miRNAs) regulate physiological and pathological reactions have recently been extensively studied (4). The specific expression patterns of miRNAs in certain diseases indicate that many of them can be used as diagnostic markers (5). Considering that many miRNAs function by suppressing their target genes, dis-regulated miRNAs can often play a regulatory role in the development of their associated diseases (6). In such cases, introduction of exogenous miRNA or suppression of endogenous miRNA can usually be used as an effective treatment approach.

Bone marrow mesenchymal stem cells (BMSCs) are multipotent cells that can differentiate into osteoblasts, chondrocytes and adipocytes (7). Infusion of BMSCs that overexpress specific miRNAs can effectively introduce exogenous miRNAs to target cells via the paracrine system and achieve therapeutic purposes (8).

In the present study, the correlation between miR-21 expression and the severity and prognosis of NSCLC patients with RILI was investigated. A rat model demonstrated that BMSCs overexpressing miR-21 could protect radiation-induced lung injury during the acute phase. Furthermore, cell-based studies showed that BMSCs overexpressing miR-21 could inhibit M1 polarization of alveolar macrophages (AMs) by suppressing TNF- α expression in these cells, and thus achieve therapeutic purposes.

Methods

Patients and samples

Totally 61 non-small cell lung cancer (NSCLC) patients before, during, and at the end of a treatment course of conventional fractionated radiotherapy were included in this study. Patients inclusion criteria, radiation therapy and classification criteria of RILI were in according with our previous description (2). The serum samples and bronchoalveolar lavage fluid (BALF) samples from all the patients (each patient was collected at various time points were collected for the subsequent assays in according with our previous reports (2). Pulmonary function markers, including diffusion capacity for carbon monoxide (DLCO), forced expiratory volume at 1 s (FEV1), and forced vital capacity (FVC), were recorded and used as baselines in all patients before radiotherapy, and would be tested

at 4, 8, 16 and 24 weeks for follow-up treatment. The prospective clinical study was performed from January 2014 to December 2018 in the Eighth Medical Center of PLA General Hospital, Beijing, China. The experimental study was performed in the laboratory of PLA General Hospital from June 2016 to March 2019.

RNA extraction and qRT-PCR assay

Serum samples were collected and centrifuged at a speed of 3,000 g for 5 min at 4 °C. Next, 200 mL serum supernatant was extracted for RNA extraction. Similarly, BALF samples were centrifuged for 10 min at a speed of 3,000 g at 4 °C, and were absorbed 200 mL supernatant for RNA extraction. Total RNA was obtained from the above samples using TRIzol reagent (Invitrogen, New York Island, USA).

Subsequently, qRT-PCR for mRNA was performed using the PrimeScript RT-PCR kit (Takara, Bio, Inc., Shiga, Japan). qRT-PCR for miRNA was performed using TaqMan miRNA kit. All the qRT-PCR experiments were performed on Eppendorf quantitative PCR detector. All the primer sequences of the target were from the previous literature (9). The parameters of reverse transcription for mRNA were set as follows: first, 65 °C reaction was set for 5 min, then 37 °C reaction was set for 15 min, and finally 98 °C reaction was set for 5 min. The parameters of qPCR for mRNA were set as follows: first 95 °C reaction for 30 s, then 40 cycles of 95 °C reaction for 5 s, and 60 °C reaction for 5 s, and finally 72 °C reaction for 30 s. The human and rat miR-21 primers were purchased from Qiagen [catalog numbers MS00009079 (Hs-miR-21-5p) and MS00013216 (Rn-miR-21-5p)]. The parameters of reverse transcription for miRNA were set as follows: first, 16 °C reaction was set for 30 min, then 42 °C reaction was set for 30 min, and finally 84 °C reaction was set for 5 min. The parameters of qPCR for miRNA were set as follows: First 95 °C reaction for 2 min, then 40 cycles of 95 °C reaction for 15 s, and 60 °C reaction for 30 s.

RILI rat model

The steps and methods of the RILI rat model constructed in this study refer to our previous publication (10). In short, female, 8-week-old Sprague-Dawley (SD) rats weighing about 240 g were used to construct RILI rat models. After being fixed, rats were anesthetized by intravenous injection of pentobarbital (40 mg/kg). Rats' heads and abdomen were protected from lead radiation. The irradiation parameters were as follows: (I) 6 mV photons (Siemens, Germany)

were generated and (II) were focused to the surface at a distance of 100 cm. (III) Single irradiation of 20 Gy (2.5 Gy/min), and (IV) irradiation area of 4.5 cm × 4.5 cm. After irradiation, rats were euthanized with carbon dioxide for a certain time (2 h and 4 weeks, 8 weeks, 16 weeks and 24 weeks). Flow rate of CO₂ displaced no more than 30% of the chamber volume per minute. The mice were observed until they were found to have stopped breathing, and then stopped injecting carbon dioxide. The rats were taken out and observed for another 5 min to confirm apnea. Between animals, the chamber is turned over to completely empty the carbon dioxide chamber. Subsequently, the serum, BALF and tissues samples from the rats were collected for the subsequent assays in according with our previous reports (10). At 28 weeks after irradiation, all the remaining mice were euthanized with CO₂ as human endpoints.

Isolation, purification, and culture of BMSCs

Carbon dioxide was used to euthanize rats, and then the bone marrow cavity of rats was washed with full culture medium. The washed medium was filtered by 200 mesh filter. The filtrate was repeatedly blown to form a uniform single cell suspension and then cultured on a 10 cm disc. After removing non-adherent cells, the medium was replaced every 1–2 days. After 10–14 days, 90% of the cells were fused and passaged. Such repeated third generation cells are used for subsequent experiments.

Stable transfection of BMSCs with miR-21 overexpressing vector

MiR-21 overexpressing vector were purchased from Ribobio, Guangzhou. According to the manufacturer's instructions, Lipofectamine[®] 2000 (Invitrogen; Thermo Fisher Science, Inc.) was used to transfer the over-expressing miRNA-21 vector into BMSC, and were select stable transfectants by G-418. BMSCs that were stable transfected with miR-21 overexpressing vector were termed BMSC-miR-21-OV and control BMSCs were termed BMSC-NC.

Transplantation of the BMSCs into the rats

BMSCs were injected into rats at the dose of 2×10⁶ by tail vein injection. The same dose of saline was injected into the control group. Every three rats were killed at 3, 7 and 14 days after BMSC injection to determine whether BMSC was successfully transplanted.

Isolation and culture of rat peritoneal macrophages

After RILI model was successfully established in rats, peritoneal macrophages were subsequently extracted. Briefly, 20 mL of PBS, together with 10% FBS, were injected into the abdominal cavity of rats. Subsequently, rat peritoneal fluid was extracted and centrifuged at a speed of 192 ×g for 10 min at 4 °C. The precipitated cells were re-dispersed and inoculated into a six-well plate and cultured in complete culture medium. After 6 hours, the uncoated cells were removed. Fresh media should be replaced every 1–2 days. Two days later, the cells were sorted by CD68 and CD11.

In vitro co-culture of peritoneal macrophages with BMSCs

Totally 1×10⁶ macrophages were inoculated into 6-well plates. Twelve hours later, the cells were replaced with a medium containing 1×10⁶ BMSCs. Macrophages and BMSC were co-cultured with Transwell inserts (Corning, Inc.).

Cytokine assay

Cytokine levels and nitric oxide production in culture supernatants or in the serum samples from both clinical patients and rat model were determined using commercial ELISA kits (R&D Systems and BD Biosciences, Franklin Lakes, NJ, USA) according to the manufacturer's protocol. Each value represents the mean of triplicate values.

Statistical analysis

All statistics were conducted using SPSS software package (version 19.0; SPSS Corporation). The clinical characteristics of the two groups were compared by Pearson Chi-square test or Fisher exact test. Student *t*-test or one-way ANOVA were used for continuous data analysis. The overall survival rate (OS) or disease-free survival rate (DFS) was assessed by Kaplan-Meier method.

Results

MiR-21 in sera and bronchoalveolar lavage fluid (BALF) supernatant is significantly increased in NSCLC patients with RILI

Before and on the day of intensity-modulated radiotherapy (IMRT), miR-21 expression characteristics in the serum of every patient were measured and considered as the baseline. Serum levels were also recorded at weeks 4,




Figure 1 Expression patterns of miR-21 in NSCLC patients with RILI. (A,B) Before IMRT, concentrations of miR-21 from (A) serum and (B) BALF supernatants in every patient were measured as baseline, and were follow-up at 4, 8, 16 and 24 weeks after IMRT. (C,D) IL-6 and TNF- α level of serum at week 4 were measured using ELISA assay. The correlation between miR-21 level and (C) IL-6; or (D) TNF- α concentrations were determined. (E,F) Concentrations of IL-6 and TNF- α from BALF supernatants at week 4 were measured using ELISA assay. The correlation between miR-21 level and (E) IL-6; or (F) TNF- α concentrations were determined.

8, 16, and 24 after treatment (Figure 1A). Serum miR-21 level has increased at the early stage after irradiation therapy in all patients, peaked at week 4 and then gradually decreased through weeks 8, 16, and 24. Similar results were also observed in BALF supernatant samples (Figure 1B). Moreover, we found negative correlations between miR-21 level and that of IL-6 and TNF- α at week 4 in both sera (Figure 1C,D) and BALF supernatants (Figure 1E,F).

MiR-21 level at week 4 could be considered a potential biomarker for predicting RILI severity

We assigned the average expression of miR-21 at week 4 as the cutoff value (cutoff value = 3.0) and then divided

the patients into miR-21 high group (32 patients) and miR-21 low group (30 patients). As shown in Table 1, at weeks 8 and 24 after treatment, patients with high miR-21 suffered RILI-related effects significantly less than that those in the low miR-21 group. This was reflected in lower RILI incidence and RILI grade in the high miR-21 group ($P=0.008$ and $P=0.031$, respectively). After comparing lung function markers between the two groups, we found that there was no difference in carbon monoxide diffusion capacity (DLCO), forced expiratory volume in one second (FEV1), and forced vital capacity (FVC) values between the high and low miR-21 groups. After IMRT, the DLCO, FEV1, and FVC values in both groups decreased relative to baseline values (Table 1). No significant differences in

Table 1 Grade of RILI and variations of pulmonary function at 8 and 24 weeks post-radiation therapy

Parameters	Post-radiation therapy 8 weeks	P value	Post-radiation therapy 24 weeks	P value
Grade of RILI ^a				
Grade 1 (n)		0.008		0.031
miR-21 low	7		10	
miR-21 high	3		8	
Grade 2 (n)				
miR-21 low	4		7	
miR-21 high	1		4	
Grade 3 (n)				
miR-21 low	3		5	
miR-21 high	1		3	
Pulmonary function ^b				
FVC (%)		0.262		0.005
miR-21 low	-3.48±0.89		-5.15 ±0.94	
miR-21 high	-3.24±0.77		-4.33 ±1.27	
FEV1 (%)		0.224		0.029
miR-21 low	-3.52±1.03		-5.88±1.23	
miR-21 high	-3.17±1.21		-5.17±1.28	
DLCO (%)		0.308		0.069
miR-21 low	-4.93±1.22		-4.65±1.53	
miR-21 high	-4.62±1.14		-4.02±1.10	

Patients in miR-21 high group (32 patients) and low group (30 patients) were examined at 8 and 24 weeks after commencement of radiation therapy for grade of RILI and pulmonary function. ^a, the values are shown as the number of patients with each grade (n). P value was calculated by Pearson Chi-Square test; ^b, Baseline values of FEV1, DLCO and FVC are set as 0 in trial and control groups. Percentage changes in FEV1, DLCO and FVC at 8 and 24 weeks are calculated from baseline. P value was calculated by student t test. FVC, forced vital capacity; FEV1, forced expiratory volume at 1 s; DLCO, diffusion capacity for carbon monoxide.

DLCO, FEV1, and FVC values were found when the two groups were evaluated on week 8. At week 24, significant decrease in FVC and FEV1 values was observed in miR-21 low group, when compared with the miR-21 high group. The above data indicate that miR-21 expression increased in NSCLC patients with RILI, and that the level of miR-21 at week 4 could be considered a potential biomarker for predicting the severity of RILI.

Treatment with miR-21-knockout BMSCs reduces mortality in rats with RILI

The use of miRNA-overexpressing stem cells is a newly developed therapeutic strategy for the treatment of

various diseases (8). Thus, we aimed to determine whether treatment with miR-21-overexpressing BMSCs was effective against RILI. Rats with RILI were randomly split into three groups. The control group received saline injection to the tail vein before and after the irradiation treatment. The miR-21-overexpressing BMSCs (BMSC-miR-21-OV) group and non-treated wildtype BMSCs (BMSC-NT) group were treated with BMSC-miR-21-OV or BMSC-NT, respectively, for 3 d before and 4 d after irradiation treatment. The rats showed signs of physical weakness, and several died within 24 weeks post-irradiation. The survival curves for the three groups are shown in *Figure 2*. Mortality rates were 45% (9/20) for the control group, 20% (4/20) for the BMSC-NT group, and 5% (1/20) for the BMSC-miR-




Figure 2 Effects of BMSCs-miR-21 or BMSCs-miR-NC on survival of RILI rats.




Figure 3 Representative rat lung histological samples of control, BMSCs-miR-21 or BMSCs-miR-NC treated RILI rats (HE, $\times 400$). (A,B,C) Alveolitis at 4 weeks; and (D,E,F) TNF- α production at 4 weeks; and (G,H,I) IL-6 production at 4 weeks in lung tissue from group BMSCs-miR-21, group BMSCs-miR-NC and control group.

21-OV group. These mortality rates differed significantly between the three groups.

Treatment with miR-21-knockout BMSCs decreases RILI-induced acute inflammation.

Studying histological sections, acute inflammation was found in all experimental groups at week 4 post-irradiation. Compared with the control group, the BMSC-NT exhibited the following weaknesses: (I) inflammatory cells throughout the lung mesenchyme, (II) collapsed alveoli with significant inflammatory exudates, and (III) pulmonary hyaline membranes in the lung tissue. These findings indicate that BMSCs treatment can significantly inhibit the development of RILI in rats. Moreover, the inhibition effect

of BMSCs was markedly enhanced in the BMSC-miR-21-OV group. In addition, immunohistochemical staining revealed that the BMSC-miR-21-OV group exhibited the lowest expression of the proinflammatory factors IL-6 and TNF- α , followed by the BMSC-NT group. Meanwhile, the control group showed the highest IL-6 and TNF- α expression levels (Figure 3).

ELISA analysis revealed significant increase in the serum level of IL-6 and expression of TNF- α in the irradiated control group. These peaked at week 4 and then decreased gradually. Compared with the BMSC-NT group, BMSC-miR-21-OV treatment was more effective in inhibiting the expression of IL-6 and TNF- α (Figure 4). The combined aforementioned data indicate that injection of miR-21-overexpressing BMSCs is an effective treatment strategy against RILI.

Co-culture of miR-21-knockout BMSCs plays an important role in inhibiting AMs M1 polarization

In RILI patients and animal model, the secretion of proinflammatory factors is closely related to polarization of lung macrophages (11). Previous reports have demonstrated that a number of important macrophage M1 polarization markers, including IL-12p35 and TNF- α , are miR-21 targets *in vitro* (12,13). We therefore explored whether BMSCs-miR21-OV could change the polarization of lung macrophages. AMs from the rats were cultured in 12-well plates. BMSCs-miR-21-OV or BMSCs-NT were then added as coculture. The cell cultures were stimulated with 200 ng/L LPS for 24 h to induce the macrophages into M1 polarization (9,14). We found that BMSCs-NT coculture could not alter the mRNA expressions of M1 markers, such as IL-6, IL-12, and TNF- α . Their expression was comparable to the control AMs. BMSCs-miR-21-OV coculture significantly inhibited the induction of the macrophages into M1 polarization (Figure 5A,B,C). BMSCs-miR-21-OV coculture also decreased production of Nitric oxide in AMs (Figure 5D). These results suggest that miR-21 may regulate RILI by controlling macrophage polarization.

Discussion

The overall prognosis of NSCLC patients after radiotherapy varies greatly among individuals. A more accurate assessment of the incidence of RILI in lung cancer patients during radiotherapy will help reduce treatment-




Figure 4 Expression of (A) IL-6 and (B) TNF- α in serum of post-irradiation rats. *P<0.05.




Figure 5 Co-cultured with the BMSCs-miR-21 trigger the M1 polarization of AMs. (A) AMs were cocultured with BMSCs-miR-21, or BMSCs-miR-NC or not, followed by 200 ng/L LPS stimulation for 24 h. (A,B,C) mRNA expression of IL-6, IL-12 and TNF- α were determined by qRT-PCR. (D) Nitric oxide production was determined by Griess assay. **P<0.01; ***P<0.001; NS, not significant.

related mortality, thereby improving the overall lung cancer survival (1,2). At present, several research groups are engaged in the search for promising RILI biomarkers.

Groves *et al.* found that CCSP/SP-D ratio may be useful as a biomarker of radiation-induced pulmonary fibrosis (15). Wang *et al.* suggested that IL-8 and TGF- β 1 could predict

RILI toxicity in NSCLC patients (16). Sun *et al.* have shown that serum miRNA signature could predict response to radiation therapy in NSCLC (17). Generally, these targets are closely related to the development of acute inflammation (11). Our study demonstrated that low miR-21 levels could predict higher RILI incidence and grade, lower pulmonary function in patients, and enhancement of IL-6 and TNF- α expressions, especially during the acute phase. These findings suggest that miR-21 can be used as a potential marker for RILI diagnosis and prognosis. Based on our findings, we also suggest that miR-21 may play a regulatory role in the pathogenesis of RILI.

Injection of miRNA-overexpressing BMSCs is considered a viable method for introducing exogenous miRNAs into the body because BMSCs can deliver miRNAs to the target cells via exosomes (18). In the present study, pretreatment with BMSC-miR-21-OV was more effective at preventing radiation-induced pulmonary injury during the acute inflammatory phase, when compared to the injection of BMSC-NT. Rats in the BMSC-miR-21-OV group exhibited more attenuated inflammation than did those in the BMSC-NT treatment group, suggesting a protective effect for miR-21. Furthermore, cell-based experiments confirmed that this therapeutic effect was achieved by targeting inflammatory factors, such as TNF- α , and regulating M1 polarization of pulmonary macrophages. These findings suggest that pretreatment with BMSC-miR-21-OV may be a novel and effective management strategy for patients undergoing radiotherapy at the early acute inflammatory stage. However, owing to the high prevalence of pulmonary interstitial fibrosis at late stages of RILI (19,20), we should further explore whether miR-21 can regulate pulmonary interstitial fibrosis formation. In addition, data on the systematic toxicity of MSC-miR-21 treatment are currently missing. Therefore, long-term analysis of the outcome of MSC-miR-21 treatment is necessary to realize its potential.

Acknowledgments

Funding: This work was supported by the Youth cultivation project of PLA (17QNP034).

Footnote

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tcr.2019.12.77>). The authors have no conflicts

of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was approved by the ethics committee of the Eighth Medical Center of PLA General Hospital and all the experiments were carried out according to principles of Helsinki Declaration. Informed consent was waived.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Bernchou U, Christiansen RL, Asmussen JT, et al. Extent and computed tomography appearance of early radiation induced lung injury for non-small cell lung cancer. *Radiother Oncol* 2017;123:93-8.
- Bao P, Zhao W, Li Y, et al. Protective effect of ulinastatin in patients with non-small cell lung cancer after radiation therapy: a randomized, placebo-controlled study. *Med Oncol* 2015;32:405.
- Hanania AN, Mainwaring W, Ghebre YT, et al. Radiation-Induced Lung Injury: Assessment and Management. *Chest* 2019;156:150-62.
- Domańska-Senderowska D, Laguet MN, Jegier A, et al. MicroRNA Profile and Adaptive Response to Exercise Training: A Review. *Int J Sports Med* 2019;40:227-35.
- Wu C, Wang X, Zhang J, et al. MicroRNA-224 Expression and Polymorphism Predict the Prognosis of Hepatitis B Virus-Related Hepatocellular Carcinoma Patients After Liver Resection. *Clin Lab* 2019;65. doi: 10.7754/Clin.Lab.2018.181025.
- Long Y, Zhu Y. Identification of FBXW7 α -regulated genes in M1-polarized macrophages in colorectal cancer by RNA sequencing. *Saudi Med J* 2019;40:766-73.
- Xu T, Zhang Y, Chang P, et al. Mesenchymal stem cell-based therapy for radiation-induced lung injury. *Stem Cell Res Ther* 2018;9:18.

8. Xu R, Shen X, Si Y, et al. MicroRNA-31a-5p from aging BMSCs links bone formation and resorption in the aged bone marrow microenvironment. *Aging Cell* 2018;17:e12794.
9. Hu Y, Xu F, Zhang R, et al. Interleukin-1 β -induced IRAK1 ubiquitination is required for TH-1GM-CSF cell differentiation in T cell-mediated inflammation. *J Autoimmun* 2019;102:50-64.
10. Bao P, Gao W, Li S, et al. Effect of pretreatment with high-dose ulinastatin in preventing radiation-induced pulmonary injury in rats. *Eur J Pharmacol* 2009;603:114-9.
11. Tian X, Wang F, Luo Y, et al. Protective Role of Nuclear Factor-Erythroid 2-Related Factor 2 Against Radiation-Induced Lung Injury and Inflammation. *Front Oncol* 2018;8:542.
12. Qiu YF, Wang MX, Meng LN, et al. MiR-21 regulates proliferation and apoptosis of oral cancer cells through TNF- α . *Eur Rev Med Pharmacol Sci* 2018;22:7735-41.
13. Yang X, Pan Y, Xu X, et al. Sialidase Deficiency in *Porphyromonas gingivalis* Increases IL-12 Secretion in Stimulated Macrophages Through Regulation of CR3, lncRNA GAS5 and miR-21. *Front Cell Infect Microbiol* 2018;8:100.
14. Jeon EJ, Kim DY, Lee NH, et al. Telmisartan induces browning of fully differentiated white adipocytes via M2 macrophage polarization. *Sci Rep* 2019;9:12336.
15. Groves AM, Williams JP, Hernady E, et al. A Potential Biomarker for Predicting the Risk of Radiation-Induced Fibrosis in the Lung. *Radiat Res* 2018;190:513-25.
16. Wang S, Campbell J, Stenmark MH, et al. Plasma Levels of IL-8 and TGF- β 1 Predict Radiation-Induced Lung Toxicity in Non-Small Cell Lung Cancer: A Validation Study. *Int J Radiat Oncol Biol Phys* 2017;98:615-21.
17. Sun Y, Hawkins PG, Bi N, et al. Serum MicroRNA Signature Predicts Response to High-Dose Radiation Therapy in Locally Advanced Non-Small Cell Lung Cancer. *Int J Radiat Oncol Biol Phys* 2018;100:107-14.
18. Tian X, Wang F, Luo Y, et al. Protective Role of Nuclear Factor-Erythroid 2-Related Factor 2 Against Radiation-Induced Lung Injury and Inflammation. *Front Oncol* 2018;8:542.
19. Chen L, Lu FB, Chen DZ, et al. BMSCs-derived miR-223-containing exosomes contribute to liver protection in experimental autoimmune hepatitis. *Mol Immunol* 2018;93:38-46.
20. Huang Y, Zhang W, Yu F, et al. The Cellular and Molecular Mechanism of Radiation-Induced Lung Injury. *Med Sci Monit* 2017;23:3446-50.

Cite this article as: Bao P, Zhao W, Mou M, Liu X. MicroRNA-21 mediates bone marrow mesenchymal stem cells protection of radiation-induced lung injury during the acute phase by regulating polarization of alveolar macrophages. *Transl Cancer Res* 2020;9(1):231-239. doi: 10.21037/tcr.2019.12.77