

PROCEEDINGS

Open Access

EXPath: a database of comparative expression analysis inferring metabolic pathways for plants

Chia-Hung Chien, Chi-Nga Chow, Nai-Yun Wu, Yi-Fan Chiang-Hsieh, Ping-Fu Hou, Wen-Chi Chang*

From The Thirteenth Asia Pacific Bioinformatics Conference (APBC 2015)
HsinChu, Taiwan. 21-23 January 2015

Abstract

Background: In general, the expression of gene alters conditionally to catalyze a specific metabolic pathway. Microarray-based datasets have been massively produced to monitor gene expression levels in parallel with numerous experimental treatments. Although several studies facilitated the linkage of gene expression data and metabolic pathways, none of them are amassed for plants. Moreover, advanced analysis such as pathways enrichment or how genes express under different conditions is not rendered.

Description: Therefore, EXPath was developed to not only comprehensively congregate the public microarray expression data from over 1000 samples in biotic stress, abiotic stress, and hormone secretion but also allow the usage of this abundant resource for coexpression analysis and differentially expression genes (DEGs) identification, finally inferring the enriched KEGG pathways and gene ontology (GO) terms of three model plants: *Arabidopsis thaliana*, *Oryza sativa*, and *Zea mays*. Users can access the gene expression patterns of interest under various conditions via five main functions (Gene Search, Pathway Search, DEGs Search, Pathways/GO Enrichment, and Coexpression analysis) in EXPath, which are presented by a user-friendly interface and valuable for further research.

Conclusions: In conclusion, EXPath, freely available at <http://expath.itps.ncku.edu.tw>, is a database resource that collects and utilizes gene expression profiles derived from microarray platforms under various conditions to infer metabolic pathways for plants.

Background

Plants, which are classified as the kingdom Plantae, provide source of energy and oxygen in ecosystems and the majority of agricultural production worldwide. To maintain the autotrophic mechanisms as well as the resistance to impacts from surroundings (e.g., extreme weather, soil salinity, and pests), the elaborate control of gene expression and collaboration under various environments or conditions at molecular level is critical and related to growth, development, and the yield of crop production in plants [1]. Since the lack of motility compels plants to be more tolerant against the threat of external stresses, genes involved in stress-related response, signal transduction pathways, and the induced transcription factors (TFs) were progressively discovered through the comparative genomics approaches [2-5].

Moreover, phytohormones, which are believed to modulate plant growth and diverse development processes, have been reported in relation to environmental variation in *Arabidopsis* and maize [6-8]. The evidence reveals that the complexity of gene regulation in significant pathways or biochemical reactions plays an important role in coping with plants survival and their self-defense mechanisms towards different circumstances.

In general, the expression of gene alters conditionally to catalyze a specific metabolic pathway [9,10]. Comprehensively investigating how genes are activated or repressed, i.e., differentially expressed genes (DEGs), in vital biological processes under various conditions are essential to understand gene functions and the coexpression manner in metabolic routes. In recent decades, microarray-based datasets have been massively produced to monitor gene expression levels in parallel with numerous experimental treatments [11]. This high-throughput

College of Biosciences and Biotechnology, Institute of Tropical Plant Sciences, National Cheng Kung University, Tainan 70101, Taiwan

detection of transcript quantity facilitates the comparative expression analysis by combining multiple microarray expression data among different samples and even different species [12]. Due to the abundance of expression datasets generated by microarray platforms for plants, a plenty of databases and resources have promptly collected gene expression data that are publicly accessible. Among them, Gene Expression Omnibus (GEO) provides most profuse microarray expression datasets presented with the function of GEO DataSets, GEO Profiles, and GEO2R Analysis [13]. Although GEO2R Analysis allows users to compare multiple expression data and then identify DEGs, GEO Profiles can only display the expression level of one gene across different samples in each dataset. Another powerful tool, eFP Browser, is easily adaptable for analyzing microarray or other large-scale datasets in plants by using pictographic representations [14]. Additionally, PLEXdb, GENEVESTIGATOR, NASCArrays, and RiceXPro are also useful repositories for microarray gene expression profiles in *Arabidopsis*, rice, and plants [15-18]. On the other hand, to gain a comprehensive insight into plant metabolic pathways that are consisted of metabolites and enzymes, relevant databases were established recently. Gramene, a comparative resource for plants, summarizes ten databases of plant metabolic pathways, e.g., AraCyc, RiceCyc, MaizeCyc, BrachyCyc and SorghumCyc [19]. Moreover, MetNet Online integrates information of metabolic pathways and regulatory networks for *Arabidopsis thaliana*, *Glycine max* and *Vitis vinifera* [20]. Other instances of similar pathway knowledge bases are *Arabidopsis* reactome and Pathway studio [21,22].

To estimate the expression level of each gene involved in significant biological processes thoroughly, it is important to integrate gene expression data with metabolic pathways. To our current knowledge, Jensen and Papin have presented a method, Metabolic Adjustment by Differential Expression (MADE), for mapping expression data onto a metabolic network model without using arbitrary expression thresholds. Unfortunately, MADE is implemented in Matlab and only supports for *Saccharomyces cerevisiae* [23]. Another web-based tool, Array2KEGG, attempted to depict up or down regulated genes in a particular KEGG pathway image of interest. However, the system were developed for human, mouse, and rat. Furthermore, although Pathway Processor 2.0 is a web resource for converting gene expression into pathway expression and identifying differentially regulated pathways in an input datasets, users have to submit their own expression data, and cannot be apply for plants [24]. Besides, advanced analysis such as pathways enrichment or how genes express under different conditions is not rendered. It is noted that AlgaePath that we published previously makes feasible to analyze metabolic pathways using transcript abundance data from next-generation sequencing in green

algae [25], but it is created for non-vascular plants. Therefore, EXPath was developed to not only comprehensively congregate the public microarray expression data from over 1000 samples in biotic stress, abiotic stress, and hormone secretion but also allow the usage of this abundant resource for coexpression analysis and DEGs identification, finally inferring the enriched KEGG pathways and gene ontology (GO) terms of three model plants: *Arabidopsis thaliana*, *Oryza sativa*, and *Zea mays*. Users can access the gene expression patterns of interest under various conditions via five main functions (Gene Search, Pathway Search, DEGs Search, Pathways/GO Enrichment, and Coexpression analysis) in EXPath, which are presented by a user-friendly interface and valuable for further research. The concept and construction of EXPath is illustrated in Figure 1.

Construction and content


Repository for microarray gene expression data

To establish overarching repository for gene expression in plants, EXPath curates 1057 samples treated with biotic stress, abiotic stress, and hormone secretion from publicly available microarray gene expression data for *Arabidopsis thaliana*, *Oryza sativa*, and *Zea mays*. Among them, expression profiles of *Arabidopsis thaliana* were retrieved from AtGenExpress [26] and NASCArrays [18], whereas others were obtained via RiceXPro (*Oryza sativa*) and GEO (*Oryza sativa* and *Zea mays*) [13,16]. Table 1 summarizes the categories and number of microarray samples collected in EXPath expression database.

Data processing and normalization

Since the platforms for inclusive microarray datasets in EXPath are either Affymetrix GeneChips (GPL189, GPL2025, and GPL432) or Agilent Technologies (GPL6864), probe set annotation data in tabular format were used to map probe set ids to detectable genes. For the purpose of optimizing input data for comparative expression analysis, ambiguous measurements of gene expression, for example, probes associated with more than two genomic loci or control probes, were discarded. In total, 20922 genes for *Arabidopsis thaliana*, 22769 genes for *Oryza sativa*, and 10724 genes for *Zea mays* were congregate respectively. Moreover, genes matched with KEGG genes and KEGG orthology (KO) were also calculated. The statistics of genes with valid expression data in EXPath is displayed in Figure 2.

After processing the raw data of microarray datasets, the normalization procedure was executed to avoid systematic biases arising from the variation between different trials (GEO series/GSEs) and samples. In this work, robust multi-array average (RMA) was performed by using the justRMA function in *affy* package, which is a part of the BioConductor project [27,28]. For those



genes with raw intensities from multiple probes and replicates, we first filtered the outliers by the interquartile range (IQR) rule and retained data between the upper and lower quartiles. Then, the average of all reserved replicates was calculated to represent the expression level for each gene in given conditions.

Collection of annotation files

EXPath offers gene general information including descriptions, cDNA and protein sequences, Pfam protein families, GO terms, and involved pathways for users' reference. The annotation files of *Arabidopsis thaliana*, *Oryza sativa*, and *Zea mays* were downloaded from TAIR10, RAP-DB, and MaizeGDB separately [29-31]. For descriptions, cDNA and protein sequences, and Pfam protein families, except the descriptions of *Arabidopsis thaliana* were from TAIR10 and the descriptions and

sequences of *Oryza sativa* were from RAP-DB, other datasets were acquired by using Ensembl BioMarts [32]. The latest GO terms and involved pathways were collected from gene ontology consortium and KEGG database [33,34].


Comparative expression analysis

Differentially expressed genes

To determine genes that are differentially expressed under given conditions, t-test statistic method was applied by using function `t.test()` of R package in EXPath. Users can specify a treatment from biotic stress, abiotic stress, or hormone secretion that are well-categorized for three model plants, and then set the time point, fold change and p-value cutoffs. Statistics of fold change and DEG lists (up-regulated and down-regulated) are also provided in EXPath.

Table 1 Categories of microarray samples in EXPath expression database.

Species	Stresses	# of samples	Resource			
<i>Arabidopsis thaliana</i>	Abiotic stress	Cold stress	298	AtGenExpress [26] NASCArrays [18]		
		Drought stress				
		Genotoxic stress				
		Heat stress				
		Osmotic stress				
		Oxidative stress				
		Salt stress				
		UV-B stress				
	Wounding stress	108				
	Biotic stress		<i>Phytophthora infestans</i> <i>P.syringae pv.tomato</i> DC3000			
	Hormones	ABA	96			
		Auxin				
		Brassinolide				
Brassinosteroids						
GA						
<i>Oryza sativa</i>	Abiotic stress	Gibberellin	94	RiceXPro [16] GEO [13]		
		Jasmonic acid				
		Zeatin (cytokinin)				
	Abiotic stress	Cold stress			94	
		Salt stress				
	Abiotic stress	Drought stress			94	
		Biotic stress				<i>Agrobacterium tumefaciens</i> <i>Magnaporthe oryzae</i> <i>Magnaporthe oryzae</i> strain Guy11 <i>Meloidogyne graminicola</i> <i>Striga hermonthica</i> <i>X. oryzae pv. oryzae</i> <i>X. oryzae pv. oryzicola</i>
	Hormones	Abscisic acid			138	
		Auxin				
		Brassinosteroid				
		Cytokinin				
		Gibberellin				
<i>Zea mays</i>	Abiotic stress	Jasmonic acid	64	GEO [13]		
		Acid soil				
		Drought				
	Abiotic stress	Waterlogging			64	
		Biotic stress				<i>Colletotrichum graminicola</i> <i>Meloidogyne incognita</i> <i>Phytophthora cinnamomi</i> <i>Sporisorium reilianum</i> f. sp. <i>Zeeae</i> (Kühn) <i>Ustilago maydis</i>
		Biotic stress				<i>Colletotrichum graminicola</i> <i>Meloidogyne incognita</i> <i>Phytophthora cinnamomi</i> <i>Sporisorium reilianum</i> f. sp. <i>Zeeae</i> (Kühn) <i>Ustilago maydis</i>
Hormones	GA3	6				
	IAA					



Coexpression gene groups

Co-expressed genes are a group of genes that express simultaneously under specific conditions. Theoretically, they tend to be controlled under similar transcriptional regulation and involve in identical biological processes or pathways. To investigate this concept, we calculated the coexpression levels of 111 KEGG pathways with number of genes more than 10 by using Pearson's correlation coefficient (PCC) in *Arabidopsis thaliana*. Among them, 92.6% of pathways are positively correlated with satisfied PCCs (most of them are between 0.6 ~ 0.9, see Figure S1, Additional file 1), which suggest that genes involved in the same pathway are generally co-expressed. In EXPath, Pearson's correlation coefficient

and Spearman's rank correlation coefficient are applied by using `cor()` functions in R package to identify genes with co-expression patterns. Normalized raw intensities of genes without log transformation were used to calculate correlation coefficient because it may alter original expression levels that we mentioned previously [35]. Users can customize positive/negative correlation and the conditions (abiotic stress, biotic stress, hormone treatment, and overall conditions) they intend to explore. The expression patterns of coexpression gene groups are illustrated based on z-score transformation:

$$z = \frac{x - \mu}{\sigma}$$

The character z denotes z -score in the above formula, whereas x , μ , and σ represent the raw intensity, mean, and standard deviation of gene expression levels respectively.

Enriched KEGG pathways and GO terms

As we know, given a gene list involved in significant biological processes, signal transduction, or metabolic pathways, the pathogenicity of diseases and the roles of these genes can be inferred. It helps researchers to determine biomarkers or treatments for specific diseases. Here, the cumulative probability (p -value) of hypergeometric distribution was calculated to evaluate the KEGG/GO enrichment of a group of input genes. The formula is as follows:

$$p(X \leq k) = \sum_{i=x}^n \frac{\binom{M}{i} \binom{N-M}{n-i}}{\binom{N}{n}}$$

where N and M denote the number of background genes and total genes involved in specific KEGG pathways or GO terms, whereas i genes out of n genes in the gene group \times belong to that KEGG pathways or GO terms. The usage of $dhyper()$ and $phyper()$ in R were applied to obtain hypergeometric p -values for each gene group.

Utility and discussion

Basic implement in EXPath

EXPath offers a user-friendly interface for exploring Gene Search, Pathway Search, DEGs Search, Pathways/GO Enrichment, and Coexpression analysis. The introduction and guideline for users can be retrieved from the main page. After selecting the EXPath function of interest, users have to specify a model plant first (Figure 3A). For Gene Search and Coexpression analysis, keywords such as HGNC symbol, description, database (pfam, KEGG,





Figure 3 The output result of "Coexpression analysis" in EXPath.

TAIR) ID, microarray probe ID, or sequence are valid as query input. Alternatively, users can utilize Gene Browser to access gene lists of three model plants categorized by chromosome. Figure 3B demonstrates a sketch of search result in Coexpression analysis. Moreover, to start a Pathway Search, users can either input a keyword or browse all available pathways classified by metabolism, genetic information processing, environmental information processing, cellular processes, and organismal systems. Prior to the display of KEGG pathway map, users are obligated to select at least one condition from sample list provided in EXPath. By following up the instructions and procedures of each function, the ultimate output is presented and interpreted systematically by users' request. Figure 3 instances an output of Coexpression analysis.

Advanced combination analysis in EXPath

In addition to explore five functions provided in EXPath separately, all of them are connected with each other using the linkage buttons or hyperlinks in output web-pages. For example, in the Gene Search result page,

EXPath not only maps the query gene to KEGG pathways to illustrate the involvement of that gene in the corresponding pathway map with its microarray expression levels under specified conditions but also furnishes the linkage button for performing coexpression analysis. Furthermore, advanced combination analysis, the most practical application in EXPath, exposes the powerful pipeline for comparative expression analysis in plants. By combining DEGs Search with Pathways/GO Enrichment, the differentially expressed genes between control and treatment samples are identified first. Then, users can designate up-regulated genes, down-regulated genes, or all DEGs to perform Pathways/GO Enrichment. The enriched KEGG pathways or GO terms of DEGs helps plant scientists to understand, for instance, the resistance to abiotic stresses, pathogenicity of microbes or viruses, and even hormone treatments. Another combination analysis of Coexpression analysis and Pathways/GO Enrichment aims to distinguish coexpressed genes for exhaustively inferring gene functions and their biological roles. A case study given below describes the details of this application.



Case study: the JAZ10

Here, we demonstrate a case study of JASMONATE ZIM-domain protein 10 (JAZ10), which belongs to the TIFY family and is one of the critical repressor in jasmonate signalling [36,37]. Jasmonate (JA) is known to be an essential phytohormone regulating defense mechanisms to pathogens, plant reproductive development, and response to stresses from various environments [38]. Since JAZ family is sensitive to JA treatment, a previous study indicated that the alternative splice variant of JAZ10 plays a significant role in repressing transcription factors that activate the expression of JA response genes in *Arabidopsis* [39]. Based on these valuable findings, we performed the combination approach of Coexpression analysis and Pathways/GO Enrichment to evaluate the usage of EXPath. First of all, 165 coexpressed genes of JAZ10 were identified by setting the parameters of Pearson's correlation coefficient, positive correlation, and hormone treatment (See Table S1, Additional file 2 for coexpressed gene list). Figure 4 shows the z-score transformed expression patterns of top 20 genes coexpressed with JAZ10. A dramatic peak was observed after the methyl jasmonate (MJ) treatment for an hour. Next, the matched genes were submitted to execute pathways and GO enrichment. As expected, both top 10 enriched

pathways and GO terms ordered by p-value are consistent with previous findings, e.g., plant-pathogen interaction (Table 2 and 3). It reveals the utility of EXPath for performing comparative expression analysis with high reliability.

Conclusions

EXPath is an overarching repository geared towards plant scientists to facilitate the retrieval of microarray gene expression data from publicly available resources and the analysis of comparative expression. As the novel database integrating gene expression data with metabolic pathways, the inferred pathways give an insight into the discovery of gene functions, pathogenicity of external invasion, and defense mechanisms for plants. By the usage of five main functions (i.e., Gene Search, Pathway Search, DEGs Search, Pathways/GO Enrichment, and Coexpression analysis) and the advanced combination analysis of them, EXPath indeed provides an effective interface for users to explore the information of interest that will be valuable for further research. Although EXPath facilitates the comparison of expression levels among genes involved in designated pathways, the limited number of plant genes recruited in KEGG database restricts the availability for comparative expression analysis. Another limitation is

Table 2 Enriched pathways of JAZ10 coexpressed gene group (partial, only shows top 10 results).

Pathway ID	Pathway name	Hit number (Query)	Percentage in query	P-value
04626	Plant-pathogen interaction	13	29.55%	3.56E-9
00592	alpha-Linolenic acid metabolism	6	13.64%	4.22E-7
00591	Linoleic acid metabolism	3	6.82%	7.45E-5
00400	Phenylalanine, tyrosine and tryptophan biosynthesis	5	11.36%	2.22E-4
00920	Sulfur metabolism	4	9.09%	6.66E-4
04075	Plant hormone signal transduction	8	18.18%	4.16E-3
00966	Glucosinolate biosynthesis	2	4.55%	0.01
00950	Isoquinoline alkaloid biosynthesis	2	4.55%	0.02
01230	Biosynthesis of amino acids	6	13.64%	0.04
00960	Tropane, piperidine and pyridine alkaloid biosynthesis	2	4.55%	0.05

Table 3 Enriched GO terms of JAZ10 coexpressed gene group (partial, only shows top 10 results).

GO ID	GO term	Hit number (Query)	Percentage in query	P-value
GO:0009753	response to jasmonic acid	46	29.11%	8.85E-55
GO:0009611	response to wounding	48	30.38%	1.09E-53
GO:0009695	jasmonic acid biosynthetic process	35	22.15%	2.11E-48
GO:0009620	response to fungus	28	17.72%	3.29E-39
GO:0009738	abscisic acid-activated signaling pathway	29	18.35%	7.19E-28
GO:0009867	jasmonic acid mediated signaling pathway	29	18.35%	1.25E-27
GO:0006950	response to stress	85	53.80%	3.71E-27
GO:0007165	signal transduction	59	37.34%	3.62E-26
GO:0009723	response to ethylene	26	16.46%	4.96E-25
GO:0042538	hyperosmotic salinity response	20	12.66%	5.86E-21

insufficient expression datasets in public for other plants rather than *Arabidopsis*, rice, and maize. For perspectives, in addition to the expectation of more available plant genes in KEGG database, we will keep surveying any relevant sample with expression profile released in public, especially for those derived from the treatments of biotic stress, abiotic stress, hormone secretion, and even development.

Availability and requirements

The EXPath database is publicly available at <http://EXPath.itsps.ncku.edu.tw>.

Additional material

Additional file 1: Figure S1.

Additional file 2: Table S1.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

WCC conceived and designed the experiments and revised the paper. CHC, CNC, and NYW developed the database and webpage. CNC, NYW, YFCH and PFH analyzed the data and performed the experiments. CHC wrote the paper.

Acknowledgements/Declarations

This research was supported by a grant from National Science Council of the Republic of China for financially supporting this research under Contract NSC 102-2313-B-006 -004 and MOST 103-2311-B-006 -001.

This article has been published as part of *BMC Genomics* Volume 16 Supplement 2, 2015: Selected articles from the Thirteenth Asia Pacific Bioinformatics Conference (APBC 2015): Genomics. The full contents of the supplement are available online at <http://www.biomedcentral.com/bmcgenomics/supplements/16/S2>

Published: 21 January 2015

References

- Atkinson NJ, Urwin PE: The interaction of plant biotic and abiotic stresses: from genes to the field. *Journal of experimental botany* 2012, **63**(10):3523-3543.
- Wang W, Vinocur B, Altman A: Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta* 2003, **218**(1):1-14.
- Cushman JC, Bohnert HJ: Genomic approaches to plant stress tolerance. *Current opinion in plant biology* 2000, **3**(2):117-124.
- Mittler R: Abiotic stress, the field environment and stress combination. *Trends in plant science* 2006, **11**(1):15-19.
- Rizhsky L, Liang H, Shuman J, Shulaev V, Davletova S, Mittler R: When defense pathways collide. The response of *Arabidopsis* to a combination of drought and heat stress. *Plant physiology* 2004, **134**(4):1683-1696.
- Ren H, Gao Z, Chen L, Wei K, Liu J, Fan Y, Davies WJ, Jia W, Zhang J: Dynamic analysis of ABA accumulation in relation to the rate of ABA catabolism in maize tissues under water deficit. *Journal of experimental botany* 2007, **58**(2):211-219.
- Gray WM: Hormonal regulation of plant growth and development. *PLoS biology* 2004, **2**(9):E311.
- Wang Y, Liu C, Li K, Sun F, Hu H, Li X, Zhao Y, Han C, Zhang W, Duan Y, et al: *Arabidopsis* EIN2 modulates stress response through abscisic acid response pathway. *Plant molecular biology* 2007, **64**(6):633-644.
- Boavida LC, Borges F, Becker JD, Feijo JA: Whole genome analysis of gene expression reveals coordinated activation of signaling and metabolic pathways during pollen-pistil interactions in *Arabidopsis*. *Plant physiology* 2011, **155**(4):2066-2080.
- Yonekura-Sakakibara K, Tohge T, Matsuda F, Nakabayashi R, Takayama H, Niida R, Watanabe-Takahashi A, Inoue E, Saito K: Comprehensive flavonol profiling and transcriptome coexpression analysis leading to decoding gene-metabolite correlations in *Arabidopsis*. *The Plant cell* 2008, **20**(8):2160-2176.
- Schena M, Shalon D, Davis RW, Brown PO: Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* 1995, **270**(5235):467-470.
- Movahedi S, Van Bel M, Heyndrickx KS, Vandepoele K: Comparative co-expression analysis in plant biology. *Plant, cell & environment* 2012, **35**(10):1787-1798.
- Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Holko M, et al: NCBI GEO: archive for functional genomics data sets—update. *Nucleic acids research* 2013, **41** Database: D991-995.
- Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ: An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets. *PLoS one* 2007, **2**(8):e718.
- Dash S, Van Hemert J, Hong L, Wise RP, Dickerson JA: PLEXdb: gene expression resources for plants and plant pathogens. *Nucleic acids research* 2012, **40** Database: D1194-1201.
- Sato Y, Takehisa H, Kamatsuki K, Minami H, Namiki N, Ikawa H, Ohyanagi H, Sugimoto K, Antonio BA, Nagamura Y: RiceXPro version 3.0: expanding the informatics resource for rice transcriptome. *Nucleic acids research* 2013, **41** Database: D1206-1213.
- Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W: GENEVESTIGATOR. *Arabidopsis* microarray database and analysis toolbox. *Plant physiology* 2004, **136**(1):2621-2632.
- Craigon DJ, James N, Okyere J, Higgins J, Jotham J, May S: NASCArrays: a repository for microarray data generated by NASC's transcriptomics service. *Nucleic acids research* 2004, **32** Database: D575-577.
- Monaco MK, Stein J, Naithani S, Wei S, Dharmawardhana P, Kumari S, Amarasinghe V, Youens-Clark K, Thomason J, Preece J, et al: Gramene 2013: comparative plant genomics resources. *Nucleic acids research* 2014, **42** Database: D1193-1199.
- Sucaet Y, Wang Y, Li J, Wurtele ES: MetNet Online: a novel integrated resource for plant systems biology. *BMC bioinformatics* 2012, **13**:267.
- Nikitin A, Egorov S, Daraselia N, Mazo I: Pathway studio—the analysis and navigation of molecular networks. *Bioinformatics* 2003, **19**(16):2155-2157.
- Tsesmetzis N, Couchman M, Higgins J, Smith A, Doonan JH, Seifert GJ, Schmidt EE, Vastrik I, Birney E, Wu G, et al: *Arabidopsis* reactome: a foundation knowledgebase for plant systems biology. *The Plant cell* 2008, **20**(6):1426-1436.
- Jensen PA, Papin JA: Functional integration of a metabolic network model and expression data without arbitrary thresholding. *Bioinformatics* 2011, **27**(4):541-547.
- Beltrame L, Bianco L, Fontana P, Cavalieri D: Pathway Processor 2.0: a web resource for pathway-based analysis of high-throughput data. *Bioinformatics* 2013, **29**(14):1825-1826.
- Zheng HQ, Chiang-Hsieh YF, Chien CH, Hsu BK, Liu TL, Chen CN, Chang WC: AlgaePath: comprehensive analysis of metabolic pathways using transcript abundance data from next-generation sequencing in green algae. *BMC genomics* 2014, **15**:196.
- Kilian J, Whitehead D, Horak J, Wanke D, Weinl S, Batistic O, D'Angelo C, Bornberg-Bauer E, Kudla J, Harter K: The AtGenExpress global stress expression data set: protocols, evaluation and model data analysis of UV-B light, drought and cold stress responses. *The Plant journal: for cell and molecular biology* 2007, **50**(2):347-363.
- Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed TP: Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 2003, **4**(2):249-264.
- Gautier L, Cope L, Bolstad BM, Irizarry RA: affy-analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics* 2004, **20**(3):307-315.
- Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher K, Alexander DL, Garcia-Hernandez M, et al: The *Arabidopsis* Information Resource (TAIR): improved gene annotation and new tools. *Nucleic acids research* 2012, **40** Database: D1202-1210.
- Sakai H, Lee SS, Tanaka T, Numa H, Kim J, Kawahara Y, Wakimoto H, Yang CC, Iwamoto M, Abe T, et al: Rice Annotation Project Database

- (RAP-DB): an integrative and interactive database for rice genomics. *Plant & cell physiology* 2013, **54**(2):e6.
31. Schaeffer ML, Harper LC, Gardiner JM, Andorf CM, Campbell DA, Cannon EK, Sen TZ, Lawrence CJ: **MaizeGDB: curation and outreach go hand-in-hand.** *Database: the journal of biological databases and curation* 2011, **2011**:bar022.
 32. Kinsella RJ, Kahari A, Haider S, Zamora J, Proctor G, Spudich G, Almeida-King J, Staines D, Derwent P, Kerhornou A, *et al*: **Ensembl BioMarts: a hub for data retrieval across taxonomic space.** *Database: the journal of biological databases and curation* 2011, **2011**:bar030.
 33. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, *et al*: **Gene ontology: tool for the unification of biology.** The Gene Ontology Consortium. *Nature genetics* 2000, **25**(1):25-29.
 34. Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M: **Data, information, knowledge and principle: back to metabolism in KEGG.** *Nucleic acids research* 2014, **42** Database: D199-205.
 35. Chien CH, Chiang-Hsieh YF, Tsou AP, Weng SL, Chang WC, Huang HD: **Large-Scale Investigation of Human TF-miRNA Relations Based on Coexpression Profiles.** *BioMed research international* 2014, **2014**:623078.
 36. Vanholme B, Grunewald W, Bateman A, Kohchi T, Gheysen G: **The tify family previously known as ZIM.** *Trends in plant science* 2007, **12**(6):239-244.
 37. Chini A, Fonseca S, Fernandez G, Adie B, Chico JM, Lorenzo O, Garcia-Casado G, Lopez-Vidriero I, Lozano FM, Ponce MR, *et al*: **The JAZ family of repressors is the missing link in jasmonate signalling.** *Nature* 2007, **448**(7154):666-671.
 38. Moreno JE, Shyu C, Campos ML, Patel LC, Chung HS, Yao J, He SY, Howe GA: **Negative feedback control of jasmonate signaling by an alternative splice variant of JAZ10.** *Plant physiology* 2013, **162**(2):1006-1017.
 39. Chung HS, Howe GA: **A critical role for the TIFY motif in repression of jasmonate signaling by a stabilized splice variant of the JASMONATE ZIM-domain protein JAZ10 in Arabidopsis.** *The Plant cell* 2009, **21**(1):131-145.

doi:10.1186/1471-2164-16-S2-S6

Cite this article as: Chien *et al*: EXPath: a database of comparative expression analysis inferring metabolic pathways for plants. *BMC Genomics* 2015 **16**(Suppl 2):S6.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

