

ORAL PRESENTATION

Open Access

Pre-clinical evaluation of novel anti-tuberculosis molecules

Maria Martha Campos

From 5th Congress of the Brazilian Biotechnology Society (SBBIOTEC)
Florianópolis, Brazil. 10-14 November 2013

Tuberculosis (TB) currently represents a major global health concern, especially when considering the emergence of multidrug-resistant strains of *Mycobacterium tuberculosis*. On this regard, the identification of new effective agents to treat this infectious disease is urgently needed. Herein, we describe the pre-clinical assays on the anti-tubercular effects of a pentacyano(isoniazid)ferrate(II) compound, namely IQG-607. Previous experimental evidence [1] clearly demonstrated the *in vivo* effectiveness of this compound in a mouse model of TB infection. Accordingly, IQG-607 (22 to and 560 $\mu\text{mol/kg}$) was able to markedly reduce the number of colony-forming units (CFU) in both spleen and lungs of *M. tuberculosis* H37Rv-infected animals, following long-term schemes of oral administration, during 28 or 56 days. In this series of experiments, the effects of IQG-607 were comparable to those observed for the reference compound isoniazid (182 $\mu\text{mol/kg}$). More recently, we demonstrated that *in vitro* incubation of IQG-607 led to a marked reduction of CFU counts in *M. tuberculosis* H37Rv-infected macrophages, being this effect similar to that displayed by either isoniazid or rifampicin [2]. Concerning the mechanism of action, IQG-607 was found able to block the biosynthesis of mycolic acids, as indicated by radiolabelling studies with acetate incorporation [2], and it displayed bactericidal effects when tested *in vivo* [1]. The acute oral administration of IQG-607 (560 to 1120 $\mu\text{mol/kg}$) was not related to any clinical sign of toxicity, whereas the treatment with isoniazid (182 $\mu\text{mol/kg}$) led to death rates of 80% in mice [3]. The same doses of IQG-607 were also devoid of toxicity, when administered acutely by oral route to rats (unpublished data). To extend evidence on the safety profile of IQG-607, this molecule was also examined in a protocol of repeated dose 90-day oral toxicity in rats. As a main conclusion, there was no evidence of severe toxic signals,

although it was not possible to accurately define the NOAEL (non-toxic adverse effect level) for IQG-607. Nevertheless, the dose of 56 $\mu\text{mol/kg}$ might be assumed, once the animals treated with this dose did not show any signal of toxicity (unpublished data). Taken together, our results provide convincing evidence on the efficacy of IQG-607. It can be assumed that IQG-607 is well absorbed when dosed by oral route, being able to reach the bacilli and killing them within the phagosome of the macrophages. Furthermore, we demonstrated a favorable toxicological outcome for this compound, even when dosed acutely, or in long-term schedules of administration, for both mice and rats. Further studies are required to determine the safety of IQG-607 in non-rodent species, as well as the efficacy of this molecule against resistant strains of *M. tuberculosis*. Although additional experiments with this molecule are still required, it is tempting to infer that IQG-607 might well represent a lead compound for development of innovative anti-TB drugs.

Published: 1 October 2014

References

1. Rodrigues-Junior VS, Santos-Junior AA, Santos AJ, Schneider CZ, Calixto JB, Sousa EHS, Lopes LGF, Souto AA, Basso LA, Santos DS, Campos MM: Activity of IQG-607, a new orally active compound in a murine model of *Mycobacterium tuberculosis* infection. *International Journal of Antimicrobial Agents* 2012, **40**(2):182-185.
2. Rodrigues-Junior VS, Santos Junior AA, Villela AD, Belardinelli JM, Morbidonid HR, Basso LA, Campos MM, Santos DS: IQG-607 abrogates the synthesis of mycolic acids and displays intracellular activity against *Mycobacterium tuberculosis*-infected macrophages. *International Journal of Antimicrobial Agents* 2013, **43**(1):82-85.
3. Basso LA, Schneider CZ, Dos Santos AJAB, Santos AA JR, Campos MM, Souto AA, Santos DS: An Inorganic Complex that Inhibits *Mycobacterium tuberculosis* Enoyl Reductase as a Prototype of a New Class of Chemotherapeutic Agents to Treat Tuberculosis. *Journal of the Brazilian Chemical Society* 2010, **21**(7):1384-1389.

doi:10.1186/1753-6561-8-S4-O17

Cite this article as: Campos: Pre-clinical evaluation of novel anti-tuberculosis molecules. *BMC Proceedings* 2014 **8**(Suppl 4):O17.

PUCRS - Pontifícia Universidade do Rio Grande do Sul, Porto Alegre, Brazil



© 2014 Campos; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.