

Establishment of a prediction model of changing trends in cardiac hypertrophy disease based on microarray data screening

CAIYAN MA, YONGJUN YING, TIANJIE ZHANG, WEI ZHANG,
HUI PENG, XUFENG CHENG, LIN XU and HONG TONG

Cardiovascular Department, Tongde Hospital of Zhejiang Province, Hangzhou, Zhejiang 310012, P.R. China

Received December 29, 2014; Accepted January 15, 2016

DOI: 10.3892/etm.2016.3105

Abstract. The aim of the present study was to construct a mathematical model to predict the changing trends of cardiac hypertrophy at gene level. Microarray data were downloaded from Gene Expression Omnibus database (accession, GSE21600), which included 35 samples harvested from the heart of Wistar rats on postoperative days 1 (D1 group), 6 (D6 group) and 42 (D42 group) following aorta ligation and sham operated Wistar rats, respectively. Each group contained six samples, with the exception of the samples harvested from the aorta ligated group after 6 days, where $n=5$. Differentially expressed genes (DEGs) were identified using a Limma package in R. Hierarchical clustering analysis was performed on common DEGs in order to construct a linear equation between the D1 and D42 groups, using linear discriminant analysis. Subsequent verification was performed using receiver operating characteristic (ROC) curve and the measurement data at day 42. A total of 319, 44 and 57 DEGs were detected in D1, D6 and D42 sample groups, respectively. *AKIP1*, *ANKRD23*, *LTBP2*, *TGF- β 2* and *TNFRSF12A* were identified as common DEGs in all groups. The predicted linear equation between D1 and D42 group was calculated to be $y=1.526x-186.671$. Assessment of the ROC curve demonstrated that the area under the curve was 0.831, with a specificity and sensitivity of 0.8. As compared with the predictive and measurement data at day 42, the consistency of the two sets of data was 76.5%. In conclusion, the present model may contribute to the early prediction of changing trends in cardiac hypertrophy disease at gene level.

Introduction

Cardiac hypertrophy is associated with the thickening of the heart muscle (1) and the risk factors of cardiac hypertrophy include hypertension, obesity, muscular dystrophy, cardiomyopathy or heart failure (2). Furthermore, it has been demonstrated that genetic factors and signaling pathways may participate in the pathogenesis of cardiac hypertrophy, which may be associated with an enhanced risk of sudden cardiac death and cardiovascular mortality (3,4). As the early symptoms of this disease are difficult to detect, it is crucial that novel molecular markers for the early therapy of cardiac hypertrophy are identified.

Molecular markers of cardiac hypertrophy have been identified (5). In particular, Kontaraki *et al* (6) identified *GATA4*, *myocardin* and *β -myosin heavy chain* as early cardiac marker genes. Furthermore, smooth muscle α -actin has been demonstrated to be a molecular marker for pressure-overload hypertrophy (7). Using mouse models, Qing *et al* (8) have previously reported that *miR-22* serves a crucial function in the regulation of cardiac hypertrophy and cardiac remodeling. *Fibroblast growth factor 21*, which is an endocrine factor, has a protective role in cardiac cells (9). As an increasing number of molecular markers are identified, mathematical models can be constructed to predict the risk of cancer (10).

Various types of mathematical models have contributed to the prediction of diseases. Flux balance models of cellular metabolism have been used to analyze and predict transcriptional regulation under certain conditions, including catabolite repression and amino acid biosynthesis pathway repression (11). Furthermore, various genes and pathways associated with differentiation, including MAOA and ADH1B metabolic genes in human pulmonary type II cells (12) and nuclear factor-kappaB pathway in a mouse model of genitourinary inflammation (13), have been identified via mathematical cluster analysis using GENECLUSTER, which is a publicly available computer package that contributed to the establishment of an effective treatment for acute promyelocytic leukemia (14). According to a previous study conducted by Kondo and Miura (15), the reaction-diffusion model is effective in biological pattern formation. Thus, these previous studies suggest the mathematical modeling is a useful tool for the prediction of disease.

Correspondence to: Mr. Caiyan Ma or Tianjie Zhang, Cardiovascular Department, Tongde Hospital of Zhejiang Province, 234 Gucui Road, Hangzhou, Zhejiang 310012, P.R. China
E-mail: caiyanmacym@163.com
E-mail: mcy-wj@163.com

Key words: cardiac hypertrophy, hierarchical clustering analysis, linear discriminant analysis, mathematical model, receiver operating characteristic curve

Using microarray data downloaded from the Gene Expression Omnibus (GEO) database (accession, GSE21600), which included 35 heart samples harvested from a Wistar rat on postoperative days 1, 6 and 42 following aorta ligation and sham-operated Wistar rats, respectively. Hellman *et al.* (16) demonstrated a correlation between hyaluronan concentration and specific gene expression levels using SPSS software. Analysis of the correlation matrix was performed according to the Principal components method (17), and orthogonal partial least squares-discrimination analysis was used to analyze the datasets of GSE21600, in which the previous clustering, including extracellular matrix and adhesion molecules were confirmed, and fatty acid metabolism, glucose metabolism, mitochondria and atherosclerosis were detected as the new clustering (18). However, these previous two studies failed to predict the changing trends of genes in this disease. Hence, the present study aimed to reanalyze the expression profiles of GSE21600 in order to construct a predictive model of cardiac hypertrophy using linear discriminant analysis (LDA) method. GSE21600 microarray data was used to identify differentially expressed genes (DEGs) using a Limma package in R (version. 3.26.5), which calculates linear models of microarray data. Common DEGs were used to construct a mathematical model in order to predict the expression levels of genes in the cardiac hypertrophy samples. The mathematical model was verified receiver operating characteristic (ROC) curve and the consistency of predictive and measurement data. The present study may be useful for the early prediction of changing trends in cardiac hypertrophy disease at the gene level.

Materials and methods

Data preprocessing and DEGs screening. GSE21600 microarray data were downloaded from the GEO database (<http://www.ncbi.nlm.nih.gov/geo/>) (16). GSE21600 included data from 35 heart samples harvested from 36 Wistar rats which were excised on postoperative days 1, 6 and 42 following aorta ligation and sham-operated groups, respectively. Each group contained six samples at each time point, with the exception of the samples harvested from the aorta ligated group at 6 days, where $n=5$. The microarray platform of GSE21600 was Illumina GPL6101 RatRef-12 expression bead chip (version 1.0; Illumina, Inc., San Diego, CA, USA).

Samples were divided into three groups: Day 1 (D1), day 6 (D6) and day 42 (D42). DEGs between the postoperative and sham-operated samples were identified in these three groups, respectively. Firstly, normalization of the microarray data was performed in the R language (19,20), and DEGs were subsequently identified using a Limma package in R (21). False discovery rate (FDR) was used to adjust the P-value, according to the method outlined by Benjamin and Hochberg (22). $FDR < 0.05$ and $> 1 \log_2$ fold change (FC) were chosen as the cut-off criteria.

Specific gene screening. In order to screen the specific expression levels of genes at each time point, DEGs were compared between the two groups. Subsequently, hierarchical clustering analysis (23) was performed on the common DEGs in the three groups.

Table I. Expression levels of five common differentially expressed genes the in aorta ligated operation group were calculated, as compared with the sham operated group.

Gene	Day 1	Day 6	Day 42
<i>AKIP1</i>	-1.24914	-1.36699	-1.80092
<i>ANKRD23</i>	-2.90253	-3.69624	-2.85077
<i>LTBP2</i>	-3.68846	-4.20566	-2.02513
<i>TGFB2</i>	-2.15313	-2.11814	-1.75841
<i>TNFRSF12A</i>	-1.99987	-2.08827	-1.54923

Sorting algorithm and construction of the mathematical model. Linear discriminant analysis (LDA) is a method that is commonly widely used in microarray classification to obtain discrimination function. LDA analysis can be performed when there are ≥ 2 groups and each group contains > 2 variables (24,25). In this method, a linear equation based on the variations in the two groups is established: $Y = a + b_{11} + b_{22} + \dots + b_n X_n$, where 'a' represents a constant and 'b₁, b₂ ... and b_n' represents the regression coefficient. In the present study, the cardiac hypertrophy samples were defined as '1' and the control samples were defined as '-1'. Based on the dynamic expression changes of the common DEGs detected in the D1 group, the expression pattern in the D42 group was predicted via the calculated mathematical model constructed using the LDA method (26).

Verification of the mathematical model. Disease classification models are typically determined using multivariate regression analysis (27,28), ROC curve (29-32) or prospective validation (33). ROC curve was used in the present study in order to evaluate the discriminant effect of the mathematical model and directly observe the accuracy of the present analysis method. Indices, including specificity and sensitivity, were calculated in order to estimate the predictive ability of LDA, in addition to area under the curve (AUC) of the ROC curve, which was also calculated to estimate accuracy. In the present study, AUC was used to distinguish non-accuracy ($AUC \leq 0.5$), low accuracy ($0.5 < AUC \leq 0.7$), moderate accuracy ($0.7 < AUC \leq 0.9$) and high accuracy ($0.9 < AUC < 1$). Furthermore, by comparing the prediction data with the measurement data in the D42 samples, the consistency of two sets of data was evaluated.

Results

Identification, comparison and feature selection of DEGs. Normalization of the microarray data is presented in Fig. 1. DEGs were identified, and the genes with $FDR < 0.05$ and $> 1 \log_2$ FC were considered as differentially expressed between the ligated samples and sham-operated samples. A total of 319, 44 and 57 DEGs were identified in the D1, D6 and D42 groups respectively.

A total of 23 DEGs were detected between the D1 and D6 groups, 14 DEGs were detected between the D1 and D42 groups, and five DEGs were identified between the D6 and D42 groups. Five common DEGs, including A kinase interacting protein 1 (*AKIP1*), ankyrin repeat domain 23

Table II. Predicted data at day 42 using a linear equation of the gene expression levels of cardiac hypertrophy.

Gene accession	State	Expression on day 1	Expression on day 42	Predicted on day 42
GSM539275	1	332.1987	337.3279	326.1898781
GSM539276	1	272.2375	126.1764	235.327208
GSM539277	1	485.7471	792.9784	558.8706386
GSM539278	1	778.9512	344.6311	1,003.179749
GSM539279	1	320.8331	108.7458	308.9669279
GSM539280	1	716.3563	479.7876	908.3260809
GSM539281	-1	85.13754	66.26252	-48.1961695
GSM539282	-1	71.55708	13.26508	-68.775425
GSM539283	-1	50.69723	41.25237	-100.385561
GSM539284	-1	23.54682	75.99313	-141.528145
GSM539285	-1	124.7012	29.73599	11.75692997
GSM539286	-1	49.61586	52.55618	-102.024223
GSM539275	1	4,201.869	6,096.354	6,190.124821
GSM539276	1	1,882.365	5,415.158	2,675.24642
GSM539277	1	3,337.275	9,621.91	4,879.955589
GSM539278	1	3,016.572	4,261.265	4,393.975807
GSM539279	1	2,658.368	3,865.638	3,851.168593
GSM539280	1	1,956.894	8,021.108	2,788.184519
GSM539281	-1	1,219.844	959.4762	1,671.290077
GSM539282	-1	1,070.036	1,546.261	1,444.277361
GSM539283	-1	1,431.854	1,145.456	1,992.561078
GSM539284	-1	1,024.116	3,023.837	1,374.692133
GSM539285	-1	988.543	1,751.745	1,320.786311
GSM539286	-1	1,213.691	2,605.091	1,661.966081
GSM539275	1	880.5447	147.3087	1,157.130248
GSM539276	1	126.5936	169.5375	14.62459301
GSM539277	1	1,011.612	281.1071	1,355.744099
GSM539278	1	1,073.774	185.5347	1,449.941769
GSM539279	1	340.023	62.10585	338.0464919
GSM539280	1	122.0065	237.4351	7.673495398
GSM539281	-1	36.33411	32.24878	-122.150826
GSM539282	-1	50.67635	24.24548	-100.417201
GSM539283	-1	36.68185	45.16885	-121.623876
GSM539284	-1	30.85578	71.55927	-130.452456
GSM539285	-1	15.40947	32.20256	-153.859142
GSM539286	-1	34.06184	51.90232	-125.594128
GSM539275	1	1,915.488	1,621.039	2,725.439616
GSM539276	1	719.9728	1,732.95	913.8063723
GSM539277	1	1,491.145	1,375.875	2,082.408155
GSM539278	1	2,425.283	3,341.205	3,497.961428
GSM539279	1	1,208.035	885.0079	1,653.395218
GSM539280	1	1,999.254	1,564.762	2,852.375074
GSM539281	-1	391.4185	495.3794	415.929062
GSM539282	-1	355.6202	400.3427	361.6818301
GSM539283	-1	437.2545	578.2272	485.3870006
GSM539284	-1	215.4102	719.1659	149.2135176
GSM539285	-1	464.529	402.5466	526.717626
GSM539286	-1	483.1193	857.3302	554.8885815
GSM539275	1	1,776.678	768.9708	2,515.092804
GSM539276	1	998.7648	732.2133	1,336.275995
GSM539277	1	2,373.809	1,362.486	3,419.959903
GSM539278	1	3,322.548	1,513.086	4,857.638915

Table II. Continued.

Gene accession	State	Expression on day 1	Expression on day 42	Predicted on day 42
GSM539279	1	879.2261	513.2602	1,155.132097
GSM539280	1	1,201.621	1,250.521	1,643.675713
GSM539281	-1	411.144	251.4373	445.8202516
GSM539282	-1	375.7809	208.7139	392.2325034
GSM539283	-1	406.536	168.1061	438.837483
GSM539284	-1	297.8341	399.4494	274.1152146
GSM539285	-1	322.7278	352.2844	311.8380763
GSM539286	-1	316.283	400.1669	302.0718985

1, the aorta ligated operation group; -1, the sham operated group.




Figure 1. Microarray data normalization. Samples were divided into three groups: days 1, 6 and 42. White, aorta ligated operation samples. Blue, sham operated samples.




Figure 2. Identification of specific differentially expressed genes. Yellow, day 1 (D1) group; green, day 6 (D6) group; purple, day 42 (D42) group.

(*ANKRD23*), latent transforming growth factor beta binding protein (*LTBP2*), transforming growth factor (*TGF*)- β 2 and tumor necrosis factor receptor superfamily member 12a

(*TNFRSF12A*), were identified among the three groups (Fig. 2).

Clustering analysis of the five common DEGs demonstrated that the sham operated and ligated samples were respectively clustered together; however, three ligated samples (16.67%; 3/18) were mixed into the operated group and two sham-operated samples (11.76%; 2/17) were mixed into the ligated group (Fig. 3). These five common DEGs were identified as downregulated genes (Table I).

Construction and verification of the mathematical model. Based on the expression levels and dynamic changes detected in the five common DEGs, a linear equation between the D1 and D42 groups was calculated as follows: $y=1.526x-186.671$; where 'y' and 'x' represent the expression levels in the D42 and D1 groups, respectively.

Assessment of the ROC curve demonstrated that AUC was 0.831, which indicated that the predictive accuracy was 83.1% and the specificity and sensitivity were 0.8, respectively (Fig. 4A). By comparing the predictive and measurement data




Figure 3. Hierarchical clustering analysis of five common differentially expressed genes in day 1 (D1), 6 (D6) and 42 (D42). Red labels represent the samples which were mixed into the false group.




Figure 4. Verification of the prediction model. (A) The model was verified by receiver operating characteristic (ROC) curve (B), which was determined by the consistency of predictive and measurement data at day 42. The area under curve (AUC) of ROC was used to assess the accuracy of data. AUC value $0.7 < AUC \leq 0.9$ implies moderate accuracy, whereas AUC value $0.9 < AUC < 1$ implies high accuracy. Cor, correlation of predicted data and measurement data.

at 42 days (Table II), the consistency of these two datasets was calculated to be 76.5% (Fig. 4B).

Discussion

In the present study, the expression profiles of sham operated and ligated heart samples harvested from a Wistar rat were analyzed and 319, 44 and 57 DEGs were subsequently identified in the D1, D6 and D42 groups, respectively. *AKIP1*, *ANKRD23*, *LTBP2*, *TGF- β 2* and *TNFRSF12A* were identified as common DEGs among the three groups, and their association with cardiac hypertrophy has previously been demonstrated (34-37). *AKIP1* was identified as a key regulator of heart function via the cAMP-dependent protein kinase signaling pathway (38). During periods of the oxidant stress, the expression of *AKIP1* is capable of protecting cardiac myocytes from the ischemic injury via enhanced mitochondrial integrity (38). Furthermore, the expression of *AKIP1* may also protect the heart via mitochondrial stress adaptation (39), and it has been demonstrated that mitochondrial DNA damage may contribute to the development of cardiac hypertrophy and heart failure (40). These results suggested that *AKIP1* may serve a crucial function in the development of cardiac hypertrophy via mitochondrial stress adaptation mechanisms. Hellman *et al* (16) have previously demonstrated that *LTBP2* and *TGF- β 2* are associated with the development of cardiac hypertrophy. *LTBP2*, which belongs to the fibrillin superfamily, regulates the release of *TGF- β 1* (41,42). Previous studies have demonstrated that *TGF- β* , including *TGF- β 1*, *TGF- β 2* and *TGF- β 3*, have an important role in the pathogenesis of cardiac hypertrophy by stimulating the proliferation of cardiomyocytes (43,44). These

results demonstrated that *LTBP2* and *TGF- β 2* are associated with the regulation of cardiac hypertrophy. However, the role of *ANKRD23* and *TNFRSF12A* in the development of cardiac hypertrophy is yet to be elucidated. As the results of the present study demonstrated that they were detected as common genes in the three groups, we hypothesize that *AKIPI*, *ANKRD23*, *LTBP2*, *TGF- β 2* and *TNFRSF12A* may contribute to the development of cardiac hypertrophy.

Numerous mathematical techniques have been developed in order to analyze large datasets, and mathematical modeling is a useful and powerful tool for the analysis of gene expression patterns (14). LDA is a well-known multivariate technique that is used for dimension reduction and classification (45). A 3-gene model, *TNFRSF8*, *BATF3* and *TMOD1*, which was obtained by LDA and leave-one-out cross-validation, was previously used to separate ALK (-) and anaplastic large-cell lymphoma from peripheral T-cell lymphoma, and the accuracy of the model was ~97% (46). Furthermore, a class-prediction model of patients with Graft-vs-host disease was previously constructed using LDA, and the accuracy was 63-80%, as estimated by reverse transcription-quantitative polymerase chain reaction (47). ROC, which directly displays the correlation of specificity and sensitivity can be used to assess the accuracy of diagnostic tests (48). In a previous study conducted by Barretina *et al* (49), Cancer Cell Line Encyclopedia, which is a predictive model, was cross-validated by specificity and sensitivity of the ROC curve and used to predict the drug response to gene expression, including topoisomerase inhibitors associated with Schlafen family member 11. Similarly, a predictions model has previously been constructed for dementia using LDA and verified by ROC curve, and the accuracy of the model was 66%; whereas the specificity and sensitivity were 73% and 64%, respectively (50). In the present study, a prediction model of cardiac hypertrophy was constructed. The assessment of ROC curve demonstrated that the predictive accuracy of the model was ~83.1% and the specificity and sensitivity were 0.8, respectively. By comparing the predictive and measurement data at 42 days, the consistency of these two datasets was calculated to be 76.5%. These results suggested that the present prediction model provides improved predictive ability, which may contribute to the early prediction of the changing trends in gene expression exhibited in patients with cardiac hypertrophy disease. However, to elevate the discrimination ability of the model, further studies with an increased number of samples and more suitable machine learning algorithm are required.

In the present study, 319, 44 and 57 DEGs were detected in D1, D6 and D42 groups, respectively. *AKIPI*, *ANKRD23*, *LTBP2*, *TGF- β 2* and *TNFRSF12A* were identified as common DEGs. A linear equation was calculated between the D1 and D42 groups, as follows: $y=1.526x-186.671$. This linear equation, which acted as a prediction model of gene expression levels, may contribute to the early prediction of the changing trends in cardiac hypertrophy disease.

Acknowledgements

The authors of the present study would like to thank Fenghe Information Technology Co., Ltd (Shanghai, China) for in-depth editing and language assistance.

References

1. Heineke J and Molkentin JD: Regulation of cardiac hypertrophy by intracellular signalling pathways. *Nat Rev Mol Cell Biol* 7: 589-600, 2006.
2. Hubert HB, Feinleib M, McNamara PM and Castelli WP: Obesity as an independent risk factor for cardiovascular disease: A 26-year follow-up of participants in the Framingham Heart Study. *Circulation* 67: 968-977, 1983.
3. Wang AY, Wang M, Woo J, Lam CW, Lui SF, Li PK and Sanderson JE: Inflammation, residual kidney function and cardiac hypertrophy are interrelated and combine adversely to enhance mortality and cardiovascular death risk of peritoneal dialysis patients. *J Am Soc Nephrol* 15: 2186-2194, 2004.
4. Dunn FG, Burns JM and Hornung RS: Left ventricular hypertrophy in hypertension. *Am Heart J* 122: 312-315, 1991.
5. Osio A, Tan L, Chen SN, Lombardi R, Nagueh SF, Shete S, Roberts R, Willerson JT and Marian AJ: Myozenin 2 is a novel gene for human hypertrophic cardiomyopathy. *Circ Res* 100: 766-768, 2007.
6. Kontaraki JE, Parthenakis FI, Patrianakos AP, Karalis IK and Vardas PE: Altered expression of early cardiac marker genes in circulating cells of patients with hypertrophic cardiomyopathy. *Cardiovasc Pathol* 16: 329-335, 2007.
7. Black FM, Packer SE, Parker TG, Michael LH, Roberts R, Schwartz RJ and Schneider MD: The vascular smooth muscle alpha-actin gene is reactivated during cardiac hypertrophy provoked by load. *J Clin Invest* 88: 1581-1588, 1991.
8. Qing YF, Zhou JG, Zhang QB, Wang DS, Li M, Yang QB, Huang CP, Yin L, Pan SY, Xie WG, *et al.*: Association of TLR4 Gene rs2149356 polymorphism with primary gouty arthritis in a case-control study. *PLoS One* 8: e64845, 2013.
9. Planavila A, Redondo I, Hondares E, Vinciguerra M, Munts C, Iglesias R, Gabrielli LA, Sitges M, Giral M, van Bilsen M and Villarroya F: Fibroblast growth factor 21 protects against cardiac hypertrophy in mice. *Nat Commun* 4: 2019, 2013.
10. Grimwade D, Walker H, Oliver F, Wheatley K, Harrison C, Harrison G, Rees J, Hann I, Stevens R, Burnett A and Goldstone A: The importance of diagnostic cytogenetics on outcome in AML: Analysis of 1,612 patients entered into the MRC AML 10 trial. The medical research council adult and children's leukaemia working parties. *Blood* 92: 2322-2333, 1998.
11. Covert MW, Schilling CH and Palsson B: Regulation of gene expression in flux balance models of metabolism. *J Theor Biol* 213: 73-88, 2001.
12. Wade KC, Guttentag SH, Gonzales LW, Maschhoff KL, Gonzales J, Kolla V, Singhal S and Ballard PL: Gene induction during differentiation of human pulmonary type II cells in vitro. *Am J Respir Cell Mol Biol* 34: 727-737, 2006.
13. Saban MR, Hellmich H, Nguyen NB, Winston J, Hammond TG and Saban R: Time course of LPS-induced gene expression in a mouse model of genitourinary inflammation. *Physiol Genomics* 5: 147-160, 2001.
14. Tamayo P, Slonim D, Mesirov J, Zhu Q, Kitareewan S, Dmitrovsky E, Lander ES and Golub TR: Interpreting patterns of gene expression with self-organizing maps: Methods and application to hematopoietic differentiation. *Proc Natl Acad Sci USA* 96: 2907-2912, 1999.
15. Kondo S and Miura T: Reaction-diffusion model as a framework for understanding biological pattern formation. *Science* 329: 1616-1620, 2010.
16. Hellman U, Mörner S, Engström-Laurent A, Samuel JL and Waldenström A: Temporal correlation between transcriptional changes and increased synthesis of hyaluronan in experimental cardiac hypertrophy. *Genomics* 96: 73-81, 2010.
17. Revell LJ: Size-correction and principal components for interspecific comparative studies. *Evolution* 12: 3258-3268, 2009.
18. Gennebäck N, Malm L, Hellman U, Waldenström A and Mörner S: Using OPLS-DA to find new hypotheses in vast amounts of gene expression data-Studying the progression of cardiac hypertrophy in the heart of aorta ligated rat. *Gene* 522: 27-36, 2013.
19. Troyanskaya O, Cantor M, Sherlock G, Brown P, Hastie T, Tibshirani R, Botstein D and Altman RB: Missing value estimation methods for DNA microarrays. *Bioinformatics* 17: 520-525, 2001.
20. Fujita A, Sato JR, Rodrigues Lde O, Ferreira CE and Sogayar MC: Evaluating different methods of microarray data normalization. *BMC Bioinformatics* 7: 469, 2006.

21. Smyth GK: Limma: Linear models for microarray data. In: Bioinformatics and computational biology solutions using R and Bioconductor. Gentleman R, Carey V, Huber W, Irizarry R and Duhot S (eds). Springer-Verlag London Ltd., London, pp397-420, 2005.
22. Benjamini Y and Hochberg Y: Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J R Statist Soc B* 57: 289-300, 1995.
23. Eisen MB, Spellman PT, Brown PO and Botstein D: Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci USA* 95: 14863-14868, 1998.
24. Liang J and Du R: Model-based fault detection and diagnosis of HVAC systems using support vector machine method. *Int J Refrig* 30: 1104-1114, 2007.
25. Polat K and Güneş S: Breast cancer diagnosis using least square support vector machine. *Digit Signal Process* 17: 694-701, 2007.
26. Chen LF, Liao HYM, Ko MT, Lin JC and Yu GJ: A new LDA-based face recognition system which can solve the small sample size problem. *Pattern Recognition* 33: 1713-1726, 2000.
27. Jiang H, Deng Y, Chen HS, Tao L, Sha Q, Chen J, Tsai CJ and Zhang S: Joint analysis of two microarray gene-expression data sets to select lung adenocarcinoma marker genes. *BMC Bioinformatics* 5: 81, 2004.
28. Imai K. Multivariate regression analysis for the item count technique. *J Amer Statist Assoc* 106: 407-416, 2011.
29. Zhou L, Cheng L, Tao L, Jia X, Lu Y and Liao P: Detection of hypopharyngeal squamous cell carcinoma using serum proteomics. *Acta Otolaryngol* 126: 853-860, 2006.
30. Navaglia F, Fogar P, Basso D, Greco E, Padoan A, Tonidandel L, Fadi E, Zambon CF, Bozzato D, Moz S, *et al*: Pancreatic cancer biomarkers discovery by surface-enhanced laser desorption and ionization time-of-flight mass spectrometry. *Clin Chem Lab Med* 47: 713-723, 2009.
31. Hewett R and Kijisanayothin P: Tumor classification ranking from microarray data. *BMC Genomics* 9 (Suppl 2): S21, 2008.
32. Roepman P, Schuurman A, Delahaye LJ, Witteveen AT, Floore AN and Glas AM: A gene expression profile for detection of sufficient tumour cells in breast tumour tissue: Microarray diagnosis eligibility. *BMC Med Genomics* 2: 52, 2009.
33. Ye QH, Qin LX, Forgues M, He P, Kim JW, Peng AC, Simon R, Li Y, Robles AI, Chen Y, *et al*: Predicting hepatitis B virus-positive metastatic hepatocellular carcinomas using gene expression profiling and supervised machine learning. *Nat Med* 9: 416-423, 2003.
34. Yu H, Tigchelaar W, Lu B, van Gilst WH, de Boer RA, Westenbrink BD and Silljé HH: AKIP1, a cardiac hypertrophy induced protein that stimulates cardiomyocyte growth via the Akt pathway. *Int J Mol Sci* 14: 21378-21393, 2013.
35. Bang ML, Gu Y, Dalton ND, Peterson KL, Chien KR and Chen J: The muscle ankyrin repeat proteins CARP, Ankrd2, and DARP are not essential for normal cardiac development and function at basal conditions and in response to pressure overload. *PloS One* 9: e93638, 2014.
36. Kuba K, Zhang L, Imai Y, Arab S, Chen M, Maekawa Y, Leschnik M, Leibbrandt A, Markovic M, Schwaighofer J, *et al*: Impaired heart contractility in Apelin gene-deficient mice associated with aging and pressure overload. *Circ Res* 101: e32-e42, 2007.
37. Dobaczewski M, Chen W and Frangogiannis NG: Transforming growth factor (TGF)- β signaling in cardiac remodeling. *J Mol Cell Cardiol* 51: 600-606, 2011.
38. Sastri M, Haushalter KJ, Panneerselvam M, Chang P, Fridolfsson H, Finley JC, Ng D, Schilling JM, Miyanojara A, Day ME, *et al*: A kinase interacting protein (AKIP1) is a key regulator of cardiac stress. *Proc Natl Acad Sci USA* 110: E387-E396, 2013.
39. Yu H, Tigchelaar W, Koonen DP, Patel HH, de Boer RA, van Gilst WH, Westenbrink BD and Silljé HH: AKIP1 expression modulates mitochondrial function in rat neonatal cardiomyocytes. *PLoS One* 8: e80815, 2013.
40. Dai DF, Johnson SC, Villarin JJ, Chin MT, Nieves-Cintrón M, Chen T, Marcinek DJ, Dorn GW II, Kang YJ, Prolla TA, *et al*: Mitochondrial oxidative stress mediates angiotensin II-induced cardiac hypertrophy and Galphaq overexpression-induced heart failure. *Circ Res* 108: 837-846, 2011.
41. Sinha S, Heagerty AM, Shuttleworth CA and Kielty CM: Expression of latent TGF-beta binding proteins and association with TGF-beta1 and fibrillin-1 following arterial injury. *Cardiovasc Res* 53: 971-983, 2002.
42. Sterner-Kock A, Thorey IS, Koli K, Wempe F, Otte J, Bangsow T, Kuhlmeier K, Kirchner T, Jin S, Keski-Oja J and von Melchner H: Disruption of the gene encoding the latent transforming growth factor-beta binding protein 4 (LTBP-4) causes abnormal lung development, cardiomyopathy, and colorectal cancer. *Genes Dev* 16: 2264-2273, 2002.
43. Bujak M and Frangogiannis NG: The role of TGF-beta signaling in myocardial infarction and cardiac remodeling. *Cardiovasc Res* 74: 184-195, 2007.
44. Dobaczewski M, Chen W and Frangogiannis NG: Transforming growth factor (TGF)- β signaling in cardiac remodeling. *J Mol Cell Cardiol* 51: 600-606, 2011.
45. Roth V and Steinhage V: Nonlinear discriminant analysis using kernel functions. In: *Advances in Neural Information Processing Systems*. Vol 12. Solla SA, Leen TK and Müller KR (eds). MIT Press, Cambridge, MA, pp568-574, 2000.
46. Agnelli L, Mereu E, Pellegrino E, Limongi T, Kwee I, Bergaggio E, Ponzoni M, Zamò A, Iqbal J, Piccaluga PP, *et al*; European T-Cell Lymphoma Study Group: Identification of a 3-gene model as a powerful diagnostic tool for the recognition of ALK-negative anaplastic large-cell lymphoma. *Blood* 120: 1274-1281, 2012.
47. Baron C, Somogyi R, Greller LD, Rineau V, Wilkinson P, Cho CR, Cameron MJ, Kelvin DJ, Chagnon P, Roy DC, *et al*: Prediction of graft-versus-host disease in humans by donor gene-expression profiling. *PLoS Med* 4: e23, 2007.
48. Beck JR and Shultz EK: The use of relative operating characteristic (ROC) curves in test performance evaluation. *Arch Pathol Lab Med* 110: 13-20, 1986.
49. Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, Wilson CJ, Lehár J, Kryukov GV, Sonkin D, *et al*: The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. *Nature* 483: 603-607, 2012.
50. Maroco J, Silva D, Rodrigues A, Guerreiro M, Santana I and de Mendonça A: Data mining methods in the prediction of Dementia: A real-data comparison of the accuracy, sensitivity and specificity of linear discriminant analysis, logistic regression, neural networks, support vector machines, classification trees and random forests. *BMC Res Notes* 4: 299, 2011.