

Zigzag Nanographene

How to cite: *Angew. Chem. Int. Ed.* **2021**, *60*, 13521–13528

International Edition: doi.org/10.1002/anie.202016437

German Edition: doi.org/10.1002/ange.202016437

Rules of Nucleophilic Additions to Zigzag Nanographene Diones**

Peter Ribar⁺, Leoš Valenta⁺, Tomáš Šolomek,^{*} and Michal Juríček^{*}

Abstract: Nucleophilic addition of carbon-centered nucleophiles to nanographene ketones represents a valuable late-stage method for the functionalization of zigzag nanographenes, but its use is rare in the chemical literature. Using two model systems, non-Kekulé triangulene-4,8-dione and Kekulé anthanthrone, we identify unexpected regioselectivities and uncover the rules that govern these reactions. Considering the large number of nanographene ketones that have been reported since the pioneering work of Eric Clar, this method enables synthesis and exploration of hitherto unknown functionalized nanographenes.

Introduction

The finite fragments of graphene, or nanographenes, are ideal molecular models for investigating the relationship between the structure and properties of semiconducting^[1] and magnetic^[2] materials based on graphene nanoribbons. The bottom-up synthetic access to these materials^[3] is key for the advancement of the molecular electronics.^[4] Routine issues that need to be addressed when synthesizing nanographenes

are stability^[5] and solubility.^[6] A classic example are acenes,^[7] the thinnest possible nanoribbons, elongation of which increases charge mobility but decreases chemical stability.^[8] Lateral extension (see anthracene and anthanthrene in Figure 1) overcomes this problem but limits the solubility, and thus processability, of such nanographenes.^[9] Conveniently, both issues can be solved by installment of peripheral substituents that prevent molecules from reacting and stacking,^[10] in addition to tuning their molecular^[11] and bulk properties.^[12] The drawback is that introduction of substituents embodies an additional hurdle on the synthetic pathway and may defy synthetic chemists to reach the target.




Figure 1. Regioselectivity of additions of carbon-centered nucleophiles to ketone precursors of Kekulé (anthracene, anthanthrene) and non-Kekulé (phenalenyl,^[13] triangulene) nanographenes.

Thanks to the recent advances in C–H activation, the cross-coupling reactions enable introduction of substituents to different positions.^[14] This method, however, cannot always be applied to all positions in zigzag^[15] nanographenes. For example, it would be difficult, if possible at all, to modify positions at the center of the edges in Clar's hydrocarbon triangulene (Figure 1). A common approach to install substituents in zigzag nanographenes is a nucleophilic addition to an aldehyde prior to the nanographene core formation by the Friedel–Crafts alkylation,^[16] which decreases modularity. A better but far less common method is to add nucleophiles to the products of the Friedel–Crafts acylation, the simplest method to construct zigzag nanographenes.^[17]

This strategy has been successful for making a variety of acenes up to pentacene by 1,2-additions of carbon-centered nucleophiles to the corresponding quinones (Figure 1),^[12,18] while only acetylide 1,2-additions have been reported for hexacene and higher acene quinones.^[10b,19] On lateral extension, acetylide 1,2-additions are common,^[20] whereas only one

[*] Dr. P. Ribar,^[†] Prof. T. Šolomek
Department of Chemistry
University of Basel
St. Johannis-Ring 19, 4056 Basel (Switzerland)
E-mail: tomas.solomek@unibas.ch

L. Valenta,^[†] Prof. M. Juríček
Department of Chemistry
University of Zurich
Winterthurerstrasse 190, 8057 Zurich (Switzerland)
E-mail: michal.juricek@chem.uzh.ch

Prof. T. Šolomek, Prof. M. Juríček
Prievdza Chemical Society
M. Hodžu 10/16, 971 01 Prievdza (Slovak Republic)

Prof. T. Šolomek
Current address: Department of Chemistry, Biochemistry and
Pharmaceutical Sciences
University of Bern
Freiestrasse 3, 3012 Bern (Switzerland)
E-mail: tomas.solomek@dcb.unibe.ch

[†] These authors contributed equally to this work.

[**] A previous version of this manuscript has been deposited on a preprint server (<https://doi.org/10.26434/chemrxiv.13348049.v1>).

Supporting information and the ORCID identification numbers for the authors of this article can be found under: <https://doi.org/10.1002/anie.202016437>.

© 2021 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.


example of heteroaryl 1,2-addition has been described on anthanthrone.^[20a] Other aryl substituents have been introduced by multistep sequences.^[14,21] Despite the examples, where 1,2-additions work, it is striking that this strategy is not generally applied to extended systems, for which the ketone precursors are known. For instance, triangulene (Figure 1) is an example of an open-shell zigzag nanographene that has never been isolated in the solid state. This non-Kekulé triplet diradical was only made twice in its neutral form: in 2001 as a short-lived tri-*tert*-butyl derivative^[5a] and in 2017 as a naked system on surface under ultra-high vacuum.^[22] It is a paradox that for almost 70 years, the ideal precursor for such task, triagulene-4,8-dione, has been known since the first synthetic attempts by Eric Clar in 1950s.^[23] All it would take to make persistent triangulene is to perform a nucleophilic 1,2-addition at the carbonyl centers of the triagulenedione core and reduce the ensuing diol. Where is the catch?

We selected two isomeric zigzag model systems, one based on triangulene and the other one on anthanthrene (Figure 1), to uncover the rules that govern nucleophilic additions on zigzag nanographene ketones and to understand why the use of this method is uncommon in the chemical literature.

Results and Discussion

Each model system is comprised of 22 carbon atoms and six hexagonal rings. While the Kekulé conjugation topology of anthanthrene is the same as that of acenes, the isomeric triangulene has a non-Kekulé topology. We improved the solubility of model diones by a minimum number of substituents distant from the reactive centers (see the SI) to get a better control over the reactions. We used two readily available reagents as carbon-centered nucleophiles, phenyl-ethylolithium (PhCCLi) and 3,5-di-*tert*-butyl-phenylmagnesium bromide (ArMgBr), which serve as models for common substituents that control solubility, aggregation, electronic structure, and solid-state morphology of nanographenes.^[24] It was important to determine the exact concentration of the reagent before each experiment to get reproducible results. In the case of double 1,2-additions and mixed 1,2-/1,4-additions, a reduction of the diol intermediate and water elimination, respectively, by SnCl₂/HCl were performed to obtain the corresponding nanographene, which simplified purification and product identification. Likewise, conjugate additions were followed by an oxidation of the enol intermediate by O₂ or I₂. The structure of all products was confirmed by 2D NMR spectroscopy to unambiguously identify the position where the nucleophilic addition took place.

Triagulenedione. We attempted a double 1,2-addition to triagulenedione by reacting its soluble derivative **T1** (Ar = 3,5-di-*tert*-butylphenyl, Scheme 1) with an excess of ArMgBr. If successful, the product would yield an ideal precursor of a persistent triangulene. Instead, we obtained a complex mixture of products, which we were unable to identify. One equivalent of ArMgBr, however, gave a clean 1,4-addition providing **T2** after oxidation in 75% yield (Scheme 1 and Table 1). A more reactive ArLi nucleophile expected to favor a 1,2-addition preserved the 1,4-selectivity, albeit with a lower



Scheme 1. Nucleophilic additions to triagulenedione. Ar = Nu = 3,5-di-*tert*-butylphenyl, M = MgBr or Li.


Table 1: Nucleophilic additions to triagulenedione and anthanthrone.^[a]

dione	Nu	Nu [equiv]	product 1 (#, yield)	product 2 (#, yield)
T1	ArMgBr	1	1,4 (T2 , 75%)	— ^[b]
T1	ArLi	1	1,4 (T2 , 44%)	— ^[b]
T1	ArLi + DMPU	1 + 2	n.r. ^[c]	n.r. ^[c]
A1	PhCCLi	2	1,2 & 1,2 (A2 , 72%)	— ^[b]
A1	PhCCLi	1	1,2 (A3 , 86%)	— ^[b]
A1	ArMgBr	10	1,2 & 1,4 (A4 , 83%)	— ^[b]
A1	ArMgBr	1	1,4 (A5 , 50%)	1,2 (A6 , 34%)
A1	ArLi	13	1,2 & 1,4 (A4 , 75%)	1,4 & 1,4 (A7 , 22%)
A1	ArLi + DMPU	10 + 20 ^[d]	n.r. ^[c]	n.r. ^[c]
A1	MesMgBr	10	1,4 & 1,4 (A11 , 91%)	— ^[b]
A8	PhCCLi	10	1,2 & 1,2 (A9 , 86%)	— ^[b]
A8	ArMgBr	10	1,4 & 1,4 (A10 , 69%)	— ^[b]
A8	ArLi	2	1,4 & 1,4 (A10 , 40%)	— ^[b]
A8	ArLi + DMPU	2 + 4	n.r. ^[c]	n.r. ^[c]

[a] Refer to Schemes 1 and 2. **A1**, **T1**: R = Ar = 3,5-di-*tert*-butylphenyl, **A8**: R = (4-octylphenyl)ethynyl, Mes = mesityl, DMPU = *N,N'*-dimethylpropyleneurea. [b] Not observed. [c] No reaction. [d] Quenching of the reagent with DMF led to > 90% ArCHO formation.

yield (44%). To our surprise, we observed no conversion in the reaction of dione **T1** with one equivalent of PhCCLi, an excess led again to a complex reaction mixture. This reactivity is in stark contrast to that observed for acene quinones, which give exclusively 1,2-additions. The most closely related system to dione **T1** that is known to undergo 1,4-additions is phenalene^[25] (Figure 1), even though it has only one carbonyl group. Nevertheless, its lowest unoccupied molecular orbital (LUMO, Figure S1) is analogous to that of triagulenedione, which suggests that conjugation topology^[13] governs the selectivity.

Anthanthrone. This dione is a lateral extension of anthraquinone. It displays an analogous LUMO (Figure S1) and one would expect it to undergo solely 1,2-additions, like anthraquinone. Indeed, the reaction of dione **A1** (R = 3,5-di-*tert*-butylphenyl) with two equivalents of PhCCLi gives the



Scheme 2. Nucleophilic additions to anthanthrone. Ar = 3,5-di-*tert*-butylphenyl, PhCC = phenylethynyl, oPhCC = (4-octylphenyl)ethynyl, Mes = mesityl. [a]–[d] Isolated after subsequent [a] protonation, [b] oxidation (O₂ or I₂), [c] reduction (SnCl₂/HCl), and [d] dehydration (SnCl₂/HCl). [e] Without protonation after the first 1,2-addition. [f] Without oxidation after the first 1,4-addition; nucleophile adds to the enol intermediate of 1,4-addition (Figure 4).

product of a double 1,2-addition (**A2**; 72% yield; Scheme 2 and Table 1); the use of one equivalent results in 1,2-product **A3** (86%). The reaction with an excess of ArMgBr, however, leads to a product of consecutive 1,2- and 1,4-additions (**A4**; 83%). To determine which addition is faster, we performed this reaction with one equivalent of ArMgBr. Unexpectedly, we isolated the product of 1,4-addition (**A5**; 50%) in a yield higher than that of 1,2-addition (**A6**; 34%). To the best of our knowledge, this has never been observed in linear acene quinones. This result suggests that each initial adduct undergoes the subsequent addition with a complementary selectivity to form one product **A4**. We tested if a more reactive nucleophile such as ArLi would suppress 1,4-addition, but we observed an opposite effect. Reacting an excess of ArLi with **A1**, we isolated the product of a double 1,4-addition (**A7**; 22%) besides **A4** (75%), the sole product of ArMgBr addition. We thought that the bulky solubilizing R groups in **A1** could steer the selectivity towards 1,4-addition due to steric hindrance. For this reason, we synthesized anthanthrone **A8** equipped with sterically non-demanding (4-octylphenyl)ethynyl substituents. The reaction of **A8** with PhCCLi proceeded as before in a 1,2-fashion (**A9**; 86%; Scheme 2 and Table 1), but ArMgBr perplexed us again—we isolated exclusively the product of a double 1,4-addition **A10** (69%).

General considerations. According to the equation of Klopman and Salem,^[26] the reactivity of electrophiles and nucleophiles is governed by three terms: Coulombic interaction, orbital overlap, and Pauli repulsion. When applied to α,β -unsaturated carbonyl systems, the Coulombic term is in favor of 1,2-additions, while orbital overlap promotes 1,4-additions. For acene quinones, exclusively 1,2-additions with nucleophiles PhCCLi and ArMgBr are documented in the

literature.^[12,18,19] In contrast, phenalenone, a compound of similar size but different conjugation topology, undergoes solely 1,4-additions^[25] (Figure 1). Clearly, the topological equivalence of triangulenedione and phenalenone leads to the same selectivity. On the other hand, the case of anthanthrone reveals that the rules, which govern selectivity, go beyond topology. Necessarily, the size of the nanographene core must be in play as well.

Indeed, we found three cases of extended Kekulé nanographene diones, where solely the products of 1,4-addition with ArMgBr were isolated.^[27] This unexpected selectivity, which has never been fully rationalized, might be the reason for the scarce use of this method in the nanographene synthesis (see sections below). In what follows, we analyze the contribution of each term of the Klopman–Salem equation and provide the basis for predicting the selectivity of nucleophilic additions to zigzag nanographene ketones.

Theoretical insight. We performed DFT calculations to determine charges (q , Hirshfeld) and LUMO coefficients (c) at the relevant positions of triangulenedione and anthanthrone. The former provides an estimate of the Coulombic term, while the latter estimates the orbital overlap term (Figure 2).




Figure 2. Hirshfeld charges (q), orbital coefficients in LUMO (c), and Fukui functions (f) for triangulenedione and anthanthrone calculated by DFT (gray table); the highest out of the values for 1,2- and 1,4-additions is shown in bold. The C–C bond lengths (Å, bold) are shown in the structures.

The carbonyl carbon atoms in both compounds bear the largest and similar positive charge and their coefficients in LUMO are smaller than those at the 1,4-positions. Compared to phenalenone and anthraquinone (Figure S2), the core extension leads to an increase of the $c_{1,4}/c_{1,2}$ ratio, while the charges remain qualitatively the same. The same trend is observed upon longitudinal extension in acene quinones (Figure S3). Based on the documented regioselectivity of phenalenone and anthraquinone, it may appear that it is dominated by orbital overlap. This observation holds true also for non-Kekulé triangulenedione but in the case of anthanthrone the situation is more complex.

The charges and the change of the $c_{1,4}/c_{1,2}$ ratio with the size of the π -system do not allow to make clear-cut predictions

in every case. For this reason, we calculated the Fukui functions (f), which express the change in electron density upon changing the number of electrons in the system. In our case, it is the change in electron density of the electrophile (ketone) upon addition of the nucleophilic electron pair. The Fukui function thus combines the Coulombic and the orbital overlap terms of the Klopman–Salem equation into one reactivity descriptor. We decomposed the Fukui function to individual positions in the ketones, an approach that has been successfully applied by others^[28] to understand the observed experimental selectivity. Note, however, that the steric repulsion (s) is not included in the Fukui function. The corresponding rate constants (k) for the respective additions (1,2 or 1,4) of a given nucleophile can be therefore thought of in terms of the simplistic relations (1) and (2).^[29]

$$k_{1,2} \approx f_{1,2} s_{1,2}^{-1} \quad (1)$$

$$k_{1,4} \approx f_{1,4} s_{1,4}^{-1} \quad (2)$$

Analysis of triangulenedione. The calculated charges (q) reflect the expected trend (Figure 2, left). They decrease monotonically with increasing distance from the carbonyl center (2–4–6–6′). Non-intuitively, the orbital coefficients (c) follow exactly the opposite trend, with the largest contribution to the LUMO at position 6′. The Fukui function (f), however, breaks the monotonous nature of these trends, with the highest values at positions 4 (0.054) and 6′ (0.060). One would then assume that the nucleophilic addition to **T1** takes place preferably at position 6′ in contrast to the experimental finding—exclusive attack at position 4 (Scheme 1, Table 1). This result suggests that metal coordination to carbonyl's oxygen atom is required for the reaction to take place. We confirmed this hypothesis by performing the addition to **T1** with ArLi with and without DMPU as a chelating agent for Li ions. While ArLi forms the expected product **T2** in 44 % yield, we observed no reaction in the presence of DMPU^[30] (Table 1). The same result was obtained with Kekulé anthanthrones **A1** and **A8** (Table 1). We can thus conclude that conjugate additions on nanographene diones require a directing group. In triangulenedione, 1,2- and two conjugate 1,4- and 1,4′-additions are therefore feasible and the Fukui function correctly predicts the experimental outcome. Indeed, a 1,4′-addition would disrupt one of the two Clar's sextets, unlike 1,4-addition (see Scheme 1). This result is further corroborated by the calculated C–C bond lengths (Figure 2).

Analysis of anthanthrone. As in triangulenedione, the Coulombic interaction clearly favors 1,2- over 1,4-addition, but the orbital overlap prefers 1,4- over 1,2-addition only moderately (Figure 2). As a result, the Fukui function at positions 2 (0.044) and 4 (0.041) has comparable values, with a slightly higher value at position 2. Accordingly, reaction of one equivalent of ArMgBr with **A1** displays both selectivities, but with a preference for 1,4-addition ($k_{1,4}/k_{1,2} \approx 3:2$, Scheme 2, Table 1). Reactions with PhCClLi, however, differ and give exclusively the products of 1,2-addition. This observation is not so surprising because the Fukui function does not include Pauli repulsive interactions and the sizes of these two nucleophiles significantly differ. Bürgi and Dunitz

showed that nucleophiles attack carbonyl centers under an ideal angle of 107°. ^[31] This means that the nucleophile can experience a steric clash with the π -electron cloud in extended nanographene ketones (Figure 3). On the other hand, the nucleophile in a conjugate addition attacks the reactive center at the edge from the outside of the nanographene core. Consequently, 1,2-additions suffer more from Pauli repulsion and in the case of bulkier nucleophiles, such as aryls, the Pauli term can outweigh the other two terms in the Klopman–Salem equation, steering the selectivity towards conjugate addition. The excess of ArMgBr gives a single product of a mixed 1,2-/1,4-addition in a very good yield (83 %),^[32] which indicates that each of the two intermediates **A12** and **A13** (Figure 4) formed after the first ArMgBr addition undergoes a selective complementary second addition (Scheme 2). That means that **A12** undergoes a 1,4-addition and **A13** a 1,2-addition. While the latter clearly follows the prediction of the Fukui functions (Figure 4, right), the former is a borderline case, where the sterics must outweigh the minor electronic preference for 1,2- in favor of 1,4-addition. If the steric repulsion has such a strong effect, then one expects that increasing the steric bulk of the




Figure 3. Illustration of the effect of the Pauli repulsion on the nucleophilic 1,2- and 1,4-addition to anthanthrone.

	1,2-addition			1,4-addition		
	A12: M = MgBr			A13: M = MgBr		
A_N	1,2	1,4	1,4'	1,2	1,4	1,4'
q	0.147	0.041	0.019	0.134	-0.006	-0.030
$ c $	0.239	0.297	0.012	0.303	0.305	0.249
f	0.054	0.053	0.015	0.065	0.050	0.044

Figure 4. Hirshfeld charges (q), orbital coefficients in LUMO (c), and Fukui functions (f) for **A12** and **A13** ($R = \text{Ar} = 3,5\text{-di-}t\text{-butylphenyl}$) calculated by DFT (gray table); the highest out of two values for 1,2- and 1,4-additions is shown in bold; calculations were carried out for $R = \text{Nu} = \text{M} = \text{H}$ (**A12**) and $R = \text{Nu} = \text{H}$, $\text{OM} = \text{O}^-$ (**A13**).

nucleophile will push the selectivity in favor of 1,4-addition even more in each of the two steps. In an extreme case, even a product of a double 1,4-addition could form. To confirm this hypothesis, we performed the reaction of **A1** with an excess of mesityl magnesium bromide (MesMgBr, Table 1) and isolated exclusively the product of a double 1,4-addition **A11** (91 %).^[33] The same change of selectivity upon increasing steric bulk of the nucleophile was observed for benzophenone.^[34] The steric argument holds true also for the case of 1,2-addition of thienyl lithium to anthanthrone,^[20a] as thienyl is less bulky than 3,5-di-*tert*-butylphenyl.

The reaction of **A1** with an excess of ArLi affords two products: mixed **A4** (75 %, Table 1) and the product of a double 1,4-addition **A7** (22 %). The formation of **A7** is unexpected considering that the size of nucleophile (Ar) did not change compared to ArMgBr and organolithiums are known to favor 1,2-additions.^[35] Because in ArLi the C–M bond is more polarized than in ArMgBr, ArLi is more basic and reactive, which might give an impression that it is also harder. In our reactions, however, ArLi appears to be softer than ArMgBr, which is in agreement with calculated average local ionization energies (PhLi: 6.0 eV, PhMgBr: 8.2 eV).^[36]

To exclude any steric effect of the R substituent on 1,2-additions, we tested the same reactions on **A8** equipped with sterically non-demanding substituents (R = (4-octylphenyl)ethynyl), Scheme 2, Table 1). Both ArMgBr and ArLi gave the same unexpected result, a single product of a double 1,4-addition. This rules out the steric effect of the R groups and points at an enhanced electronic effect. Indeed, the ethynyl groups perturb the frontier molecular orbitals and affect the Fukui functions. While in **A1** and **A13**, the Fukui functions favor 1,2-addition, this preference diminishes with ethynyl side groups (Figure S5). Even though the change is not large, it has a profound consequence on selectivity.

Rules of thumb. Our observations can be summarized into the following basic set of rules for nucleophilic additions to zigzag nanographene ketones:

1. A directing group is required.
2. Conjugation topology affects the electronic preference for 1,2- versus 1,4-selectivity, expressed by $f_{1,4}/f_{1,2}$ ratio (Table 2, vertical).
3. Increasing the extent of π -electron cloud and size of the nucleophile steers the selectivity towards 1,4-addition (Table 2, row 2).

Table 2: Rules of thumb.

Nu (\rightarrow) dione (\downarrow)	non-bulky (e.g., PhCCLi)	bulky (e.g., ArMgBr)
$f_{1,4}/f_{1,2} > 1$ ^[a]	1,4 (phenaleneone)	1,4 (triangulenedione)
$f_{1,4}/f_{1,2} \approx 1$ ^[b]	1,2 (anthanthrone)	1,2 and 1,4 (anthanthrone)
$f_{1,4}/f_{1,2} < 1$ ^[c]	1,2 (anthraquinone)	1,2 (anthraquinone)


[a] Typical for non-Kekulé conjugation topology (possible also for Kekulé, see Applications section). [b] Possible for both non-Kekulé and Kekulé conjugation topology. [c] Typical for Kekulé conjugation topology.

4. Steric bulk imposed by additional substituents at or close to the reactive position or non-planar geometry may alter the expected selectivity (see Applications section).


Applications. In the chemical literature, nucleophilic addition is uncommon as a method to functionalize nanographene ketones. Our results show that the reason for this might be that a mixture of products is often formed, in particular if an excess of nucleophile is used. Indeed, we found three reports describing this issue.^[27a,37] Although in some cases, products can be formed selectively and in good yields, understanding the principles that govern these reactions is crucial for controlling the selectivity. Below, we test the predictive power of our analysis on three examples from the literature.

In two reported reactions of extended Kekulé nanographene diones with ArMgBr, 1,4-addition products were obtained in good yields (Scheme 3). The “butterfly” system from Zhang et al. (2010)^[27c] is analogous to anthanthrone, the second one^[27b] to anthraquinone. In the former case, Fukui functions show a preference for 1,2-addition, like in anthanthrone (rule no. 2). We would, however, expect that the steric clash between nucleophile and the “butterfly arms” of the nanographene dione significantly increases the magnitude of the steric repulsion compared to anthanthrone, which should outweigh the electronic preference (rule no. 4). Indeed, the unexpected Michael addition is the experimental outcome.^[27c] The Fukui functions calculated for the unsymmetrically extended hexacenequinone system from Li et al. (2011)^[27b] favor 1,4-addition, unlike in anthraquinone or anthanthrone. In this scenario, 1,4-addition should be the unequivocal result, which is the case (rule no. 2).

An intriguing observation was made during the synthesis of persistent [7]uthrene, another non-Kekulé triplet diradical (Li et al., 2014).^[38] The first attempted synthetic route relied on a double 1,2-addition to [7]uthredione (**U1**, Scheme 4).




Scheme 3. Application to examples from literature. [a] Note that only a partial oxidation in air of the intermediate hydroquinone obtained after the work-up takes place. Further dehydrogenation did not occur even upon heating the product in the presence of *p*-chloranil at reflux.



Scheme 4. Nucleophilic additions to [7]uthrenedione.

The use of an excess of MesMgBr led to a complex reaction mixture and no desired product could be detected. It led the authors to re-design the synthesis and to install the substituents by a nucleophilic addition to a dialdehyde prior to the nanographene core formation by the Friedel–Crafts alkylation. Once we calculated the Fukui functions for **U1** (Figure S6), this system caught our attention. Analogously to triangulenedione, no 1,2- but 1,4-addition is predicted (rule no. 2), despite the twisted backbone on account of the methyl groups. We therefore re-synthesized **U1** and tested the selectivity of additions with ArMgBr. Stepwise addition/oxidation provided 1,4-addition products **U2** (22%, Scheme 4) and **U3** (26%). The reactions with one equivalent of ArMgBr were selective but slow and the unreacted starting material was recovered. Interestingly, the addition of a large excess of ArMgBr to **U3** afforded **U4** (55%), the product of 1,2-addition, despite a lower f value ($f_{1,4}/f_{1,2} \approx 1.2$). We attribute this change in the selectivity to a steric hindrance imposed by the substituent present in position 4 that must be more significant than that of the π -electron cloud (rule no. 4), the extent of which is intermediate between those of acene quinones and anthanthrone.

To test if triangulenedione gives a similar result as [7]uthrenedione, we performed the second and the third addition with ArMgBr to **T2** (Scheme 5). The second addition/oxidation proceeded as expected and the Ar group was installed at the second, unsubstituted position 4, affording **T3** in an excellent yield (98%). Interestingly, the third addition also gave a product of 1,4-addition **T5** (53%), even though the addition occurred at a position already bearing one Ar substituent (Table 2, row 1). Compared to [7]uthrenedione, the steric hindrance of the Ar substituent does not outweigh the more pronounced electronic preference of triangulenedione for 1,4-addition ($f_{1,4}/f_{1,2} \approx 2$, rule no. 2) combined with the steric hindrance imposed by the larger π -electron cloud disfavoring 1,2-addition (rule no. 3). When an excess of ArMgBr was reacted with **T3**, a product of 1,4- and subsequent 1,6-addition **T6** was isolated (37%). To the best of our knowledge, this is the first example of 1,6-addition on a nanographene dione. It can be rationalized by the fact that 1) the Fukui function at position 6'' of intermediate **T4**



Scheme 5. Nucleophilic additions to **T2**. Calculations of the Fukui functions (f) were carried out for triangulenedione (**T3**) and OM=O (**T4**).

(Figure S7) formed upon 1,4-addition is comparable to those at positions 2 and 4, hindered by the Ar group, and larger than that in position 4'; and 2) metal coordination is possible with the oxygen atom of the oxy group, which acts as the directing group (rule no. 1). This result demonstrates the versatility of this method by judicious placement of a suitable directing group.

Conclusion

We explored the regioselectivity of nucleophilic additions of carbon-centered nucleophiles ArMgBr and PhCClLi to zigzag nanographene ketones, namely, non-Kekulé triangulene-4,8-dione and Kekulé anthanthrone. We observed a number of unexpected results and unequivocally rationalized them as an interplay of electronic effects, expressed by Fukui functions, and sterics related to the size of nanographene and nucleophile. We compiled our findings into a basic set of rules that govern these reactions (Table 2), the use of which is rare in the chemical literature. These principles helped us explain the few known examples with unanticipated selectivity.

The synthesis of new nanographenes aids and abets understanding of physical phenomena in graphene-based materials, important for the future development of organic electronics. Functionalization of their periphery is crucial for controlling their electronic structure, solubility, stability, and self-assembly behavior. In this respect, the selectivity rules of nucleophilic additions to zigzag nanographene ketones developed in this work do not only allow to install substituents at a late stage of the synthesis, but make it possible to address positions which would not be accessible otherwise. In addition, this method will facilitate synthesis and exploration of new functionalized nanographenes as a large number of

nanographene ketones have been described since the pioneering work of Eric Clar.

Acknowledgements

This project received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant Agreement No. 716139), the Swiss National Science Foundation (SNSF, T.Š./PZ00P2_174175, M.J./PZ00P2_148043, PP00P2_170534), and Experientia Foundation (T.Š.). We would like to thank Prof. Marcel Mayor (University of Basel) for his generous support of our research and the reviewers for their useful suggestions to improve our manuscript.

Conflict of interest

The authors declare no conflict of interest.

Keywords: Fukui function analysis · Michael addition · nucleophilic addition · regioselectivity · zigzag nanographene

- [1] a) J. Yamaguchi, H. Hayashi, H. Jippo, A. Shiotari, M. Ohtomo, M. Sakakura, N. Hieda, N. Aratani, M. Ohfuchi, Y. Sugimoto, H. Yamada, S. Sato, *Commun. Mater.* **2020**, *1*, 36; b) Y.-W. Son, M. L. Cohen, S. G. Louie, *Phys. Rev. Lett.* **2006**, *97*, 216803.
- [2] a) O. V. Yazyev, *Rep. Prog. Phys.* **2010**, *73*, 056501; b) J. Jung, T. Pereg-Barnea, A. H. MacDonald, *Phys. Rev. Lett.* **2009**, *102*, 227205; c) W. Y. Kim, K. S. Kim, *Nat. Nanotechnol.* **2008**, *3*, 408–412.
- [3] a) M. Grzybowski, B. Sadowski, H. Butenschön, D. T. Gryko, *Angew. Chem. Int. Ed.* **2020**, *59*, 2998–3027; *Angew. Chem.* **2020**, *132*, 3020–3050; b) A. Senese, W. Chalifoux, *Molecules* **2018**, *24*, 118; c) A. Narita, X.-Y. Wang, X. Feng, K. Müllen, *Chem. Soc. Rev.* **2015**, *44*, 6616–6643.
- [4] a) X.-Y. Wang, X. Yao, K. Müllen, *Sci. China Chem.* **2019**, *62*, 1099–1144; b) J. Wu, W. Pisula, K. Müllen, *Chem. Rev.* **2007**, *107*, 718–747.
- [5] a) J. Inoue, K. Fukui, T. Kubo, S. Nakazawa, K. Sato, D. Shiomi, Y. Morita, K. Yamamoto, T. Takui, K. Nakasuji, *J. Am. Chem. Soc.* **2001**, *123*, 12702–12703; b) K. Goto, T. Kubo, K. Yamamoto, K. Nakasuji, K. Sato, D. Shiomi, T. Takui, M. Kubota, T. Kobayashi, K. Yakusi, J. Ouyang, *J. Am. Chem. Soc.* **1999**, *121*, 1619–1620.
- [6] a) C. D. Simpson, J. D. Brand, A. J. Berresheim, L. Przybilla, H. J. Räder, K. Müllen, *Chem. Eur. J.* **2002**, *8*, 1424–1429.
- [7] a) J. E. Anthony, *Angew. Chem. Int. Ed.* **2008**, *47*, 452–483; *Angew. Chem.* **2008**, *120*, 460–492; b) J. E. Anthony, *Chem. Rev.* **2006**, *106*, 5028–5048.
- [8] a) J. Mei, Y. Diao, A. L. Appleton, L. Fang, Z. Bao, *J. Am. Chem. Soc.* **2013**, *135*, 6724–6746; b) M. Watanabe, Y. J. Chang, S.-W. Liu, T.-H. Chao, K. Goto, Md. M. Islam, C.-H. Yuan, Y.-T. Tao, T. Shinmyozu, T. J. Chow, *Nat. Chem.* **2012**, *4*, 574–578.
- [9] a) R. Rieger, K. Müllen, *J. Phys. Org. Chem.* **2010**, *23*, 315–325; b) K.-H. Koch, K. Müllen, *Chem. Ber.* **1991**, *124*, 2091–2100.
- [10] a) B. Purushothaman, M. Bruzek, S. R. Parkin, A.-F. Miller, J. E. Anthony, *Angew. Chem. Int. Ed.* **2011**, *50*, 7013–7017; *Angew. Chem.* **2011**, *123*, 7151–7155; b) M. M. Payne, S. R. Parkin, J. E. Anthony, *J. Am. Chem. Soc.* **2005**, *127*, 8028–8029.
- [11] a) I. Kaur, M. Jazdyk, N. N. Stein, P. Prusevich, G. P. Miller, *J. Am. Chem. Soc.* **2010**, *132*, 1261–1263; b) H. Qu, C. Chi, *Org. Lett.* **2010**, *12*, 3360–3363.
- [12] J. E. Anthony, J. S. Brooks, D. L. Eaton, S. R. Parkin, *J. Am. Chem. Soc.* **2001**, *123*, 9482–9483.
- [13] Phenalenyl has an odd number of carbon atoms and is not, strictly speaking, a non-Kekulé system. Nevertheless, it is generally considered to have non-Kekulé topology because of a similar distribution of spin density compared to non-Kekulé systems such as triangulene. For IUPAC definition of non-Kekulé systems, see: V. I. Minkin, *Pure Appl. Chem.* **1999**, *71*, 1919.
- [14] a) H. Ito, Y. Segawa, K. Murakami, K. Itami, *J. Am. Chem. Soc.* **2019**, *141*, 3–10; b) B. P. Mathew, M. R. Kuram, *Inorg. Chim. Acta* **2019**, *490*, 112–129; c) H. Ito, K. Ozaki, K. Itami, *Angew. Chem. Int. Ed.* **2017**, *56*, 11144–11164; *Angew. Chem.* **2017**, *129*, 11296–11317.
- [15] Polycyclic aromatic hydrocarbons, or nanographenes, have two basic types of edges, armchair or zigzag (see ref. [9a]). The term “zigzag nanographene” refers to systems in which all edges are zigzag. For example, triangulene and anthanthrene have three and four zigzag edges, respectively.
- [16] a) Y. Gu, Y. G. Tullimilli, J. Feng, H. Phan, W. Zeng, J. Wu, *Chem. Commun.* **2019**, *55*, 5567–5570; b) W. Zeng, T. Y. Gopalakrishna, H. Phan, T. Tanaka, T. S. Herng, J. Ding, A. Osuka, J. Wu, *J. Am. Chem. Soc.* **2018**, *140*, 14054–14058; c) Y. Gu, X. Wu, T. Y. Gopalakrishna, H. Phan, J. Wu, *Angew. Chem. Int. Ed.* **2018**, *57*, 6541–6545; *Angew. Chem.* **2018**, *130*, 6651–6655.
- [17] a) E. Clar, R. Schoental, *Polycyclic Hydrocarbons, Vol. 1*, Springer, Berlin, **1964**; b) E. Clar, *Polycyclic Hydrocarbons, Vol. 2*, Springer, Berlin, **1964**.
- [18] Examples with lithium acetylide: a) D. Lehnher, J. Gao, F. A. Hegmann, R. R. Tykwinski, *Org. Lett.* **2008**, *10*, 4779–4782; b) M. S. Taylor, T. M. Swager, *Org. Lett.* **2007**, *9*, 3695–3697; c) D. P. Lydon, L. Porrès, A. Beeby, T. B. Marder, P. J. Low, *New J. Chem.* **2005**, *29*, 972–976; d) C.-C. A. Voll, T. M. Swager, *J. Am. Chem. Soc.* **2018**, *140*, 17962–17967; e) V. H. Gessner, T. D. Tilley, *Org. Lett.* **2011**, *13*, 1154–1157; f) J. Wang, K. Liu, Y.-Y. Liu, C.-L. Song, Z.-F. Shi, J.-B. Peng, H.-L. Zhang, X.-P. Cao, *Org. Lett.* **2009**, *11*, 2563–2566; g) I. Kaur, W. Jia, R. P. Kopsreski, S. Selvarasah, M. R. Dokmeci, C. Pramanik, N. E. McGruer, G. P. Miller, *J. Am. Chem. Soc.* **2008**, *130*, 16274–16286; Examples with (hetero)aryl magnesium bromide: h) R. M. Chalke, V. R. Patil, *Polymer* **2017**, *123*, 355–365; i) D.-S. Kim, J. E. Jung, N. S. Baek, T.-D. Kim, *Org. Electron.* **2015**, *17*, 319–324.
- [19] B. Purushothaman, S. R. Parkin, J. E. Anthony, *Org. Lett.* **2010**, *12*, 2060–2063.
- [20] a) J.-B. Giguère, J. Boismenu-Lavoie, J.-F. Morin, *J. Org. Chem.* **2014**, *79*, 2404–2418; b) T. Matsuno, S. Kamata, S. Hitosugi, H. Isobe, *Chem. Sci.* **2013**, *4*, 3179–3183; c) J.-B. Giguère, Q. Verolet, J.-F. Morin, *Chem. Eur. J.* **2013**, *19*, 372–381; d) L. Zhang, B. Walker, F. Liu, N. S. Colella, S. C. B. Mannsfeld, J. J. Watkins, T.-Q. Nguyen, A. L. Briseno, *J. Mater. Chem.* **2012**, *22*, 4266–4268.
- [21] B. K. Shah, D. C. Neckers, J. Shi, E. W. Forsythe, D. Morton, *J. Phys. Chem. A* **2005**, *109*, 7677–7681.
- [22] N. Pavliček, A. Mistry, Z. Majzik, N. Moll, G. Meyer, D. J. Fox, L. Gross, *Nat. Nanotechnol.* **2017**, *12*, 308–311.
- [23] E. Clar, D. G. Stewart, *J. Am. Chem. Soc.* **1953**, *75*, 2667–2672.
- [24] Nucleophilic additions with MeMgBr and MeLi as the model sp³-carbon nucleophiles were also performed. All attempts, however, led either to a decomposition of the starting material or a mixture of many products that could not be identified, except for the reaction of **T1** with one equivalent of MeMgBr, which gave a product of 1,4-addition.

- [25] a) C. M. Wehrmann, R. T. Charlton, M. S. Chen, *J. Am. Chem. Soc.* **2019**, *141*, 3240–3248; b) L. I. Rosquete, M. G. Cabrera-Serra, J. E. Piñero, P. Martín-Rodríguez, L. Fernández-Pérez, J. G. Luis, G. McNaughton-Smith, T. Abad-Grillo, *Bioorg. Med. Chem.* **2010**, *18*, 4530–4534; c) M. S. Newman, L.-F. Lee, *J. Org. Chem.* **1975**, *40*, 2650–2652.
- [26] a) I. Fleming, *Ionic Reactions*, in *Frontier Orbitals and Organic Chemical Reactions*, Wiley, Chichester, **1976**, pp. 33–85; b) G. Klopman, *J. Am. Chem. Soc.* **1968**, *90*, 223–234; c) L. Salem, *J. Am. Chem. Soc.* **1968**, *90*, 543–552.
- [27] a) C. Reus, M. P. Lechner, M. Schulze, D. Lungerich, C. Diner, M. Gruber, J. M. Stryker, F. Hampel, N. Jux, R. R. Tykwinski, *Chem. Eur. J.* **2016**, *22*, 9097–9101; b) J. Li, C. Jiao, K.-W. Huang, J. Wu, *Chem. Eur. J.* **2011**, *17*, 14672–14680; c) X. Zhang, J. Li, H. Qu, C. Chi, J. Wu, *Org. Lett.* **2010**, *12*, 3946–3949.
- [28] a) B. Wang, C. Rong, P. K. Chattaraj, S. Liu, *Theor. Chem. Acc.* **2019**, *138*, 124; b) S. Liu, C. Rong, T. Lu, *J. Phys. Chem. A* **2014**, *118*, 3698–3704; c) R. K. Roy, *J. Phys. Chem. A* **2003**, *107*, 10428–10434.
- [29] To support the view that these reactions occur under kinetic control, the following crossover experiment suggested by a reviewer was carried out: The reaction of **A1** with 10 equivalents of ArMgBr (Table 1) was performed as usual, but before quenching, the reaction mixture was cooled to -78°C and 20 equivalents of *p*-tolylmagnesium bromide (TolMgBr) were added. Upon warming to room temperature and work-up, no product of TolMgBr addition was observed, only **A4** was formed.
- [30] Chelation of Li ions by DMPU was shown to increase the reactivity of the nucleophile (which we did not observe in our case) and to alter the selectivity of the nucleophilic addition to cyclohexenone from 1,2 to 1,4: T. Mukhopadhyay, D. Seebach, *Helv. Chim. Acta* **1982**, *65*, 385–391.
- [31] H. B. Bürgi, J. D. Dunitz, J. M. Lehn, G. Wipff, *Tetrahedron* **1974**, *30*, 1563–1572.
- [32] Variable-concentration experiments suggested by a reviewer revealed that the decrease of concentration from $c(\text{ArMgBr}) = 10 \times c(\mathbf{A1}) \approx 80 \text{ mM}$ (reaching the solubility limit of **A1**) by a factor of three leads to a decrease of the yield by approx. 20%, while no other product is observed. The addition of LiCl (1.2 equivalent with respect to ArMgBr) at $c(\text{ArMgBr}) \approx 80 \text{ mM}$ does not change the reaction outcome. These results indicate that in the range $c(\text{ArMgBr}) \approx 26\text{--}80 \text{ mM}$, the aggregation effects of ArMgBr can be neglected. Typically, the dione concentration was approx. 5 mM.
- [33] After the submission of this manuscript, Du et al. reported a double 1,4-addition of MesMgBr to 4,10-dibromoanthanthrone. Interestingly, the addition of 4-dodecylphenyllithium to the same dione gave a product of a double 1,2-addition. Both results are in agreement with our hypothesis, as *para*-substituted phenyl is less bulky than 3,5-di-*tert*-butylphenyl. Y. Du, H. B. Lovell, F. Lirette, J.-F. Morin, K. N. Plunkett, *J. Org. Chem.* **2021**, *86*, 1456–1461.
- [34] M. Debeaux, K. Brandhorst, P. G. Jones, H. Hopf, J. Grunenberg, W. Kowalsky, H.-H. Johannes, *Beilstein J. Org. Chem.* **2009**, *5*, 31.
- [35] E. Amoah, R. K. Dieter, *J. Org. Chem.* **2017**, *82*, 2870–2888.
- [36] J. J. Brown, S. L. Cockroft, *Chem. Sci.* **2013**, *4*, 1772–1780.
- [37] a) P. Hu, S. Lee, K. H. Park, S. Das, T. S. Heng, T. P. Gonçalves, K.-W. Huang, J. Ding, D. Kim, J. Wu, *J. Org. Chem.* **2016**, *81*, 2911–2919; b) Q. Ye, J. Chang, K.-W. Huang, G. Dai, J. Zhang, Z.-K. Chen, J. Wu, C. Chi, *Org. Lett.* **2012**, *14*, 2786–2789.
- [38] Y. Li, K.-W. Huang, Z. Sun, R. D. Webster, Z. Zeng, W. Zeng, C. Chi, K. Furukawa, J. Wu, *Chem. Sci.* **2014**, *5*, 1908–1914.

Manuscript received: December 14, 2020

Accepted manuscript online: March 1, 2021

Version of record online: May 1, 2021