

RESEARCH ARTICLE

# First-trimester smoking cessation in pregnancy did not increase the risk of preeclampsia/eclampsia: A Murmansk County Birth Registry study

Olga A. Kharkova<sup>1,2\*</sup>, Andrej M. Grjibovski<sup>2,3,4</sup>, Alexandra Krettek<sup>1,5,6</sup>, Evert Nieboer<sup>7</sup>, Jon Ø. Odland<sup>1,8</sup>

**1** Department of Community Medicine, Faculty of Health Sciences, UiT The Arctic University of Norway, Tromsø, Norway, **2** International School of Public Health, Northern State Medical University, Arkhangelsk, Russia, **3** Department of International Public Health, Norwegian Institute of Public Health, Oslo, Norway, **4** Department of Preventive Medicine, International Kazakh-Turkish University, Turkestan, Kazakhstan, **5** Department of Biomedicine and Public Health, School of Health and Education, University of Skövde, Skövde, Sweden, **6** Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden, **7** Department of Biochemistry and Biomedical Sciences, Hamilton, ON, Canada, **8** School of Health Systems and Public Health, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

\* [okh000@post.uit.no](mailto:okh000@post.uit.no)



**OPEN ACCESS**

**Citation:** Kharkova OA, Grjibovski AM, Krettek A, Nieboer E, Odland JØ (2017) First-trimester smoking cessation in pregnancy did not increase the risk of preeclampsia/eclampsia: A Murmansk County Birth Registry study. PLoS ONE 12(8): e0179354. <https://doi.org/10.1371/journal.pone.0179354>

**Editor:** Kelli K. Ryckman, University of Iowa, UNITED STATES

**Received:** March 30, 2017

**Accepted:** May 26, 2017

**Published:** August 10, 2017

**Copyright:** © 2017 Kharkova et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** According to the Russian legislation (Federal Law №152 from 27th of July 2006 “Personal data”), cross-border data transfer may be prohibited or restricted in order to protect constitutional rights of the Russian citizens. However existing Russian legislation allows the attachment of the dataset from the corresponding author on the basis of properly submitted requests. Data requests may also be sent to Anton Kovalenko ([anton.a.kovalenko@uit.no](mailto:anton.a.kovalenko@uit.no)). For additional information, please contact Andrej B. Gudkov,

## Abstract

### Background

Although prior studies have shown that smoking reduces preeclampsia/eclampsia risk, the consequence of giving up this habit during pregnancy should be assessed. The aims of the current study were threefold: (i) describe maternal characteristics of women with preeclampsia/eclampsia; (ii) examine a possible association between the number of cigarettes smoked daily during pregnancy and the development of this affliction; and (iii) determine if first-trimester discontinuation of smoking during pregnancy influences the risk.

### Methods

A registry-based study was conducted using data from the Murmansk County Birth Registry (MCBR). It included women without pre-existing hypertension, who delivered a singleton infant during 2006–2011 and had attended the first antenatal visit before 12 week of gestation. We adjusted for potential confounders using logistic regression.

### Results

The prevalence of preeclampsia/eclampsia was 8.3% (95%CI: 8.0–8.6). Preeclampsia/eclampsia associated with maternal age, education, marital status, parity, excessive weight gain and body mass index at the first antenatal visit. There was a dose-response relationship between the number of smoked cigarettes per day during pregnancy and the risk of preeclampsia/eclampsia (adjusted  $OR_{1-5 \text{ cig/day}} = 0.69$  with 95%CI: 0.56–0.87;  $OR_{6-10 \text{ cig/day}} = 0.65$  with 95%CI: 0.51–0.82; and  $OR_{\geq 11 \text{ cig/day}} = 0.49$  with 95%CI: 0.30–0.81). There was no

Chairman of the Ethical Committee of Northern State Medical University, Arkhangelsk, Russia at 163000 Russia Arkhangelsk, Troitsky ave 51, Tel.: (8182)28-57-91.

**Funding:** The authors received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

**Abbreviations:** MCBR, Murmansk County Birth Registry; WHO, World health organization; EMRO, Eastern Mediterranean region; AFRO, African region; EURO, European region; ICD, International classification of diseases; BMI, Body mass index; CO, carbon monoxide.

difference in this risk among women who smoked before and during pregnancy and those who did so before but not during pregnancy (adjusted OR = 1.10 with 95%CI: 0.91–1.32).

## Conclusions

Preeclampsia/eclampsia was associated with maternal age, education, marital status, parity, excessive weight gain, and body mass index at the first antenatal visit. There was a negative dose-response relationship between the number of smoked cigarettes per day during pregnancy and the odds of preeclampsia/eclampsia. However, women who gave up smoking during the first trimester of gestation had the same risk of preeclampsia/eclampsia as those who smoked while pregnant. Consequently, antenatal clinic specialists are advised to take these various observations into account when counselling women on smoking cessation during pregnancy.

## Introduction

Preeclampsia and eclampsia are common complications of pregnancy that annually affect 8,370,000 women worldwide [1, 2]. Preeclampsia in a woman who before 20 weeks' gestation was previously normotensive is defined by the presence of hypertension (systolic or diastolic blood pressures of >140mm and >90 mm Hg) in combination with proteinuria with or without oedema [3–5]. Eclampsia constitutes the onset of seizures in a woman with preeclampsia [6].

The crude incidences of preeclampsia between 2002 and 2010 in the Eastern Mediterranean (EMRO) and Western Pacific (WPRO) WHO regions, respectively, were 1.2 to 4.2% [7]. When a logistic model using bootstrapped data was employed to estimate the incidence of preeclampsia using country macroeconomic indicators of health care and population characteristics for adjustment, the overall estimate for all WHO regions was 4.6% (with a 95% uncertainty range of 2.7–8.2); in the EMRO and AFRO WHO regions the incidences were 1.0 and 5.6%, respectively. In the EURO WHO region excluding Russia (due to missing data), the crude incidence of preeclampsia was 3.8%, with a corresponding model-based incidence (in %) of 5.3 (1.8–9.3) [7]. The overall prevalence of preeclampsia/eclampsia in Russia depends on the year of measurement and the federal district [8–10]. For example, during 2005–2010 the prevalence ranged from 2.1 to 1.4% in the Southern and Northern Caucasians, 0.1–0.6% in Far Eastern federal districts and 2.4–2.6% in the Northwestern and Ural districts.

We recently demonstrated a high prevalence of smoking before and during pregnancy in Northwest Russia, with only one of four smokers quitting during pregnancy [11]. This raised concern since tobacco smoking during pregnancy increases the risk of placental complications including placenta previa [12] and placental abruption [13], as well as preterm birth [14] and low birth weight [15, 16]. Maternal smoking also associates with increased perinatal morbidity and mortality, while in the general population maternal smoking constitutes a risk factor for many chronic diseases including cardiovascular disease, diabetes, and inflammatory diseases [17, 18].

Cigarette smoking appears to reduce the risk of preeclampsia and eclampsia [19–21]. However, there is no consistent evidence to determine if giving up smoking during pregnancy influences this protective effect. Some studies suggest a reduced risk [22–24], while others show no effect [21, 25, 26]. This discrepancy could be a consequence of variation in study

design, sample selection or its size. Moreover, most studies do not adjust for potential confounders such as socioeconomic status [22, 23] and weight gain during pregnancy [21, 22, 24, 26]. In the present study, we employ registry data to examine potential modifiers of the impact maternal smoking has on the risk of preeclampsia/eclampsia, including putative confounding factors. Our aims are to: (i) describe maternal characteristics of women with preeclampsia/eclampsia; (ii) examine a potential association between the number of smoked cigarettes per day during pregnancy and the development of preeclampsia/eclampsia; and (iii), determine if first-trimester smoking cessation during pregnancy influences the risk of preeclampsia/eclampsia.

## Materials methods

### Study setting, design and sample size

The study was conducted in Murmansk County, which is a federal subject of the north-western part of Russia. The County borders with the Republic of Karelia in Russia, Lapland Region in Finland and Finnmark County in Norway. Murmansk County is surrounded in part by the Barents Sea and White Sea.

The study population consisted of all women who were registered in the Murmansk County Birth Registry (MCBR). Its background details are described elsewhere [11], including implementation and quality control details [27]. For the purposes of this study we excluded women if they had a multiple pregnancy, pre-existing hypertension, or had their first antenatal visit after week 12 of gestation. Three specific issues are the focus of the current publication: (i) maternal characteristics of the study participants (N = 39 566); (ii) the association between the numbers of smoked cigarettes per day and incidence of preeclampsia/eclampsia (N = 36 376); and (iii) and the impact on the latter of the first-trimester smoking cessation during pregnancy (N = 39 566). Sampling and analysis details are provided in Fig 1.


### Data collection

Based on medical records and personal interviews with the expectant mothers, the MCBR contained information on maternal characteristics including age, ethnicity, residence, educational level, marital status, parity, alcohol abuse as diagnosed by a doctor, self-reported smoking before and during pregnancy, and maternal weight and height measured at the first antenatal visit. Information in the MCBR on preeclampsia/eclampsia occurrence, excessive weight gain during pregnancy and year of delivery were derived from individual obstetric journals.

Preeclampsia and eclampsia were classified according to the International Classification of Diseases, tenth revision (ICD-10) [28]. Preeclampsia (ICD-10 codes O14.0 “mild to moderate preeclampsia”; O14.1 “severe preeclampsia”) is a pregnancy-induced hypertensive state that occurs after 20 weeks of gestation. It is characterized by hypertension (blood pressure of 140/90 or higher), along with oedema and proteinuria (300 mg of protein in a 24-hour urine sample) [8, 29]. Eclampsia (ICD-10 code O15.0) involves convulsions and coma in pregnant or puerperal women along with hypertension, oedema, and proteinuria.

We analysed preeclampsia (N = 3276) and eclampsia (N = 5) cases together because of the limited number of cases of eclampsia. The variable “preeclampsia/eclampsia” (N = 3281) was treated as binary variable.

In terms of smoking status during pregnancy, women were grouped as smokers (did so before and during pregnancy), quitters (smoked before but not during pregnancy), or non-smokers (did not smoke before nor during pregnancy). Smoking status was assessed during the first antenatal visit. Number of smoked cigarettes per day during pregnancy was taken as a categorical variable, specifically as 0, 1–5, 6–10, and  $\geq 11$ .



**Fig 1. Sampling and analysis details.**

<https://doi.org/10.1371/journal.pone.0179354.g001>

Maternal age was also treated as a categorical variable, namely:  $\leq 19$  years, 20–24 years, 25–29 years, 30–34 years and  $\geq 35$  years. Place of residence was registered as urban or rural. Ethnicity was dichotomized into Russian and other. Maternal education was divided into three categories: *as less than university* that included primary (0–9 years of schooling), secondary (10–11 years of schooling) and vocational training, *university* and *unknown*. Marital status was classified as married, cohabitating and single, with the latter including divorced and widows. Parity was categorized as 0, 1, and  $\geq 2$  previous deliveries. Alcohol abuse was recorded as either yes or no. Year of delivery was denoted by the exact calendar year.

Body mass index (BMI) was calculated for the women’s weight at the first antenatal visit (kg) divided by height (m<sup>2</sup>). By BMI, women were classified into five groups: underweight ( $\leq 18.4$  kg/m<sup>2</sup>), normal weight (18.5–24.9 kg/m<sup>2</sup>), overweight (25.0–29.9 kg/m<sup>2</sup>), obese ( $\geq 30.0$  kg/m<sup>2</sup>), and unknown.

Excessive weight gain was defined as weight gain during pregnancy of  $> 18$  kg in underweight women,  $> 16$  kg in normal weight women,  $> 11.5$  kg in overweight women, and  $\geq 6$  kg in obese women. Excessive weight gain in pregnancy (ICD-10 code O26.0) was dichotomized as yes and no.

## Data analysis

Pearson's chi-squared tests were used to analyse the categorical variables for differences. By logistic regression we examined the effects of giving up smoking during pregnancy on the risk of preeclampsia/eclampsia, and any association between preeclampsia/eclampsia and daily numbers of smoked cigarettes while pregnant. Both crude and adjusted odds ratios (ORs) and their 95% confidence intervals (CI) are reported. All statistical analyses were conducted using SPSS, version 22 (SPSS Inc., Chicago, IL).

## Ethical considerations

The study was approved by the Ethical Committee of Northern State Medical University, Arkhangelsk, Russia, and the Norwegian Regional Committee for Medical and Health Research Ethics (REC-North), Tromsø, Norway.

## Results

### Maternal characteristics of women with preeclampsia/eclampsia

Of the 39 566 participants in our study, 8.3% (95% CI: 8.0–8.6) had preeclampsia/eclampsia during their current pregnancy. Occurrence was more prevalent in women  $\geq 35$  years old, having less than university education, and who were nulliparous and single (Table 1). Preeclampsia/eclampsia developed more frequently among women who were obese at the first antenatal visit and those with normal weight. Furthermore, risk of preeclampsia/eclampsia was 6.7% among women who smoked in pregnancy and 8.7% among those who did not ( $p < 0.001$ ). The proportion of women with preeclampsia/eclampsia decreased with the number of cigarettes smoked per day ( $p < 0.001$ ), while ethnicity, residence (urban *versus* rural) and alcohol abuse had no impact.

### Association between daily numbers of smoked cigarettes during pregnancy and development of preeclampsia/eclampsia

Non-smokers both before and during pregnancy had a greater risk of preeclampsia/eclampsia compared to smokers (Table 2). A dose-response relationship was evident between the number of cigarettes smoked per day during pregnancy and the risk of preeclampsia/eclampsia ( $p_{\text{trend}} < 0.001$ ). Note that pregnant women who smoked 1–5, 6–10 or  $\geq 11$  cigarettes per day during pregnancy had decreased odds of having preeclampsia/eclampsia compared to non-smokers (crude  $OR_{1-5\text{cig.}}$  of 0.72 with 95% CI: 0.58–0.90; for  $OR_{6-10\text{cig.}}$  of 0.68 with 95% CI: 0.54–0.85; and for  $OR_{\geq 11\text{cig.}}$  of 0.46 with 95% CI: 0.28–0.74, respectively). Adjustment for potential confounders did not change the association.

### Quitting smoking during pregnancy and risk of preeclampsia/eclampsia

Women who smoked before but not during pregnancy had lower risk of having preeclampsia/eclampsia compared to those who did not smoke before and during pregnancy [adjusted OR of 0.80 (95% CI: 0.68–0.94)]. However, there was no significant difference in the occurrence of preeclampsia/eclampsia among women smoking before and during pregnancy and those who smoked before, but not during pregnancy—either before or after adjustment for other maternal characteristics (Table 3).

**Table 1. Maternal characteristics of the study participants (N = 39 566).**

Maternal characteristics	Total number	Women without preeclampsia/eclampsia		Women with preeclampsia/eclampsia		P-level
		N	%	N	%	
<i>Maternal age (years)</i>						0.001
≤ 19	2112	1943	92.0	169	8.0	
20–24	11 532	10 601	91.9	931	8.1	
25–29	13 801	12 723	92.2	1078	7.8	
30–34	8724	7955	91.2	769	8.8	
≥ 35	3397	3063	90.2	334	9.8	
<i>Residence</i>						0.223
Urban	34 349	31 478	91.6	2871	8.4	
Rural	5217	4807	92.1	410	7.9	
<i>Ethnicity</i>						0.329
Russian	38 043	34 878	91.7	3165	8.3	
Other	1523	1407	92.4	116	7.6	
<i>Education</i>						<0.001
Less than university	24 897	22 797	91.6	2100	8.4	
University	14 488	13 340	92.1	1148	7.9	
Unknown	181	148	81.8	33	18.2	
<i>Marital status</i>						<0.001
Married	30 402	27872	91.7	2530	8.3	
Cohabitation	6107	5705	93.4	402	6.6	
Single	3057	2708	88.6	349	11.4	
<i>Parity</i>						<0.001
0	22 489	20 481	91.1	2008	8.9	
1	14 742	13 649	92.6	1093	7.4	
≥2	2335	2155	92.3	180	7.7	
<i>Alcohol abuse</i>						0.844
No	39 499	36 224	91.7	3275	8.3	
Yes	67	61	91.0	6	9.0	
<i>BMI</i>						<0.001
Underweight	2478	2404	97.0	74	3.0	
Normal weight	25 836	24 165	93.5	1671	6.5	
Overweight	7331	6432	87.7	899	12.3	
Obese	2918	2374	81.4	544	18.6	
Unknown	1003	910	90.7	93	9.3	
<i>Excessive weight gain</i>						<0.001
No	37 148	34001	91.5	3147	8.5	
Yes	2418	2284	94.5	134	5.5	
<i>Smoking status during pregnancy</i>						<0.001
Non-smoker	30 690	28 019	91.3	2671	8.7	
Quitter	2534	2350	92.7	184	7.3	
Smoker	6342	5916	93.3	426	6.7	
<i>Number of smoked cigarettes per day during pregnancy*</i>						<0.001
0	33 219	30 364	91.4	2855	8.6	
1–5	1411	1321	93.6	90	6.4	
6–10	1333	1253	94.0	80	6.0	
≥ 11	413	396	95.9	17	4.1	

\* Total number is 36 376 because of missing data  
 Calculated using chi-squared test

<https://doi.org/10.1371/journal.pone.0179354.t001>

**Table 2. Association between daily numbers of smoked cigarettes during pregnancy and preeclampsia/eclampsia among women with singleton pregnancies in Murmansk County, Northwest Russia (N = 36 376).**

Variable	Crude			Adjusted <sup>1</sup>		
	OR	95% CI	P <sub>trend</sub>	OR	95% CI	P <sub>trend</sub>
<i>Number of smoked cigarettes per day during pregnancy</i>			<0.001			<0.001
0	1.00			1.00		
1–5	0.72	0.58–0.90		0.69	0.56–0.87	
6–10	0.68	0.54–0.85		0.65	0.51–0.82	
≥ 11	0.46	0.28–0.74		0.49	0.30–0.81	

<sup>1</sup> OR adjusted for variables maternal age, residence, ethnicity, marital status, parity, alcohol abuse, year of delivery, body mass index and excessive weight gain

<https://doi.org/10.1371/journal.pone.0179354.t002>

## Discussion

### Main findings

Preeclampsia/eclampsia was more common in older, single, less educated and primiparae women and those who were overweight or obese at the first antenatal visit and had no excessive weight gain during pregnancy. Non-smokers both before and during pregnancy had a greater risk of preeclampsia/eclampsia compared to smokers. A dose-response relationship was evident between the number of cigarettes smoked per day during pregnancy and the risk of preeclampsia/eclampsia. Furthermore, discontinuance of smoking during pregnancy did not increase the risk of preeclampsia/eclampsia.

### Data interpretation and comparisons with previous studies

The observed preeclampsia/eclampsia prevalence in Murmansk County of 8.3% is higher than previous estimates [7–10, 23, 30]. This could reflect different definitions and differing proportions of primiparae [31]. Preeclampsia is more common in primiparae than in multiparae, which is a potential reason for discrepancies in parity among countries. Furthermore, regional data are often different from national figures, as they reflect the variation of preeclampsia/eclampsia within one country. For example in St. Petersburg (located in the Northwest federal district of Russia), the prevalence of preeclampsia/eclampsia in 2005 was 7.1%, while it was 8.6% in Orenburg County (Volga federal district), 10.5% in Kurgan County (Ural federal district) and <0.1% in Vladimir County (Central federal district) and Vologda (Northwest federal district) [10].

**Table 3. Association between smoking status during pregnancy and preeclampsia/eclampsia among women with singleton pregnancies in Murmansk County, Northwest Russia (N = 39 566).**

Variable	Crude		Adjusted <sup>1</sup>	
	OR	95% CI	OR	95% CI
Smoking before and during pregnancy	1.00		1.00	
Smoked before, but not during pregnancy	1.09	0.91–1.30	1.10	0.91–1.32
Did not smoke both before and during pregnancy	1.32	1.19–1.47	1.37	1.23–1.54

<sup>1</sup> OR adjusted for variables maternal age, residence, ethnicity, marital status, parity, alcohol abuse, year of delivery, body mass index and excessive weight gain

<https://doi.org/10.1371/journal.pone.0179354.t003>



Our observations that the risk of preeclampsia/eclampsia increased in women who are  $\geq 35$  years old, have less education than university, are single, primiparae and overweight or obese at the first antenatal visit are consistent with earlier studies [21, 31, 32].

Our finding of a 2-fold protective effect for the development of preeclampsia/eclampsia in women who smoked more than  $\geq 11$  cigarettes per day relative to non-smokers supports previous reports [19–21]. According to Linqvist et al [32], moderate smokers (1–9 cigarettes per day) are characterized by a lower incidence of preeclampsia compared to non-smokers. Similarly, Yang et al [33] report an inverse exposure-response association as does Bainbridge et al [34]. Venditti et al [35] state that the use of carbon monoxide (CO) could prevent the development of hypertension and proteinuria in a rodent model of preeclampsia. Bainbridge et al [34] suggest that CO, a combustible product in cigarettes, may be the active agent. More recently Zhai et al [36] demonstrated an inverse correlation between increased environmental ambient CO and preeclampsia. However, any interpretations must consider that the pathogenesis of preeclampsia is complex as it appears to involve genetic, immunological and environmental factors [37].

Tobacco smoking during pregnancy can potentially impact angiogenic factors, endothelial function and the immune system, which could lead to a lower risk of preeclampsia. However, this protective role is most likely explained by CO's biological role in heme-degradation processes including the promotion of anti-inflammatory and pro-angiogenic effects [38–40]. On the other hand, the mechanisms underlying the increased risk of preeclampsia among previous and passive smokers remain unclear. Luo et al [41] suggest that this could be due to adverse chronic effects of low tobacco exposure in the absence of significant exposure to a transient protective factor such as CO in association with current smoking.

Our data suggest that discontinuing smoking after pregnancy awareness did not alter the risk of preeclampsia/eclampsia statistically speaking. By contrast, some studies demonstrate a lower incidence of preeclampsia among women who stop smoking at the beginning of pregnancy compared to those who never smoked [22, 23]. Neither do our findings align with those of England et al [26] in their randomized clinical trial “Calcium for Preeclampsia Prevention” (N = 4,589). They observed that the incidence of preeclampsia among women who stopped smoking 13–21 weeks before pregnancy was similar to that among women who never smoked. This difference is likely related to whether cessation of smoking occurred after pregnancy recognition rather than well before pregnancy.

Studies based on the measurements of biomarkers of smoking such as plasma or salivary cotinine demonstrate diverse findings as well [41, 42]. A prospective pregnancy cohort study defined smoking status according to plasma cotinine, and found that previous and passive smokers compared to non-smokers were almost six-fold more likely to exhibit preeclampsia [41]. However, women who smoked during their current pregnancy had almost the same risk of preeclampsia as non-smokers. Another study did not show significant differences in preeclampsia rates using lower cutoffs of cotinine exposure [42].

Mainstream smoke contains multiple toxic chemicals in addition to nicotine and CO that are volatile, e.g., acetaldehyde, hydrogen cyanide, nitric acid, acetone, ammonia, hydrogen sulfide, hydrocarbons, nitrosamines, and carbonyl compound [17]. The smoke particulate phase also contains multiple toxicants such as carboxylic acids, phenols, terpenoids, paraffin waxes, tobacco-specific nitrosamines, polyaromatic hydrocarbons. Clearly smoking during pregnancy is not recommended in the context of reported detrimental concerns that include increases in perinatal mortality, abruptio placenta, intrauterine growth retardation [43, 44] and birth defects (e.g., oral clefts) [45].



## Limitations and strengths

All data regarding smoking status was self-reported, which may have contributed to misclassification, and thus would constitute measurement bias. If exposure misclassification did occur in our study it most likely was among smoking women who falsely reported that they stopped after pregnancy recognition, or among those who gave up smoking in the first trimester during pregnancy but subsequently resumed this practise. This type of misclassification would have decreased the risk of preeclampsia/eclampsia among those who gave up smoking while pregnant. However, we found that women who reported that they gave up smoking after pregnancy recognition had the same risk of preeclampsia/eclampsia as women who indicated they smoked before and during pregnancy. Our information about smoking behaviour was collected during the first antenatal visit. Räisänen et al [46] consider that gathering smoking status information during the first antenatal visit is more reliable than assessing it at the time of birth. Although we did not have data about the duration of tobacco smoke exposure, we did control for the number of cigarettes smoked daily during pregnancy.

The major strength of our study is that the data represent almost the total population of pregnant women in Murmansk County who delivered a singleton infant during 2006–2011 and had the first antenatal visit before 12 weeks of gestation. The registry data were collected in clinics and the number of births registered in the MCBR comprised 98.8% of the official number of births recorded by the Health Department in Murmansk County [27]. This allows generalizing of the results at the population level. Furthermore, quality controls in 2006–2007 suggested that the validity of data in MCBR is sufficient for epidemiological studies [27]. In contrast to previous studies [19–21], we had the ability to control for the influence of possible confounding factors such as social-demographic characteristics of pregnant women, body mass index at the first antenatal visit and excessive weight gain in pregnancy.

Our findings and perspective may provide clinicians with a better understanding of the necessity of promoting women to discontinue smoking during pregnancy, as well as rationale for the benefits of doing so early in the pregnancy. Clinicians might communicate that the risk of preeclampsia/eclampsia will not be affected by giving up smoking during pregnancy.

## Conclusions

In summary, we found that preeclampsia/eclampsia was associated with some maternal characteristics, such as maternal age, education, marital status, parity, excessive weight gain, and body mass index at the first antenatal visit. Our study demonstrates that maternal smoking was inversely associated with preeclampsia/eclampsia. Moreover, increased number of daily smoked cigarettes during pregnancy decreased odds of preeclampsia/eclampsia. Interesting, if women quit smoking during pregnancy they had the same risk of preeclampsia/eclampsia as those who smoked while pregnant. Even though our findings imply that giving up smoking does not alter the reduced risk of preeclampsia/eclampsia, it would most likely mitigate known smoking-related risks to the mother and the unborn. This advice might be shared during the first antenatal visit.

## Acknowledgments

We thank the office staff at the Murmansk County Birth Registry for their help in accessing the data.

## Author Contributions

**Conceptualization:** OAKh.

**Formal analysis:** OAKh AMG.

**Methodology:** OAKh AMG AK.

**Project administration:** JØO.

**Supervision:** JØO.

**Validation:** OAKh AMG AK EN.

**Writing – original draft:** OAKh EN.

**Writing – review & editing:** OAKh AMG.

## References

- Chappell LC, Enye S, Seed P, Briley AL, Poston L, Shennan AH: Adverse perinatal outcomes and risk factors for preeclampsia in women with chronic hypertension: a prospective study. *Hypertension* 2008, 51(4):1002–1009. <https://doi.org/10.1161/HYPERTENSIONAHA.107.107565> PMID: 18259010
- Villar J, Say L, Gulmezoglu AM, Merialdi M, Lindheimer MD, Betran AP et al. Eclampsia and preeclampsia: a worldwide health problem for 2000 years. In Critchley H, MacLean A, Poston L, Walker J, eds. *Preeclampsia*. London (UK) RCOG Press 2003: 189–207
- WHO recommendations for Prevention and treatment of pre-eclampsia and eclampsia. 2011.
- Tranquilli AL, Dekker G, Magee L, Roberts J, Sibai BM, Steyn W et al: The classification, diagnosis and management of the hypertensive disorders of pregnancy: A revised statement from the ISSHP. *Pregnancy Hypertens* 2014, 4(2):97–104. <https://doi.org/10.1016/j.preghy.2014.02.001> PMID: 26104417
- RCOG. Guideline No. 10(A). The Management of Severe Pre- eclampsia/Eclampsia 2010.
- Vest AR, Cho LS: Hypertension in pregnancy. *Curr Atheroscler Rep* 2014, 16(3):395. <https://doi.org/10.1007/s11883-013-0395-8> PMID: 24477794
- Abalos E, Cuesta C, Grosso AL, Chou D, Say L: Global and regional estimates of preeclampsia and eclampsia: a systematic review. *Eur J Obstet Gynecol Reprod Biol* 2013, 170(1):1–7. <https://doi.org/10.1016/j.ejogrb.2013.05.005> PMID: 23746796
- Federal clinical guidelines. Hypertensive disorders in pregnancy, childbirth and the postpartum period. Preeclampsia. Eclampsia (in Russian). 2013, 85p. Available from: <http://www.ncagip.ru/upload/obrazovanie/4.pdf> Accessed 3 May 2017
- Frolova OG, Pavlovich SV, Grebennik TK: Statistics of preeclampsia and eclampsia with contemporary parameters of registration of births (in Russian). *Bulletin PFUR* 2014, 2:70–74.
- Main indicators of mother and child health, activities of childhood service and obstetrics in RF. The Ministry of Health and Social Development Russian Federation. Moscow, 2011. Available from: [http://www.mednet.ru/images/stories/files/statistika/for\\_miac/materinstvo\\_i\\_detstvo\\_2010.pdf](http://www.mednet.ru/images/stories/files/statistika/for_miac/materinstvo_i_detstvo_2010.pdf). Accessed 2 Aug 2016.
- Kharkova OA, Krettek A, Grjibovski AM, Nieboer E, Odland JO: Prevalence of smoking before and during pregnancy and changes in this habit during pregnancy in Northwest Russia: a Murmansk county birth registry study. *Reprod Health* 2016, 13:18. <https://doi.org/10.1186/s12978-016-0144-x> PMID: 26952100
- Tikkanen M, Nuutila M, Hiilesmaa V, Paavonen J, Ylikorkala O: Prepregnancy risk factors for placental abruption. *Acta Obstet Gynecol Scand* 2006, 85:40–44. PMID: 16521678
- Oyelese Y, Smulian JC: Placenta Previa, placenta accreta, and vasa Previa. *Obstet Gynecol* 2006, 107:927–941. <https://doi.org/10.1097/01.AOG.0000207559.15715.98> PMID: 16582134
- Jaddoe VW, Troe EJ, Hofman A, Mackenbach JP, Moll HA, Steegers EA et al: Active and passive maternal smoking during pregnancy and the risks of low birthweight and preterm birth: the Generation R Study. *Paediatric and perinatal epidemiology* 2008, 22(2):162–171. <https://doi.org/10.1111/j.1365-3016.2007.00916.x> PMID: 18298691
- Stojanovic M, Bojanic V, Musovic D, Milosevic Z, Stojanovic D, Visujic A et al: Maternal smoking during pregnancy and socioeconomic factors as predictors of low birth weight in term pregnancies in Nis. *Vojnosanitetski pregled Military-medical and pharmaceutical review* 2010, 67(2):145–150. PMID: 20337097
- Aagaard-Tillery KM, Porter TF, Lane RH, Varner MW, Lacoursiere DY: In utero tobacco exposure is associated with modified effects of maternal factors on fetal growth. *Am J Obstet Gynecol* 2008, 198:661–666.

17. U.S. Department of Health and Human Services. How tobacco smoke causes disease: the biology and behavioral basis for smoking-attributable disease: a report of the surgeon general. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health;2010. 706 p. Available from: [http://www.ncbi.nlm.nih.gov/books/NBK53017/pdf/Bookshelf\\_NBK53017.pdf](http://www.ncbi.nlm.nih.gov/books/NBK53017/pdf/Bookshelf_NBK53017.pdf). Accessed 5 Sep 2016
18. Campos Tda S, Richter KP, Cupertino AP, Galil AG, Banhato EF, Colugnati FA et al. Cigarette smoking among patients with chronic diseases. *Int J Cardiol* 2014, 174(3):808–810. <https://doi.org/10.1016/j.ijcard.2014.04.150> PMID: 24801077
19. Engel SM, Janevic TM, Stein CR, Savitz DA: Maternal smoking, preeclampsia, and infant health outcomes in New York City, 1995–2003. *Am J Epidemiol* 2009, 169(1):33–40. <https://doi.org/10.1093/aje/kwn293> PMID: 19001134
20. Conde-Agudelo A, Althabe F, Belizan JM, Kafury-Goeta AC: Cigarette smoking during pregnancy and risk of preeclampsia: a systematic review. *Am J Obstet Gynecol* 1999, 181(4):1026–1035. PMID: 10521771
21. Wikstrom AK, Stephansson O, Cnattingius S: Tobacco use during pregnancy and preeclampsia risk: effects of cigarette smoking and snuff. *Hypertension* 2010, 55(5):1254–1259. <https://doi.org/10.1161/HYPERTENSIONAHA.109.147082> PMID: 20231527
22. Pipkin FB, Genetics of Preeclampsia C: Smoking in moderate/severe preeclampsia worsens pregnancy outcome, but smoking cessation limits the damage. *Hypertension* 2008, 51(4):1042–1046. <https://doi.org/10.1161/HYPERTENSIONAHA.107.106559> PMID: 18259022
23. Sibai BM, Gordon T, Thom E, Caritis SN, Klebanoff M, McNellis D et al: Risk factors for preeclampsia in healthy nulliparous women: a prospective multicenter study. The National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. *Am J Obstet Gynecol* 1995, 172(2 Pt 1):642–648.
24. Perni UC, Wikstrom AK, Cnattingius S, Villamor E: Interpregnancy change in smoking habits and risk of preeclampsia: a population-based study. *Am J Hypertens* 2012, 25(3):372–378. <https://doi.org/10.1038/ajh.2011.225> PMID: 22113171
25. Xiong X, Zhang J, Fraser WD: Quitting smoking during early versus late pregnancy: the risk of preeclampsia and adverse birth outcomes. *J Obstet Gynaecol Can* 2009, 31(8):702–707. PMID: 19772701
26. England LJ, Levine RJ, Qian C, Morris CD, Sibai BM, Catalano PM et al: Smoking before pregnancy and risk of gestational hypertension and preeclampsia. *Am J Obstet Gynecol* 2002, 186(5):1035–1040. PMID: 12015533
27. Anda EE, Nieboer E, Voitov AV, Kovalenko AA, Lapina YM, Voitova EA et al: Implementation, quality control and selected pregnancy outcomes of the Murmansk County Birth Registry in Russia. *Int J Circumpolar Health* 2008, 67:318–334. PMID: 19024802
28. [ICD10Data.com](http://ICD10Data.com) (Available from: [www.icd10data.com](http://www.icd10data.com)). Accessed 12 Sep 2016.
29. Bhide A, Rana R, Dhavilkar M, Amodio-Hernandez M, Deshpande D, Caric V: The value of the urinary protein:creatinine ratio for the detection of significant proteinuria in women with suspected preeclampsia. *Acta Obstet Gynecol Scand* 2015, 94(5):542–546. <https://doi.org/10.1111/aogs.12624> PMID: 25737188
30. Roberts JM, Cooper DW: Pathogenesis and genetics of pre-eclampsia. *Lancet* 2001, 357(9249):53–56. PMID: 11197372
31. Martin CL, Hall MH, Campbell DM: The effect of smoking on pre-eclampsia in twin pregnancy. *BJOG* 2000, 107(6):745–749. PMID: 10847230
32. Lindqvist PG, Marsal K: Moderate smoking during pregnancy is associated with a reduced risk of preeclampsia. *Acta Obstet Gynecol Scand* 1999, 78(8):693–697. PMID: 10468061
33. Yang Q, Wen SW, Smith GN, Chen Y, Krewski D, Chen XK et al: Maternal cigarette smoking and the risk of pregnancy-induced hypertension and eclampsia. *Int J Epidemiol* 2006, 35(2):288–293. <https://doi.org/10.1093/ije/dyi247> PMID: 16303811
34. Bainbridge SA, Sidle EH, Smith GN: Direct placental effects of cigarette smoke protect women from pre-eclampsia: the specific roles of carbon monoxide and antioxidant systems in the placenta. *Med Hypotheses* 2005, 64(1):17–27. <https://doi.org/10.1016/j.mehy.2004.06.019> PMID: 15533604
35. Venditti CC, Casselman R, Young I, Karumanchi SA, Smith GN: Carbon monoxide prevents hypertension and proteinuria in an adenovirus sFlt-1 preeclampsia-like mouse model. *PLoS One* 2014, 9(9): e106502. <https://doi.org/10.1371/journal.pone.0106502> PMID: 25202912
36. Zhai D, Guo Y, Smith G, Krewski D, Walker M, Wen SW: Maternal exposure to moderate ambient carbon monoxide is associated with decreased risk of preeclampsia. *Am J Obstet Gynecol* 2012, 207(1):57 e51–59.

37. Naljayan MV, Karumanchi SA: New developments in the pathogenesis of preeclampsia. *Adv Chronic Kidney Dis* 2013, 20(3):265–270. <https://doi.org/10.1053/j.ackd.2013.02.003> PMID: 23928392
38. Karumanchi SA, Levine RJ: How does smoking reduce the risk of preeclampsia? *Hypertension* 2010, 55(5):1100–1101. <https://doi.org/10.1161/HYPERTENSIONAHA.109.148973> PMID: 20231524
39. Loboda A, Jozkowicz A, Dulak J: HO-1/CO system in tumor growth, angiogenesis and metabolism—Targeting HO-1 as an anti-tumor therapy. *Vascul Pharmacol* 2015, 74:11–22. <https://doi.org/10.1016/j.vph.2015.09.004> PMID: 26392237
40. Loboda A, Jozkowicz A, Dulak J: Carbon monoxide: pro- or anti-angiogenic agent? Comment on Ahmad et al. (*Thromb Haemost* 2015; 113: 329–337). *Thromb Haemost* 2015, 114(2):432–433. <https://doi.org/10.1160/TH15-01-0082> PMID: 25904352
41. Luo ZC, Julien P, Wei SQ, Audibert F, Smith GN, Fraser WD et al: Plasma cotinine indicates an increased risk of preeclampsia in previous and passive smokers. *Am J Obstet Gynecol* 2014, 210(3):232 e231–235.
42. Janakiraman V, Gantz M, Maynard S, El-Mohandes A: Association of cotinine levels and preeclampsia among African-American women. *Nicotine Tob Res* 2009, 11(6):679–684. <https://doi.org/10.1093/ntr/ntp049> PMID: 19395687
43. Cnattingius S, Mills JL, Yuen J, Eriksson O, Salonen H: The paradoxical effect of smoking in preeclamptic pregnancies: smoking reduces the incidence but increases the rates of perinatal mortality, abruptio placentae, and intrauterine growth restriction. *Am J Obstet Gynecol* 1997, 177(1):156–161. PMID: 9240600
44. Baba S, Wikstrom AK, Stephansson O, Cnattingius S: Changes in snuff and smoking habits in Swedish pregnant women and risk for small for gestational age births. *BJOG* 2013, 120(4):456–462. <https://doi.org/10.1111/1471-0528.12067> PMID: 23190416
45. Meyer KA, Williams P, Hernandez-Diaz S, Cnattingius S: Smoking and the risk of oral clefts: exploring the impact of study designs. *Epidemiology* 2004, 15(6):671–678. PMID: 15475715
46. Raisanen S, Kramer MR, Gissler M, Saari J, Hakulinen-Viitanen T, Heinonen S: Smoking during pregnancy was up to 70% more common in the most deprived municipalities—a multilevel analysis of all singleton births during 2005–2010 in Finland. *Prev Med* 2014, 67:6–11. <https://doi.org/10.1016/j.ypmed.2014.06.026> PMID: 24983887