

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(1*S*,2*R*,3*S*,6*S*,7*R*)-3,7,11,11-Tetramethyl-6,7-epoxybicyclo[5.4.0]undecane-2-olMohamed Loubidi,^{a*} Ahmed Benharref,^a Lahcen El Ammari,^b Mohamed Saadi^b and Moha Berraho^a^aLaboratoire de Chimie des Substances Naturelles, "Unité Associé au CNRST (URAC16)", Faculté des Sciences Semlalia, BP 2390 Bd My Abdellah, 40000 Marrakech, Morocco, and ^bLaboratoire de Chimie du Solide Appliquée, Faculté des Sciences, Université Mohammed V-Agdal, BP 1014, Avenue Ibn Battouta, Rabat, Morocco

Correspondence e-mail: loubidim@gmail.com


Received 14 April 2014; accepted 16 April 2014

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.045; wR factor = 0.131; data-to-parameter ratio = 18.7.

The title compound, $\text{C}_{15}\text{H}_{26}\text{O}_2$, was synthesized from β -himachalene (3,5,5,9-tetramethyl-2,4a,5,6,7,8-hexahydro-1*H*-benzocycloheptene), which was isolated from the Atlas cedar (*cedrus atlantica*). The molecule is built up from a seven-membered ring to which a six- and a three-membered ring are fused. The seven- and six-membered rings each have a twist-boat conformation. In the crystal, $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into zigzag chains running along the b -axis direction.

Related literature

For background to β -himachalene, see: El Haib *et al.* (2011). For the reactivity of this sesquiterpene and its derivatives, see: El Jamili *et al.* (2002); Benharref *et al.* (2013); Ourhriss *et al.* (2013). For their potential antifungal activity against the phytopathogen *Botrytis cinerea*, see: Daoubi *et al.* (2004). For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{26}\text{O}_2$
 $M_r = 238.36$
 Monoclinic, $P2_1$
 $a = 5.9617$ (10) Å
 $b = 12.068$ (2) Å
 $c = 9.5909$ (15) Å
 $\beta = 95.789$ (8)°
 $V = 686.48$ (19) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.07$ mm⁻¹
 $T = 298$ K
 $0.38 \times 0.11 \times 0.10$ mm

Data collection

Bruker X8 APEX diffractometer
 11541 measured reflections
 3036 independent reflections
 2698 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.131$
 $S = 1.06$
 3036 reflections
 162 parameters
 1 restraint
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.34$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.16$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H11}\cdots\text{O2}^i$	0.72 (3)	2.11 (3)	2.820 (2)	171 (3)

Symmetry code: (i) $-x, y + \frac{1}{2}, -z$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: PLATON (Spek, 2009) and publCIF (Westrip, 2010).

The authors thank the Unit of Support for Technical and Scientific Research (UATRS, CNRST) for the X-ray measurements

Supporting information for this paper is available from the IUCr electronic archives (Reference: BT6976).

References

- Benharref, A., Ourhriss, N., El Ammari, L., Saadi, M. & Berraho, M. (2013). *Acta Cryst.* **E69**, o933–o934.
 Bruker (2009). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Daoubi, M., Duran-Patron, R., Hmamouchi, M., Hernandez-Galan, R., Benharref, A. & Isidro, G. C. (2004). *Pest Manag. Sci.* **60**, 927–932.
 El Haib, A., Benharref, A., Parrès-Maynadié, S., Manoury, E., Urrutigoity, M. & Gouygou, M. (2011). *Tetrahedron Asymmetry*, **22**, 101–108.
 El Jamili, H., Auhmani, A., Dakir, M., Lassaba, E., Benharref, A., Pierrot, M., Chiaroni, A. & Riche, C. (2002). *Tetrahedron Lett.* **43**, 6645–6648.
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Ourhriss, N., Benharref, A., Saadi, M., El Ammari, L. & Berraho, M. (2013). *Acta Cryst.* **E69**, o275.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2014). E70, o595 [doi:10.1107/S1600536814008642]

(1*S*,2*R*,3*S*,6*S*,7*R*)-3,7,11,11-Tetramethyl-6,7-epoxybicyclo[5.4.0]undecane-2-ol

Mohamed Loubidi, Ahmed Benharref, Lahcen El Ammari, Mohamed Saadi and Moha Berraho

1. Comment


This work is a part of our ongoing program concerning the valorization of the most abundant essential oils in Morocco, such as *Cedrus atlantica*. This oil is made up mainly (75%) of bicyclic sesquiterpene hydrocarbons, among which is found the compound β -himachalene (El Haib *et al.*, 2011). The reactivity of this sesquiterpene and its derivatives has been studied extensively by our team in order to prepare new products having biological properties (El Jamili *et al.*, 2002; Benharref *et al.*, 2013; Ourhriss *et al.*, 2013). Indeed, these compounds were tested, using the food poisoning technique, for their potential antifungal activity against phytopathogen *Botrytis cinerea* (Daoubi *et al.*, 2004). The molecule of the title compound (Fig.1) is built up from two fused six- and seven-membered rings, both rings have a twist-boat conformation as indicated by the total puckering amplitude $QT = 0.7821(19)$ Å and spherical polar angle $\theta = 92.53(15)^\circ$ with $\varphi = 272.91(14)^\circ$, for the six-membered ring, and $QT = 1.1479(21)$, $\theta = 87.76(10)$, $\varphi_2 = -148.25(21)$, $\varphi_3 = 27.36(3)$ for the seven-membered ring (Cremer & Pople, 1975). In the crystal, O—H \cdots O hydrogen bonds links the molecules into zigzag chains running along the *b* axis (Table 1, Fig. 1).

2. Experimental

Diborane is prepared by addition at 0 °C of 2.5 g (17 mmol) of boron trifluoride etherate in 0.5 g (12.6 mmol) of sodium borohydride in 30 ml of diglyme. Diborane formed is driven by a stream of dry nitrogen in 2 g (10 mmol) of β -himachalene dissolved in 20 ml of tetrahydrofuran at 273K. This takes about 4 h. 2 ml of sodium hydroxide 3 N is then added carefully between 263K and 273K in 15 minutes and then 2 ml of 30% hydrogen peroxide in the vicinity of 298K. The reaction mixture was then extracted with diethyl ether, the organic phase is washed to neutrality and the solvent was evaporated under vacuum. The residue obtained is chromatographed on a column of silica gel with hexane-ethyl acetate (95:5), which allowed the isolation of pure (1*S*, 2*R*, 3*S*, 6*S*, 7*R*)-6,7-epoxy-3,7,11,11- tetramethylbicyclo[5.4.0]undecane-2-ol with a yield of 25% (600 mg 2.5 mmol). The title compound was recrystallized from its cyclohexane solution.

3. Refinement

Except H11, which was freely refined, all H atoms were fixed geometrically and treated as riding with C—H = 0.96 Å (methyl), 0.97 Å (methylene), 0.98 Å (methine) with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{methylene, methine})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{methyl})$. The methyl groups were allowed to rotate, but not to tip. In the absence of significant anomalous scattering, the absolute configuration could not be reliably determined and any references to the Flack parameter were removed. The absolute configuration of the chiral centres was arbitrarily set.

**Figure 1**

Molecular structure of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.




Figure 2

Packing view showing the O–H...O hydrogen bonds as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity.

(1*S*,2*R*,3*S*,6*S*,7*R*)-3,7,11,11-Tetramethyl-6,7-epoxybicyclo[5.4.0]undecane-2-ol

Crystal data

$C_{15}H_{26}O_2$
 $M_r = 238.36$
 Monoclinic, $P2_1$
 Hall symbol: P 2yb
 $a = 5.9617(10) \text{ \AA}$
 $b = 12.068(2) \text{ \AA}$
 $c = 9.5909(15) \text{ \AA}$
 $\beta = 95.789(8)^\circ$
 $V = 686.48(19) \text{ \AA}^3$
 $Z = 2$

$F(000) = 264$
 $D_x = 1.153 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 3036 reflections
 $\theta = 2.7\text{--}27.1^\circ$
 $\mu = 0.07 \text{ mm}^{-1}$
 $T = 298 \text{ K}$
 Needle, colourless
 $0.38 \times 0.11 \times 0.10 \text{ mm}$

Data collection

Bruker X8 APEX
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator

φ and ω scans
 11541 measured reflections
 3036 independent reflections
 2698 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$
 $\theta_{\text{max}} = 27.1^\circ$, $\theta_{\text{min}} = 2.7^\circ$
 $h = -7 \rightarrow 7$

$k = -15 \rightarrow 15$
 $l = -11 \rightarrow 12$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.131$
 $S = 1.06$
 3036 reflections
 162 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0862P)^2 + 0.0329P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.16 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C2	0.0588 (3)	0.50820 (14)	0.02419 (18)	0.0314 (4)
H2	0.1734	0.5611	0.0000	0.038*
C4	0.2591 (3)	0.37310 (19)	-0.11371 (19)	0.0453 (5)
H4A	0.3279	0.4210	-0.1786	0.054*
H4B	0.2399	0.3005	-0.1567	0.054*
C9	0.3831 (5)	0.3909 (3)	0.4739 (2)	0.0657 (7)
H9A	0.5201	0.3606	0.5226	0.079*
H9B	0.2942	0.4216	0.5439	0.079*
H11	-0.153 (4)	0.608 (2)	-0.012 (3)	0.051 (8)*
C1	0.1382 (3)	0.45493 (13)	0.16735 (17)	0.0289 (3)
H1	0.0019	0.4272	0.2047	0.035*
C3	0.0279 (3)	0.41938 (15)	-0.09006 (19)	0.0372 (4)
H3	-0.0630	0.3592	-0.0565	0.045*
C5	0.4180 (3)	0.36301 (17)	0.02114 (19)	0.0395 (4)
H5A	0.5132	0.2982	0.0162	0.047*
H5B	0.5143	0.4279	0.0323	0.047*
C6	0.2821 (3)	0.35300 (14)	0.14433 (17)	0.0304 (3)
C7	0.3361 (3)	0.27224 (16)	0.2586 (2)	0.0412 (4)
C8	0.2506 (4)	0.2968 (2)	0.3985 (2)	0.0545 (6)
H8A	0.2637	0.2308	0.4564	0.065*
H8B	0.0925	0.3170	0.3840	0.065*
C10	0.4472 (4)	0.4851 (2)	0.3774 (2)	0.0531 (6)

H10A	0.5572	0.4567	0.3186	0.064*
H10B	0.5204	0.5431	0.4355	0.064*
C11	0.2506 (3)	0.53852 (15)	0.28073 (19)	0.0372 (4)
C12	0.5456 (4)	0.2016 (2)	0.2643 (3)	0.0612 (6)
H12A	0.5204	0.1334	0.3119	0.092*
H12B	0.6694	0.2405	0.3140	0.092*
H12C	0.5803	0.1859	0.1708	0.092*
C13	-0.0927 (4)	0.4641 (2)	-0.2266 (2)	0.0575 (6)
H13A	-0.0141	0.5281	-0.2558	0.086*
H13B	-0.2443	0.4842	-0.2120	0.086*
H13C	-0.0954	0.4080	-0.2977	0.086*
C14	0.3584 (4)	0.64005 (19)	0.2140 (3)	0.0574 (6)
H14A	0.2429	0.6824	0.1611	0.086*
H14B	0.4661	0.6150	0.1529	0.086*
H14C	0.4329	0.6856	0.2865	0.086*
C15	0.0719 (4)	0.5824 (2)	0.3696 (3)	0.0636 (7)
H15A	0.1416	0.6300	0.4416	0.095*
H15B	0.0005	0.5214	0.4118	0.095*
H15C	-0.0391	0.6235	0.3113	0.095*
O1	-0.1435 (2)	0.56718 (13)	0.04280 (17)	0.0465 (4)
O2	0.1605 (2)	0.24617 (10)	0.14695 (15)	0.0414 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C2	0.0338 (8)	0.0274 (8)	0.0338 (9)	-0.0005 (7)	0.0068 (6)	0.0023 (7)
C4	0.0540 (11)	0.0512 (12)	0.0321 (9)	0.0069 (9)	0.0119 (8)	-0.0066 (8)
C9	0.0770 (16)	0.0825 (18)	0.0347 (11)	0.0019 (14)	-0.0090 (10)	0.0007 (12)
C1	0.0297 (7)	0.0277 (7)	0.0300 (8)	-0.0025 (6)	0.0062 (6)	-0.0001 (6)
C3	0.0429 (10)	0.0379 (9)	0.0304 (9)	-0.0010 (7)	0.0020 (7)	-0.0009 (7)
C5	0.0379 (9)	0.0398 (9)	0.0419 (10)	0.0050 (8)	0.0100 (7)	-0.0051 (8)
C6	0.0321 (7)	0.0265 (7)	0.0317 (8)	-0.0036 (6)	-0.0003 (6)	-0.0027 (7)
C7	0.0462 (10)	0.0351 (10)	0.0403 (10)	-0.0033 (8)	-0.0060 (8)	0.0039 (8)
C8	0.0599 (13)	0.0623 (14)	0.0407 (11)	-0.0036 (11)	0.0022 (9)	0.0197 (10)
C10	0.0487 (11)	0.0572 (14)	0.0514 (13)	-0.0088 (9)	-0.0045 (9)	-0.0138 (10)
C11	0.0398 (9)	0.0374 (9)	0.0347 (9)	-0.0056 (7)	0.0052 (7)	-0.0098 (7)
C12	0.0595 (14)	0.0520 (13)	0.0681 (15)	0.0150 (11)	-0.0130 (11)	0.0080 (11)
C13	0.0698 (14)	0.0646 (14)	0.0352 (11)	0.0089 (12)	-0.0088 (9)	-0.0009 (10)
C14	0.0689 (14)	0.0398 (11)	0.0640 (14)	-0.0125 (11)	0.0086 (11)	-0.0075 (10)
C15	0.0596 (13)	0.0766 (17)	0.0561 (14)	0.0007 (12)	0.0141 (11)	-0.0262 (13)
O1	0.0504 (8)	0.0417 (8)	0.0490 (9)	0.0166 (6)	0.0121 (6)	0.0105 (7)
O2	0.0490 (8)	0.0270 (6)	0.0459 (7)	-0.0068 (5)	-0.0067 (6)	0.0019 (6)

Geometric parameters (\AA , $^\circ$)

C2—O1	1.427 (2)	C7—C12	1.508 (3)
C2—C3	1.531 (2)	C7—C8	1.512 (3)
C2—C1	1.546 (2)	C8—H8A	0.9700
C2—H2	0.9800	C8—H8B	0.9700
C4—C3	1.526 (3)	C10—C11	1.558 (3)

C4—C5	1.529 (3)	C10—H10A	0.9700
C4—H4A	0.9700	C10—H10B	0.9700
C4—H4B	0.9700	C11—C15	1.525 (3)
C9—C8	1.523 (4)	C11—C14	1.552 (3)
C9—C10	1.539 (4)	C12—H12A	0.9600
C9—H9A	0.9700	C12—H12B	0.9600
C9—H9B	0.9700	C12—H12C	0.9600
C1—C6	1.528 (2)	C13—H13A	0.9600
C1—C11	1.582 (2)	C13—H13B	0.9600
C1—H1	0.9800	C13—H13C	0.9600
C3—C13	1.527 (3)	C14—H14A	0.9600
C3—H3	0.9800	C14—H14B	0.9600
C5—C6	1.503 (2)	C14—H14C	0.9600
C5—H5A	0.9700	C15—H15A	0.9600
C5—H5B	0.9700	C15—H15B	0.9600
C6—C7	1.478 (3)	C15—H15C	0.9600
C6—O2	1.481 (2)	O1—H11	0.72 (3)
C7—O2	1.455 (2)		
O1—C2—C3	113.36 (15)	C6—C7—C8	117.44 (17)
O1—C2—C1	106.45 (14)	C12—C7—C8	115.52 (19)
C3—C2—C1	110.40 (13)	C7—C8—C9	111.26 (19)
O1—C2—H2	108.8	C7—C8—H8A	109.4
C3—C2—H2	108.8	C9—C8—H8A	109.4
C1—C2—H2	108.8	C7—C8—H8B	109.4
C3—C4—C5	113.30 (14)	C9—C8—H8B	109.4
C3—C4—H4A	108.9	H8A—C8—H8B	108.0
C5—C4—H4A	108.9	C9—C10—C11	116.50 (19)
C3—C4—H4B	108.9	C9—C10—H10A	108.2
C5—C4—H4B	108.9	C11—C10—H10A	108.2
H4A—C4—H4B	107.7	C9—C10—H10B	108.2
C8—C9—C10	114.41 (18)	C11—C10—H10B	108.2
C8—C9—H9A	108.7	H10A—C10—H10B	107.3
C10—C9—H9A	108.7	C15—C11—C14	107.35 (18)
C8—C9—H9B	108.7	C15—C11—C10	109.69 (18)
C10—C9—H9B	108.7	C14—C11—C10	104.72 (17)
H9A—C9—H9B	107.6	C15—C11—C1	109.53 (15)
C6—C1—C2	109.41 (13)	C14—C11—C1	112.59 (15)
C6—C1—C11	114.05 (14)	C10—C11—C1	112.74 (15)
C2—C1—C11	114.57 (14)	C7—C12—H12A	109.5
C6—C1—H1	106.0	C7—C12—H12B	109.5
C2—C1—H1	106.0	H12A—C12—H12B	109.5
C11—C1—H1	106.0	C7—C12—H12C	109.5
C4—C3—C13	110.86 (16)	H12A—C12—H12C	109.5
C4—C3—C2	108.51 (15)	H12B—C12—H12C	109.5
C13—C3—C2	112.26 (17)	C3—C13—H13A	109.5
C4—C3—H3	108.4	C3—C13—H13B	109.5
C13—C3—H3	108.4	H13A—C13—H13B	109.5
C2—C3—H3	108.4	C3—C13—H13C	109.5

C6—C5—C4	109.51 (14)	H13A—C13—H13C	109.5
C6—C5—H5A	109.8	H13B—C13—H13C	109.5
C4—C5—H5A	109.8	C11—C14—H14A	109.5
C6—C5—H5B	109.8	C11—C14—H14B	109.5
C4—C5—H5B	109.8	H14A—C14—H14B	109.5
H5A—C5—H5B	108.2	C11—C14—H14C	109.5
C7—C6—O2	58.92 (11)	H14A—C14—H14C	109.5
C7—C6—C5	122.77 (16)	H14B—C14—H14C	109.5
O2—C6—C5	112.77 (14)	C11—C15—H15A	109.5
C7—C6—C1	120.48 (15)	C11—C15—H15B	109.5
O2—C6—C1	114.54 (12)	H15A—C15—H15B	109.5
C5—C6—C1	113.81 (14)	C11—C15—H15C	109.5
O2—C7—C6	60.64 (11)	H15A—C15—H15C	109.5
O2—C7—C12	115.87 (18)	H15B—C15—H15C	109.5
C6—C7—C12	121.17 (19)	C2—O1—H11	105 (2)
O2—C7—C8	114.42 (17)	C7—O2—C6	60.45 (11)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O1—H11...O2 ⁱ	0.72 (3)	2.11 (3)	2.820 (2)	171 (3)

Symmetry code: (i) $-x, y+1/2, -z$.