



Draft Genome Sequences of *Dickeya* Species Associated with Soft Rot Diseases in Pineapple (*Ananas comosus*) and Banana (*Musa* spp.)

 Anthony J. Young,^a Henry W. G. Birt,^b Niloofer Vaghefi,^c Robert D. Hoelzle,^b Paul G. Dennis^b

^aSchool of Agriculture and Food Sciences, The University of Queensland, Gatton, Queensland, Australia

^bSchool of Earth and Environmental Sciences, The University of Queensland, Brisbane, Queensland, Australia

^cSchool of Agriculture and Food, Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Parkville, Victoria, Australia

ABSTRACT *Dickeya* species cause soft rots on many commercial crops. Here, we present the draft genomes of *Dickeya oryzae* (BRIP 64262) and *Dickeya zae* (BRIP 64263) isolates causing soft rot on banana (*Musa* spp.) and pineapple (*Ananas comosus*) plants, respectively. This expands the range of available genomes from plant-pathogenic *Dickeya* species.

D *ickeya* species cause soft rots on a wide range of plants, including potato, maize, rice, pineapple, and banana (1–5). While draft and complete genome sequences are available for *Dickeya zae* and *Dickeya oryzae* strains from North America, Europe, and Asia (6–9), none has been reported from Australia. Here, we present draft genome sequences for two isolates, *D. oryzae* BRIP 64262, sourced from banana (*Musa* spp.), and *D. zae* BRIP 64263, sourced from pineapple (*Ananas comosus*).

Banana samples exhibiting corm rot were collected from a field at South Johnstone, Queensland, Australia (17°36'19.0"S, 145°59'54.6"E). Corms were split to reveal infected tissue, and streaks onto Kelman's medium were made from the margin of infection. Bacteria were isolated in the same way from the necrotic margins of symptomatic pineapple plants (1). Plates were incubated aerobically at 28°C for 3 days until individual colonies appeared, which were then purified by three successive streaks from single colonies. DNA was extracted using the Isolate II plant DNA extraction kit (Bioline). Briefly, ~250 mg of biomass was transferred from solid medium into a 2-mL screw-cap tube containing extraction buffer and a 4-mm-diameter steel ball and then was processed in a FastPrep bead beater for 40 s at a setting of 4.5 m s⁻¹ before purification according to the manufacturer's instructions. DNA libraries were prepared using the Nextera XT DNA library preparation kit (Illumina, Inc., San Diego, CA, USA) and sequenced on the MiSeq platform with 150-bp paired-end reads (Illumina, Inc.), generating 296 Mbp and 333 Mbp of data from 2.00 million (BRIP 64262) and 2.25 million (BRIP 64263) read pairs, respectively.

Raw reads were trimmed and quality filtered with Trimmomatic v0.36 (10) using the following settings: ILLUMINACLIP:NexteraPE-PE.fa:2:30:10, TRAILING:10, SLIDINGWINDOW:4:15, and MINLEN:75. Paired-end trimmed reads were *de novo* assembled using SPAdes v3.14.1 (11) in isolate mode. The assembled contigs were checked for completeness and contamination using CheckM v1.1.2 (12) and assigned taxonomy using GTDB-Tk v1.3.0 and the Genome Taxonomy Database (GTDB) 07-RS207 release (13). The NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v6.1 was used to identify genes (14). Average nucleotide identity (ANI) was assessed with FastANI v1.3 (15). Default parameters were used unless otherwise specified.

The BRIP 64262 and BRIP 64263 draft genomes were of similar sizes (4.7 and 4.8 Mbp, respectively), compared to neighboring species, according to phylogenetic

Editor Leighton Pritchard, SIPBS, University of Strathclyde

Copyright © 2022 Young et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Anthony J. Young, a.young@uq.edu.au, or Paul G. Dennis, p.dennis@uq.edu.au.

The authors declare no conflict of interest.

Received 14 July 2022

Accepted 16 September 2022

Published 5 October 2022

analysis, and were alike in composition, with 94.7% ANI. BRIP 64262 had 97.2% ANI to the *D. oryzae* type strain (GenBank assembly accession number [GCF_009372235.1](https://doi.org/10.1093/genbank/GCF_009372235.1)), while BRIP 64263 had 96.3% ANI to the *D. zeae* type strain (GenBank assembly accession number [GCF_002887555.1](https://doi.org/10.1093/genbank/GCF_002887555.1)). The BRIP 64262 and BRIP 64263 draft genomes were assembled into 191 and 173 contigs, with N_{50} values of 402 and 194 kbp, respectively, and both had GC contents of 58%. According to CheckM, the draft genomes of BRIP 64262 and BRIP 64263 were both 99.97% complete, with 0.61% and 1.00% contamination, respectively. Both isolates had 3 copies of the 16S rRNA gene, while PGAP annotated 4,234 genes for BRIP 64262 and 4,293 genes for BRIP 64263, similar to comparable genomes of *D. zeae* (GenBank assembly accession number [GCF_002887555.1](https://doi.org/10.1093/genbank/GCF_002887555.1)) (4,216 genes) and *D. oryzae* (GenBank assembly accession number [GCF_009372235.1](https://doi.org/10.1093/genbank/GCF_009372235.1)) (4,031 genes).

Data availability. The data from this project have been deposited under NCBI BioProject accession number [PRJNA737524](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA737524). The raw sequencing data have been deposited in the Sequence Read Archive (SRA) under accession numbers [SRR14876326](https://www.ncbi.nlm.nih.gov/sra/SRR14876326) (BRIP 64262) and [SRR14876325](https://www.ncbi.nlm.nih.gov/sra/SRR14876325) (BRIP 64263), and the draft genome assemblies have been deposited in GenBank (BRIP 64262, BioSample accession number [SAMN19735995](https://www.ncbi.nlm.nih.gov/biosample/SAMN19735995); BRIP 64263, BioSample accession number [SAMN19735996](https://www.ncbi.nlm.nih.gov/biosample/SAMN19735996)). The versions described in this paper are the first versions (BioSample accession numbers [SAMN19735995](https://www.ncbi.nlm.nih.gov/biosample/SAMN19735995) and [SAMN19735996](https://www.ncbi.nlm.nih.gov/biosample/SAMN19735996)). The annotated genomes are available in GenBank under the following accession numbers: BRIP 64262, [JAMXSN000000000](https://www.ncbi.nlm.nih.gov/genbank/JAMXSN000000000); BRIP 64263, [JAMXSO000000000](https://www.ncbi.nlm.nih.gov/genbank/JAMXSO000000000).

ACKNOWLEDGMENTS

We would like to acknowledge Nandita Pathania, Andrew Manners and Ken Pegg for assistance with the cultures.

REFERENCES

- Young AJ, Pathania N, Manners A, Pegg K. 2022. Heart rot of Australian pineapples caused by *Dickeya zeae*. *Australas Plant Pathol* <https://doi.org/10.1007/s13313-022-00880-x>.
- Samson R, Legendre JB, Christen R, Fischer-Le Saux M, Achouak W, Gardan L. 2005. Transfer of *Pectobacterium chrysanthemi* (Burkholder et al., 1953) Brenner et al. 1973 and *Brenneria paradisiaca* to the genus *Dickeya* gen. nov. 60 as *Dickeya chrysanthemi* comb. nov. and *Dickeya paradisiaca* comb. nov. and delineation of four novel species. *Int J Syst Evol Microbiol* 55:1415–1427. <https://doi.org/10.1099/ijs.0.02791-0>.
- Sueno WSK, Marrero G, de Silva AS, Sether DM, Alvarez AM. 2014. Diversity of *Dickeya* strains collected from pineapple plants and irrigation water in Hawaii. *Plant Dis* 98:817–824. <https://doi.org/10.1094/PDIS-03-13-0219-RE>.
- Prokić A, Zlatković N, Kuzmanović N, Ivanović M, Gašić K, Pavlović Ž, Obradović A. 2020. Identification and characterization of *Dickeya zeae* strains associated with maize stalk soft-rot in northern Serbia. *Eur J Plant Pathol* 157:685–691. <https://doi.org/10.1007/s10658-020-02019-4>.
- Toth IK, Van Der Wolf JM, Saddler G, Lojkowska E, Hélias V, Pirhonen M, Tsror L, Elphinstone JG. 2011. *Dickeya* species: an emerging problem for potato production in Europe. *Plant Pathol* 60:385–399. <https://doi.org/10.1111/j.1365-3059.2011.02427.x>.
- Bertani I, da Silva DP, Abbruscato P, Piffanelli P, Venturi V. 2013. Draft genome sequence of the plant pathogen *Dickeya zeae* DZ2Q, isolated from rice in Italy. *Genome Announc* 1:e00905-13. <https://doi.org/10.1128/genomeA.00905-13>.
- Pritchard L, Humphris S, Saddler GS, Elphinstone JG, Pirhonen M, Toth IK. 2013. Draft genome sequences of 17 isolates of the plant pathogenic bacterium *Dickeya*. *Genome Announc* 1:e00978-13. <https://doi.org/10.1128/genomeA.00978-13>.
- Li B, Shi Y, Ibrahim M, Liu H, Shan C, Wang Y, Kube M, Xie GL, Sun G. 2012. Genome sequence of the rice pathogen *Dickeya zeae* strain ZJU1202. *J Bacteriol* 194:4452–4453. <https://doi.org/10.1128/JB.00819-12>.
- Wang X, He SW, Guo HB, Han JG, Thin KK, Gao JS, Wang Y, Zhang XX. 2020. *Dickeya oryzae* sp. nov., isolated from the roots of rice. *Int J Syst Evol Microbiol* 70:4171–4178. <https://doi.org/10.1099/ijsem.0.004265>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* 36:1925–1927. <https://doi.org/10.1093/bioinformatics/btz848>.
- Tatusova T, Dicuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 9:5114. <https://doi.org/10.1038/s41467-018-07641-9>.