



Review

Effects of Taijiquan and Qigong exercises on depression and anxiety levels in patients with substance use disorders: A systematic review and meta-analysis

Peng Zhang^a, Zaimin Li^a, Qing Yang^b, Jiali Zhou^a, Xiujie Ma^{a,c,*}

^a School of Wushu, Chengdu Sport University, Chengdu, 610041, China

^b Department of Martial Arts, Shanghai University of Sport, Shanghai, 200438, China

^c Chinese Guoshu Academy, Chengdu Sport University, Chengdu, 610041, China

ARTICLE INFO

Keywords:

Depression
Anxiety
Substance abuse disorder
Exercise
Taijiquan
Qigong

ABSTRACT

Previous studies have shown that Taijiquan and Qigong exercise can effectively reduce depression and anxiety in healthy and clinical populations. At present, only a few reviews have elaborated on the premise of different types of interventions, and there is still a lack of studies that systematically summarize the clinical evidence of patients with substance use disorders. The purpose of this meta-analysis was to assess the impact of the Taijiquan and Qigong exercise on the mood of individuals with substance use disorders (SUD); Articles were searched in Pubmed, Embase, Cochrane Library, Web of Science, and CNKI from their inception to the May 24, 2021. All randomized controlled trials (RCT) using Taijiquan and Qigong intervention were included. With the Review Manager software to determine the effect (standardized mean difference, SMD), subgroup analysis was conducted to intervention type, exercise time, and exercise duration. Nine studies totaling 823 participants were included in the study. Overall results indicated that Taijiquan and Qigong exercise had significant benefits in reducing depression ($SMD = -0.35$, 95% $CI: -0.53$ to -0.18 , $Z = 4.00$, $p < 0.01$) and anxiety ($SMD = -0.31$, 95% $CI: -0.50$ to -0.11 , $Z = 3.06$, $p < 0.01$) symptoms in SUD. When depression and anxiety were examined separately, subgroup analysis demonstrated that Taijiquan intervention ($SMD = -0.45$, 95% $CI: -0.71$ to -0.19 , $Z = 3.39$, $p < 0.01$) and Qigong intervention ($SMD = -0.28$, 95% $CI: -0.51$ to -0.04 , $Z = 2.33$, $p = 0.02$), 30–60 min exercise time ($SMD = -0.41$, 95% $CI: -0.61$ to -0.20 , $Z = 3.82$, $p < 0.01$) and duration of 9–12 weeks ($SMD = -0.57$, 95% $CI: -1.10$ to -0.04 , $Z = 2.10$, $p = 0.04$) more than 12 weeks ($SMD = -0.28$, 95% $CI: -0.50$ to -0.07 , $Z = 2.59$, $p = 0.01$) could effectively alleviate the symptoms of depression. Meanwhile, Qigong intervention ($SMD = -0.31$, 95% $CI: -0.52$ to -0.10 , $Z = 2.85$, $p < 0.01$), 30–60 min exercise time ($SMD = -0.29$, 95% $CI: -0.53$ to -0.04 , $Z = 2.28$, $p = 0.02$), more than 60 min exercise time ($SMD = -0.54$, 95% $CI: -1.06$ to -0.02 , $Z = 2.05$, $p = 0.04$), and duration of less than 9 weeks ($SMD = -0.35$, 95% $CI: -0.68$ to -0.02 , $Z = 2.07$, $p = 0.04$) had positive effects on alleviating anxiety symptoms with SUD. Taijiquan and Qigong exercise could reduce levels of anxiety and depression in those with SUD. Considering the small number of included studies, more reliable RCTs are needed on this topic.

Introduction

Substance abuse is a global problem that encompasses the misuse of alcohol, tobacco, heroin, and synthetic drugs. According to the World Health Organization, in 2016, approximately 275 million people worldwide used drugs at least once, accounting for 5.6% of the world's population in the age range of 15–64 years.¹ In 2017, about 585,000 people died from drug abuse.² Numerous studies have shown that

drug-dependent people generally suffer from affective disturbances. Drug abuse can cause chronic illness³; long-term abuse may also lead to symptoms such as anxiety, depression, insomnia, and impaired social functioning.^{4–8} Opioid dependence is one of the most severe types of drug dependency.⁹ Currently, the treatment of drug dependency relies on drug replacement therapy, in which either methadone or buprenorphine is used to treat opioid addiction.^{10,11} However, methadone or buprenorphine use is a risk factor for addiction, and overuse of either substances

* Corresponding author. School of Wushu/Chinese Guoshu Academy, Chengdu Sport University, Chengdu, 610041, China.

E-mail address: maxiujie@cdsu.edu.cn (X. Ma).

<https://doi.org/10.1016/j.smhs.2021.12.004>

Received 16 October 2021; Received in revised form 30 November 2021; Accepted 14 December 2021

Available online 30 December 2021

2666-3376/© 2022 Chengdu Sport University. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd.

may lead to serious health and social problems.^{12,13} There is currently no effective way to manage the toxic effects of drugs, including those of synthetic drugs.¹⁴ Increasing physical activity levels has been shown to improve mental health by reducing the risk of depression,^{15–17} anxiety, and other stress-related disorders.^{18,19} Physical exercise has recently been used to treat mental illness,²⁰ such as anorexia nervosa,²¹ bulimia, and alcohol and substance abuse disorders.²² Exercise may be a complementary approach to pharmacological and psychological interventions because it is non-discriminatory.²³ Exercise and movement may aid smoking cessation, enhancing mood, reducing negative affect, and decreasing the severity of withdrawal symptoms.^{24–26}

Qigong and Taijiquan are two of the most popular traditional Chinese exercise and martial arts modalities, respectively, known as mind-body exercise therapy. Qigong is older than Taijiquan and is considered a kind of “medical” movement in the field of Chinese medicine.²⁷ Recently, both modalities have attracted research attention due to their efficacy in treating mental diseases, mood disorders, and other secondary diseases. Both modalities represent mind-body exercises, helping achieve physical and mental health harmony through body movements, regular breathing, and adjustments of the mind. Taijiquan and Qigong exercise can effectively relieve depression and anxiety symptoms among drug abusers.^{28,29} Previous studies have shown promising effects of Taijiquan or Qigong practice on improving depression and anxiety in otherwise healthy adults,³⁰ young people with major depressive disorder,³¹ obese women,³² cancer patients,³³ and chronic obstructive pulmonary disease patients.^{34,35} However, previous studies have used both modalities at varied frequencies and over different periods; therefore, their effects on the levels of anxiety, depression, and overall mood in patients with substance use disorders (SUD) remain unclear. The present meta-analysis aimed to clarify the impact of these exercise modalities on the mood of patients with SUD, accounting for the differences in exercise type, and session and practice duration.

Materials and methods

Search strategy

The literature search was conducted in PUBMED, Web of Science, EMBASE, Cochrane Library, and China National Knowledge Infrastructure databases, covering the period from database inception to the May 24, 2021. There were no restrictions on the publication date or language. Databases were searched using the following terms: Tai Ji, Tai Chi, Tai Ji Quan, Taijiquan, Tai Chi, Tai Chi Chuan, Qigong, Qi Gong, Qi-gong, Chi Kung, Chi Chung, Qi-training, Qigong Massage, Health Qigong, Meditation, Meditative Movement, Liuzijue, Wuqinxi, Yijinjing, Baduanjin, Traditional Exercise, Chinese Traditional Exercise, Substance-Related Disorders, Alcohol, Tobacco, Nicotine, Heroin, Morphine, Marijuana, Cannabis, Opioid, Cocaine, Methamphetamine, Drug Abuse, Drug Addiction, Drug Dependence, Drug Use Disorder, Substance Use, Substance Abuse, Depression, Anxiety, Mental Health, Mental Problem, Psychological Health, and Psychological Problem. Chinese translations of these terms were used for Chinese database searches. A complete record of search strings is provided in the supplementary file. We manually searched the reference lists of all included studies and relevant reviews to identify any additional potentially relevant studies. If an article was incomplete or its full text was unavailable, we contacted the corresponding author by email to obtain detailed information. For literature tracing, based on the retrieved literature or related references listed in the review, we used Baidu Scholar and Google Scholar to search for them retrospectively.

Eligibility criteria

Only randomized clinical trials (RCT) of interventions for depression and anxiety in individuals aged ≥ 18 years and diagnosed with SUD were included. Reviews, comments, conference abstracts, and book chapters were excluded.

To be included in the meta-analysis, a study must have included both an experimental group that received Taijiquan or Qigong exercise, and a control condition that included hospital routine care, no treatment, or other types of intervention (e.g., medication).

The main outcomes were depression and anxiety symptoms. Outcome measures included the Hamilton Anxiety Scale, Beck Depression Inventory, Profile of Mood States, State-Trait Anxiety Inventory, Hamilton Rating Scale for Depression, Self-Rating Anxiety Scale, Hospital Anxiety and Depression Scale, Symptom Checklist-90, and Self-Rating Depression Scale scores.

Study selection and data extraction

Two reviewers (Zhang and Yang, respectively) extracted and cross-checked trial data. The decision to include a study was made primarily based on its title and abstract; in cases where title and abstract were insufficient to determine study eligibility, full-text reading was performed. Inconsistencies were resolved by a third independent reviewer and group discussion, as required (Ma). We extracted data on the following variables: publication year, first author name, study design (e.g., intervention model, practice time, persistence time, and outcome measurements), and participant characteristics (e.g. age, sex, and sample size). Risk of bias evaluation details are presented in Table 1.

Study quality assessment

Two authors (Zhang and Yang) evaluated the risk of bias in each study using a tool provided by the Cochrane Collaboration.³⁶ We evaluated seven key domains: allocation concealment, sequence generation, blinding of participants and personnel, incomplete outcome data, blinding of outcome assessment, selective outcome reporting, and other sources of bias. The risk of bias was classified as low, unclear, or high for each included study. Discrepancies in assessments were resolved by a third party.

Study analysis method

The Review Manager (version: 5.4.1, Cochrane, UK) was used for statistical analysis. For continuous outcomes, we calculated the standardized mean difference (SMD) and 95% confidence intervals (95% CI). The I^2 statistic was used to assess heterogeneity; fixed- and random-effects models were used to evaluate studies with low ($I^2 < 50\%$) and high ($I^2 \geq 50\%$) heterogeneity, respectively. The p -values of < 0.05 indicated statistically significant findings. Subgroup analyses were performed to minimize the impact of heterogeneity on the reported estimates. The included studies varied in terms of exercise type, time, and duration, which were the factors considered in subgroup analysis. The variables of interest were: (a) intervention type (two groups): Taijiquan and Qigong; (b) exercise time (three groups): less than 30 min, 30–60 min, and more than 60 min; and (c) exercise duration (three groups): shorter than 9 weeks, 9–12 weeks, and more than 12 weeks.

Results

A total of 752 articles were identified. In preliminary screening, we excluded 721 studies based on their title and abstract for reasons including duplications ($n = 214$) and irrelevance ($n = 507$). The remaining 31 articles were further screened by reading the full text. Twenty-two studies were removed owing to no body-exercise ($n = 5$), reviews ($n = 3$), no control group ($n = 1$), no outcomes of depression or anxiety ($n = 11$), incomplete data ($n = 2$). Finally, the meta-analysis included five and four studies on Taijiquan and Qigong exercises, respectively (Fig. 1).

Table 1
Characteristics of included studies.

Study name	Location (Language)	Substance	Sample Size (Attrition Rate)	Sample size (Mean age \pm SD)	Intervention Program	Training			Outcome measures
						Frequency (Week)	Time (Min)	Duration (Week)	
Li et al. (2002) ³⁷	Guangzhou, China (English)	Heroin	86,0%	E:n = 34 (33.3 \pm 6.5) C 1:n = 26 (31.9 \pm 5.9) C 2:n = 26 (31.7 \pm 6.1)	E = Pangu Qigong C1 = Medication C2 = No treatment	7	120–150	1 week +3 days	HAS
Li et al. (2013) ³⁸	Yunnan, China (English)	Heroin	70,53%	E: n = 36 (30.7 \pm 6.3) C:n = 34 (30.7 \pm 6.3)	E = Taijiquan C = Treatment as usual	14	60	24 weeks	HRSD
Zhu et al. (2018) ³⁹	Shanghai, China (English)	Amphetamine-drugs	80,39%	E:n = 40 (33.74 \pm 7.1) C:n = 40 (37.76 \pm 9.85)	E = 24–posture Taijiquan routine C = the 9th Guang Bo Ti Cao	1–3 months =5 4–6 months =3	60	24 weeks	SDS
Oh et al. (2016) ⁴⁰	Jeungpyeong, Korea (English)	Alcohol	50,24%	E:n = 25 (45.4 \pm 7.8) C:n = 25 (48.1 \pm 11.5)	E = 24–posture Taijiquan routine C = Hospital–rehabilitation program	3	50	8 weeks	BDI
Geng et al. (2016) ⁴¹	Shanghai, China (Chinese)	Amphetamine-drugs	60,5%	E:n = 30 (34 \pm 7) C:n = 30 (37.76 \pm 9.85)	E = Yang–style Taijiquan C = the 9th Guang Bo Ti Cao	5	45	12 weeks	SCL-90
Huang et al. (2015) ⁴²	Guangdong, China (Chinese)	Heroin	100,13%	E:n = 50 (35.26 \pm 12.22) C:n = 50 (35.21 \pm 12.12)	E = Qigong (Baduanjin) +Medication C = Medication	14	30	20 weeks	SAS
Zhang et al. (2020) ⁴³	Heze, China (English)	Amphetamine-drugs	76,5%	E:n = 38 (41.08 \pm 9.94) C:n = 38 (39.11 \pm 8.90)	E = 24–posture Taijiquan routine C = the 9th Guang Bo Ti Cao	5	80	24 weeks	BDI
David et al. (2013) ⁴⁴	Bedford, USA (English)	Cocaine	101,17%	E:n = 51 (36.0 \pm 9.4) C:n = 50 (40.4 \pm 11.9)	E = Qigong C = Sham Qigong	2–3	15	2 weeks	BDI STAIT
Fu et al. (2016) ⁴⁵	Anhui, China (Chinese)	Amphetamine-drugs	200,0%	E:n = 100 (28.3 \pm 7.83) C:n = 100 (27.99 \pm 8.17)	E = Qigong (Wuqinxi) C = No treatment	7	30	20 weeks	SAS SDS

¹ E-experiment group; C-control group; HAS-Hamilton Anxiety Scales; BDI-Beck Depression Inventory; STAIT-Trait Anxiety Inventory; POMS-Profile of Mood States; HRSD-Hamilton Rating Scale for Depression; SCL–90-Symptom Checklist 90; SAS-Self–Rating Anxiety Scale; SDS Self-Rating Depression Scale.

Study characteristics

Selected studies are shown in Table 1. Our systematic review included nine articles.^{37–45} Table 1 presents the characteristics of the included studies. These studies used Taijiquan or Qigong exercise modalities as an intervention in the experimental groups, and “hospital routine care”, “no treatment”, and “other types of intervention” in the control groups. Of all included studies, six were published in English^{37–40,43,44} and three were published in Chinese.^{41,42,45} In addition, one study was performed in the United States,⁴⁴ one was performed in Korea,⁴⁰ and seven were performed in China.^{37–39,41–43,45} The included participants were dependent on heroin, methamphetamine, amphetamine-type stimulant, K powder, ecstasy, and synthetic drugs. The sample sizes ranged from 50 to 200 participants, with a total of 823 participants (404 and 419 participants in the Taijiquan and Qigong, and control groups, respectively). Four studies included only female participants^{38–41,45} and one study included only male participants.³⁷

Selected studies included three styles of Taijiquan. Three of the included articles involved the style of 24-posture,^{39,40,43} one used the Yang style,⁴¹ and one did not specify the exercise style used. Among the studies involving Qigong, one used Pangu Qigong,³⁷ created by Chinese Qigong experts and considered easy to learn and practice; one study used the Baduanjin style,⁴² one used the Wuqinxi style,⁴⁵ and one study did not specify the style used.⁴⁴ One trial compared the effect of Qigong versus that of no treatment and that of medication.³⁷ Intervention duration ranged from 10 days to 24 weeks. All studies examined the effects of Taijiquan and Qigong exercise immediately after the intervention, but few studies reported follow-up outcomes. Seven studies measured depression-related variables,^{37–41,43–45} and five studies measured anxiety-related variables.^{37,41,42,44,45} The anxiety scales used included the Spielberger State-Trait Anxiety Inventory-State, Hamilton Anxiety Scale, Self-rating Anxiety Scale, and Symptom Checklist-90. Depression scales used included the Hamilton Rating Scale for




Fig. 1. Flow of study selection.

Depression, Center for Epidemiological Studies Depression Scale, Self-rating Depression Scale, and Symptom Checklist-90.

Methodological quality assessment

Methodological details of the included studies are shown in Table 2. Most included articles failed to report the process of sequence generation and allocation of concealment and were thus classified as “unclear” in the risk of bias domain. Six articles described study drop-outs, and the

dropout rate was within reasonable limits (Table 1). Blinding of instructors and participants is impossible in this type of exercise intervention. Therefore, the risk of bias was judged as low for group allocation and result blinding. All studies reported their randomization protocols, including four studies that used random sequence generation processes,^{37,39,40,45} two studies that used the order of admission to hospital,^{37,45} and two studies that used digital random tables and computer randomization.^{39,40} All studies described interventions used in the control group, but only three articles accounted for blinding.^{39,43,44} One

Table 2
Quality assessment of the included articles.

Study name	Random sequence generation	Allocation concealment	Blinding of Participants and personnel	Blinding of outcome assessments	Incomplete outcome data	Selective reporting	Other bias
Li et al.(2002) ³⁷	High	Unclear	Unclear	Low	Low	Low	Unclear
Li et al.(2013) ³⁸	Unclear	Unclear	Unclear	Unclear	Low	Low	Unclear
Zhu et al.(2018) ³⁹	Low	Low	Low	Unclear	Low	Low	Unclear
Oh et al.(2016) ⁴⁰	Low	Unclear	Unclear	Unclear	Low	Low	Unclear
Geng et al.(2016) ⁴¹	Unclear	Unclear	Unclear	Unclear	Low	Low	Unclear
Huang et al.(2015) ⁴²	Unclear	Unclear	Unclear	Unclear	Low	Low	Unclear
Zhang et al.(2020) ⁴³	Unclear	Low	Low	Low	Low	Low	Unclear
David et al.(2013) ⁴⁴	Low	Low	Low	Low	Low	Low	Unclear
Fu et al.(2016) ⁴⁵	High	Unclear	Unclear	Unclear	Low	Low	Unclear

article failed to report the reason for participant loss to follow-up and withdrawal.^{43,44} Overall, the quality of the included studies was relatively high.

Meta-analysis of outcome indicators

Seven studies ($n = 535$) were included in the meta-analysis of the impact of Taijiquan and Qigong exercise on depression in SUD patients. Heterogeneity was low ($I^2 = 0\%$, $p < 0.05$) and the fixed-effects model was used for analysis, revealing that, compared with the control intervention, Taijiquan and Qigong exercise practice may alleviate depressive symptoms of SUD patients ($SMD = -0.35$, 95% $CI: -0.53$ to -0.18 , $Z = 4.00$, $p < 0.01$; Fig. 2). A subgroup analysis was conducted to further explore the relationship between moderating variables and depression in drug users, confirming that Taijiquan and Qigong exercises can alleviate depressive symptoms in this group ($SMD = -0.45$, 95% $CI: -0.71$ to -0.19 , $Z = 3.39$, $p < 0.01$, $I^2 = 0\%$, and $SMD = -0.28$, 95% $CI -0.51$ to -0.04 , $Z = 2.33$, $p = 0.02$, $I^2 = 0\%$, respectively) (Figs. 2 and 3). In addition, 30–60 min of exercise ($SMD = -0.41$, 95% $CI: -0.61$ to -0.20 , $Z = 3.82$, $p < 0.01$; Fig. 4) was more effective at reducing depressive symptoms than the other factors tested. In addition, 9–12 weeks ($SMD = -0.57$, 95% $CI: -1.10$ to -0.04 , $Z = 2.10$, $p = 0.04$) and more than 12 weeks ($SMD = -0.28$, 95% $CI: -0.50$ to -0.07 , $Z = 2.59$, $p = 0.01$) of exercise alleviated symptoms of depression. However, subgroup analysis of two studies with exercise duration of less than 9 weeks revealed high heterogeneity ($I^2 = 65\%$); thus, the random-effects model was used for meta-analysis, which revealed some non-significant effects ($SMD = -0.56$, 95% $CI: -1.22$ to -0.11 , $Z = 1.64$, $p = 0.1$; Fig. 5).

We included 5 ($n = 491$) studies in the meta-analysis of the impact of Taijiquan and Qigong exercise on anxiety in SUD patients were as follows. After the heterogeneity test, ($I^2 = 54\%$, $p = 0.001$). It suggests that the 5 groups of data selected in this study have high heterogeneity and heterogeneity lookup is necessary. Through sensitivity analysis of the 5 groups of data, the article written by Huang et al.⁴² affected heterogeneity after removing the study, which showed that the remaining 4 groups of data ($I^2 = 0\%$, $p = 0.80$) and Effect Size relative risk (RR) could be subjected to meta-analysis with fixed effects. The results showed that compared with the control group, the intervention of Taijiquan and Qigong exercise practice could significantly alleviate anxiety symptoms among SUD patients ($SMD = -0.31$, 95% $CI: -0.50$ to -0.11 , $Z = 3.06$, $p = 0.002$; Fig. 6).

Subgroup analyses were performed based on exercise types. Heterogeneity among the four studies on Qigong exercise was high ($I^2 = 64\%$, $p = 0.04$) suggesting between the data selected in this study. The heterogeneity is statistically significant and requires a heterogeneous lookup. Sensitivity analysis of four literature found that the article written by Huang et al.⁴² had a great impact on heterogeneity, removed again for heterogeneity test analysis, and the results show the remaining 3 articles There was no heterogeneity ($I^2 = 0\%$, $p = 0.61$), and Effect Size RR could be subjected to meta-analysis with fixed effects ($SMD = -0.31$, 95% $CI: -0.52$ to -0.10) and were statistically significant ($Z = 2.85$, $p = 0.004$). Subgroup analysis showed that the practice of Qigong

exercise can significantly alleviate symptoms of anxiety among SUD patients. In the subgroup analysis of Tai Chi exercise intervention events, since there is only one study on the intervention of anxiety, there is no statistical significance (Fig. 7). About the effect of exercise time on anxiety, 3 groups of data involving exercise 30–60 min, tested for heterogeneity, ($I^2 = 73\%$, $p = 0.04$), suggesting between the data selected in this study. The heterogeneity is statistically significant and requires a heterogeneous lookup. Another heterogeneity test after removing the article of Huang et al.⁴² showed no heterogeneity between the remaining 2 articles ($I^2 = 0\%$, $p = 0.96$) fixed-effects model ($SMD = -0.29$, 95% $CI: -0.53$ to -0.04) and were statistically significant ($Z = 2.28$, $p = 0.02$). The results showed more than 60 min exercise time ($SMD = -0.54$, 95% $CI: -1.06$ to -0.02 , $Z = 2.05$, $p = 0.04$) and 30–60 min exercise time ($SMD = -0.29$, 95% $CI: -0.53$ to -0.04 , $Z = 2.28$, $p = 0.02$) were more effective than the other factors tested (Fig. 8). In the 2 articles on the effect of exercise duration on the duration of anxiety for more than 12 weeks, after the heterogeneity test, ($I^2 = 86\%$, $p = 0.008$), indicating that the heterogeneity between the two studies is high, So when choosing random-effects for meta-analysis. The results showed a duration of longer than 12 weeks that are not statistically significant ($SMD = -0.61$, 95% $CI: -1.31$ to -0.08 , $Z = 1.74$, $p = 0.08$). In addition, about the effect of exercise duration on anxiety the duration of shorter than 9 weeks ($SMD = -0.35$, 95% $CI: -0.68$ to -0.02 , $Z = 2.07$, $p = 0.04$), could reduce anxiety symptoms, which were more effective than the other factors tested (Fig. 9).

Discussion

This meta-analysis aimed to summarize the existing evidence on the impact of Taijiquan and Qigong exercise on the levels of depression and anxiety in SUD patients. The present findings suggest that Taijiquan and Qigong exercise modalities may alleviate depression and anxiety symptoms. Given the safety of these modalities, clinicians may prescribe Taijiquan and Qigong exercise to support treatment for depression and anxiety in SUD patients. The present findings are consistent with those of previous systematic reviews on the effects of Taijiquan and Qigong exercise on the levels of anxiety in SUD patients.⁴⁶ Overall, Taijiquan and Qigong exercise reduced depression ($SMD = -0.35$, 95% $CI: -0.53$ to -0.18 , $Z = 4.00$, $p < 0.01$) and anxiety ($SMD = -0.45$, 95% $CI: -0.50$ to -0.11 , $Z = 3.06$, $p < 0.01$) levels in SUD patients. Meanwhile, there may be significant heterogeneity in the research of Huang et al.⁴² compared with the effect size of other studies, they proved to have a great impact on SUD anxiety, Through the analysis of this study, from the event intervention aspect, it was found that the intervention (Qigong + Medication) for SUD patients was different from other studies. Subgroup analysis revealed that Taijiquan and Qigong, 30–60 min exercise time had significant benefits in anxiety symptoms in SUD patients. In addition, exercise duration of more than 9 weeks improved depression symptoms and that of less than 9 weeks improved depression symptoms; these findings are inconsistent with those of some individual studies. Further RCTs on Taijiquan and Qigong in the treatment of anxiety and depression are required to elucidate the underlying mechanisms. Studies that showed




Fig. 2. The forest plot about the effect of Taijiquan and Qigong exercise on depression.




Fig. 3. The forest plot about the effect of intervention type on depression.




Fig. 4. The forest plot about the effect of exercise time on depression.

the impact of less than 9 weeks of exercise practice on anxiety levels all involved Qigong exercise, suggesting the suitability of this intervention in the short term. Meanwhile, the impact of Taijiquan exercise on anxiety symptoms of SUD patients remains unclear due to the limited number of studies available, suggesting a need for further research.

Substance dependence involves complex mechanisms, including the brain's adaptation to exogenous opioids,^{47,48} which act on the same receptors as endogenous neurotransmitters such as endorphins and dopamine.⁴⁹ Endorphins and dopamine are key signaling molecules in the brain's reward circuits, associated with habit and long-term memory formation.⁵⁰ Studies have reported that exercise improves the mood of drug-dependent individuals mainly by regulating norepinephrine (NE) and glucocorticoid pathways. The sympathetic nervous system and hypothalamic-pituitary-adrenal (HPA) system are both activated in response to stress.⁵¹ The adrenal medulla secretes a large amount of NE, increasing NE levels in the ventral tegmental area and nucleus accumbens.⁵² Elevated NE levels increase CRF secretion in the hypothalamus, activate HPA, and increase corticosterone (CORT) production. CORT crosses the blood-brain barrier and enters the brain,^{53,54} increasing the levels of serotonin (5-HT) and its metabolite 5-hydroxyindoleacetic acid

in the frontal cortex, nucleus accumbens, hypothalamus, and CA. In general, 5-HT acts in synergy with NE. These changes affect the limbic system.⁵⁵

Moderate-intensity exercise affects the HPA activation–CORT-mediated pathway. The relative reduction in the rate of NE synthesis and activation of the sympathetic nervous system can be achieved through stress management techniques.⁵⁶ Liu et al.⁵¹ used a 12-week Taijiquan training intervention, showing it may increase the physiological adaptability and modulate the reactivity of the sympathetic nervous system, helping reduce the fluctuation of stress-related hormones and levels of depression and anxiety. Taijiquan is characterized by continuous slow movement,⁵⁷ deep breathing, and mental concentration. Qigong involves Qi and Yi to help achieve inner balance. It consists of meditation, relaxation, guided imagery, deep tranquility, and breathing exercises. Taijiquan and Qigong are both moderate-intensity exercise modalities. The smooth rhythmic motions of Taijiquan and Qigong exercise are usually experienced as relatively pleasurable.^{58,59} Several studies have assessed the effects of Taijiquan and Qigong on psychological responses, such as depression, stress, and life satisfaction.^{60–63} The effects of these modalities may be mediated by moderate rhythmic movement, which




Fig. 5. The forest plot about the effect of exercise duration on depression.




Fig. 6. The forest plot about the effect of Taijiquan and Qigong exercise on anxiety.




Fig. 7. The forest plot about the effect of intervention type on anxiety.

increases the parasympathetic tone.³⁹ In addition, both modalities help patients achieve relaxation, and reconnect with their mind, body, and spirit, making them more mindful and internally focused.⁶⁴ Finally, practicing Taijiquan and Qigong exercises in a group can promote social interaction in interpersonal relationships, providing social support, and benefiting patients' physical and mental health.⁶³

Cognitive behavioral therapy and other non-drug therapies, such as exercise, acupuncture, and psychotherapy, have become important complementary modalities in treating SUD patients. Although exercise is

a low-risk and low-cost intervention, further research is needed to develop evidence-based exercise prescriptions in this patient group.

Our study has several strengths. Relatively few relevant studies have been published, precluding subgroup analysis. However, we examined the impact of the type of intervention, exercise session time, and practice duration on the outcomes of interest. Our review methods were systematic and described in detail. We used extensive search strategies with no constraints on publication date or language. Furthermore, study eligibility criteria were clearly defined, and only RCTs were included in




Fig. 8. The forest plot about the effect of exercise time on anxiety.




Fig. 9. The forest plot about the effect of exercise duration on anxiety.

this review. At least two independent reviewers were responsible for study selection, quality assessment, and data extraction.

This study has some limitations. First, only nine studies were included (five on Taijiquan and four on Qigong) and the overall sample size was small. Second, outcomes were difficult to measure; therefore, the presented results should be approached with caution. Moreover, different levels of exercise intensity may differently affect hormonal regulation and the resulting mood; however, exercise intensity was not reported in the examined studies.

Conclusions

Our meta-analysis revealed that Taijiquan and Qigong exercise may improve depression and anxiety symptoms in patients with SUD. In addition, this meta-analysis showed that Qigong exercise duration of less than 9 weeks may reduce anxiety levels in SUD patients. The present findings may provide a foundation for future research and clinical applications. However, the number of studies included was small. Future studies are required to validate the present findings.

Submission statement

All authors have read and agreed to the content of the manuscript, which is not yet published and is not considered for publication elsewhere.

Authors' contributions

M.X. conceived of and designed the review; M.X., P.Z., Y.Q., L.Z., Z.J. performed the searches and conducted the meta-analysis; all authors analyzed the data including screening of studies, quality assessment, data extraction; P.Z. and M.X. wrote the paper with all authors contributing to the drafting and finalization of the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgement

This research was funded by Sichuan Social Science Fund, grant number SC21ZW003.

References

- UNODC. *World Drug Report -Executive Summary—Conclusions and Policy Implications*. 1. New York: United Nations Office on Drugs and Crime (UNODC); 2018.
- UNODC. *World Drug Report -Executive Summary—Conclusions and Policy Implications*. 1. New York: United Nations Office on Drugs and Crime (UNODC); 2019.
- Zhang Y, Lu C, Zhang J, et al. Gender differences in abusers of amphetamine-type stimulants and ketamine in southwestern China. *Addict Behav*. 2013;38(1):1424–1430. <https://doi.org/10.1016/j.addbeh.2012.06.024>.
- Anglin MD, Burke C, Perrochet B, Stamper E, Dawud-Noursi S. History of the methamphetamine problem. *J Psychoactive Drugs*. 2000;32(2):137–141. <https://doi.org/10.1080/02791072.2000.10400221>.
- Homer BD, Solomon TM, Moeller RW, Mascia A, DeRaleau L, Halkitis PN. Methamphetamine abuse and impairment of social functioning: a review of the underlying neurophysiological causes and behavioral implications. *Psychol Bull*. 2008;134(2):301–310. <https://doi.org/10.1037/0033-2909.134.2.301>.
- Beadell NC, Thompson EM, Delashaw JB, Cetas JS. The deleterious effects of methamphetamine use on initial presentation and clinical outcomes in aneurysmal subarachnoid hemorrhage. *J Neurosurg*. 2012;117(4):781–786. <https://doi.org/10.3171/2012.7.JNS12396>.
- Cohen JB, Dickow A, Horner K, et al. Abuse and violence history of men and women in treatment for methamphetamine dependence. *Am J Addict*. 2003;12(5):377–385.
- Pocuca N, Young JW, MacQueen DA, et al. Sustained attention and vigilance deficits associated with HIV and a history of methamphetamine dependence. *Drug Alcohol Depend*. 2020;215:108245. <https://doi.org/10.1016/j.drugaldep.2020.108245>.
- Krupitsky E, Zvartau E, Woody G. Use of naltrexone to treat opioid addiction in a country in which methadone and buprenorphine are not available. *Curr Psychiatry Rep*. 2010;12(5):448–453. <https://doi.org/10.1007/s11920-010-0135-5>.
- Deng Q, Tang Q, Schottenfeld RS, Hao W, Chawarski MC. Drug use in rural China: a preliminary investigation in Hunan Province. *Addiction*. 2012;107(3):610–613. <https://doi.org/10.1111/j.1360-0443.2011.03648.x>.
- Joseph H, Stancliff S, Langrod J. *Methadone maintenance treatment (MMT): a review of historical and clinical issues*. *Mt Sinai J Med*. 2000;67(5-6):347–364.
- Huhn AS, Strain EC, Tompkins DA, Dunn KE. A hidden aspect of the U.S. opioid crisis: Rise in first-time treatment admissions for older adults with opioid use disorder. *Drug Alcohol Depend*. 2018;193:142–147. <https://doi.org/10.1016/j.drugaldep.2018.10.002>.
- Nielsen S, Dietze P, Lee N, Dunlop A, Taylor D. Concurrent buprenorphine and benzodiazepines use and self-reported opioid toxicity in opioid substitution treatment. *Addiction*. 2007;102(4):616–622. <https://doi.org/10.1111/j.1360-0443.2006.01731.x>.
- Rawson RA. Current research on the epidemiology, medical and psychiatric effects, and treatment of methamphetamine use. *J Food Drug Anal*. 2013;21(4):S77–S81. <https://doi.org/10.1016/j.jfda.2013.09.039>.
- Schuch FB, Vancampfort D, Firth J, et al. Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. *Am J Psychiatr*. 2018;175(7):631–648. <https://doi.org/10.1176/appi.ajp.2018.17111194>.
- Harvey SB, Øverland S, Hatch SL, Wessely S, Mykletun A, Hotopf M. Exercise and the Prevention of Depression: Results of the HUNT Cohort Study. *Am J Psychiatr*. 2018;175(1):28–36. <https://doi.org/10.1176/appi.ajp.2017.16111223>.
- Vieira J, Porcu M, Rocha P. The regular physical exercises practice as complementary therapy to the treatment of depressed women. *Jornal Brasileiro De Psiquiatria [in Portuguese]*. 2007;56(1):23–28. <https://doi.org/10.1590/S0047-20852007000100007>.
- McDowell CP, Dishman RK, Vancampfort D, et al. Physical activity and generalized anxiety disorder: results from The Irish Longitudinal Study on Ageing (TILDA). *Int J Epidemiol*. 2018;47(5):1443–1453. <https://doi.org/10.1093/ije/dyy141>.
- LeardMann CA, Kelton ML, Smith B, et al. Prospectively assessed posttraumatic stress disorder and associated physical activity. *Public Health Rep*. 2011;126(3):371–383. <https://doi.org/10.1177/003335491112600311>.
- Torales J, Almirón M, Gonzalez I, et al. Physical exercise as an adjuvant treatment of mental disorders. *A narrative review*. *Anales de la Facultad de Ciencias Médicas (Asunción) [in Spanish]*. 2018;51(3):27–32. [https://doi.org/10.18004/anales/2018.051\(03\)27-032](https://doi.org/10.18004/anales/2018.051(03)27-032).
- Bassoe HH. Anorexia/bulimia nervosa: the development of anorexia nervosa and of mental symptoms. Treatment and the outcome of the disease. *Acta Psychiatr Scand Suppl*. 1990;361:7–13. <https://doi.org/10.1111/j.1600-0447.1990.tb11076.x>.
- Zschucke E, Heinz A, Ströhle A. Exercise and physical activity in the therapy of substance use disorders. *Sci World J*. 2012;2012:901741. <https://doi.org/10.1100/2012/901741>.
- Hallgren M, Vancampfort D, Giesen ES, Lundin A, Stubbs B. Exercise as treatment for alcohol use disorders: systematic review and meta-analysis. *Br J Sports Med*. 2017;51(14):1058–1064. <https://doi.org/10.1136/bjsports-2016-096814>.
- Bock BC, Marcus BH, King TK, Borrelli B, Roberts MR. Exercise effects on withdrawal and mood among women attempting smoking cessation. *Addict Behav*. 1999;24(3):399–410. [https://doi.org/10.1016/s0306-4603\(98\)00088-4](https://doi.org/10.1016/s0306-4603(98)00088-4).
- Scully D, Kremer J, Meade MM, Graham R, Dudgeon K. Physical exercise and psychological well being: a critical review. *Br J Sports Med*. 1998;32(2):111–120. <https://doi.org/10.1136/bjism.32.2.111>.
- Ussher MH, Taylor AH, West R, McEwen A. Does exercise aid smoking cessation? A systematic review. *Addiction*. 2000;95(2):199–208. <https://doi.org/10.1046/j.1360-0443.2000.9521996.x>.
- LG X. *Chinese Medical Qigong*. Harbin: Heilongjiang Science and Technology Publisher; 1990.
- Zhu D, Xu D, Dai G, Wang F, Xu X, Zhou D. Beneficial effects of Tai Chi for amphetamine-type stimulant dependence: a pilot study. *Am J Drug Alcohol Abuse*. 2016;42(4):469–478. <https://doi.org/10.3109/00952990.2016.1153646>.
- Wang D, Wang Y, Wang Y, Li R, Zhou C. Impact of physical exercise on substance use disorders: a meta-analysis. *PLoS One*. 2014;9(10):e110728. <https://doi.org/10.1371/journal.pone.0110728>.
- Wang CW, Chan CH, Ho RT, Chan JS, Ng SM, Chan CL. Managing stress and anxiety through qigong exercise in healthy adults: a systematic review and meta-analysis of randomized controlled trials. *BMC Complement Altern Med*. 2014;14:8. <https://doi.org/10.1186/1472-6882-14-8>.
- Giles A, Nasstasia Y, Baker A, et al. Exercise as Treatment for Youth With Major Depression: The Healthy Body Healthy Mind Feasibility Study. *J Psychiatr Pract*. 2020;26(6):444–460. <https://doi.org/10.1097/PRA.0000000000000516>.
- McCain NL, Gray DP, Elswick RK, et al. A randomized clinical trial of alternative stress management interventions in persons with HIV infection. *J Consult Clin Psychol*. 2008;76(3):431–441. <https://doi.org/10.1037/0022-006X.76.3.431>.
- Zeng Y, Luo T, Xie H, Huang M, Cheng AS. Health benefits of qigong or tai chi for cancer patients: a systematic review and meta-analysis. *Complement Ther Med*. 2014;22(1):173–186. <https://doi.org/10.1016/j.ctim.2013.11.010>.
- Yeh GY, Roberts DH, Wayne PM, Davids RB, Quilty MT, Phillips RS. Tai Chi Exercise for Patients With Chronic Obstructive Pulmonary Disease: A Pilot Study. *Respir care*. 2010;55(11):1475–1482.
- Wu JJ, Zhang YX, Du WS, et al. Effect of Qigong on self-rating depression and anxiety scale scores of COPD patients: A meta-analysis. *Medicine (Baltimore)*. 2019;98(22):e15776. <https://doi.org/10.1097/MD.00000000000015776>.
- Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. The Cochrane Collaboration; 2011 [updated March 2011] Version 5.1.0. <http://www.cochrane-handbook.org>.
- Li M, Chen K, Mo Z. Use of qigong therapy in the detoxification of heroin addicts. *Altern Ther Health Med*. 2002;8(1):50–59.
- Li DX, Zhuang XY, Zhang YP, et al. Effects of Tai Chi on the protracted abstinence syndrome: a time trial analysis. *Am J Chin Med*. 2013;41(1):43–57. <https://doi.org/10.1142/S0192415X13500043>.
- Zhu D, Dai G, Xu D, et al. Long-Term Effects of Tai Chi Intervention on Sleep and Mental Health of Female Individuals With Dependence on Amphetamine-Type Stimulants. *Front Psychol*. 2018;9:1476. <https://doi.org/10.3389/fpsyg.2018.01476>.
- Oh CU, Kim NC. Effects of Tai Chi on Serotonin, Nicotine Dependency, Depression, and Anger in Hospitalized Alcohol-Dependent Patients. *J Altern Complement Med*. 2016;22(12):957–963. <https://doi.org/10.1089/acm.2016.0246>.
- Geng J, Zhu D, Xu D. Rehabilitation effect of Taiji rehabilitation exercise on compulsory isolation of female drug abusers. *Chin J Sports Med [in Chinese]*. 2016;35(11):1048–1051. <https://doi.org/10.16038/j.1000-6710.2016.11.011>.
- Huang X, Wu S, Xu H. Clinical observation of modified Ba Duan Jin combined with methadone in improving anxiety symptoms of heroin addicts. *Shenzhen J Integrated Tradit Chin West Med [in Chinese]*. 2015;25(13):98–99. <https://doi.org/10.16458/j.cnki.1007-0893.2015.13.052>.
- Zhang Z, Zhu D. Effect of Taijiquan exercise on rehabilitation of male amphetamine-type addicts. *Evid Based Complement Alternat Med*. 2020;2020:8886562. <https://doi.org/10.1155/2020/8886562>.
- Smelson D, Chen KW, Ziedonis D, et al. A pilot study of Qigong for reducing cocaine craving early in recovery. *J Altern Complement Med*. 2013;19(2):97–101. <https://doi.org/10.1089/acm.2012.0052>.
- Fu G, Huang S, Tang C, Liu Y, Gao Y, et al. Effect of Wuqinxi on emotion in patients abstaining from drug. *J Anhui Univ Chin Med [in Chinese]*. 2016;35(05):26–29. <https://doi.org/10.3969/j.issn.2095-7246.2016.05.008>.
- Liu F, Cui J, Liu X, Chen KW, Chen X, Li R. The effect of tai chi and Qigong exercise on depression and anxiety of individuals with substance use disorders: a systematic review and meta-analysis. *BMC Complement Med Ther*. 2020;20(1):161. <https://doi.org/10.1186/s12906-020-02967-8>.
- Belcheva MM, Ho MT, Ignatova EG, et al. Buprenorphine differentially alters opioid receptor adaptation in rat brain regions. *J Pharmacol Exp Ther*. 1996;277(3):1322–1327.
- Swain JE, Ho SS. Opioids and maternal brain-behavior adaptation. *Neuropsychopharmacology*. 2021;46(1):265–266. <https://doi.org/10.1038/s41386-020-00858-7>.
- Martin JA, Campbell A, Killip T, et al. QT interval screening in methadone maintenance treatment: report of a SAMHSA expert panel [published correction appears in *J Addict Dis*. *J Addict Dis*. 2012 Jan;31(1):91. Haigney, Mark C P [added]]. *J Addict Dis*. 2011;30(4):283–306. <https://doi.org/10.1080/10550887.2011.610710>.
- Leshner AI. Addiction is a brain disease, and it matters. *Science*. 1997;278(5335):45–47. <https://doi.org/10.1126/science.278.5335.45>.
- Liu YP, Yuan MZ, H C, et al. Effect of 12-week taichi on catecholamines and angiotensin-II in young people. *Journal of Fujian Normal University (Natural Science Edition) [in Chinese]*. 2015;31(5):115–119.
- Shaham Y, Erb S, Stewart J. Stress-induced relapse to heroin and cocaine seeking in rats: a review. *Brain Res Brain Res Rev*. 2000;33(1):13–33. [https://doi.org/10.1016/S0165-0173\(00\)00024-2](https://doi.org/10.1016/S0165-0173(00)00024-2).
- Sairanen M, O'Leary OF, Knuutila JE, Castrén E. Chronic antidepressant treatment selectively increases expression of plasticity-related proteins in the hippocampus and medial prefrontal cortex of the rat. *Neuroscience*. 2007;144(1):368–374. <https://doi.org/10.1016/j.neuroscience.2006.08.069>.

54. Lapmanee S, Charoenphandhu J, Teerapornpantakit J, Krishnamra N, Charoenphandhu N. Agomelatine, venlafaxine, and running exercise effectively prevent anxiety- and depression-like behaviors and memory impairment in restraint stressed rats. *PLoS One*. 2017;12(11):e0187671. <https://doi.org/10.1371/journal.pone.0187671>.
55. Müller CP, Homberg JR. The role of serotonin in drug use and addiction. *Behav Brain Res*. 2015;277:146–192. <https://doi.org/10.1016/j.bbr.2014.04.007>.
56. Fink G. *Encyclopedia of stress*. Academic Press; 2000.
57. Zou L, Han J, Li C, et al. Effects of Tai Chi on lower limb proprioception in adults aged over 55: A systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2019; 100(6):1102–1113. <https://doi.org/10.1016/j.apmr.2018.07.425>.
58. Pa J, Goodson W, Bloch A, King AC, Yaffe K, Barnes DE. Effect of exercise and cognitive activity on self-reported sleep quality in community-dwelling older adults with cognitive complaints: a randomized controlled trial. *J Am Geriatr Soc*. 2014; 62(12):2319–2326. <https://doi.org/10.1111/jgs.13158>.
59. Kim TH, Pascual-Leone J, Johnson J, Tamim H. The mental-attention Tai Chi effect with older adults. *BMC Psychol*. 2016;4(1):29. <https://doi.org/10.1186/s40359-016-0137-0>.
60. Wang C, Collet JP, Lau J. The effect of Tai Chi on health outcomes in patients with chronic conditions: a systematic review. *Arch Intern Med*. 2004;164(5):493–501. <https://doi.org/10.1001/archinte.164.5.493>.
61. Gillespie LD, Robertson MC, Gillespie WJ, et al. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev*. 2009;2:CD007146. <https://doi.org/10.1002/14651858.CD007146.pub2>.
62. Garber CE, Blissmer B, Deschenes MR, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011;43(7): 1334–1359. <https://doi.org/10.1249/MSS.0b013e318213febf>.
63. Wang F, Man JK, Lee EK, et al. The effects of qigong on anxiety, depression, and psychological well-being: a systematic review and meta-analysis. *Evid Based Complement Alternat Med*. 2013;2013:152738. <https://doi.org/10.1155/2013/152738>.
64. Breslin KT, Reed MR, Malone SB. An holistic approach to substance abuse treatment. *J Psychoactive Drugs*. 2003;35(2):247–251. <https://doi.org/10.1080/02791072.2003.10400006>.