

# Draft Genome Sequence of the Organophosphorus Compound-Degrading *Burkholderia zhejiangensis* Strain CEIB S4-3

Armando Hernández-Mendoza,<sup>a</sup> Fernando Martínez-Ocampo,<sup>b</sup> Luis Fernando Lozano-Aguirre Beltrán,<sup>c</sup> Elida Carolina Popoca-Ursino,<sup>b</sup> Laura Ortiz-Hernández,<sup>b</sup> Enrique Sánchez-Salinas,<sup>b</sup>  Edgar Dantán-González<sup>b</sup>

Departamento de Bioquímica y Biología Molecular, Facultad de Ciencias, Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos, Mexico<sup>a</sup>; Laboratorio de Investigaciones Ambientales, Centro de Investigación en Biotecnología, Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos, Mexico<sup>b</sup>; Programa de Genómica Evolutiva, Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico<sup>c</sup>

***Burkholderia* species are widely distributed in the environment. A *Burkholderia zhejiangensis* strain was isolated from pesticide-contaminated soil from an agricultural field in Mexico and identified as an organophosphorus compound-degrading bacterium. In this study, we report the draft genome sequence of *Burkholderia zhejiangensis* strain CEIB S4-3.**

Received 7 November 2014 Accepted 10 November 2014 Published 18 December 2014

**Citation** Hernández-Mendoza A, Martínez-Ocampo F, Lozano-Aguirre Beltrán LF, Popoca-Ursino EC, Ortiz-Hernández L, Sánchez-Salinas E, Dantán-González E. 2014. Draft genome sequence of the organophosphorus compound-degrading *Burkholderia zhejiangensis* strain CEIB S4-3. *Genome Announc.* 2(6):e01323-14. doi:10.1128/genomeA.01323-14.

**Copyright** © 2014 Hernández-Mendoza et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Edgar Dantán-González, edantan@uaem.mx.

Bacteria of the genus *Burkholderia* are broadly distributed in several ecological niches. Pesticides that are applied to agricultural soil control pests that disrupt crop production, but when pesticides persist and accumulate in soils, they can alter microbial processes. Methyl parathion (MP) is a toxic organophosphate insecticide that irreversibly inhibits acetylcholinesterase (1). The hydrolysis of MP generates *p*-nitrophenol (PNP) (2), a toxic environmental pollutant (3, 4). In 2012, Popoca-Ursino (5) reported three bacterial strains of the genus *Burkholderia* isolated from crop soils in Morelos, Mexico. Here, we present a bacterium identified as *Burkholderia zhejiangensis* strain CEIB S4-3 that has the capability to completely degrade PNP and use it as a carbon source. These observations indicate that this organism potentially has genes that simultaneously hydrolyze MP and degrade PNP.

Strain CEIB S4-3 was grown at 30°C with constant agitation for 12 h. Genomic DNA was obtained using the AxyPrep bacterial genomic DNA miniprep kit (Axygen). The concentration was calculated in a UV-Vis NanoDrop 2000 spectrophotometer (Thermo Scientific), and 5 µg of genomic DNA was sequenced in the Genome Analyzer IIX system (Illumina). We obtained a random data set of 34,641,784 paired-end reads 72 bases in length. Quality-based trimming was performed with a DynamicTrim (SolexaQA++) perl script, and genome assembly was accomplished using the SPAdes (version 3.1.1) program. The draft genome has 154 contigs, with a calculated 7,666,843 bp total length and an  $N_{50}$  contig size of 156,081 bp. We aligned the draft genome with 41 complete genomes (chromosomes and plasmids) from members of the genus *Burkholderia* using the NUCmer program. The draft genome has 86.95 and 87.6% identity and 4.57 and 5.31 Mb alignment coverage with *Burkholderia* sp. strain RPE64 (3 chromosomes, 2 plasmids, and 6,964,487 bp total length) (6) and *Burkholderia* sp. strain YI23 (3 chromosomes, 3 plasmids, and 8,896,411 bp total length) (7), respectively. This analysis suggests that *B. zhejiangensis* CEIB S4-3 has at least 2 chromosomes.

The contigs were analyzed on the RAST server, which identi-

fied 7,228 coding sequences (CDS). We identified the 16S rRNA gene of *B. zhejiangensis* CEIB S4-3 using the RNAmmer 1.2 server. We compared 101 16S rRNA genes from the genus *Burkholderia* and two 16S rRNA genes from the genus *Ralstonia* (outgroup). The sequences were aligned in the MUSCLE server, and a phylogenetic analysis was performed in the MEGA (version 6.1) program with a neighbor-joining algorithm using 1,000 replicates for bootstrapping.

The 7,228 predicted CDS were compared with the COG database, and 4,391 proteins (60.8%) were identified: 1,545 open reading frames (ORFs) (21.4%) had an identified function and 2,846 ORFs (39.4%) had an uncharacterized function. We also identified a methyl parathion-degrading (*mpd*) gene (99% identity with the MpdB protein from *Burkholderia cepacia*) (8) and two PNP catabolic gene clusters (*pnpABA'E1E2FDC* and *pnpE1E2FDC*; identities ranging from 67 to 100% with PnpABE1E2FDC proteins from *Burkholderia* sp. strain SJ98) (9, 10). These catabolic genes will explain the capabilities of this bacterium to hydrolyze MP and degrade PNP completely.

**Nucleotide sequence accession number.** The draft genome sequence of *B. zhejiangensis* CEIB S4-3 (including 154 scaffolds) has been deposited in GenBank database under accession no. [JSBM000000000](https://www.ncbi.nlm.nih.gov/nuccore/JSBM000000000).

## ACKNOWLEDGMENTS

This work was supported by the Consejo Nacional de Ciencia y Tecnología (CONACyT) project CB-2012-182225 and UAEM project UAEMOR-CA-118 and a scholarship from CONACyT to F.M.-O. (293552) and E.C.P.-U. (351686).

## REFERENCES

- Singh BK, Walker A. 2006. Microbial degradation of organophosphorus compounds. *FEMS Microbiol. Rev.* 30:428–471. <http://dx.doi.org/10.1111/j.1574-6976.2006.00018.x>.
- Ramanathan MP, Lalithakumari D. 1999. Complete mineralization of methylparathion by *Pseudomonas* sp. A3. *Appl. Biotechnol.* 80:1–12.

3. Spain JC. 1995. Biodegradation of nitroaromatic compounds. *Annu. Rev. Microbiol.* 49:523–555. <http://dx.doi.org/10.1146/annurev.mi.49.100195.002515>.
4. Bhushan B, Chauhan A, Samanta SK, Jain RK. 2000. Kinetics of biodegradation of *p*-nitrophenol by different bacteria. *Biochem. Biophys. Res. Commun.* 274:626–630. <http://dx.doi.org/10.1006/bbrc.2000.3193>.
5. Popoca-Ursino EC. 2012. Aislamiento de bacterias con capacidad de degradación de plaguicidas organofosforados, evaluación molecular de sus actividades. Tesis de la Maestría en Biotecnología. Centro de Investigación en Biotecnología (CEIB), Universidad Autónoma del Estado de Morelos (UAEM), Toluca, Mexico.
6. Shibata TF, Maeda T, Nikoh N, Yamaguchi K, Oshima K, Hattori M, Nishiyama T, Hasebe M, Fukatsu T, Kikuchi Y, Shigenobu S. 2013. Complete Genome sequence of *Burkholderia* sp. strain RPE64, bacterial symbiont of the bean bug *Riptortus pedestris*. *Genome Announc.* 1(4): e00441-13. <http://dx.doi.org/10.1128/genomeA.00441-13>.
7. Lim JS, Choi BS, Choi AY, Kim KD, Kim DI, Choi IY, Kaa JO. 2011. Complete genome sequence of the fenitrothion-degrading *Burkholderia* sp. strain YI23. *J. Bacteriol.* 193:896–908.
8. Ekkhunnatham A, Jongsareejit B, Yamkunthong W, Wichitwechkarn J. 2012. Purification and characterization of methyl parathion hydrolase from *Burkholderia cepacia* capable of degrading organophosphate insecticides. *World J. Microbiol. Biotechnol.* 28:1739–1746. <http://dx.doi.org/10.1007/s11274-011-0985-y>.
9. Vikram S, Pandey J, Bhalla N, Pandey G, Ghosh A, Khan F, Jain RK, Raghava GPS. 2012. Branching of the *p*-nitrophenol (PNP) degradation pathway in *Burkholderia* sp. strain SJ98: evidences from genetic characterization of PNP gene cluster. *AMB Express* 2:30. <http://dx.doi.org/10.1186/2191-0855-2-30>.
10. Vikram S, Pandey J, Kumar S, Raghava GPS. 2013. Genes involved in degradation of *para*-nitrophenol are differentially arranged in form of non-contiguous gene clusters in *Burkholderia* sp. strain SJ98. *PLoS One* 8:e84766. <http://dx.doi.org/10.1371/journal.pone.0084766>.