



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Reconocimientos

Ninguno.

Bibliografía

1. Legido-Quigley H, Mateos-García JT, Campos VR, Gea-Sánchez M, Muntaner C, McKee M. The resilience of the Spanish health system against the COVID-19 pandemic. *Lancet Public Health*. 2020;5:e251–2.
2. Vieta E, Pérez V, Arango C. Psychiatry in the aftermath of COVID-19. *Rev Psiquiatr Salud Ment*. 2020;13:105–10.
3. Ellaway R, Tworek J. The Net generation illusion: Challenging conformance to social expectations. En: Ferris SP, editor. *Teaching, learning and the Net generation: Concepts and tools for reaching digital learners*. 1.^a ed. IGI Global; 2012. p. 324–39.
4. Bauman Z. *Education in Liquid Modernity*. *Rev Educ Pedagog Cult Stud*. 2005;27:303–17.

Santiago Ovejero^{a,b}, Maria Luisa Barrigon^{a,c}, Héctor Guadalajara^d, Eloísa Pérez-Santos^b y Enrique Baca-García^{a,e,f,g,h,i,j,*}

^a *Department of Psychiatry, Jimenez Diaz Foundation Hospital, Madrid, España*

- ^b *School of Psychology, University Complutense of Madrid, Madrid, España*
^c *Department of Psychiatry, Madrid Autonomous University, Madrid, España*
^d *Department of General and Digestive Surgery, Fundación Jiménez Díaz University Hospital, Madrid, España*
^e *Department of Psychiatry, Rey Juan Carlos University Hospital, Móstoles, España*
^f *Department of Psychiatry, General Hospital of Villalba, Madrid, España*
^g *Department of Psychiatry, Infanta Elena University Hospital, Valdemoro, España*
^h *Department of Psychiatry, Centre Hospitalier Universitaire de Nîmes, Francia*
ⁱ *Universidad Católica del Maule, Talca, Chile*
^j *Centro de Investigación Biomédica en Red Salud Mental (CIBERSAM), Madrid, España*

* Autor para correspondencia.
 Correo electrónico: EBaca@quironsalud.es
 (E. Baca-García).

<https://doi.org/10.1016/j.rpsm.2021.01.001>
 1888-9891/ © 2021 SEP y SEPB. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Psychiatrists as internists: Some considerations following a COVID-19 redeployment experience



Psiquiatras como internistas: algunas reflexiones tras una experiencia de reconversión en COVID-19

Dear Editor,

We have read with enthusiasm the recently published article about the changes that we can expect in psychiatry after the pandemic caused by the novel coronavirus SARS-CoV-2.¹ In particular, we believe that the authors are fully correct when, in first place, they highlight the medical nature of psychiatry. Since the first outbreak in China, several authors have warned about the mental health problems that the pandemic might cause. Also, they have pointed out the need for psychiatric care networks to adapt to this new reality.² However, few have mentioned the fact that, in the most critical moments of the pandemic, many psychiatrists had to put aside their common clinical practice to act as general practitioners and directly treat patients with COVID-19. Fortunately or unfortunately, this was the fate of the authors of this article when, one day in mid-March, we were told that a few hours later we should join as doctors a COVID-19 hospitalization facility in our general hospital in Madrid, Spain.

It is important to briefly put in context the place of psychiatry in modern medicine. To be honest, psychiatrists are not especially popular among other medical special-

lists. Psychiatry is one of the most reviled specialties within medicine.³ Although this is probably largely based on prejudice, sometimes we ourselves contribute to this bad reputation. For example, when we overlook the physical comorbidities of our patients, disregarding the risk that this may imply.⁴ Or when we insist on ruling out “organic” causes of mental disorders, apparently ignoring their obvious biological bases.⁵ Thus, in recent years psychiatry has gone through an identity crisis, and its scientific nature has even been questioned.⁶

The authors of this article fit pretty well with these stereotypes about psychiatrists. Currently working in community mental health, our last professional contact with general or internal medicine was many years ago, as first-year psychiatry residents. We had not touched a stethoscope (figuratively and almost literally) for a long time. All these shortcomings made the experience of the first days stressful and overwhelming. Whereas our fellow infectionists faced the novelty of an unknown disease and its dramatic consequences, for us everything was new. The critical situation of the hospital, on the verge of collapse, and the general climate of catastrophe possibly also influenced us. However, after a few days, we recovered from the initial shock. Stress began to shape us and our routines, in parallel to the society that, hit by COVID-19, initiated its own adaptive effort.⁷ Then we remembered that psychiatrists are also doctors, and we understood that the distinction between “medical” and “psychiatric” patients is useless. When we were able to return to our regular practice at the end of April, the redeployed psychiatrists shared the feeling of having achieved considerable clinical autonomy and that the experience, brief but very intense, had contributed to our professional development. Undoub-

tedly, the high protocolization of the clinical management of COVID-19 and, above all, being part of a multidisciplinary medical team, increased our perception of clinical accountability.⁸

Although, as mentioned, the balance of our experience was not negative, the SARS-Cov-2 pandemic has exposed gaps in the medical skills of the psychiatric community. It must be said, in our defense, that this is largely due to the high degree of specialization achieved in modern medicine. In fact, psychiatrists are not the only who have experienced difficulties during the COVID-19 outbreak, other specialties have also suffered from their lack of medical training when responding to the crisis.⁹ However, psychiatrists are often prone to despise the more medical part of the biopsychosocial model, perhaps because of the mistaken belief that a biological approach is not compatible with a humane, patient-centered management. Several proposals have been made to try to enhance general medicine skills during residency in psychiatry. Nevertheless, experienced psychiatrists should also continue to update regularly, and not lose touch with general medicine. Ideally, psychiatrists should acquire enough skills to be able to lead integrated health teams.¹⁰ Otherwise, the psychiatric community runs the risk of becoming irrelevant in major public health decisions, especially if it continues to strive to distance itself from major "physical" health problems. It is time for psychiatrists to react and prepare for an uncertain future of health crises, in which our knowledge and skills cannot be restricted to the field of our specialty.

Bibliografía

- Vieta E, Pérez V, Arango C. Psychiatry in the aftermath of COVID-19. *Rev Psiquiatr Salud Ment.* 2020, <http://dx.doi.org/10.1016/j.rpsm.2020.04.004>. S1888-9891(20)30029-X.
- Xiang YT, Yang Y, Li W, Zhang L, Zhang Q, Cheung T, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry.* 2020;7:228–9, [http://dx.doi.org/10.1016/S2215-0366\(20\)30046-8](http://dx.doi.org/10.1016/S2215-0366(20)30046-8).
- Ajaz A, David R, Brown D, Smuk M, Korszun A. BASH: bad-mouthing, attitudes and stigmatisation in healthcare as experienced by medical students. *BJPsych Bull.* 2016;40:97–102, <http://dx.doi.org/10.1192/pb.bp.115.053140>.
- Dornquast C, Tomzik J, Reinhold T, Walle M, Mönter N, Berghöfer A. To what extent are psychiatrists aware of the comorbid somatic illnesses of their patients with serious mental illnesses? A cross-sectional secondary data analysis. *BMC Health Serv Res.* 2017;17:162, <http://dx.doi.org/10.1186/s12913-017-2106-6>.
- Benrimoh D, Pomerleau VJ, Demoustier A, Poulin S, Maltais JR, Brouillette J, et al. Why we still use "organic causes": results from a survey of psychiatrists and residents. *J Neuropsychiatry Clin Neurosci.* 2019;31:57–64, <http://dx.doi.org/10.1176/appi.neuropsych.18050099>.
- de Leon J. Is psychiatry scientific? A letter to a 21st century psychiatry resident. *Psychiatry Investig.* 2013;10:205–17, <http://dx.doi.org/10.4306/pi.2013.10.3.205>.
- Crespo-Facorro B. Mental health and the SARS-CoV-2 pandemic. *Rev Psiquiatr Salud Ment.* 2020, <http://dx.doi.org/10.1016/j.rpsm.2020.04.010>. S1888-9891(20)30035-5.
- Molleman E, Broekhuis M, Stoffels R, Jaspers F. Consequences of participating in multidisciplinary medical team meetings for surgical, nonsurgical, and supporting specialties. *Med Care Res Rev.* 2010;67:173–93, <http://dx.doi.org/10.1177/1077558709347379>.
- Plachouri KM, Georgiou S. How well prepared are dermatologists redeployed to COVID-19 wards? *Int J Dermatol.* 2020;10, <http://dx.doi.org/10.1111/ijd.14978>, 1111/ijd.14978.
- Annamalai A, Rohrbaugh RM, Sernyak MJ. General medicine training in psychiatry residency. *Acad Psychiatry.* 2015; 39:437–41, <http://dx.doi.org/10.1007/s40596-015-0344-7>.

Jorge Gómez-Arnau*, Raúl González-Lucas,
Patricia Sánchez-Páez

Department of Psychiatry. Hospital Universitario Ramón y Cajal, Madrid, Spain

* Corresponding author.

E-mail address: jorge.gomezarnau@salud.madrid.org
(J. Gómez-Arnau).

<https://doi.org/10.1016/j.rpsm.2020.07.001>
1888-9891/ © 2020 SEP y SEPB. Published by Elsevier España, S.L.U. All rights reserved.