

## Clinical Study

# Vitamin D and Serum Cytokines in a Randomized Clinical Trial

**Eleanor Yusupov, Melissa Li-Ng, Simcha Pollack, James K. Yeh, Mageda Mikhail, and John F. Aloia**

Winthrop University Hospital, Bone Mineral Research Center, 222 Station Plaza North, Suite 350A, Mineola, NY 11501, USA

Correspondence should be addressed to Eleanor Yusupov, eleanor.rx@gmail.com

Received 27 March 2010; Revised 18 June 2010; Accepted 7 July 2010

Academic Editor: Vin Tangpricha

Copyright © 2010 Eleanor Yusupov et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** The role of vitamin D in the body's ability to fight influenza and URI's may be dependent on regulation of specific cytokines that participate in the host inflammatory response. The aim of this study was to test the hypothesis that vitamin D can influence intracellular signaling to regulate the production of cytokines. **Subjects and Methods.** This study was a 3-month prospective placebo-controlled trial of vitamin D3 supplementation in ambulatory adults [Li-Ng et al., 2009]. 162 volunteers were randomized to receive either 50  $\mu\text{g}/\text{d}$  (2000 IU) of vitamin D3 or matching placebo. 25(OH)D and the levels of 10 different cytokines (IL-2, 4, 5, 6, 8, 10, 13, GM-CSF, IFN- $\gamma$ , TNF- $\alpha$ ) were measured in the serum of participants at baseline and the final visit. There were 6 drop-outs from the active vitamin D group and 8 from the placebo group. **Results.** In the active vitamin D group, we found a significant median percent decline in levels of GM-CSF ( $-62.9\%$ ,  $P < .0001$ ), IFN- $\gamma$  ( $-38.9\%$ ,  $P < .0001$ ), IL-4 ( $-50.8\%$ ,  $P = .001$ ), IL-8 ( $-48.4\%$ ,  $P < .0001$ ), and IL-10 ( $-70.4\%$ ,  $P < .0001$ ). In the placebo group, there were significant declines for GM-CSF ( $-53.2\%$ ,  $P = .0007$ ) and IFN- $\gamma$  ( $-34.4\%$ ,  $P = .0011$ ). For each cytokine, there was no significant difference in the rate of decline between the two groups. 25(OH)D levels increased in the active vitamin D group from a mean of  $64.3 \pm 25.4$  nmol/L to  $88.5 \pm 23.2$  nmol/L. **Conclusions.** The present study did not show that vitamin D3 supplementation changed circulating cytokine levels among healthy adults.

## 1. Introduction

Vitamin D is produced in the skin when sunlight is absorbed. Thus, vitamin D levels, or serum 25-hydroxyvitamin D (25(OH)D), fluctuate seasonally. 25(OH)D levels are low during the winter in northern latitudes because of decreased amounts of sunlight. A conventional diet usually does not provide adequate amounts of vitamin D. Vitamin D insufficiency results in a number of skeletal and extraskeletal complications. It has been associated with decreased muscle strength [1], breast cancer [2, 3], colon cancer [4], cardiovascular disease [5], and autoimmune disorders such as type 1 diabetes [6], rheumatoid arthritis [7] and systemic lupus erythematosus [7]. Based on vitamin D's role in the basic innate immune defense mechanisms, it has been suggested as an adjuvant treatment of tuberculosis [8]. Levels of 25(OH)D, below 17.8 ng/ml are associated with a 26% increased risk of all-cause mortality in the general population even after adjusting for known CVD

risk factors, socioeconomic status, and other characteristics [9].

In vitro and in vivo studies show a role for vitamin D as an important component of the innate immune system. The innate immune system provides front-line protection against infectious agents. Expression of vitamin D receptor occurs in different cells of the myeloid and lymphoid lineage. The active form of vitamin D, 1,25-dihydroxyvitamin D (1,25(OH)<sub>2</sub>D), increases the production of endogenous antibiotics called antimicrobial peptides (AMP) in human monocytes, neutrophils, and epithelial cells [10]. AMPs such as defensin and cathelicidin have a broad range of actions against microorganisms, including bacteria, fungi, and viruses. Liu et al. showed that stimulation of toll-like receptors (TLR) 2/1 engages a vitamin D-dependent intracellular circuit that results in the expression of cathelicidin, enhancing the microbicidal capability of the monocyte [8]. Remarkably, the authors also observed that sera from African-American individuals, who are known

to have substantially lower serum 25(OH)D levels than whites [11], were inefficient in inducing genetic expression of cathelicidin. When the sera was supplemented with 25(OH)D, cathelicidin levels increased to levels observed in monocytes collected from whites. This suggests that vitamin D insufficiency during the winter may increase susceptibility to infections, particularly viral respiratory infections.

It is estimated that 72% of adults experience at least one URI per year and those adults experience an average of 2.5 URIs per year [12]. Every year 5% to 20% of the U.S. population get the flu [13]. Vitamin D may also play a role in reducing the severity of URI symptoms by regulating specific cytokines that participate in the host inflammatory response. Vitamin D's role in fighting influenza may be dependent on suppression of pro-inflammatory cytokines Interferon gamma, TNF- $\alpha$ , and IL-12 in the macrophage [14].

1,25(OH)<sub>2</sub>D has been shown to inhibit mononuclear and T lymphocyte cell proliferation by decreasing the production of IL-1 $\beta$ , IL-2, IL-6, interferon  $\gamma$  (IFN- $\gamma$ ), and TNF- $\alpha$  [15]. A randomized controlled trial by Schleithoff et al. showed that 50  $\mu$ g/day (2000 IU/day) of vitamin D3 reduced the inflammatory milieu in CHF patients [16].

1,25(OH)<sub>2</sub>D may play a protective role against inflammatory bowel disease as was shown in a mouse model of colitis. Tissue-specific alterations in the expression of cytokines IL-1 $\beta$ , IL-10, and IL-17 were demonstrated in Cyp27b1 knockout mice [17].

When reviewing the adverse events from our prior study of vitamin D3 supplementation in postmenopausal African American women [18], we noticed a significant difference in the reported incidence of URI symptoms between the active vitamin D group and the placebo group. This finding led us to conclude that higher doses of vitamin D supplementation may protect against viral URI's [19].

Current recommendations for vitamin D intake are based on amounts required to sustain optimal skeletal health. Schleithoff's study and our prior study suggest that optimal function of the innate immune system might require higher doses of vitamin D. In this study, we administered 50  $\mu$ g/day (2000 IU) of vitamin D3 which is the tolerable upper intake level (UL) for vitamin D for children and adults set by the Food and Nutrition Board of the Institute of Medicine [20]. Experts in the U.S. believe that higher intakes of vitamin D are necessary and that these higher intakes are safe [21, 22].

The role of vitamin D in the body's ability to fight influenza and URI's may be dependent on regulation of specific cytokines that participate in the host inflammatory response. The aim of this study was to test the hypothesis that vitamin D can influence intracellular signaling to regulate the production of cytokines. The levels of 10 different cytokines produced by T cells (Th1, Th2, regulatory T cells, NK cells) and macrophages were measured in the serum of participants at baseline and the final visit.

## 2. Patients and Methods

**2.1. Subjects.** Study participants were recruited from the Long Island, New York community (latitude, 40.7 degrees

N) between December 2006 and March 2007 (Figure 1 Flowchart). Volunteers were recruited from local newspaper advertisements, mailing of brochures to community residents, and flyers posted at Winthrop University Hospital medical offices. Patients were eligible for the study if they met the following criteria: ambulatory adult age 18–80 and stable medical condition with no change in medications for 6 months prior to study entry. Exclusion criteria included morbid obesity (body mass index >35 kg/m<sup>2</sup>); current tobacco use; history of hypercalcemia, nephrolithiasis or sarcoidosis; pregnancy; recent hospitalization; current liver or kidney disorders, malignancy and malabsorption; and use of immunosuppressants or medications that interfere with vitamin D metabolism such as phenytoin and carbamazepine. Race determination was by self-declaration. All participants provided written informed consent and the trial was approved by the institutional review board of Winthrop University Hospital. The results of this study have been recently reported [19].

**2.2. Study Design.** This study was a 3-month prospective, randomized, double-blind, placebo-controlled trial of vitamin D3 supplementation in ambulatory adults [19]. Recruitment began from December 2006 to March 2007 and the study was completed in June 2007. The participants were randomly assigned using a computer-generated randomization sequence to receive either 50  $\mu$ g/d of vitamin D3 or matching placebo. Each subject was sequentially assigned a number upon study entry and the investigators dispensed the corresponding sequentially numbered container of study medication to the subject. All participants and investigators were blinded throughout the study except for the research pharmacist and the statistician. Neither the statistician nor the research pharmacist had any contact with study participants. Eligible subjects underwent a baseline medical history, height and weight measurements, and blood tests. Subjects were seen at 6-weeks and 12-weeks postrandomization. Blood was collected again at the 12-week visit. 25(OH)D and the levels of 10 different cytokines (IL-2, 4, 5, 6, 8, 10, 13, GM-CSF, IFN- $\gamma$ , TNF- $\alpha$ ) were measured in the serum of participants at baseline and the final visit.

**2.3. Laboratory Tests.** Serum samples were analyzed by Ray Biotech Quantibody Human Th1/Th2 Array 1 kit using glass-chip-based multiplexed sandwich ELISA system. The signals are detected using fluorescence-based detection method. Baseline and study-end serum samples were analyzed in one assay. The well-to-well CV is <20%.

Serum 25(OH)D was measured by a radio-receptor assay from DiaSiorin, Inc (Stillwater, MN). The intraassay variability in our laboratory is 4.1% and interassay variability is 7.0%. Our laboratory participates in the international Vitamin D External Quality Assessment Program (<http://www.deqas.org/>). Vitamin D3 content was analyzed in an independent laboratory (Vitamin D, Skin, and Bone Research Laboratory, Department of Medicine, Boston University School of Medicine, Boston, MA).




FIGURE 1: Epidemiology and Infection, Volume 137, Issue 3, 2009 “A randomized controlled trial of vitamin D3 supplementation for the prevention of symptomatic upper respiratory tract infections” by John Aloia. © 2009 Cambridge University Press.

**2.4. Statistical Analysis.** The signed rank test was used to assess the median percent change of each cytokine level within the active vitamin D or placebo group during the 12-week period. The rank-sum test was used to compare median percent change between the active vitamin D and placebo groups. An estimate of the difference of the two medians (active vitamin D-placebo) and the corresponding 95% confidence interval was calculated using the software Confidence Interval Analysis (CIA), Version 2.0.0. We also pooled both groups together and calculated percent changes in 25(OH)D over 12 weeks. Spearman correlations were then calculated between these percent changes in 25(OH)D and percent changes of each cytokine level. Since analysis was done for 10 cytokines, we used the Bonferroni adjustment and results were considered significant at the  $P = .05/10 = .005$  level of significance. For all calculations (except for confidence interval analysis), we utilized SAS 9.2 for Windows, SAS Institute Inc., Cary, N.C.

### 3. Results

**3.1. Baseline Characteristics.** The baseline characteristics and laboratory values of the study population are summarized in Table 1 [19]. *T*-tests comparing active to placebo patients at baseline did not reveal any differences between the groups. The baseline 25(OH)D levels ranged from 16 to 156 nmol/L with a mean level of  $63.7 \pm 28.7$  nmol/L in the study population. At baseline, 23% of the active patients exceeded 75 nmol/L.

**3.2. Adherence.** Adherence (defined as the ratio of the number of pills consumed to the number of days in the study) ranged from 59% to 100%. Mean compliance was  $94\% \pm 9\%$ . Adherence did not significantly differ between the active and placebo groups.

**3.3. Cytokine Levels.** There were no statistically significant differences between the two groups with respect to median percent differences for each cytokine. The only exception was for IL-10, where the percent change for the active vitamin D group was  $-70.4\%$  versus only  $-49.6\%$  for placebo ( $P = .02$ ). However, the Bonferroni adjustment would need to be removed in order to obtain significance at  $P < .05$ .

In the active vitamin D group, we found a significant median percent decline in levels of GM-CSF ( $-62.9\%$ ,  $P < .0001$ ), IFN- $\gamma$  ( $-38.9\%$ ,  $P < .0001$ ), IL-4 ( $-50.8\%$ ,  $P = .001$ ), IL-8 ( $-48.4\%$ ,  $P < .0001$ ), and IL-10 ( $-70.4\%$ ,  $P < .0001$ ). In the placebo group, there were significant declines (but slightly smaller compared to the active vitamin D group) for GM-CSF ( $-53.2\%$ ,  $P = .0007$ ) and IFN- $\gamma$  ( $-34.4\%$ ,  $P = .0011$ ). For each cytokine, there was no significant difference in the rate of decline between the two groups. 25(OH)D levels increased in the active vitamin D group from a mean of  $64.3 \pm 25.4$  nmol/L to  $88.5 \pm 23.2$  nmol/L. At the end of the study, 73% of the active vitamin D group patients exceeded 75 nmol/L. Spearman correlation analysis showed no significant correlations for any of the 10 cytokines: the strongest correlations were  $r = -0.096$  ( $P = .30$ ) for IL-8 and  $r = 0.079$  ( $P = .39$ ) for IL-5.

TABLE 1: Baseline characteristics of study participants. There were no significant differences between the study groups at baseline.

Characteristic	Active ( <i>n</i> = 78)	Placebo ( <i>n</i> = 70)
*Age, y	59.3 ± 13.0	58.1 ± 13.4
BMI, kg/m <sup>2</sup>	26.1 ± 4.5	26.6 ± 4.1
**Male	17 (21.8%)	13 (18.6%)
Female	61 (78.2%)	57 (81.4%)
Race		
Caucasian	70 (89.7%)	61 (87.1%)
African American	3 (3.8%)	3 (4.3%)
Asian	2 (2.6%)	6 (8.6%)
Other	3 (3.8%)	0 (0.0%)
*25-OHD, nmol/L	64.3 ± 25.4	63.0 ± 25.8
PTH, pg/mL	29.2 ± 13.6	28.4 ± 12.6
History of tobacco use	26 (33.3%)	28 (40%)
History of asthma	6 (7.7%)	2 (2.9%)
History of COPD	3 (3.8%)	2 (2.9%)
Received flu vaccine	44 (56.4%)	45 (64.3%)
Dietary calcium intake (mg/d)	762.8 ± 375.7	854.6 ± 518.6
Dietary vitamin D intake (IU/d)	147.3 ± 182.3	168 ± 146.5

\*Values are expressed as mean ± SD.

\*\*Values are expressed as *n* (%).

Epidemiology and Infection, see [19] © 2009 Cambridge University Press.

Cytokine results as described above are summarized in Table 2. We provide in Table 3, for each cytokine, the difference of the median percent change (Active vitamin D-Placebo) and the corresponding 95% confidence interval. We note that for nine of the ten cytokines, the estimated difference of medians between the two groups was within ±10%, which would likely represent a clinically insignificant difference. The only exception was for cytokine IL-10, where the group difference was statistically significant ( $P = .02$  before the Bonferroni adjustment as previously mentioned). We also considered a subanalysis consisting of placebo group who are vitamin D insufficient (<75 nmol/l) both at the start and the end of the study versus. active vitamin D group who were insufficient at the start and became sufficient at the end of the study. These “modified” placebo and active vitamin D groups consisted of 33 and 30 patients, respectively. When we repeat the analysis done in Table 2, the rank sum  $P$ -values (comparing active vitamin D with placebo) are all nonsignificant, with  $P > .30$  for all 10 parameters.

**3.4. Adverse Events.** A total of 72 adverse events were reported in the study over 3 months, 38 in the active vitamin D group and 34 in the placebo group,  $P = .99$ . There was no significant difference in the adverse events between the study groups. There were 3 serious adverse events, one in the active vitamin D group and two in the placebo group. None of the serious adverse events were considered to be related to the study medication. There were no episodes of nephrolithiasis or hypercalcemia.

## 4. Discussion

The present study did not show an effect of vitamin D3 supplementation on the circulating levels of certain

cytokines. We predicted that the levels of IL-2, 6, IFN- $\gamma$ , TNF- $\alpha$ , and GM-CSF will decrease under the influence of vitamin D, while the levels of IL-4, 5, 10, and 13 were expected to rise [23–25]. We did not find that cytokine levels or changes in cytokine levels were responsive to vitamin D supplementation or related to serum 25(OH)D levels. However, there is accumulating evidence that vitamin D does significantly alter the immune system [26]. In gene chip experiments, the active form of vitamin D has been shown to downregulate gene expression of T-helper 1 cells, at the same time up regulating gene expression of T-helper 2 cells [24]. Decreased production of INF-gamma and IL-12 by T-helper 1 cells, and induction of regulatory T cells influenced by 1,25(OH)<sub>2</sub>D regulates immune responses to infections [23]. 1,25(OH)<sub>2</sub>D has been shown to induce the expression of IL-10 by CD40/IL-4-activated B-cells by binding of the VDR to the promoter of IL-10 [25].

The role of 1,25(OH)<sub>2</sub>D in innate and adaptive immunity has also been demonstrated in vivo. Following in vivo stimulation of murine dendritic cells by lipopolysaccharide (bacterial-derived endotoxin), they were found to produce 1-alpha-hydroxylase, resulting in production of active form of vitamin D [27]. Hansdottir et al. proposed a mechanism by which vitamin D plays a role in host defense by demonstrating the conversion of inactive vitamin D (25(OH)D) to active 1,25(OH)<sub>2</sub>D by enzyme 1alpha-hydroxylase in respiratory epithelial cells [28]. Impaired production of 1,25(OH)<sub>2</sub>D due to low-circulating 25(OH)D levels results in inhibition of VDR-dependent innate immune response [8].

1,25(OH)<sub>2</sub>D inhibits the production of IL-12, IFN- $\gamma$  and promotes production of IL-4 following endotoxin stimulation in rats [24]. Human B-cells experiments demonstrated the ability of 1,25(OH)<sub>2</sub>D to induce the expression of

TABLE 2: Laboratory values.

Cytokine	Median baseline value (in pg/ml)	Active ( <i>n</i> = 63)		Placebo ( <i>n</i> = 57)		Rank-sum <i>P</i> -value	
		Median percent change (12 weeks)	<i>P</i> -value (signed-rank test)	Median baseline value (in pg/ml)	Median percent change (12 weeks)		<i>P</i> -value (signed-rank test)
GM-CSF	5.91	-63.0	<.0001	4.78	-53.2	.0007	.53
IFN- $\gamma$	130.8	-38.9	<.0001	115.2	-34.4	.0011	.41
IL-2	2.13	-17.6	.70	2.15	-23.8	.52	.96
IL-4	6.10	-50.8	.001	8.26	-39.4	.006	.47
IL-5	4.88	-8.32	.49	4.53	-30.0	.66	.50
IL-6	2.19	-43.2	.37	2.75	-52.9	.0963	.61
IL-8	16.3	-48.4	<.0001	24.1	-33.0	.0210	.21
IL-10	6.82	-70.4	<.0001	5.12	-49.6	.0401	.02
IL-13	0.604	-59.4	.0104	0.500	-14.4	.22	.21
TNF- $\alpha$	5.64	-12.0	.21	7.60	-21.8	.0542	.64

TABLE 3: Median Difference of Percent Changes (Active vitamin D-Placebo).

Cytokine	Difference of MedianPercent Change (Active vitamin D-Placebo)	
		95% CI of Difference of Medians
GM-CSF	-3.86%	(-19.9%, 9.69%)
IFN- $\gamma$	-5.24%	(-18.6%, 8.47%)
IL-2	0.21%	(-31.4%, 26.9%)
IL-4	-5.59%	(-27.2%, 9.10%)
IL-5	8.62%	(-17.6%, 41.0%)
IL-6	4.14%	(-12.5%, 24.4%)
IL-8	-9.43%	(-27.2%, 5.15%)
IL-10	-17.0%	(-38.6%, -1.15%)
IL-13	-9.08%	(-48.7%, 4.71%)
TNF- $\alpha$	-5.67%	(-38.1%, 23.9%)

chemokine (C-C motif) receptor 10 (CCR10), which plays an important role in the mucosal immune system [29].

In patients with severe congenital neutropenia, 1,25(OH)<sub>2</sub>D induced the expression of the human cathelicidin antimicrobial peptide 18 precursor protein in myeloid precursors [30]. The process of activation of vitamin D in human tracheobronchial cells causes upregulation of cathelicidin antimicrobial peptide gene and the TLR coreceptor CD14, contributing to innate immunity in the lungs [28].

Data on vitamin D regulating serum cytokines in placebo-controlled clinical trials is very limited. In a 6-month double-blind, placebo-controlled, randomized trial in vitamin D-insufficient patients with MS, 25  $\mu$ g (1000 IU) vitamin D significantly increased serum levels of TGF-beta 1, while having no effect on TNF- $\alpha$ , IFN- $\gamma$ , and IL-13 [31]. Another randomized, placebo-controlled clinical trial in 123 CHF patients demonstrated that 50  $\mu$ g (2000 IU) of vitamin D, administered for 9 months, increased median serum levels of anti-inflammatory IL-10 by 43%, and suppressed the levels of proinflammatory TNF- $\alpha$  [16]. The median baseline 25(OH)D levels in that trial were significantly lower

compared to our study: 35.87 nmol/l in vitamin D group and 38.09 nmol/l in the placebo group. Seasonal differences in cytokine levels were observed by Stewart et al. among healthy controls, who noted lower winter levels of IL-4, IL-10 and TNF- $\alpha$ , and a winter excess of IFN- $\gamma$  [32]. These findings further support the hypothesis that seasonal differences in vitamin D influence the levels of cytokines and may help explain our finding of declining serum cytokines levels during winter months in the active vitamin D, as well as the control group.

The strengths of this study include the high compliance rate of medication intake (94%) and low-dropout rate. The mean 25(OH)D level of the subjects at baseline (63.7  $\pm$  28.7 nmol/L) was in accord with the mean 25(OH)D level in NHANES III (64.8 nmol/L) measured during the winter [33]. Our laboratory participates in the international Vitamin D External Quality Assessment Program which ensures the analytical reliability of 25(OH)D assays. Vitamin D content in the tablets was verified by an independent laboratory.

We recognize that this study has several limitations. 50  $\mu$ g/day may not be enough to stimulate innate immunity,

and/or the length of the study may not have been sufficient to observe an effect. Another reason why we may not have observed a benefit is that the mean 25(OH)D level at the beginning of the study was not that low but it was not different from the average level seen in NHANES III. Vitamin D repletion in subjects who were deficient at baseline may have resulted in meaningful changes in cytokine levels however, our study was not designed to address this question. 73% of patients in the vitamin D group achieved 25(OH)D levels greater than 75 nmol/L but the difference between baseline and end-of-study levels may not be enough to confer a benefit.

Although there is an increasing amount of evidence from animal experiments and human case-control studies that demonstrate the role of vitamin D in host defense, additional randomized clinical trials are needed [34].

## Acknowledgments

This research was partially funded by the Empire Clinical Research Investigator Program (ECRIP). The authors thank Jane Greensher for her expertise as the Nurse Coordinator and Martin Feuerman for contributing to the data and statistical analyses. They also thank Lynn Maier for preparation of the typescript.

## Conflict of Interests

The authors declare that there is no conflict of interests.

## References

- [1] M. Visser, D. J. H. Deeg, and P. Lips, "Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): the Longitudinal Aging Study Amsterdam," *Journal of Clinical Endocrinology and Metabolism*, vol. 88, no. 12, pp. 5766–5772, 2003.
- [2] E. R. Bertone-Johnson, W. Y. Chen, M. F. Holick et al., "Plasma 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D and risk of breast cancer," *Cancer Epidemiology Biomarkers and Prevention*, vol. 14, no. 8, pp. 1991–1997, 2005.
- [3] J. M. Lappe, D. Travers-Gustafson, K. M. Davies, R. R. Recker, and R. P. Heaney, "Vitamin D and calcium supplementation reduces cancer risk: results of a randomized trial," *American Journal of Clinical Nutrition*, vol. 85, no. 6, pp. 1586–1591, 2007.
- [4] D. M. Freedman, M. Dosemeci, and K. McGlynn, "Sunlight and mortality from breast, ovarian, colon, prostate, and non-melanoma skin cancer: a composite death certificate based case-control study," *Occupational and Environmental Medicine*, vol. 59, no. 4, pp. 257–262, 2002.
- [5] D. H. Kim, S. Sabour, U. N. Sagar, S. Adams, and D. J. Whellan, "Prevalence of hypovitaminosis D in cardiovascular diseases (from the National Health and Nutrition Examination Survey 2001 to 2004)," *American Journal of Cardiology*, vol. 102, no. 11, pp. 1540–1544, 2008.
- [6] E. Hypponen, E. Laara, A. Reunanen, M.-R. Järvelin, and S. M. Virtanen, "Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study," *The Lancet*, vol. 358, no. 9292, pp. 1500–1503, 2001.
- [7] M. Cutolo, K. Otsa, M. Uprus, S. Paolino, and B. Seriolo, "Vitamin D in rheumatoid arthritis," *Autoimmunity Reviews*, vol. 7, no. 1, pp. 59–64, 2007.
- [8] P. Liu, M. Hewison, and J. S. Adams, "Vitamin D and Innate Immunity," *Clinical Reviews in Bone and Mineral Metabolism*, vol. 7, pp. 176–184, 2009.
- [9] M. L. Melamed, E. D. Michos, W. Post, and B. Astor, "25-hydroxyvitamin D levels and the risk of mortality in the general population," *Archives of Internal Medicine*, vol. 168, no. 15, pp. 1629–1637, 2008.
- [10] T.-T. Wang, F. P. Nestel, V. Bourdeau et al., "Cutting edge: 1,25-dihydroxyvitamin D3 is a direct inducer of antimicrobial peptide gene expression," *Journal of Immunology*, vol. 173, no. 5, pp. 2909–2912, 2004.
- [11] S. S. Harris and B. Dawson-Hughes, "Seasonal changes in plasma 25-hydroxyvitamin D concentrations of young American black and white women," *American Journal of Clinical Nutrition*, vol. 67, no. 6, pp. 1232–1236, 1998.
- [12] A. M. Fendrick, A. S. Monto, B. Nightengale, and M. Sarnes, "The economic burden of non-influenza-related viral respiratory tract infection in the United States," *Archives of Internal Medicine*, vol. 163, no. 4, pp. 487–494, 2003.
- [13] Centers for Disease Control and Prevention, "Key facts about influenza and influenza vaccine," April 2009, <http://www.cdc.gov/flu/keyfacts.htm>.
- [14] J. J. Cannell, M. Zaslavoff, C. F. Garland, R. Scragg, and E. Giovannucci, "On the epidemiology of influenza," *Virology Journal*, vol. 5, article 29, 2008.
- [15] J. M. Lemire, "Immunomodulatory role of 1,25-dihydroxyvitamin D3," *Journal of Cellular Biochemistry*, vol. 49, no. 1, pp. 26–31, 1992.
- [16] S. S. Schleithoff, A. Zittermann, G. Tenderich, H. K. Berthold, P. Stehle, and R. Koerfer, "Vitamin D supplementation improves cytokine profiles in patients with congestive heart failure: a double-blind, randomized, placebo-controlled trial," *American Journal of Clinical Nutrition*, vol. 83, no. 4, pp. 754–759, 2006.
- [17] N. Liu, L. Nguyen, R. F. Chun et al., "Altered endocrine and autocrine metabolism of vitamin D in a mouse model of gastrointestinal inflammation," *Endocrinology*, vol. 149, no. 10, pp. 4799–4808, 2008.
- [18] J. F. Aloia, S. A. Talwar, S. Pollack, and J. Yeh, "A randomized controlled trial of vitamin D3 supplementation in African American women," *Archives of Internal Medicine*, vol. 165, no. 14, pp. 1618–1623, 2005.
- [19] M. Li-Ng, J. F. Aloia, S. Pollack et al., "A randomized controlled trial of vitamin D3 supplementation for the prevention of symptomatic upper respiratory tract infections," *Epidemiology and Infection*, vol. 137, no. 10, pp. 1396–1404, 2009.
- [20] Institute of Medicine, Food and Nutrition Board, *Dietary Reference Intakes: Calcium, Phosphorus, Magnesium, Vitamin D and Fluoride*, National Academy Press, Washington, DC, USA, 1999.
- [21] R. P. Heaney, K. M. Davies, T. C. Chen, M. F. Holick, and M. J. Barger-Lux, "Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol," *American Journal of Clinical Nutrition*, vol. 77, no. 1, pp. 204–210, 2003.
- [22] R. Vieth, H. Bischoff-Ferrari, B. J. Boucher et al., "The urgent need to recommend an intake of vitamin D that is effective," *American Journal of Clinical Nutrition*, vol. 85, no. 3, pp. 649–650, 2007.

- [23] M. T. Cantorna, S. Yu, and D. Bruce, "The paradoxical effects of vitamin D on type 1 mediated immunity," *Molecular Aspects of Medicine*, vol. 29, no. 6, pp. 369–375, 2008.
- [24] X.-P. Qi, P. Li, G. Li, Z. Sun, and J.-S. Li, "1,25-dihydroxyvitamin D3 regulates LPS-induced cytokine production and reduces mortality in rats," *World Journal of Gastroenterology*, vol. 14, no. 24, pp. 3897–3902, 2008.
- [25] G. Heine, U. Niesner, H.-D. Chang et al., "1,25-dihydroxyvitamin D3 promotes IL-10 production in human B cells," *European Journal of Immunology*, vol. 38, no. 8, pp. 2210–2218, 2008.
- [26] D. L. Kamen and V. Tangpricha, "Vitamin D and molecular actions on the immune system: modulation of innate and autoimmunity," *Journal of Molecular Medicine*, vol. 88, no. 5, pp. 441–450, 2010.
- [27] E. Y. Enioutina, D. Bareyan, and R. A. Daynes, "TLR ligands that stimulate the metabolism of vitamin D3 in activated murine dendritic cells can function as effective mucosal adjuvants to subcutaneously administered vaccines," *Vaccine*, vol. 26, no. 5, pp. 601–613, 2008.
- [28] S. Hansdottir, M. M. Monick, S. L. Hinde, N. Lovan, D. C. Look, and G. W. Hunninghake, "Respiratory epithelial cells convert inactive vitamin D to its active form: potential effects on host defense," *Journal of Immunology*, vol. 181, no. 10, pp. 7090–7099, 2008.
- [29] A.-K. Shirakawa, D. Nagakubo, K. Hieshima, T. Nakayama, Z. Jin, and O. Yoshie, "1,25-dihydroxyvitamin D3 induces CCR10 expression in terminally differentiating human B cells," *Journal of Immunology*, vol. 180, no. 5, pp. 2786–2795, 2008.
- [30] J. Karlsson, G. Carlsson, O. Larne, M. Andersson, and K. Pütsep, "Vitamin D3 induces pro-LL-37 expression in myeloid precursors from patients with severe congenital neutropenia," *Journal of Leukocyte Biology*, vol. 84, no. 5, pp. 1279–1286, 2008.
- [31] B. D. Mahon, S. A. Gordon, J. Cruz, F. Cosman, and M. T. Cantorna, "Cytokine profile in patients with multiple sclerosis following vitamin D supplementation," *Journal of Neuroimmunology*, vol. 134, no. 1-2, pp. 128–132, 2003.
- [32] N. Stewart, B. Taylor, A.-L. Ponsonby et al., "The effect of season on cytokine expression in multiple sclerosis and healthy subjects," *Journal of Neuroimmunology*, vol. 188, no. 1-2, pp. 181–186, 2007.
- [33] A. C. Looker, B. Dawson-Hughes, M. S. Calvo, E. W. Gunter, and N. R. Sahyoun, "Serum 25-hydroxyvitamin D status of adolescents and adults in two seasonal subpopulations from NHANES III," *Bone*, vol. 30, no. 5, pp. 771–777, 2002.
- [34] N. M. Maalouf, "The noncalcitropic actions of vitamin D: recent clinical developments," *Current Opinion in Nephrology and Hypertension*, vol. 17, no. 4, pp. 408–415, 2008.