



Omega-3 Fatty Acids Could Alleviate the Risks of Traumatic Brain Injury – A Mini Review

Parvathy R. Kumar^{1,2}, Musthafa Mohamed Essa^{1,2}, Samir Al-Adawi^{2,3}, Ghazi Dradekh^{1,2},
Mushtaq A. Memon⁴, Mohammed Akbar⁵, Thamilarasan Manivasagam⁶

¹Department of Food Science and Nutrition, College of Agricultural and Marine Sciences, Sultan Qaboos University, Sultanate of Oman.

²Ageing and Dementia Research Group, Sultan Qaboos University, Sultanate of Oman.

³College of Medicine and Health Sciences, Sultan Qaboos University, Sultanate of Oman.

⁴College of Veterinary Medicine, Washington State University, Pullman, WA, USA.

⁵Laboratory of Molecular Signaling, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Rockville, MD, USA.

⁶Department of Biochemistry and Biotechnology, Annamalai University, Tamil Nadu, India.

ABSTRACT

Traumatic brain injury (TBI) is an acquired brain trauma that occurs when any sudden trauma/injury causes damage to the brain. TBI is characterized by tissue damage and imbalance in the cerebral blood flow and metabolism. It has been established through laboratory experiments that the dietary supplementation of omega-3 fatty acids (FAs) could reduce the oxidative stress developed in brain due to TBI. The inclusion of omega-3 FA in diet could normalize the levels of brain-derived neurotrophic factor (BDNF), and thus, it could restore the survival of neuronal cells. BDNF improves the synaptic transmission by regulating synapsin 1 and cyclic adenosine monophosphate (cAMP) response element binding protein. The brain tissue analysis of TBI models supplemented with omega-3 polyunsaturated fatty acids (PUFAs) showed significantly reduced lipid peroxidation, nucleic acid and protein oxidation, thereby promoting neuronal and glial cell survival. Thus, omega-3 FA intake could be considered as a therapeutic option to reduce the secondary neuronal damages initiated by TBI.

Key words: Brain trauma, Neuronal damage, Omega-3 fatty acids, Oman, Traumatic brain injury

INTRODUCTION

Traumatic brain injury (TBI) is an acquired brain trauma that occurs when any sudden trauma/injury causes damage to the brain. An external mechanical force could displace the brain inside the skull, causing injury against the dura or neurocranium. Even the blood vessels in brain and nervous tissues could get disrupted. The impact of trauma ranges from mild to moderate to severe TBI.^[1]

The mortality indices vary from 1% in mild TBI to 2-5% in moderate TBI and up to 30-50% after suffering a severe head injury. According to the report of Centers for Disease Control and Prevention, about 50,000 individuals die every year from traumatic brain and twice this number suffer from permanent disabilities in the US.^[2] The treatment opportunities are much limited in reversing the effects of TBI. It has been reported that omega-3 fatty acids (FAs) are effective natural agents in reducing the neuronal damage, and reducing the brain oxidative stress.

Correspondence to:

Dr. Musthafa Mohamed Essa, Department of Food Science and Nutrition, P. O. 34, CAMS, Sultan Qaboos University, Al-Khoud, Muscat, P. C. 123, Sultanate of Oman. Tel: +968 2414 3604; Fax: +968 2441 3418; E-mail: drmdessa@gmail.com

DOI: 10.4103/2225-4110.130374

FATAL SECONDARY EFFECTS OF TBI

The initial stages of cerebral injury after TBI are indicated by tissue damage and imbalance in the cerebral blood flow and metabolism. These irregular patterns can lead to increased accumulation of lactic acid due to anaerobic glycolysis, with increased membrane permeability causing more fluid retention and edema formation.^[3] As a result of these events, there is depletion of ATP reserves, as the cellular energy requirements become inadequate with the anaerobic glycolysis. This ATP depletion, in turn, disrupts the membrane ion pumps. Therefore, the secondary stage of the TBI is initiated, which is characterized by the depolarization of terminal membrane leading to the release of excitatory neurotransmitters like glutamate. The Ca^{2+} and Na^+ influx becomes high due to the impaired ion pumps. The excess Ca^{2+} causes activation of lipases, proteases, and lipid peroxidases, and produces free FA and free radicals. Moreover, the activated caspases and endonucleases initiate the degradation of biomembranes and the nucleosomal DNA. This eventually leads to programmed cell death of neuronal cells. The oxidative stress formed in response to TBI generates superoxides, NO, H_2O_2 , and peroxynitrite. These events lead to the peroxidation of cellular structures, proteins, and vascular systems in the brain, and the cleavage of DNA, and finally inhibition of electron transport chain (ETC) in the mitochondria. All these factors sum up for the apoptosis.^[3-5] All the above factors are given in Figure 1 as




Figure 1. Schematic representation of possible damage occurring after TBI

a flowchart. The therapeutic options are very much limited in reversing the impairments initiated as a result of TBI.

IMPORTANCE OF OMEGA-3 FAs

Natural products and their active nutrients/components often cause possible improvement in most of the brain-related damages.^[6] Omega-3 FAs are polyunsaturated fatty acids (PUFAs) having double bond at the third position from the end of the FA chain. They are highly essential for maintaining the membrane fluidity and are present in all tissues, but are rich particularly in retina, brain, and spermatozoa. There are three important physiological omega-3 FAs available, including alpha linolenic acid (ALA), eicosapentaenoic acid (EPA), and docosahexaenoic acid (DHA). Dietary omega-3 FAs are the precursors for the FA components of neuronal membrane and are essential for maintaining the structural balance of cell membranes, signaling, neurotransmission, and modulation of enzymatic activity.^[7] Inadequate intake of omega-3 FAs results in inefficiency in learning and memory.^[8,9] Brain tissue analysis of TBI models supplemented with omega-3 PUFAs showed significant reduction in the lipid peroxidation and nucleic acid and protein oxidation.^[10]

POSSIBLE EFFECTS OF OMEGA-3 FAs IN RESCUING THE TBI SURVIVORS

As described earlier, brain injury may cause severe neuronal damage and consequent impairment in memory, learning, and motor coordination, and supplementation of omega-3 FAs might be able to reduce the neuronal damage. Neural trauma is usually associated with irregular phospholipid metabolism of neuronal membrane.^[11,12] Omega-3FAs are the major constituents of the membrane phospholipids, which suggests that supplementation of PUFAs could help in reducing the irregular phospholipid metabolism that occurs during neuronal damage. Supplementation of DHA could enhance the gene expression, release of neurotransmitters, and restoration of synaptic activity, which are all deleteriously affected by the TBI.^[13,14] The striatal dopamine levels reduced due to TBI could be normalized with the oral administration of fish oil.^[15] The PUFAs are capable of inhibiting arachidonic acid catabolism and maintaining the levels of cytokines, and have a strong antioxidant potential, and thus, they are effective as strong anti-inflammatory factors.^[16] Ying *et al.*, from Pinnilla's group have manifested that induced Omega-3 deficiency during gestation enhances the susceptibility to TBI in rat pups.^[17] Pinnilla's lab has also studied the effects of Omega-3FA supplementation and the action of progesterone in early life in the fluid percussion injury induced TBI rat models. They have demonstrated that Omega-3FA and progesterone are good candidates in counteracting the neuronal plasticity impairments.^[18] A recent report showed that omega-3 FA intake could normalize the levels of brain-derived neurotrophic factor (BDNF), synapsin 1, and cyclic adenosine monophosphate (cAMP)-responsive element binding protein, reduce the oxidative damage, and improve memory and learn-

ing abilities.¹¹⁹ BDNF is required for the synaptic transmission and the survival of neuronal cells. The authors of this study have demonstrated that omega-3 FAs could increase the levels of BDNF in TBI experimental models, thereby possibly activating the downstream targets. Furthermore, could reduce the deleterious effects of oxidative stress generated due to trauma.¹¹⁹ Neuronal survival is regulated by the phosphatidylinositol kinase and Akt signaling pathway.¹²⁰ It has been demonstrated that DHA could activate Akt and helps in the nuclear translocation. Thus, it initiates the Akt signaling pathway. So, DHA intake could have a positive effect on neuronal survival.¹²¹ Another group have established through their experiments that the EPA and DHA supplementation after the injury could ameliorate the traumatic axonal injury.¹²² Eady *et al.*¹²³ established through their work that DHA could induce cell survival, modulate neuro-inflammation, and restore synaptic circuits. A recent case study reported that a critical traumatic brain injured survivor showed improvement from its fatal effects by the enteral ingestion of omega-3 FAs.¹²⁴ All the effects are presented as a flowchart in Figure 2 and the experimental evidences are summarized in Table 1.




Figure 2. Possible support offered by omega-3 fatty acids in TBI

Table 1. Summary of beneficial effects of omega-3 FAs in Experimental TBI models

Omega-3FA dosage	Models	Effect	Reference
DHA enriched diet (1.2% DHA)	Rats	Improved cognition and synaptic plasticity	Wu <i>et al.</i> , 2011
Diet with 0.9% DHA +1%EPA	Rats	Improved neurotransmission and reduced oxidative stress	Wu <i>et al.</i> , 2004
Diet with 3.1% ALA	Rats	Neuronal cell survival	Akbar <i>et al.</i> , 2005
Regular diet + 1.5 ml fish oil daily (360 mg EPA and 240 mg DHA).	Rats	Maintains the neurotransmitter, dopamine levels	Samuel <i>et al.</i> , 2011
Diet+EPA+DHA (24 mg each/Kg of bodyweight)	Rats	Reduce axonal injury	James <i>et al.</i> , 2011
Diet with DHA (5 mg/kg of bodyweight)	Rats	Neuronal cell survival and neuroprotection	Eady <i>et al.</i> , 2012

TBI: Traumatic brain injury, DHA: Docosahexaenoic acid, ALA: Alpha linolenic acid

CONCLUSION

The role of PUFAs in the neuronal growth and maintenance is well documented. The omega-3 FA intake could lessen the secondary effects of the trauma in brain. The beneficial effects of omega-3 FAs suggest the importance of considering their supplementation as a therapeutic option for TBI survivors. Further studies are warranted to determine the effective dose for positive effects against TBI.

ACKNOWLEDGMENTS

The project was supported by a research grant from Research Council, Oman (Grant # RC/AGR/FOOD/11/01) to EMM, which is gratefully acknowledged. The PhD student fellowship offered to PVR by The Research Council, Oman (Grant # RC/AGR/FOOD/11/01) is highly acknowledged.

REFERENCES

- Thurman D, Guerrero J. Trends in hospitalization traumatic brain injury. *JAMA* 1999;282:954-7.
- Available from: <http://www.cdc.gov/traumaticbraininjury/Page> last reviewed: March 6, 2014.
- Shao C, Roberts KN, Markesbery WR, Scheffs W, Lovell MA. Oxidative stress in head trauma imaging. *Free Radic Biol Med* 2006;41:77-85.
- Bayir H, Kagan VE, Borisenko GG, Tyurina YY, Janesko KL, Vagni VA, *et al.* Enhanced oxidative stress in iNOS-deficient mice after traumatic brain injury: Support for a neuroprotective role of iNOS. *J Cereb Blood Flow Metab* 2005;25:673-84.
- Chong ZZ, Li F, Maiese K. Oxidative stress in the brain: Novel cellular targets that govern survival during neurodegenerative disease. *Prog Neurobiol* 2005;75:207-46.
- Essa MM, Vijayan RK, Castellano-Gonzalez G, Memon MA, Braidy N, Guillemin GJ. Neuroprotective effect of natural products against Alzheimer's disease. *Neurochem Res* 2012;37:1829-42.
- du Bois TM, Deng C, Bell W, Huang XF. Fatty acids differentially affect serotonin receptor and transporter binding in the rat brain. *Neuroscience* 2006;139:1397-403.
- Yamamoto N, Saitoh M, Moriuchi A, Nomura M, Okuyama H. Effect of dietary α -linolenate/linoleate balance on brain lipid compositions and learning ability of rats. *J Lipid Res* 1987;28:144-51.
- Lamprey MS, Walker BL. A possible essential role for dietary linolenic acid in the development of the young rat. *J Nutr* 1976;106:86-93.
- Michael-Titus AT. Omega-3 fatty acids and neurological injury. *Prostaglandins Leukot Essent Fatty Acids* 2007;77:295-300.
- Florent S, Malaplate-Armand C, Youssef I, Kriem B, Koziel V, Escanyé MC, *et al.* Docosahexaenoic acid prevents neuronal apoptosis induced by soluble amyloid- β oligomers. *J Neurochem* 2006;96:385-395.
- Klein J. Membrane breakdown in acute and chronic neurodegeneration: Focus on choline-containing phospholipids. *J Neural Transm* 2000;107:1027-63.
- Sullivan PG, Rabchevsky AG, Waldmeier PC, Springer JE. Mitochondrial permeability transition in CNS trauma: Cause or effect of neuronal cell death? *J Neurosci Res* 2005;79:231-9.
- Wu A, Ying Z, Gomez-Pinilla F. The salutary effects of DHA dietary supplementation on cognition, neuroplasticity, and membrane homeostasis after brain trauma. *J Neurotrauma* 2011;28:2113-22.
- Shin SS, Dixon CE. Oral fish oil restores striatal dopamine release after traumatic brain injury. *Neuroscience Letters*.2011;496:168-71.
- Farooqui AA, Horrocks LA, Farooqui T. Modulation of inflammation in

- brain: A matter of fat. *J Neurochem* 2007;101:577-99.
17. Ying Z, Agrawal R, Zhuang Y, Gomez-Pinilla F. Dietary Omega-3 Deficiency from Gestation Increases Spinal Cord Vulnerability to Traumatic Brain Injury-Induced Damage. *Plos One*. 2012; 7:e52998.
 18. Tyagi E, Agrawal R, Ying Z, Gomez-Pinilla F. TBI and sex : Crucial role of progesterone protecting the brain in an omega-3 deficient condition. *Experimental Neurology*. 2014;253:41-51.
 19. Wu A, Ying Z, Gomez-Pinilla F. Dietary omega-3 fatty acids normalize BDNF levels, reduce oxidative damage, and counteract learning disability after traumatic brain injury in rats. *J Neurotrauma* 2004;21:1457-67.
 20. Dudek H, Datta SR, Franke TF, Birnbaum MJ, Yao R, Cooper GM, *et al.* Akt. *Science* 1997;275:661-5.
 21. Akbar M, Calderon F, Wen Z, Kim KY. Docosahexaenoic acid: A positive modulator of Akt signaling in neuronal survival. *Proc Natl Acad Sci* 2005;102:10858-63.
 22. Mills JD, Bailes JE, Sedney CL, Hutchins H, Sears B. Omega-3 fatty acid supplementation and reduction of traumatic axonal injury in a rodent head injury model. *J Neurosurg*. 2011;114:77-84.
 23. Eady TN, Belayev L, Khoutorova L, Atkins KD, Zhang C, Bazan NG. Docosahexaenoic acid signaling modulates cell survival in experimental ischemic stroke penumbra and initiates long-term repair in young and aged rats. *PLoS One* 2012;7:e46151.
 24. Lewis M, Ghassemi P, Hibbeln J. Therapeutic use of omega-3 fatty acids in severe head trauma. *Am J Emerg Med* 2013;31:273.e5-8.