



MEETING ABSTRACT

Open Access

Reproducibility of neutrophil percentages in pooled and non-pooled nasal lavage samples

Dominik A Nowak^{1*}, Penelope Ferrie², Paul K Keith²

From Canadian Society of Allergy and Clinical Immunology Annual Scientific Meeting 2011
Quebec, Canada. 20-23 October 2011

Background

Neutrophils form a host's main immediate immune response. COPD, ozone, infection and other factors have been shown to increase neutrophils found in blood, sputum and nasal lavage. Nasal lavage is used to collect cells and inflammatory mediators from the nasal cavity, and can be used to quantify neutrophilic response to inhaled stimuli.

Objective

To compare the reproducibility of neutrophil percentage in a single sample lavage, SSL, versus multiple sample lavage, MSL (3 lavages performed 15 minutes apart and pooled).

Methods

Randomized crossover trial of nasal lavage performed on 4 visits 7-10 days apart, alternating between SSL and MSL done in 7 subjects with perennial allergic rhinitis, 7 with bilateral nasal polyposis and 7 controls.

Results

The mean (\pm SEM) neutrophil percentage was not significantly different between the two methods (80 ± 12 for SSL, 86 ± 8 for MSL, $p=0.2$). At a minimum total cell count (TCC) cutoff of ≥ 20 , the neutrophil percentage intraclass correlation (ICC) was similar but not sufficiently reproducible for SSL at 0.588 compared to MSL at 0.641. For samples with $TCC \geq 100$, the neutrophil percentage ICC of SSL was 0.93 (excellent correlation) and that of MSL was 0.676 (satisfactory). The evaluable samples for $TCC \geq 100$ were 60/84 for SSL and 67/84 for MSL. We previously demonstrated in the same experiment that a minimal TCC cutoff of ≥ 100 cells gave excellent eosinophil percentage ICC for both methods (>0.8).

Conclusion

SSL ICC was superior to MSL in measuring nasal lavage neutrophil percentage. This method could be used to assess the effects of inhaled stimuli on the nasal cavity.

Author details

¹University of Toronto Mississauga, Mississauga, Ontario, Canada, L5L 1C6.

²McMaster University, Hamilton, Ontario, Canada, L8S 4L8.

Published: 14 November 2011

doi:10.1186/1710-1492-7-S2-A26

Cite this article as: Nowak et al.: Reproducibility of neutrophil percentages in pooled and non-pooled nasal lavage samples. *Allergy, Asthma & Clinical Immunology* 2011 **7**(Suppl 2):A26.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹University of Toronto Mississauga, Mississauga, Ontario, Canada, L5L 1C6
Full list of author information is available at the end of the article