


Article

Convergent evolution of kidney sizes and supraorbital salt glands for birds living in saline habitats



Chi-Cheng Chiu,
Cheng-Te Yao,
Ben-Yang Liao,
Shou-Hsien Li

liaooby@nhri.edu.tw (B.-Y.L.)
t43028@ntnu.edu.tw (S.-H.L.)

Highlights

Birds have independently evolved larger kidneys and salt glands multiple times

Kidney sizes and salt glands in birds correlate positively with habitat salinity

Habitat salinity constrains the evolution of bird kidney sizes and salt glands

Chiu et al., iScience 27, 109169
March 15, 2024 © 2024
<https://doi.org/10.1016/j.isci.2024.109169>



Article

Convergent evolution of kidney sizes and supraorbital salt glands for birds living in saline habitats

Chi-Cheng Chiu,¹ Cheng-Te Yao,² Ben-Yang Liao,^{3,*} and Shou-Hsien Li^{1,4,*}

SUMMARY

Only a small number of avian species inhabit salty environments. To understand how they adapted, we examined the evolution of kidney sizes, supraorbital salt glands (SSGs), and the utilization of salty habitats across 230 species spanning 25 avian orders. Phylogenetic analysis indicates that SSGs, large kidneys, and thriving in salty habitats emerged convergently in birds. Transition rate analysis reveals that species possessing SSGs and large kidneys tended to move from low-to high-salinity environments, while others moved in the opposite direction. However, habitat salinity also influenced kidney evolution; lineages residing in high-salinity environments tended to develop larger kidneys than those in low-salinity environments. Our findings suggest that SSGs and large kidneys may have evolved through adaptation to high salinity. Overall, habitat conditions and physiological traits influenced avian adaptation to salty environments in a reciprocal manner. These results shed the new light on the evolutionary mechanisms underlying functional diversity in birds.

INTRODUCTION

Oceans provide vast space and rich resources for organisms. However, organisms must overcome the osmotic challenge of saltwater intake to live in such environments. Therefore, less than 3% of avian species (i.e., seabirds) live entirely or partially in high-osmolarity environments.¹

Supraorbital salt glands (SSGs, salt glands) in birds, also known as glandulae nasalis,^{2,3} produce concentrated solutions primarily consisting of NaCl or KCl, which help maintain proper internal solute levels while minimizing water loss.^{4–6} Glands with a similar function might have evolved, independently and convergently in reptiles such as turtles, sea snakes, lizards, and crocodiles.^{7,8} These glands are widespread and found in various parts of the skull across different evolutionary lineages, although their homology remains to be examined. Moreover, the unique positioning of the salt glands above the eye is a distinctive feature of birds.^{3,4} Supraorbital salt glands have been discovered in at least 40 bird families, which cover almost all traditional bird orders except for the Passeriformes.⁸ These glands are particularly common in marine birds such as gulls, petrels, albatrosses, auks, and penguins but can also be found in some freshwater species, such as dabbling ducks, mallards, and rails.⁹ Desert-dwelling birds such as ostriches and North African partridges^{1,8} and some carnivorous birds with high-protein diets such as Tawny Eagles¹⁰ also possess supraorbital salt glands. Avian supraorbital salt glands display significant levels of plasticity in both their structure and function,^{1,6,11,12,44} but their size and excretory capacity largely depend on the species' environmental salinity and diet.^{13,14}

Birds also use their kidneys to maintain the osmotic balance of their blood.¹⁵ The avian kidney comprises units of lobule with a renal medulla containing loops of Henle able to extract water from urine with higher salt concentrations than in the plasma. It has been shown that the mass of the kidney is positively correlated with the number of medullary cones, the smaller units of a renal medulla, which are associated with its ability to regulate osmotic pressure.¹⁵ How birds physiologically maintain their osmotic balance has been studied intensively.¹⁴ However, we know little about how the avian physiological traits associated with osmoregulation, such as the SSGs and large kidneys, have evolved in response to the types of habitats they use.

Non-passerine birds with SSGs tend to have larger kidneys (compared with their body sizes) than others.¹ Hughes, therefore, proposed that birds with high-salinity tolerance have larger kidneys for better osmoregulation. However, shared ancestry among species can lead to a significant association between two traits evolving independently.¹⁹ Without controlling for such “phylogenetic signals” (as Hughes did not¹), we do not know whether high-salinity environments could have driven these two physiological traits to evolve in birds. Independent evolution of similar traits, convergent evolution, in the same environmental conditions, has long been considered convincing evidence of adaptation.⁵⁶ If a salty environment could drive the evolution of a larger kidney and the presence of an SSG, could it lead to the convergent evolution of the

¹School of Life Science, National Taiwan Normal University, Taipei 11677, Taiwan

²High altitude research station, Taiwan Endemic Species Research Institute, Nantou 55244, Taiwan

³Institute of Population Health Sciences, National Health Research Institutes, Zhunan 350, Taiwan

⁴Lead contact

*Correspondence: liaoby@nhri.edu.tw (B.-Y.L.), t43028@ntnu.edu.tw (S.-H.L.)

<https://doi.org/10.1016/j.isci.2024.109169>






Figure 1. Estimating ancestral states and reconstructing traits in avian species: habitat salinity, presence of SSGs, and kidney sizes

Estimated ancestral states for the level of habitat salinity (A) and the presence of SSGs (B) for the 230 avian species studied and kidney sizes for the 167 avian species studied (C). The characteristics of each species are shown at the tips of the phylogenetic tree. The posterior probability of the reconstructed characteristic state of each trait is shown in the pie on each ancestral node. The gray and black bars next to the phylogenetic tree indicate the order of the species. The color patterns at the rightmost indicate the characteristic state of the extant species.

larger kidneys and SSGs in different avian lineages? It is also unclear whether the evolution of the two physiological traits was correlated or independent in birds.

RESULTS

Convergent evolution of larger kidney sizes, presence of supraorbital salt glands, and the use of high salinity habitat

In our dataset, species in completely or mainly saline habitats usually have large kidneys (residual kidney weight ≥ 0.13 , 67%) and SSGs (91%); in contrast, few species in mainly and completely freshwater habitats have “large kidneys” (88% had residual kidney weight < 0.13), and most (80%) do not have SSGs. Ancestral characteristics reconstruction suggested that the ancestor of birds probably lived in freshwater environments, had no SSG, and had medium-sized kidneys (the probabilities of having “non-large kidneys” and of having “large-kidneys” are roughly equal) (Figure 1).

Interestingly, our results suggest that the birds’ utilization of high-salinity habitats evolved together with the growth of their kidneys and the development of SSGs. The use of high-salinity habitats (sum of posterior probability for completely or mainly saline habitats > 0.5) occurred independently at least 14 times in 12 orders of birds (Anseriformes, Charadriiformes, Coraciiformes, Gaviiformes, Gruiformes, Phoenicopteriformes, Podicipediformes, Pelecaniformes, Passeriformes, Suliformes and Sphenisciformes); SSGs evolved independently 8 times in 14 orders of birds (Accipitriformes, Anseriformes, Charadriiformes, Falconiformes, Galliformes, Gaviiformes, Gruiformes, Pelecaniformes, Phoenicopteriformes, Podicipediformes, Procellariiformes, Sphenisciformes, Struthioniformes and Suliformes) (Figure 1). Intriguingly, both the utilization of high-salinity habitats and the presence of SSGs evolved in the same nodes for at least four avian lineages (Columbea, Aequornithes, and part of Galloanserae and Charadriiformes). Furthermore, SSGs are present in the common ostrich (*Struthio camelus*), the 13 Accipitriformes species, and 5 Falconiformes species that do not utilize high-salinity habitats. In contrast, four species in Passeriformes, the ash-throated flycatcher (*Myiarchus cinerascens*), the marsh wren (*Cistothorus palustris*), the snow bunting (*Plectrophenax nivalis*), and the Savannah sparrow (*Passerculus sandwichensis*) and one species in Coraciiformes, the belted kingfisher (*Megaceryle alcyon*), have evolved to utilize high-salinity habitats without having SSGs.

The use of high-salinity habitats and the presence of SSGs are evolutionarily inert in birds. Only five lineages, the pin-tailed duck (*Anas acuta*), the redhead (*Aythya Americana*), the greater scaup (*Aythya marila*), the upland sandpiper (*Bartramia longicauda*), and the long-billed dowitcher (*Limnodromus scolopaceus*), have evolved from ancestors utilizing high salinity habitats to descendants using freshwater habitats. Only four lineages, the white stork (*Ciconia Ciconia*), the African fish eagle (*Haliaeetus vocifer*), the American kestrel (*Falco sparverius*), and the Robert Falco (*Falco jugger*) have lost SSGs possessed by their ancestors.




Figure 2. The proportion of avian species living in completely saline, mainly saline, mainly freshwater, and completely freshwater habitats with and without supraorbital salt glands

By contrast, the evolution of birds' kidney size has been relatively labile. Large kidneys evolved at least 25 times in twelve orders of birds (Struthioniformes, Anseriformes, Galliformes, Phoenicopteriformes, Podicipediformes, Gaviiformes, Gruiformes, Pelecaniformes, Suliformes, Charadriiformes, Coraciiformes, and Passeriformes). In addition, lineages derived from those with large kidneys have 16 times reverted to having residual kidney weight <0.13 , namely the mallard duck (*Anus platyrhynchos*), the pin-tailed duck (*A. acuta*), the redhead (*Aythya americana*) in the non-Passeriformes, and the horned lark (*Eremophila alpestris*), the American bushtit (*Psaltriparus minimus*), the cedar waxwing (*Bombycilla cedrorum*), the cactus wren (*Campylorhynchus brunneicapillus*), the northern mockingbird (*Mimus polyglottos*), the Bendire's thrasher (*Toxostoma bendirei*), the zebra finch (*Taeniopygia guttata*), the chipping sparrow (*Spizella passerina*), the grey-cheeked thrush (*Catharus minimus*), the russet nightingale-thrush (*C. occidentalis*), the hermit thrush (*C. guttatus*), the ancestors of corvoidea, and all species of passeroidea in Passeriformes. However, birds in other avian lineages re-acquired large kidneys. These include the Eurasian magpie (*Pica pica*) in corvoidea, the painted redstart (*Myioborus pictus*), Bullock's oriole (*Icterus bullockii*), Brewer's blackbird (*Euphagus cyanocephalus*), the rufous-crowned sparrow (*Aimophila ruficeps*), the Savannah sparrow, and Lincoln's sparrow (*Melospiza lincolni*) in passeroidea, Carduelinae and a "masked" clade (*Piranga*, *Cardinalis*, and *Caryothraustes*) in Cardinalidae. It is worth noting that birds adapted to high-salinity habitats have all evolved at least one characteristic assumed to be related to osmosis except the snow bunting (*Plectrophenax nivalis*).

In addition, the results of search.conv clarify that when birds inhabit saline environments, the kidney size and SSGs in different lineages tend to be similar (found highly significant under "state," $p = 0.001$). Furthermore, birds' habitat diversity within each clade is diverse. Different clades do not exhibit convergent evolution toward the same direction (found no significance under "clade," $p = 0.117$).

Correlated evolution of large kidneys and functional supraorbital salt glands

Our results show high phylogenetic signals in levels of habitat salinity ($\lambda = 0.78$, $p \ll 0.001$; $K = 0.26$, $p = 0.001$; null hypothesis λ and $K = 0$), the presence of SSGs ($\lambda = 1$, $p \ll 0.001$; $K = 1.373$, $p = 0.001$; null hypothesis λ and $K = 0$), and residual kidney weight ($\lambda = 0.7$, $p \ll 0.001$; $K = 0.2$, p value = 0.012; null hypothesis λ and $K = 0$). After controlling for the phylogenetic signal, the PGLS results show that birds in high-salinity environments tend to have larger kidneys; the residual kidney weight is significantly positively correlated with levels of habitat salinity ($\lambda = 0.54$, $K = 0$, $\delta = 0.26$, estimate = 0.07, s.e. = 0.01, $t = 7.38$, $p \ll 0.001$, $R^2 = 0.25$). The evolutionary dependency between residual kidney weight and habitat types is also supported by the results of the correlated evolution analysis, except at the 25th percentile cut-off point (logBF for the set of cut-off points: 25th = 0.67; 50th = 10.872; 75th = 23.041).

The results of our PGLM analysis show that the presence of SSGs is significantly positively correlated with levels of habitat salinity ($\lambda = 0.91$, $K = 0.42$, $\delta = 2.09$, estimate = 0.488, s.e. = 0.226, $t = 2.16$, $p = 0.03$, $R^2 = 0.099$) (Figure 2). The results of the correlated evolutionary analyses also support such an association; the salt gland characteristic and habitat types are highly evolutionarily dependent on each other (logBF = 16.012). Furthermore, the presence of SSGs is positively correlated with kidney size ($\lambda = 0.46$, $K = 0$, $\delta = 0.3$, estimate = 0.21, s.e. = 0.03, $t = 6.4$, $p \ll 0.001$, $R^2 = 0.19$) (Figure 3). The results of the correlated evolutionary analyses show that the residual kidney weight and salt gland characteristics are strongly dependent on each other (logBF in the set of the cut-offs point 25th = 8.85; 50th = 16.618; 75th = 16.305).

Supraorbital salt glands and larger kidneys drove the evolution of salinity adaptation

Transition rate analysis indicates that the presence of SSGs drove birds to shift from low-to high-salinity habitats, while the lack of SSGs drove the evolution of habitat use in the other direction. For species with SSGs, the transition rate from low to high-salinity habitats was 4.56, while the reverse was 1.73; in contrast, for species without SSGs, the transition rate was biased toward low-salinity habitats (18.38 vs. 1.09 for low-to




Figure 3. The allometric effect of body weight on kidney weights

Linear regression between the $\log(\text{body weight})$ and $\log(\text{kidney size})$ is $\log(\text{kidney size}) = 1.164994 \pm 0.072970 + (0.928436 \pm 0.019287) \log(\text{body weight})$ ($p < 2.2\text{e-}16$; adjusted $r^2 = 0.933$).

high-salinity habitats transition; Figure 4A). Similarly, kidney size probably drove the evolution of birds' habitat use; for species with large kidneys, the transition rate from high-to low-salinity habitats was about 13.4 times higher than that from low-to high-salinity habitats (16.59 vs. 1.24 for low-to high-salinity habitats transition, Figure 4B).

Results of the transition rate analysis also support correlated evolution between the presence of SSGs and kidney sizes. For species without SSGs, the transition rate from large to non-large kidneys was 12.57, and the reverse was 7.68, whereas for species with SSGs, the transition rate from non-large to large kidneys was 17.81 and the reverse was 2.45 (Figure 4C).

Salinity is a force to fine-tune the evolution of avian kidney size

Our results suggest that different levels of environmental salinity might have modulated the evolution of kidney size (Figure 4B). For species living in low-salinity habitats, the transition rate from large to non-large kidneys was 12.09, and the reverse was 7.75, whereas, for species in high-salinity habitats, the transition rate from non-large to large kidneys (1.08) was about five times higher than that from large to non-large kidneys (0.18).

DISCUSSION

Our ancestral state reconstruction suggests that high-salinity habitats evolved independently in different lineages of birds (Figure 1), as it did with sea snakes.²⁹ Furthermore, we found that this ecological trait was quite evolutionarily inert; among the 230 species we sampled, only five species whose ancestors had used saline habitats reverted to using freshwater habitats. Similarly, we found that the two physiological traits related to salt tolerance, namely the presence of SSGs and large kidneys, also evolved convergently in birds. Of these, the presence of SSGs was also highly evolutionarily conserved (Figure 1). Therefore, our results suggest that the evolutionary potential of organisms might be constrained by the cost of niche shift.^{30,31} The "cost" of niche shift may arise because of antagonistic pleiotropy, in which a mutation in a single gene controlling multiple traits increases fitness in one trait while also reducing fitness in another one.³³ Alternatively, such "cost" may be rendered by a reduction in evolvability, the capacity to generate variation useful for adaptive change, after shifting to a new ecological niche.^{34,35}

Shifting to high salinity habitats might have allowed these avian lineages to escape from the intensive competition with other species using freshwater habitats and paved the way for further adaptive radiations.³⁶ It allowed avian lineages (such as Charadriiformes³⁷) to radiate by filling various empty niches in such habitats. However, ecological niche shift may constrain changes in other directions and thus limit niche diversity and evolvability^{38,39} later. Therefore, birds using high-salinity habitats might become an "evolutionary dead-end"^{40,41} after the empty salty niche is filled. The speciation rate for these birds could be reduced or even followed by an increasing extinction rate.




Figure 4. Evolutionary transition rates of salinity adaptation traits in avian species

Evolutionary transition rates between the use of high salinity habitats and (A) the presence of SSGs, (B) larger kidneys, and (C) between the larger kidneys and the presence of SSGs. For each characteristic, the transition rates between the three different combinations of characteristic categories were estimated by the dependent model in BayesTraits^{13,14} with the assumption that the two characteristics evolved interdependently. The dashed arrows are the estimated minimum transition rates ($\cong 0$). The thin black arrow indicates the estimated lower transition rates (< 10). The thick red arrows indicate the estimated higher transition rates (> 10).

Convergent evolution (the independent evolution of similar phenotypes⁸) to use high-salinity habitats also implies that different lineages of birds might have adopted different genetic routes to develop complex traits when adapting to high-salinity habitats.^{42,43} For instance, the supraorbital salt glands of birds have been considered to be derived from the nasal gland, which can also be found in reptiles and most mammals.⁸ The convergent evolution of salt glands in the tetrapod has been suggested to be the result of the cooption of an existing nasal gland in the ancestor of tetrapods.⁸ The convergent evolution of SSGs we found in this study implies that the development of this complex phenotype could have been modulated by the same set of developmental toolkits for the same existing gland in different lineages of birds independently. Therefore, the evolution of salt tolerance could have been driven by mutations in different compartments of the same genetic

pathways, or genetic toolkit, that regulates the developmental process of SSGs, kidney size, and other physiological traits related to salt tolerance or osmotic pressure. Convergent phenotypic evolution with divergent genetic causes has been shown in the cases of four Passerellidae sparrows,⁴⁵ the Savannah sparrow (*P. sandwichensis beldingi*), Nelson's sparrow (*Ammospiza nelsoni subvirgatus*), the song sparrow (*Melospiza melodia pusillula*), and the swamp sparrow (*M. georgiana nigrescens*); fewer than 50% of genes that were considered to contribute to adaptation to salty coastal habitats were shared among them. Hence the convergent evolution of salt-tolerance phenotypes suggests that birds should provide a rich opportunity to investigate the diverse genetic mechanisms underpinning their salt-tolerant phenotype.

In 1970, Hughes¹ found that species in avian orders composed mainly of non-saline species have relatively smaller kidneys whereas species associated with saline habitats have larger kidneys. However, he neither tested for the positive associations between kidney size and environmental salinity nor controlled for the effect of phylogenetic constraints on the evolution of kidney size and habitat use. Therefore, our results provide the first modern evolutionary test to demonstrate that avian species living in high-salinity environments have larger kidneys. Such an evolutionary trend can be illustrated by avian orders whose species mainly use freshwater habitats. For example, species specialized to high-salt habitats, such as the black-faced spoonbill of Pelecaniformes, the American coot (*Fulica americana*) of Gruiformes, and the Savannah sparrow of Passeriformes, all have the largest kidneys recorded in their order (Table 1). The importance of kidney size in osmoregulation is also supported by the result that lineages without large kidneys tend to shift their habitats from high-to low-salinity habitats (Figure 4B).

In addition to kidney size, our results suggest that the presence of SSGs in species also correlates with their habitats' salinity. We found that species with SSGs tend to become high-salinity specialists; in contrast, species without SSGs shift from high-salinity habitats to low-salinity habitats (Figure 4). Therefore, the transition rate between the SSG and environmental salinity (from low-to high-salinity habitats was 16.47, while the reverse was 3.35). This not only supports the important role of the SSG in osmoregulation in high-salinity environments^{13,46} but also suggests that the presence of SSGs could be a physiological prerequisite for birds in some lineages to use the harsh high-salinity environment.

Hughes proposed that SSGs should be associated with increased renal mass.¹ According to our PGLS results, the presence of SSGs is positively correlated with kidney size; our evolutionary dependency analysis results also suggest that SSGs tend to be highly correlated with the evolution of kidney sizes. This is also supported by the results of transition rate analysis; kidney size tended to decrease in species without SSGs and increase in species with SSGs (Figure 4). Our results support the intimate interactions between the kidneys and the salt glands for the osmoregulation suggested by Hughes.¹

The correlated evolution of two traits could be caused either by the linkage between the genes underpinning the two traits (genetic correlation) or because the two traits respond to the same selection pressure.⁴⁷ Genetic correlation is the proportion of variance that two traits share due to genetic causes.⁴⁸ Such correlations may arise from linkage disequilibrium (i.e., the non-random association of alleles at two or more loci) or pleiotropy, which occurs when a single gene influences two or more characteristics.⁴⁸ However, if the correlated evolution between the kidney size and functional salt glands had been caused by genetic linkage, we would have found a perfect (or nearly perfect) correlation between these two traits. Although the result of the correlated evolutionary analysis was highly significant ($\log_{10} BF > 8.0$) for the kidney size and presence of SSGs, the r^2 between relative kidney size and the functional salt gland was only 0.19. Therefore, our results suggest that the association between the large kidneys and the presence of SSGs was probably driven by the selection force of environmental salinity influencing both of them independently. This result implies that birds may need large kidneys and salt glands working synergistically to regulate the osmotic pressure in high-salinity habitats effectively. Or if multiple genes control each of the two traits, and some of the associated genes are linked, the correlations between the traits could be imperfectly correlated. Or perhaps the evolution of the traits was genetically correlated, and initially led to a perfect or near-perfect correlation between them which has to some extent broken down more recently, but has not they could still drift apart recently.

We also found that the correlation between kidney size and habitat type used ($r^2 = 0.25$) is much higher than that between the presence of supraorbital salt glands and habitat type used ($r^2 = 0.02$). The lower correlation between the presence of SSGs and habitat use might be because birds that can produce hypertonic salt secretions from SSGs are not restricted to species living in high-osmotic environments. For instance, the salt glands of raptor species (e.g., the tawny eagle), were found to produce secretions in response to a high protein diet.¹⁰ The common ostrich also produces salt secretions from the salt gland when facing heat stress.^{49,50}

Our analysis suggests that larger kidneys and the presence of SSGs are highly associated with the level of habitat salinity in birds. Almost all birds adapted to high-salinity habitats have evolved at least one characteristic that is assumed to be related to osmotic pressure. There are two lines of evidence to support the conclusion that salty environments drive the evolution of SSG and large kidneys. (1) Transition rate analysis indicates that species living in high salinity habitats tend to develop larger kidneys and SSGs; in contrast, species living in low-salinity habitats tend to develop smaller kidneys and to lose their SSGs (Figures 4B and 4C). (2) The results of our investigation of morphological convergence reveal that birds' kidney size and SSGs tend to undergo convergent evolution when they occupy saline water environments. Therefore, the salty environment might not only play a role in facilitating the emergence of physiological traits that participate in osmoregulation but could also further fine-tune the evolutionary dynamics of these physiological traits.

Our results suggest that environmental salinity could be a powerful driving force for large kidneys' convergent and correlated evolution and the presence of SSGs in birds. The strong association between the presence of SSGs and large kidneys suggests that birds might need both to handle the strong selection pressure posed by high-salinity environments. Because it is highly energetically demanding to maintain osmotic balance through SSGs and large kidneys in high-salinity environments,^{51,52} evolution to utilize salty habitats may be constrained in birds. Such physiological constraints might explain why less than 3% of avian species live entirely or partially in such high-osmolarity environments.⁵³ Our results provide tantalizing evidence of correlated evolution between different physiological traits to handle high osmotic pressure in high-salinity environments and suggest that such osmotic pressure has independently contributed to the expansion of avian physiological and functional diversity.

Table 1. List of avian species with information on the sample size, presence/absence (+/–) of supraorbital salt glands (SSGs), kidney weight in grams, body weight in grams, the residual of kidney weight/body weight (K/W), and habitat type (completely freshwater, mainly freshwater, mainly saline, completely saline)

Species	Sample size	SSG	Kidney (g)	Body (g)	K/W residuals	Habitat type
Accipitriformes						
<i>Accipiter gentilis</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
<i>Accipiter striatus</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
<i>Aquila rapax</i> ²	3	+	13.9	2532	–0.187	Completely freshwater
<i>Aquila verreauxii</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
<i>Buteo rufofuscus</i> ¹⁰	1	+	NA	NA	NA	Mainly freshwater
<i>Buteo jamaicensis</i> ¹⁵	1	–	7.6	1225	–0.155	Completely freshwater
<i>Gyps africanus</i> ²	1	–	35.8	5270	–0.073	Completely freshwater
<i>Haliaeetus vocifer</i> ²	1	–	18.4	3500	–0.197	Completely freshwater
<i>Gyps coprotheres</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
<i>Polemaetus bellicosus</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
<i>Terathopus ecaudatus</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
<i>Torgos tracheliotos</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
Anseriformes						
<i>Anas platyrhynchos</i> ¹	1	+	6.5	1305	–0.249	Mainly saline
<i>Anas acuta</i> ¹⁵	1	+	5.1	862	–0.186	Mainly freshwater
<i>Anas rubripes</i> ⁵⁴	1	+	NA	NA	NA	Mainly freshwater
<i>Anas crecca</i> ⁵⁵	1	+	NA	NA	NA	Mainly saline
<i>Anas clypeata</i> ⁵⁵	1	+	NA	NA	NA	Mainly saline
<i>Aythya valisineria</i> ⁵⁵	1	+	NA	NA	NA	Mainly saline
<i>Anser anser</i> ⁵⁷	1	+	NA	NA	NA	Mainly saline
<i>Cygnus atratus</i> ⁵⁸	1	+	NA	NA	NA	Mainly saline
<i>Aythya americana</i> ¹⁵	2	+	8.4	1055	–0.051	Mainly freshwater
<i>Aythya marila</i> ²	1	+	9.1	787	0.102	Mainly freshwater
<i>Aythya affinis</i> ²	1	+	18.1	1041	0.288	Mainly saline
<i>Oxyura jamaicensis</i> ¹	1	+	5.4	411	0.139	Mainly saline
<i>Mergus serrator</i> ²	1	+	9.7	700	0.178	Mainly saline
<i>Somateria mollissima</i> ⁵⁹	1	+	NA	NA	NA	Completely saline
Caprimulgiformes						
<i>Chordeiles minor</i> ¹⁵	3	–	0.6	80	–0.152	Completely freshwater
<i>Amazilia tzacatl</i> ²	1	–	0.04	5	–0.204	Completely freshwater
<i>Selasphorus platycercus</i> ¹⁵	2	–	0.03	3	–0.129	Completely freshwater
<i>Selasphorus rufus</i> ¹⁵	1	–	0.04	4	–0.114	Completely freshwater
Cathartiformes						
<i>Cathartes aura</i> ¹	1	–	12.6	1761	–0.083	Completely freshwater
Charadriiformes						
<i>Charadrius alexandrinus</i> ¹	1	+	0.6	37	0.161	Completely saline
<i>Charadrius vociferus</i> ¹⁵	3	+	1.3	88	0.145	Mainly saline
<i>Arenaria melanocephala</i> ¹	1	+	2	118	0.214	Completely saline
<i>Bartramia longicauda</i> ¹⁵	1	+	1.6	141	0.045	Completely freshwater
<i>Actitis macularius</i> ¹⁵	2	+	0.7	41	0.186	Mainly saline
<i>Tringa melanoleuca</i> ¹⁵	2	+	2.9	212	0.137	Mainly saline
<i>Calidris melanotos</i> ¹⁵	2	+	1.1	85	0.087	Mainly saline

(Continued on next page)

Table 1. Continued

Species	Sample size	SSG	Kidney (g)	Body (g)	K/W residuals	Habitat type
<i>Calidris minutilla</i> ¹	1	+	0.3	19	0.13	Mainly saline
<i>Calidris ruficollis</i> ¹	1	+	0.8	46	0.198	Mainly saline
<i>Limnodromus scolopaceus</i> ¹⁵	2	+	1.8	130	0.129	Mainly freshwater
<i>Calidris pusilla</i> ¹⁵	2	+	0.4	23	0.177	Mainly saline
<i>Limosa fedoa</i> ¹	1	+	6.4	388	0.236	Mainly saline
<i>Larus californicus</i> ⁶⁰	1	+	NA	NA	NA	Mainly saline
<i>Larus marinus</i> ⁵⁹	1	+	NA	NA	NA	Completely saline
<i>Larus hyperboreus</i> ⁶¹	1	+	NA	NA	NA	Completely saline
<i>Larus delawarensis</i> ¹⁵	4	+	7.2	488	0.194	Mainly saline
<i>Larus heermanni</i> ¹	1	+	7.8	544	0.185	Completely saline
<i>Larus glaucescens</i> ¹	1	+	7.4	618	0.111	Mainly saline
<i>Larus occidentalis</i> ¹	2	+	15.5	1283	0.136	Completely saline
<i>Chlidonias niger</i> ¹⁵	3	+	1	60	0.187	Mainly saline
<i>Aethia cristatella</i> ¹	2	+	6	289	0.328	Completely saline
<i>Aethia pusilla</i> ¹	2	+	2.3	96	0.358	Completely saline
<i>Fratercula cirrhata</i> ¹	1	+	11.7	734	0.24	Completely saline
<i>Uria aalge</i> ¹	1	+	17.5	1031	0.277	Completely saline
<i>Alle alle</i> ²	2	+	2.1	103	0.29	Completely saline
<i>Cephus grylle</i> ⁶¹	1	+	NA	NA	NA	Completely saline
<i>Uria lomvia</i> ⁶¹	1	+	NA	NA	NA	Completely saline
<i>Alca torda</i> ⁶¹	1	+	NA	NA	NA	Completely saline
<i>Recurvirostra americana</i> ⁶²	1	+	NA	NA	NA	Mainly saline
<i>Rynchops niger</i> ⁵⁹	1	+	NA	NA	NA	Mainly saline
<i>Fratercula corniculata</i> ¹	2	+	9.7	572	0.26	Completely saline
<i>Fratercula arctica</i> ⁶¹	1	+	NA	NA	NA	Completely saline
<i>Charadrius hiaticula</i> ⁶¹	1	+	NA	NA	NA	Mainly saline
<i>Pluvialis apricaria</i> ⁶¹	1	+	NA	NA	NA	Mainly saline
<i>Rissa tridactyla</i> ⁶¹	1	+	NA	NA	NA	Completely saline
<i>Actitis hypoleucos</i> ⁶¹	1	+	NA	NA	NA	Mainly saline
<i>Calidris alpina</i> ⁶¹	1	+	NA	NA	NA	Mainly saline
<i>Calidris canutus</i> ⁶¹	1	+	NA	NA	NA	Completely saline
<i>Calidris minuta</i> ⁶¹	1	+	NA	NA	NA	Mainly saline
<i>Calidris alba</i> ⁶¹	1	+	NA	NA	NA	Mainly saline
<i>Gallinago gallinago</i> ⁶¹	1	+	NA	NA	NA	Mainly saline
<i>Limosa lapponica</i> ⁶¹	1	+	NA	NA	NA	Mainly saline
<i>Tringa glareola</i> ⁶¹	1	+	NA	NA	NA	Completely freshwater
<i>Tringa ochropus</i> ⁶¹	1	+	NA	NA	NA	Completely freshwater
Ciconiiformes						
<i>Ciconia ciconia</i> ²	3	-	23	3350	-0.082	Completely freshwater
Columbiformes						
<i>Columba livia</i> ¹⁵	1	-	2.8	286	0.001	Completely freshwater
<i>Zenaida asiatica</i> ¹⁵	3	-	0.9	141	-0.205	Mainly freshwater
<i>Zenaida macroura</i> ¹	1	-	0.7	110	-0.214	Completely freshwater
<i>Columbina inca</i> ¹⁵	2	-	0.3	43	-0.201	Completely freshwater

(Continued on next page)

Table 1. Continued

Species	Sample size	SSG	Kidney (g)	Body (g)	K/W residuals	Habitat type
Coraciiformes						
<i>Megaceryle alcyon</i> ¹⁵	2	–	2.2	173	0.1	Mainly saline
Cuculiformes						
<i>Geococcyx californianus</i> ¹⁵	2	–	1.7	190	–0.05	Completely freshwater
Falconiformes						
<i>Falco sparverius</i> ²	1	–	1.1	112	–0.025	Completely freshwater
<i>Falco biarmicus</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
<i>Falco cherrug</i> ¹⁰	1	+	NA	NA	NA	Mainly freshwater
<i>Falco chicquera</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
<i>Falco jugger</i> ¹⁰	1	–	NA	NA	NA	Complete freshwater
<i>Falco peregrinus</i> ¹⁰	1	+	NA	NA	NA	Mainly saline
<i>Polihierax semitorquatus</i> ¹⁰	1	+	NA	NA	NA	Complete freshwater
Galliformes						
<i>Bonasa umbellus</i> ¹⁵	2	–	4.9	577	–0.041	Completely freshwater
<i>Lagopus lagopus</i> ¹⁵	5	–	6	602	0.03	Completely freshwater
<i>Tympanuchus phasianellus</i> ¹⁵	1	–	5.1	791	–0.151	Completely freshwater
<i>Centrocercus urophasianus</i> ¹⁵	2	–	22.3	2013	0.111	Completely freshwater
<i>Callipepla squamata</i> ¹⁵	2	–	1.4	168	–0.084	Completely freshwater
<i>Callipepla gambelii</i> ¹⁵	3	–	1.2	150	–0.105	Completely freshwater
<i>Cyrtonyx montezumae</i> ¹⁵	2	–	1.3	212	–0.211	Completely freshwater
<i>Phasianus colchicus</i> ¹⁵	4	–	8.2	1283	–0.14	Completely freshwater
<i>Perdix perdix</i> ¹⁵	2	–	3.8	521	–0.11	Completely freshwater
<i>Numida meleagris</i> ²	1	–	7.3	1620	–0.286	Completely freshwater
<i>Coturnix coturnix</i> ⁶³	18	+	NA	NA	NA	Mainly freshwater
<i>Ammoperdix heyi</i> ⁴⁹	1	+	NA	NA	NA	Completely freshwater
Gaviiformes						
<i>Gavia stellata</i> ²	3	+	25.2	1549	0.27	Mainly saline
Gruiformes						
<i>Gallirallus owstoni</i> ⁶⁴	4	+	NA	NA	NA	Completely freshwater
<i>Balearica pavonina</i> ²	2	–	26	4448	–0.144	Completely freshwater
<i>Fulica americana</i> ¹	1	+	9	699	0.146	Mainly saline
Otidiformes						
<i>Ardeotis kori</i> ²	2	–	44.3	7770	–0.138	Completely freshwater
Pelecaniformes						
<i>Pelecanus erythrorhynchos</i> ¹⁵	2	+	66	6777	0.09	Mainly saline
<i>Ardea herodias</i> ¹⁵	1	–	15.8	1840	–0.002	Mainly saline
<i>Nycticorax nycticorax</i> ¹⁵	4	–	4.6	623	–0.099	Mainly freshwater
<i>Botaurus lentiginosus</i> ¹⁵	2	–	4.7	625	–0.091	Mainly saline
<i>Platalea minor</i> ^c	4	+	21.52	1573	0.196	Completely saline
<i>Threskiornis aethiopicus</i> ^c	3	+	9.08	1068	–0.022	Mainly saline
<i>Gorsachius melanolophus</i> ^c	3	–	2.38	397	–0.203	Completely freshwater
<i>Bubulcus ibis</i> ^c	3	–	1.26	238	–0.271	Completely freshwater
<i>Pelecanus occidentalis</i> ⁵⁹		+	NA	NA	NA	Completely saline

(Continued on next page)

Table 1. Continued

Species	Sample size	SSG	Kidney (g)	Body (g)	K/W residuals	Habitat type
Phoenicopteriformes						
<i>Phoeniconaias minor</i> ²	5	+	18	1541	0.126	Completely saline
Piciformes						
<i>Colaptes auratus</i> ¹⁵	1	–	1	126	–0.114	Completely freshwater
<i>Colaptes chrysoides</i> ¹⁵	2	–	1	116	–0.08	Completely freshwater
<i>Melanerpes uropygialis</i> ¹⁵	3	–	0.7	68	–0.019	Completely freshwater
<i>Melanerpes formicivorus</i> ¹⁵	2	–	0.5	63	–0.134	Completely freshwater
<i>Sphyrapicus varius</i> ¹⁵	2	–	0.5	45	0.002	Completely freshwater
<i>Sphyrapicus thyroideus</i> ¹⁵	3	–	0.5	43	0.021	Completely freshwater
Podicipediformes						
<i>Podiceps nigricollis</i> ¹⁵	2	+	3.7	330	0.064	Mainly saline
<i>Podilymbus podiceps</i> ¹⁵	2	+	5.9	496	0.101	Mainly saline
Procellariiformes						
<i>Phoebastria immutabilis</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Phoebastria nigripes</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Phoenicopterus ruber</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Phoebastria irrorata</i> ⁶⁶	1	+	NA	NA	NA	Completely saline
Psittaciformes						
<i>Melopsittacus undulatus</i> ¹⁵	6	–	0.3	39	–0.162	Completely freshwater
Sphenisciformes						
<i>Eudyptula minor</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Pygoscelis adeliae</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Pygoscelis papua</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Spheniscus mendiculus</i> ⁴⁴	1	+	NA	NA	NA	Completely saline
<i>Spheniscus humboldti</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
Strigiformes						
<i>Bubo virginianus</i> ¹	1	–	3.7	635	–0.201	Completely freshwater
<i>Micrathene whitneyi</i> ¹⁵	2	–	0.4	37	–0.015	Completely freshwater
Struthioniformes						
<i>Struthio camelus</i> ²	1	+	920	123000	0.06	Completely freshwater
Suliformes						
<i>Urile pelagicus</i> ¹	1	+	16.9	1300	0.168	Completely saline
<i>Fregata minor</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Aptenodytes patagonicus</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Sula nebowxii</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
<i>Sula sula</i> ⁶⁵	1	+	NA	NA	NA	Completely saline
Passeriformes						
<i>Pachyrhamphus aglaiae</i> ¹⁵	1	–	0.295	29	–0.049	Completely freshwater
<i>Tyrannus tyrannus</i> ¹⁵	3	–	0.604	49	0.05	Completely freshwater
<i>Tyrannus verticalis</i> ¹⁵	1	–	0.388	38.8	–0.048	Completely freshwater
<i>Tyrannus vociferans</i> ¹⁵	2	–	0.557	46.4	0.037	Completely freshwater
<i>Pitangus sulphuratus</i> ¹⁵	1	–	0.657	79	–0.107	Completely freshwater
<i>Contopus sordidulus</i> ¹⁵	3	–	0.157	14.3	–0.036	Completely freshwater

(Continued on next page)

Table 1. Continued

Species	Sample size	SSG	Kidney (g)	Body (g)	K/W residuals	Habitat type
<i>Contopus cooperi</i> ¹⁵	1	–	0.515	36.5	0.1	Completely freshwater
<i>Pyrocephalus rubinus</i> ¹⁵	2	–	0.211	13.9	0.104	Completely freshwater
<i>Empidonax oberholseri</i> ¹⁵	1	–	0.204	12.1	0.145	Completely freshwater
<i>Myiarchus cinerascens</i> ¹⁵	4	–	0.299	26.3	–0.003	Mainly freshwater
<i>Eremophila alpestris</i> ¹⁵	4	–	0.284	27.5	–0.044	Mainly freshwater
<i>Tachycineta thalassina</i> ¹⁵	2	–	0.282	15.7	0.18	Mainly freshwater
<i>Tachycineta bicolor</i> ¹⁵	3	–	0.315	18	0.173	Mainly freshwater
<i>Hirundo rustica</i> ¹⁵	3	–	0.325	18	0.187	Completely freshwater
<i>Petrochelidon pyrrhonota</i> ¹⁵	5	–	0.374	22.2	0.163	Completely freshwater
<i>Progne subis</i> ¹⁵	2	–	0.544	46.6	0.025	Mainly freshwater
<i>Cyanocitta stelleri</i> ¹⁵	2	–	1.101	103.7	0.007	Completely freshwater
<i>Aphelocoma ultramarina</i> ¹⁵	2	–	1.274	120.2	0.01	Completely freshwater
<i>Pica pica</i> ¹⁵	4	–	2.556	198.2	0.11	Completely freshwater
<i>Baeolophus wollweberi</i> ¹⁵	4	–	0.144	10.4	0.055	Completely freshwater
<i>Auriparus flaviceps</i> ¹⁵	4	–	0.107	7	0.087	Mainly freshwater
<i>Psaltriparus minimus</i> ¹⁵	1	–	0.102	8	0.012	Mainly freshwater
<i>Sitta carolinensis</i> ¹⁵	1	–	0.225	15.8	0.08	Completely freshwater
<i>Certhia familiaris</i> ¹⁵	2	–	0.145	8.4	0.145	Mainly freshwater
<i>Troglodytes aedon</i> ¹⁵	3	–	0.175	11	0.117	Completely freshwater
<i>Cistothorus palustris</i> ¹⁵	2	–	0.278	13	0.255	Mainly saline
<i>Thryomanes bewickii</i> ¹⁵	1	–	0.212	12.2	0.159	Completely freshwater
<i>Campylorhynchus brunneicapillus</i> ¹⁵	5	–	0.424	38.4	–0.005	Completely freshwater
<i>Catherpes mexicanus</i> ¹⁵	1	–	0.18	11.2	0.122	Completely freshwater
<i>Salpinctes obsoletus</i> ¹⁵	2	–	0.249	15.5	0.132	Mainly freshwater
<i>Mimus polyglottos</i> ¹⁵	3	–	0.508	49.1	–0.026	Completely freshwater
<i>Dumetella carolinensis</i> ¹⁵	1	–	0.452	36.8	0.04	Completely freshwater
<i>Toxostoma bendirei</i> ¹⁵	1	–	0.66	63.6	–0.017	Completely freshwater
<i>Toxostoma curvirostre</i> ¹⁵	3	–	0.875	72.4	0.053	Completely freshwater
<i>Toxostoma crissale</i> ¹⁵	2	–	0.677	58.8	0.026	Completely freshwater
<i>Oreoscoptes montanus</i> ¹⁵	3	–	0.57	40	0.107	Completely freshwater
<i>Turdus migratorius</i> ¹⁵	3	–	1.172	86.6	0.107	Completely freshwater
<i>Catharus guttatus</i> ¹⁵	1	–	0.359	31	0.009	Completely freshwater
<i>Catharus ustulatus</i> ¹⁵	1	–	0.461	29.8	0.134	Mainly freshwater
<i>Catharus minimus</i> ¹⁵	1	–	0.396	33.2	0.024	Completely freshwater
<i>Catharus occidentalis</i> ¹⁵	1	–	0.252	31.2	–0.147	Completely freshwater
<i>Catharus aurantirostris</i> ¹⁵	1	–	0.405	33.8	0.027	Completely freshwater
<i>Sialia mexicana</i> ¹⁵	4	–	0.408	29.3	0.088	Mainly freshwater
<i>Polioptila caerulea</i> ¹⁵	2	–	0.078	5.1	0.078	Completely freshwater
<i>Polioptila melanura</i> ¹⁵	3	–	0.086	5.4	0.097	Completely freshwater
<i>Bombycilla cedrorum</i> ¹⁵	4	–	0.471	40.1	0.023	Completely freshwater
<i>Phainopepla nitens</i> ¹⁵	3	–	0.368	25.3	0.103	Completely freshwater
<i>Sturnus vulgaris</i> ¹⁵	4	–	1.133	82.2	0.113	Mainly freshwater
<i>Vireo solitarius</i> ¹⁵	1	–	0.169	16	–0.05	Completely freshwater
<i>Setophaga ruticilla</i> ¹⁵	1	–	0.1	9.5	–0.066	Mainly freshwater
<i>Myioborus pictus</i> ¹⁵	1	–	0.163	9.1	0.163	Completely freshwater

(Continued on next page)

Table 1. Continued

Species	Sample size	SSG	Kidney (g)	Body (g)	K/W residuals	Habitat type
<i>Myioborus miniatus</i> ¹⁵	1	–	0.097	9.5	–0.079	Completely freshwater
<i>Passer domesticus</i> ¹⁵	5	–	0.301	29.3	–0.044	Completely freshwater
<i>Taeniopygia guttata</i> ¹⁵	2	–	0.09	11.9	–0.203	Completely freshwater
<i>Dolichonyx oryzivorus</i> ¹⁵	3	–	0.396	35.2	0.001	Mainly freshwater
<i>Sturnella neglecta</i> ¹⁵	2	–	1.247	113.4	0.025	Completely freshwater
<i>Agelaius phoeniceus</i> ¹⁵	4	–	0.668	67.5	–0.036	Mainly freshwater
<i>Icterus cucullatus</i> ¹⁵	2	–	0.294	25.2	0.007	Completely freshwater
<i>Icterus bullockii</i> ¹⁵	3	–	0.41	33.1	0.041	Completely freshwater
<i>Euphagus carolinus</i> ¹⁵	1	–	0.687	66.2	–0.016	Mainly freshwater
<i>Euphagus cyanocephalus</i> ¹⁵	3	–	0.863	70.9	0.055	Mainly freshwater
<i>Quiscalus quiscula</i> ¹⁵	3	–	1.037	94.3	0.019	Completely freshwater
<i>Molothrus ater</i> ¹⁵	2	–	0.398	49.7	–0.137	Completely freshwater
<i>Piranga ludoviciana</i> ¹⁵	2	–	0.514	33.2	0.138	Completely freshwater
<i>Cardinalis cardinalis</i> ¹⁵	3	–	0.59	47	0.057	Completely freshwater
<i>Cardinalis sinuatus</i> ¹⁵	1	–	0.574	35.1	0.163	Completely freshwater
<i>Passerina caerulea</i> ¹⁵	1	–	0.273	28.1	–0.07	Completely freshwater
<i>Coccothraustes vespertinus</i> ¹⁵	1	–	0.86	52.3	0.177	Completely freshwater
<i>Pinicola enucleator</i> ¹⁵	1	–	0.833	53.5	0.154	Completely freshwater
<i>Passerculus sandwichensis</i> ¹⁵	3	–	0.261	17.7	0.254	Completely saline
<i>Pipilo erythrophthalmus</i> ¹⁵	1	–	0.397	39.6	–0.046	Mainly freshwater
<i>Poocetes gramineus</i> ¹⁵	2	–	0.285	26.3	–0.024	Completely freshwater
<i>Chondestes grammacus</i> ¹⁵	1	–	0.166	26.2	–0.257	Completely freshwater
<i>Aimophila ruficeps</i> ¹⁵	1	–	0.264	17.9	0.099	Completely freshwater
<i>Amphispiza bilineata</i> ¹⁵	6	–	0.154	12.9	–0.003	Completely freshwater
<i>Junco hyemalis</i> ¹⁵	2	–	0.216	21.7	–0.067	Completely freshwater
<i>Spizella passerina</i> ¹⁵	1	–	0.125	13.3	–0.106	Completely freshwater
<i>Spizella pallida</i> ¹⁵	1	–	0.17	13	0.037	Completely freshwater
<i>Spizella breweri</i> ¹⁵	4	–	0.178	10.6	0.14	Completely freshwater
<i>Zonotrichia albicollis</i> ¹⁵	3	–	0.279	26.5	–0.036	Completely freshwater
<i>Melospiza lincolni</i> ¹⁵	5	–	0.228	18.1	0.03	Mainly freshwater
<i>Melospiza melodia</i> ¹⁵	2	–	0.247	21.1	0.003	Completely freshwater
<i>Calcarius lapponicus</i> ¹⁵	2	–	0.261	26.7	–0.068	Mainly freshwater
<i>Plectrophenax nivalis</i> ¹⁵	1	–	0.337	31.1	–0.019	Mainly saline

Data sources are shown in superscript after the species name; a superscript C indicates data collected in the current study.

Limitations of the study

Since kidney weight measurements require fresh samples, they are not easy to obtain. Only about 56% of the orders of birds were included in this study. However, such limitations probably only affect the estimation of the numbers of species having adapted to high-salinity habitats, changed kidney size, or developed or lost SSGs in the evolutionary process, rather than the trend of the results from the phylogenetic analysis, the ancestral state reconstruction, and the correlated evolution analysis.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- RESOURCE AVAILABILITY
 - Lead contact

- Materials availability
- Data and code availability
- **METHOD DETAILS**
 - Collection of characteristics
 - Standardization of kidney sizes
 - Ancestral state reconstruction
 - Phylogenetic signals in kidney sizes, possession of SSGs, and habitat salinity
 - Evolutionary transitions in kidney size, SSGs and levels of habitat salinity
 - Searching for morphological convergence among species in diverse saline habitat types
- **QUANTIFICATION AND STATISTICAL ANALYSIS**

ACKNOWLEDGMENTS

We greatly thank two anonymous reviewers who provide numerous insightful comments to improve the quality of this article. We thank the Taiwan Biodiversity Research Institute for providing the specimens they archived. We are grateful to Yao-Hua Chen, Jun-Ling Yao, and Yi-Hua Guo who assisted with sample preparation and anatomy. We are in debt to Chih-Ming Hung who had provided many insightful comments to the early draft of this article. We thank Alan Watson for greatly improving the readability of this article. We thank the Ministry of Science and Technology of Taiwan (MOST 111-2621-B-003 -002 -MY3) for their financial support.

AUTHOR CONTRIBUTIONS

Shou-Hsien Li, Chi-Cheng Chiu, and Ben-Yang Liao designed the research; Chi-Cheng Chiu, and Cheng-Te Yao performed the research; Chi-Cheng Chiu analyzed data; and Shou-Hsien Li, Chi-Cheng Chiu, Ben-Yang Liao, and Cheng-Te Yao wrote the article. All authors approved the final submission.

DECLARATION OF INTERESTS

None of the authors of this article have any affiliations with or involvement in any organization or entity with any financial or non-financial interest (such as patent or stock ownership, membership of a company board of directors, membership of an advisory board or committee for a company, and consultancy for or receipt of speaker's fees from a company) in the subject matter or materials discussed in this article.

Received: November 23, 2022

Revised: October 30, 2023

Accepted: February 6, 2024

Published: February 9, 2024

REFERENCES

1. Hughes, M.R. (1970). Relative Kidney Size in Nonpasserine Birds with Functional Salt Glands. *Condor* 72, 164–168. <https://doi.org/10.2307/1366626>.
2. Quiring, D.P. (1950). *Functional Anatomy of the Vertebrates* (McGraw-Hill).
3. Baumel, J.J. (1993). *Handbook of Avian Anatomy: Nomina Anatomica Avium* (Publications of the Nuttall Ornithological Club).
4. Schmidt-Nielsen, K., Jørgensen, C.B., and Osaki, H. (1958). Extrarenal salt excretion in birds. *Am. J. Physiol.* 193, 101–107. <https://doi.org/10.1152/ajplegacy.1958.193.1.101>.
5. Hughes, M.R. (2003). Regulation of salt gland, gut and kidney interactions. *Comp. Biochem. Physiol. Mol. Integr. Physiol.* 136, 507–524. <https://doi.org/10.1016/j.cbpb.2003.09.005>.
6. Gutiérrez, J.S., Dietz, M.W., Masero, J.A., Gill Jr, R.E., Dekinga, A., Battley, P.F., Sánchez-Guzmán, J.M., and Piersma, T. (2012). Functional ecology of saltglands in shorebirds: flexible responses to variable environmental conditions. *Funct. Ecol.* 26, 236–244. <https://doi.org/10.1111/j.1365-2435.2011.01929.x>.
7. Vidal, N., and Hedges, S.B. (2009). The molecular evolutionary tree of lizards, snakes, and amphisbaenians. *C. R. Biol.* 332, 129–139. <https://doi.org/10.1016/j.crv.2008.07.010>.
8. Babonis, L.S., and Brischoux, F. (2012). Perspectives on the Convergent Evolution of Tetrapod Salt Glands. *Integr. Comp. Biol.* 52, 245–256. <https://doi.org/10.1093/icb/ics073>.
9. Carpenter, R.E., and Stafford, M.A. (1970). The secretory rates and the chemical stimulus for secretion of the nasal salt glands in the Rallidae. *Condor* 72, 316–324. <https://doi.org/10.2307/1366010>.
10. Cade, T.J., and Greenwald, L. (1966). Nasal Salt Secretion in Falconiform Birds. *Condor* 68, 338–350. <https://doi.org/10.2307/1365449>.
11. Schmidt-Nielsen, K., and Kim, Y.T. (1964). The effect of salt intake on the size and function of the salt gland of ducks. *Auk* 81, 160–172. <https://doi.org/10.2307/4082766>.
12. Shuttleworth, T.J., and Hildebrandt, J.P. (1999). Vertebrate salt glands: Short-and long-term regulation of function. *J. Exp. Zool.* 283, 689–701. [https://doi.org/10.1002/\(sici\)1097-010x\(19990601\)283:7<689::aid-jez7>3.0.co;2-t](https://doi.org/10.1002/(sici)1097-010x(19990601)283:7<689::aid-jez7>3.0.co;2-t).
13. Peaker, M., and Linzell, J.L. (1975). *Salt Glands in Birds and Reptiles* (Cambridge University Press).
14. Skadhauge, E. (1982). A quantitative survey of salt and water excretion. *Comp. Biochem. Physiol. A Comp. Physiol.* 71, 481–483. [https://doi.org/10.1016/0300-9629\(82\)90196-7](https://doi.org/10.1016/0300-9629(82)90196-7).
15. Johnson, O.W. (1968). Some Morphological Features of Avian Kidneys. *Auk* 85, 216–228. <https://doi.org/10.2307/4083582>.
16. S.M. Billerman, B.K. Keeney, P.G. Rodewald, and T.S. Schulenberg, eds. (2020). *Birds of the World* (Cornell Laboratory of Ornithology).
17. Jetz, W., Thomas, G.H., Joy, J.B., Redding, D.W., Hartmann, K., and Mooers, A.O. (2014). Global Distribution and Conservation of Evolutionary Distinctness in Birds. *Curr. Biol.* 24, 919–930. <https://doi.org/10.1016/j.cub.2014.03.011>.
18. Hackett, S.J., Kimball, R.T., Reddy, S., Bowie, R.C.K., Braun, E.L., Braun, M.J., Chojnowski, J.L., Cox, W.A., Han, K.-L., Harshman, J., et al. (2008). A Phylogenomic Study of Birds Reveals Their Evolutionary History. *Science* 320, 1763–1768. <https://doi.org/10.1126/science.1157704>.
19. Revell, L.J. (2012). phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* 3, 217–223. <https://doi.org/10.1111/j.2041-210X.2011.00169.x>.

20. Pagel, M. (1999). The Maximum Likelihood Approach to Reconstructing Ancestral Character States of Discrete Characters on Phylogenies. *Syst. Biol.* 48, 612–622.
21. Pagel, M. (1994). Detecting correlated evolution on phylogenies: a general method for the comparative analysis of discrete characters. *Proc. R. Soc. B. Biol. Sci.* 255, 37–45. <https://doi.org/10.1098/rspb.1994.0006>.
22. Pagel, M., and Lutzoni, F. (2002). Accounting for phylogenetic uncertainty in comparative studies of evolution and adaptation. In *Biological Evolution and Statistical Physics Lecture Notes in Physics*, M. Lässig and A. Valleriani, eds. (Springer), pp. 148–161. https://doi.org/10.1007/3-540-45692-9_8.
23. Orme, D., Freckleton, R., Thomas, G., Petzoldt, T., Fritz, S., Isaac, N., and Pearse, W. (2018). *Caper: Comparative Analyses of Phylogenetics and Evolution in R*.
24. Ho, L.S.T., Ane, C., Lachlan, R., Tarpinian, K., Feldman, R., Yu, Q., and van der Bijl, W. (2016). *Phyloim*.
25. Pagel, M., Meade, A., and Barker, D. (2004). Bayesian Estimation of Ancestral Character States on Phylogenies. *Syst. Biol.* 53, 673–684. <https://doi.org/10.1080/10635150490522232>.
26. Pagel, M., and Meade, A. (2006). Bayesian Analysis of Correlated Evolution of Discrete Characters by Reversible-Jump Markov Chain Monte Carlo. *Am. Nat.* 167, 808–825. <https://doi.org/10.1086/503444>.
27. Meade, A., and Pagel, M. (2014). *BayesTraits*.
28. Fristoe, T.S., Iwaniuk, A.N., and Botero, C.A. (2017). Big brains stabilize populations and facilitate colonization of variable habitats in birds. *Nat. Ecol. Evol.* 1, 1706–1715. <https://doi.org/10.1038/s41559-017-0316-2>.
29. Gearty, W., Carrillo, E., and Payne, J.L. (2021). Ecological Filtering and Exaptation in the Evolution of Marine Snakes. *Am. Nat.* 198, 506–521. <https://doi.org/10.1086/716015>.
30. Futuyma, D.J., and Moreno, G. (1988). The Evolution of Ecological Specialization. *Annu. Rev. Ecol. Syst.* 19, 207–233.
31. Schluter, D. (2000). Ecological Character Displacement in Adaptive Radiation. *Am. Nat.* 156, S4–S16. <https://doi.org/10.1086/303412>.
32. Castiglione, S., Tesone, G., Piccolo, M., Melchionna, M., Mondanaro, A., Serio, C., Di Febbraro, M., and Raia, P. (2018). A new method for testing evolutionary rate variation and shifts in phenotypic evolution. *Methods Ecol. Evol.* 9, 974–983.
33. MacArthur, R.H. (1984). *Geographical Ecology: Patterns in the Distribution of Species* (Princeton University Press).
34. Colegrave, N., and Collins, S. (2008). Experimental evolution: experimental evolution and evolvability. *Heredity* 100, 464–470. <https://doi.org/10.1038/sj.hdy.6801095>.
35. Woods, R.J., Barrick, J.E., Cooper, T.F., Shrestha, U., Kauth, M.R., and Lenski, R.E. (2011). Second-Order Selection for Evolvability in a Large *Escherichia coli* Population. *Science* 331, 1433–1436. <https://doi.org/10.1126/science.1198914>.
36. Ackerly, D.D., Schwilk, D.W., and Webb, C.O. (2006). Niche Evolution and Adaptive Radiation: Testing the Order of Trait Divergence. *Ecology* 87, S50–S61. [https://doi.org/10.1890/0012-9658\(2006\)87\[50:NEAART\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[50:NEAART]2.0.CO;2).
37. Baker, A.J., Pereira, S.L., and Paton, T.A. (2007). Phylogenetic relationships and divergence times of Charadriiformes genera: multigene evidence for the Cretaceous origin of at least 14 clades of shorebirds. *Biol. Lett.* 3, 205–209. <https://doi.org/10.1098/rsbl.2006.0606>.
38. Buckling, A., Wills, M.A., and Colegrave, N. (2003). Adaptation Limits Diversification of Experimental Bacterial Populations. *Science* 302, 2107–2109. <https://doi.org/10.1126/science.1088848>.
39. Collar, D.C., O'Meara, B.C., Wainwright, P.C., and Near, T.J. (2009). Piscivory Limits Diversification of Feeding Morphology in Centrarchid Fishes. *Evolution* 63, 1557–1573. <https://doi.org/10.1111/j.1558-5646.2009.00626.x>.
40. Colles, A., Liow, L.H., and Prinzing, A. (2009). Are specialists at risk under environmental change? Neoeological, paleoecological and phylogenetic approaches. *Ecol. Lett.* 12, 849–863. <https://doi.org/10.1111/j.1461-0248.2009.01336.x>.
41. Simpson, G.G. (1944). *Tempo and Mode in Evolution* (Columbia University Press).
42. Stern, D.L. (2013). The genetic causes of convergent evolution. *Nat. Rev. Genet.* 14, 751–764. <https://doi.org/10.1038/nrg3483>.
43. Arendt, J., and Reznick, D. (2008). Convergence and parallelism reconsidered: what have we learned about the genetics of adaptation? *Trends Ecol. Evol.* 23, 26–32. <https://doi.org/10.1016/j.tree.2007.09.011>.
44. Fänge, R., Schmidt-Nielsen, K., and Robinson, M. (1958). Control of Secretion From the Avian Salt Gland. *Am. J. Physiol.* 195, 321–326. <https://doi.org/10.1152/ajplegacy.1958.195.2.321>.
45. Walsh, J., Benham, P.M., Deane-Coe, P.E., Arcese, P., Butcher, B.G., Chan, Y.L., Cheviron, Z.A., Elphick, C.S., Kovach, A.I., Olsen, B.J., et al. (2019). Genomics of rapid ecological divergence and parallel adaptation in four tidal marsh sparrows. *Evol. Lett.* 3, 324–338. <https://doi.org/10.1002/evl3.126>.
46. Holmes, W.N., and Phillips, J.G. (1985). The Avian Salt Gland. *Biol. Rev.* 60, 213–256. <https://doi.org/10.1111/j.1469-185X.1985.tb00715.x>.
47. Price, T., and Langen, T. (1992). Evolution of correlated characters. *Trends Ecol. Evol.* 7, 307–310. [https://doi.org/10.1016/0169-5347\(92\)90229-5](https://doi.org/10.1016/0169-5347(92)90229-5).
48. Falconer, D.S. (1996). *Introduction to Quantitative Genetics* (Pearson Education India).
49. Schmidt-Nielsen, K., Borut, A., Lee, P., and Crawford, E. (1963). Nasal Salt Excretion and the Possible Function of the Cloaca in Water Conservation. *Science* 142, 1300–1301. <https://doi.org/10.1126/science.142.3597.1300>.
50. Sheldon, F.H., and Whittingham, L.A. (1997). Phylogeny in studies of bird ecology, behavior, and morphology. In *Avian Molecular Evolution and Systematics*, D.P. Mindell, ed. (Academic Press).
51. Peña-Villalobos, I., Valdés-Ferranty, F., and Sabat, P. (2013). Osmoregulatory and metabolic costs of salt excretion in the Rufous-collared sparrow *Zonotrichia capensis*. *Comp. Biochem. Physiol. Mol. Integr. Physiol.* 164, 314–318. <https://doi.org/10.1016/j.cbpa.2012.10.027>.
52. Sabat, P., Narváez, C., Peña-Villalobos, I., Contreras, C., Maldonado, K., Sanchez-Hernandez, J.C., Newsome, S.D., Nespolo, R., and Bozinovic, F. (2017). Coping with Salt Water Habitats: Metabolic and Oxidative Responses to Salt Intake in the Rufous-Collared Sparrow. *Front. Physiol.* 8, 654.
53. Rajpar, M.N., Ozdemir, I., Zakaria, M., Sheryar, S., and Rab, A. (2018). Seabirds as bioindicators of marine ecosystems. In *Intechopen, H.M. Seabirds*, ed., pp. 47–65.
54. Barnes, G.G., and Nudds, T.D. (1991). Salt tolerance in American black ducks, mallards, and their F1-hybrids. *Auk* 108, 89–98. <https://doi.org/10.1093/auk/108.1.89>.
55. Cooch, F.G. (1964). A preliminary study of the survival value of a functional salt gland in prairie Anatidae. *Auk* 81, 380–393. <https://doi.org/10.2307/4082692>.
56. Losos, J.B. (2011). Convergence, adaptation, and constraint. *Evolution* 65, 1827–1840. <https://doi.org/10.1111/j.1558-5646.2011.01289.x>.
57. Hanwell, A., Linzell, J.L., and Peaker, M. (1971). Salt-gland secretion and blood flow in the goose. *J. Physiol.* 213, 373–387. <https://doi.org/10.1113/jphysiol.1971.sp009387>.
58. Hughes, M.R. (1976). The effects of salt water adaptation on the Australian black swan, *Cygnus atratus* (Latham). *Comp. Biochem. Physiol. A Comp. Physiol.* 55, 271–277. [https://doi.org/10.1016/0300-9629\(76\)90143-2](https://doi.org/10.1016/0300-9629(76)90143-2).
59. Schmidt-Nielsen, K. (1960). The salt-secreting gland of marine birds. *Circulation* 21, 955–967. <https://doi.org/10.1161/01.CIR.21.5.955>.
60. McFARLAND, L.Z. (1964). Minimal salt load required to induce secretion from the nasal salt glands of sea gulls. *Nature* 204, 1202–1203.
61. Staaland, H. (1967). Anatomical and physiological adaptations of the nasal glands in Charadriiformes birds. *Comp. Biochem. Physiol.* 23, 933–944. [https://doi.org/10.1016/0010-406X\(67\)90354-4](https://doi.org/10.1016/0010-406X(67)90354-4).
62. Rubega, M.A., and Oring, L.W. (2004). Excretory organ growth and implications for salt tolerance in hatchling American avocets *Recurvirostra americana*. *J. Avian Biol.* 35, 13–15. <https://doi.org/10.1111/j.0908-8857.2004.03126.x>.
63. Dunson, W.A., Dunson, M.K., and Ohmart, R.D. (1976). Evidence for the presence of nasal salt glands in the roadrunner and the cottontail quail. *J. Exp. Zool.* 198, 209–216. <https://doi.org/10.1002/jez.1401980210>.
64. Carpenter, R.E., and Stafford, M.A. (1970). The secretory rates and the chemical stimulus for secretion of the nasal salt glands in the Rallidae. *Condor* 72, 316–324. <https://doi.org/10.2307/1366010>.
65. McFARLAND, L.Z. (1959). Captive marine birds possessing a functional lateral nasal gland (salt gland). *Nature* 184, 2030–2031.
66. Watson, G.E., and Divoky, G.J. (1971). Identification of *Diomedea leptorhyncha* Coues 1866, an albatross with remarkably small salt glands. *Condor* 73, 487–489.
67. Team, R.C. (2022). R: A language and environment for statistical computing.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Biological samples		
Pelicaniforme samples	Taiwan Biodiversity Research Institute	SPBorgan001
		SPBorgan002
		SPBorgan003
		SPBorgan004
		ASibis001
		ASibis002
		ASibis003
		BcNH001
		BcNH002
		MNH001
		MNH002
		MNH003
		Bubibis001
		Bubibis002
		Bubibis003
		GE001
GE002		
Deposited data		
Analyzed data	This paper	https://doi.org/10.17632/bkvjx52nsr.1
Software and algorithms		
R software	Team, R.C. ⁶⁷	https://www.r-project.org/
phytools	Revell ¹⁹	https://github.com/liamrevell/phytools
caper	Orme et al. ²³	https://github.com/cran/caper/blob/master/R/brunch.R
phylolm	Ho et al. ²⁴	https://github.com/lamho86/phylolm
RRphylo	Castiglione et al. ³²	https://github.com/cran/RRphylo
BayesTraits 3.0	Pagel and Meade ²⁶	http://www.evolution.reading.ac.uk/BayesTraitsV3/BayesTraitsV3.html

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the lead contact, Shou-Hsien Li (e-mail: t43028@ntnu.edu.tw).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- Data: All the raw data enrolled in this study have been deposited to the Digital Commons Data and are publicly available as of the date of publication. The accession number is listed in the [key resources table](#).
- Code: This paper does not report original code.
- Any additional information required to reanalyze the data reported in this paper is available from the [lead contact](#) upon request.

METHOD DETAILS

Collection of characteristics

Data on kidney and body weight were collected for 161 avian species and on SSGs for 224 species, both featuring known phylogenetic information from the literature.^{1,2,10,15,44,54–66} In 2023, we conducted searches using Google Scholar, utilizing various combinations of search terms such as 'salt gland', 'kidney size', and 'osmotic regulation' to obtain the most pertinent data. Because information on kidney weight was only published for two species of Pelicaniformes, we measured kidney and body weights from six additional Pelicaniformes species, namely the great egret (*Ardea alba*; n= 2), cattle egret (*Bubulcus ibis*; n= 3), black-crowned night heron (*Nycticorax nycticorax*; n= 2), Malayan night heron (*Gorsachius melanolophus*; n= 3), black-faced spoonbill (*Platalea minor*; n= 4), and African sacred ibis (*Threskiornis aethiopicus*; n= 3) which live in habitats with various levels of salinity (Table 1). All Pelicaniforme samples measured in this study were from carcasses of roadkill or of rescued individuals archived in the Taiwan Biodiversity Research Institute (TBRI). Before measurement, carcasses were stored in sealed plastic bags to prevent desiccation, and frozen at -20°C. After thawing, we followed the method of Maryanne (1970) to weigh the birds to the nearest 0.1 g; then, we took out the kidneys, dried them with absorbent paper, and weighed them to the nearest 0.001 g. Our subsequent analyses related to kidney size included 167 species and those relating to SSGs covered 230 species representing 25 avian orders.

We categorized the habitat type used by these birds into different levels of salinity: (1) completely freshwater (completely living in freshwater habitats, such as forests, grasslands, streams, lakes, and reservoirs.), (2) mainly freshwater (almost exclusively living in freshwater habitats; only a few records appear in saline habitats such as saltmarshes, salt pans, mangroves, seacoasts, beaches, and marine environments), (3) mainly saline (including the ocean and habitats associated with high salinity; sometimes appearing in freshwater habitats or just for parts of the year), and (4) completely saline habitats (exclusively living in habitats with high salinity) by assessing the habitat information on these species described in Birds of the World.¹⁶ The completely saline habitat has the highest level of environmental salinity, followed by mainly saline, mainly freshwater, and then completely freshwater.

Standardization of kidney sizes

Since the data obtained from the literature mostly only provides the average values of multiple individuals for each species, this study uses the average values of each species for subsequent analysis. Because both the body and kidney sizes of the avian species included in this study differ by up to five orders of magnitude (3 ~ 123,000 g for body weight; 0.03 ~ 920 g for kidney weight), we log-transformed body and kidney weights to better meet the statistical assumptions of a regression test. We used the residual from linear regressions ($\text{Log}(\text{kidney size}) = 1.165 \pm 0.073 + (0.928 \pm 0.0193) \log(\text{body weight})$ ($p < 2.2e-16$; adjusted r^2 : 0.933; Figure 3) of kidney weight on body weight as a proxy for relative kidney sizes to control the allometric effect.

Ancestral state reconstruction

To infer the macroevolutionary history of the ecological and physiological traits associated with salt tolerance, we extracted 10,000 avian phylogenetic trees of all 230 species based on the complete Bayesian maximum clade credibility (MCC) species-level avian phylogeny from <http://birdtree.org>¹⁷ with the Hackett constraint.¹⁸ Using the R package, phytools,¹⁹ we constructed a consensus tree based on the majority rule of the 10,000 trees we extracted. We estimated the branch length with a least square method. In the consensus tree of the 167 avian species the values were: Minimum: 0.188, 1st Quartile: 3.565, Median: 7.831, 3rd Quartile: 17.514, Maximum: 108.376. And in the consensus tree of the 230 avian species the values were: Minimum: 0.12, 1st Quartile: 3.252, Median: 7.337, 3rd Quartile: 15.979, Maximum: 108.376. Then, we used a Bayesian inference method (stochastic characteristic mapping with the "make.simmap" function) implemented in the phytools package for R²⁶ to reconstruct the ancestral state of the kidney size of the 167 avian species and SSG and habitat type of the 230 avian species. To compare the continuous data of residual kidney weight with the discrete data of SSGs and habitat type, we recoded the residual kidney weight data as a binary variable using the fourth quartile (the 75th percentile) of residual kidney weights as a reference. Species with residual kidney weights equal to or greater than 0.13 (the 75th percentile) were considered to have 'large kidneys', while those with residual kidney weights less than 0.13 were considered to have 'non-large' kidneys. The parameters were set with 1,000 MCMC generations and the ARD ("All Rates Different") model, which assumes that rates of trait evolution were not the same among all branch. The transition rates between different characteristic states were calculated by simulations, and posterior probabilities were mapped to the phylogeny using the "densityMap" function in phytools.¹⁹ The simulation involves two steps: first, simulating ancestral states at each internal node by sampling the posterior distribution of states; second, generating a substitution (mutational) history by sampling the posterior distribution conditioned on the reconstructions and observed states at the tips of the topology. Posterior probability values $\geq 50\%$ indicated a characteristic state being ancestral to a clade.

Phylogenetic signals in kidney sizes, possession of SSGs, and habitat salinity

After using the Shapiro-Wilk test to check the normality of the data, we used the function 'phylosig' in the package 'phytools'¹⁹ to calculate the Pagel's λ ²⁰ and Blomberg et al.'s K ²¹ for the phylogenetic signal in the variation of relative kidney sizes, the presence of SSGs, and the category of habitat. Assuming a Brownian motion model of trait evolution, a λ of 0 indicates that trait correlation between species is independent of their shared evolutionary history; by contrast, a λ of 1 suggests that trait correlation between species is constrained by their shared evolutionary history.²⁰ Also, assuming a Brownian motion model of trait evolution, K values greater than 1 indicate a higher trait variance among

clades than expected, whereas K values smaller than 1 imply a lower trait variance within clades than expected.²² Therefore, low K values suggest that their shared evolutionary history constrains the trait variance. We used phylogenetic generalized least-squares (PGLS) in the R package 'caper'²³ to control the phylogenetic effect and infer the association between variations in relative kidney size and habitat types with different levels of salinity and between variations in relative kidney size and the presence of SSGs. We used the phylogenetic generalized linear model (PGLM) in R package 'phylolm'²⁴ to infer the linear association between the possession of SSGs and habitat salinity.

Evolutionary transitions in kidney size, SSGs and levels of habitat salinity

We used the Discrete module of BayesTraits 3.0^{25,26} to estimate the transition rates of correlated evolution between habitat salinity, relative kidney sizes, and the possession of SSGs on a phylogeny. We took the dependent model approach, which assumes that the rate of change in one trait depends on the state of the other trait. We estimated the log Bayes factor (logBF) for the dependent model (allowing correlation between variables) against the independent model (null model, which fixes all correlations to be zero) as twice the difference between the estimated log marginal likelihoods using the formula $\log BF = 2 * (\log \text{marginal likelihood dependent model} - \log \text{marginal likelihood, independent model})$. Then, we interpreted comparisons where $\log BF > 2$ as having weak support, $\log BF > 5$ as having moderate support, and $\log BF > 10$ as having strong support to reject the null model.²⁷

Because the transition rate of correlated evolutionary analyses can only be calculated from binary data, residual kidney weight was transformed into a binary variable by reference to the fourth quartile of residual kidney weights; species with residual kidney weights equal to or larger than 0.13 (the 75th percentile) were considered to have 'large kidneys,' and those with residual kidney weights less than 0.13 were supposed to have 'non-large kidneys.' To test the sensibility of this cutoff value for our analysis, we repeated these analyses setting the cut-off points to the 25th and 50th percentiles of residual kidney weight, as suggested by Fristoe et al.²⁸ We similarly re-grouped the habitat categories into two by combining the entirely freshwater habitat and the mainly freshwater habitat to be 0, and the mainly saline habitat and the completely saline habitat to be 1. We ran an MCMC chain with 5.05 million iterations and a burn-in of 50,000 iterations and sampled every 1000 iterations. We scaled the branch length of the phylogenetic trees by 0.001 and used an exponential prior with a mean of 10 for all parameters.

Searching for morphological convergence among species in diverse saline habitat types

We used the RRphylo³² method to perform phylogenetic ridge regression on trees and data, yielding branch-wise evolutionary rates and ancestral character estimates (ACEs) at each node. This process is applied independently to each phenotype component for multivariate data, using a normalization factor to prevent extreme rate values and multicollinearity while assuming minimized rate variation within clades. We also employed search.conv,³² a fast and effective method for identifying phenotypic convergence among clades or species groups within specific categories. With search.conv, the phenotypic distance between species is quantified as the angle between their phenotypic vectors (i.e., multivariate phenotypes for each species). Under a Brownian motion model of evolution, this angle should increase proportionally to the patristic distance, the sum of the lengths of the branches that link two nodes in a tree, between species. However, when morphological convergence is present, the angle (per unit time) becomes smaller than the expected.

When comparing clades, the function calculates the mean angle over all possible combinations of species pairs, taking one species per clade, and divides this value by the patristic distance between the nodes subtending the clades (i.e., the phylogenetic distance between the most recent common ancestors, MRCAs, of the clades). This value is contrasted with a random distribution of 1,000 angle-by-time values to assess significance. When comparing species, the function randomly samples two species within the tree, computes the angle between them, and divides it by the patristic distance between their immediate ancestors.

Given two clades presumed to evolve under convergence, search.conv derives the ACEs for the MRCAs for both clades from the RRphylo results and calculates the angle between them. This angle is added to the mean angle between species and divided by the patristic distance between the MRCAs. If the "summed" angle is smaller than expected by chance, it indicates that the clades converged since their origin and subsequently followed parallel phenotypic evolutionary trajectories. The significance level is assessed as described above by randomizing phenotypes across the tree tips. Finally, if no specific hypothesis about converging clades is available, the function automatically scans the phylogeny to identify instances of convergence.

Under the "state case", search.conv computes the mean angle over all possible combinations of species pairs using one species per state. Each angle is divided by the patristic distance between the species. Significance is assessed by contrasting this value with a family of 1,000 random angles obtained by shuffling the state across the species.

In this study, we employed search.conv for both "clade" and "state" scenarios, by categorizing species based on their usage of saline and non-saline habitats. We further conducted the analyses by functional variable values (kidney sizes and presence of SSGs) for each species, utilizing these vectors as input for search.conv.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical Analysis Methods and Software: Used R v 4.2.3 language and related packages (phytools, caper, phylolm, RRphylo) and BayesTraits V3.0, to perform various statistical analyses, including ancestral state reconstruction, correlated evolutionary transition rate estimation, phylogenetic signal testing, phylogenetically adjusted least squares regression, phylogenetically adjusted generalized linear models, phylogenetic ridge regression, and detection of morphological convergence.

Location of Statistical Details: All statistical details of this research, such as the types of statistical tests, the exact value and meaning of sample size n , the definition and measurement of central tendency and dispersion, the setting of significance level, etc., were found in [Table 1](#) and [Method details](#).

Premises and Strategies of Statistical Analysis: Before performing statistical analysis, we tested the normality of the data and log-transformed or binarized the data to meet the assumptions of the statistical model. This study obtained all species with available characteristic values and phylogenetic information, stratified them based on their habitat types and physiological characteristics, and did not exclude any data or species during the analysis process.