

# Effects of *Lecanicillium lecanii* strain JMC-01 on the physiology, biochemistry, and mortality of *Bemisia tabaci* Q-biotype nymphs

Ting Xie, Ling Jiang, Jianshe Li, Bo Hong, Xinpu Wang and Yanxia Jia

School of Agriculture, Ningxia University, Yinchuan, Ningxia, China

## ABSTRACT

**Background:** *Lecanicillium lecanii* is an entomopathogenic fungi, which was isolated from insects suffering from disease. Now, it is an effective bio-control resource that can control agricultural pests such as whitefly and aphids. There are many studies on the control of various agricultural pests by *L. lecanii*, but no report on its control of *Bemisia tabaci* biotype-Q exists. In this work, we studied the susceptibility of *B. tabaci* Q-biotype (from Ningxia, China) to *L. lecanii* JMC-01 in terms of nymph mortality and the changes in detoxifying protective enzymes activities.

**Methods:** *B. tabaci* nymphs were exposed to *L. lecanii* JMC-01 conidia by immersion with the host culture. Mortality was assessed daily for all nymph stages. The detoxifying and protective enzyme activity changes, weight changes, and fat, and water contents of the nymphs were determined spectrophotometrically.

**Results:** All instars of *B. tabaci* died after being infested with  $1 \times 10^8$  conidia/mL. The 2nd-instar nymphs were the most susceptible, followed by the 3rd-instar nymphs. The corrected cumulative mortality of the 2nd- and 3rd-instar nymphs was 82.22% and 75.55%, respectively. The levels of detoxifying and protective enzymes initially increased and then decreased. The highest activities of carboxylesterase, acetylcholinesterase, peroxidase, and catalase occurred on the 3rd day, reaching 10.5, 0.32, 20, and 6.3 U/mg prot, respectively. These levels were 2.2-, 4.3-, 2.4-, and 1.4-fold the control levels, respectively. The highest activities of glutathione-S transferase and superoxide dismutase on the 2nd day were, respectively, 64 and 43.5 U/mg prot. These levels were, respectively, 2.7 and 1.1-fold that of the control level. The water and fat content in the infested *B. tabaci* nymphs decreased and differed significantly from the control levels. The weight increased continuously in the first 24 h, decreasing thereafter. At 72 h, the infestation level was about 0.78-fold that of the control level.

**Conclusions:** The studied *L. lecanii* JMC-01 strain is pathogenic to the *B. tabaci* Q-biotype. This strain interferes with the normal functioning of detoxifying and protective enzymes, and is also involved in the disruption of normal physiological metabolism in *B. tabaci*.

Submitted 9 February 2019  
Accepted 19 August 2019  
Published 16 September 2019

Corresponding author  
Yanxia Jia,  
helenjia\_2006@nxu.edu.cn

Academic editor  
Jasmine Janes

Additional Information and  
Declarations can be found on  
page 18

DOI [10.7717/peerj.7690](https://doi.org/10.7717/peerj.7690)

© Copyright  
2019 Xie et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Agricultural Science, Entomology, Mycology

**Keywords** *Lecanicillium lecanii* JMC-01, *Bemisia tabaci*, Protective enzymes, Detoxification enzymes, Physiological and biochemical metabolism, Mortality

## INTRODUCTION

The whitefly or tobacco whitefly *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) is a cosmopolitan insect pest with more than 900 documented host plant species. This species is considered to belong to a cryptic species complex with more than 40 morphotypes distributed across the biotypes, with the B- and Q-biotypes being the most important (Tang et al., 2018). The whitefly is of economic importance due to its direct (by sapping plant fluids and vectoring plant pathogens) and indirect (phytosanitary and quarantine measures) damage to crops (De Barro, 2011; Xu et al., 2014). Its control mainly relies on chemical pesticide application, which has resulted in the development of insecticide resistance. In addition to the emergence of resistant strains, farms and other stakeholders are challenged by safety concerns. Pesticide application causes environmental pollution, alters the abundance of natural enemies, increases pest resistance, and promotes secondary pest population resurgence (Liu et al., 2009). Environmentally-friendly pest management methods such as biological control using natural enemies and entomopathogen microorganisms (bacteria, fungi, and viruses) are being established worldwide in response to this.

Entomopathogenic fungi were the first microorganisms identified as insect pathogens, whereas entomopathogenic bacteria were the first to be commercialized (Lacey et al., 2001). *Lecanicillium lecanii* (= *Verticillium lecanii* (Zimmerman) Viegas) belongs to Deuteromycotina, Hyphomycetes, Moniliales, Moniliaceae, that is widely used as entomopathogenic fungi in bio-control up to now. And the entomopathogenic fungal species described and commercialized, *L. lecanii* (Zare & Gams, 2001) deserves further consideration as a broad range commercial biopesticide, due to its wide range of hosts and wide geographical distribution (Xie et al., 2015). Indeed, this species can infect the diamondback moth *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) (Ravindran et al., 2018), aphids (Hemiptera: Aphididae) (Askary, Benhamou & Brodeur, 1999), the citrus mealybug *Planococcus citri* Risso (Hemiptera: Pseudococcidae) (Ghaffari et al., 2017), and the soybean cyst nematode *Heterodera glycines* Ichinohe (Tylenchida: Heteroceridae) (Shinya et al., 2008), and has also been documented to infect *Bemisia tabaci* (Zhu & Kim, 2011). In insects, the spores of entomopathogenic fungi germinate, and the fungal hyphae penetrate the epidermis and invade the tissues and organs until reaching the hemocoel (Duan et al., 2017). When the hyphae come into contact with the hemolymph, the defense system of the insects, which includes detoxifying and protective enzymes, is induced (Liu et al., 2013).

Physiological and biochemical approaches have been used to describe the chronological events leading to fungal infestation success in an insect host. Reactive oxygen species (ROS) are forms of atmospheric oxygen (Tian et al., 2016b) produced in the mitochondria that are equilibrated by cellular antioxidative mechanisms (Esmail et al., 2018). In many instances, microbial pathogens are associated with an increase in ROS, which induces an oxidative stress response in the host (Foyer & Noctor, 2013). The antioxidative mechanism of the cells includes antioxidant enzymes, such as catalase (CAT), superoxide dismutase (SOD), and peroxidase (POD), which degrade H<sub>2</sub>O<sub>2</sub> to reduce oxidative damage

(*Felton & Summers, 1995*). In addition to this antioxidative mechanism, insects also harbor detoxifying enzymes, such as carboxylesterase (CarE), glutathione-S transferase (GST), and acetylcholinesterase (AChE), which are able to metabolize exogenous toxicants (*Xu et al., 2006*), and have been the target of insecticide synergist research (*Wang et al., 2016*).

The effects of these insect detoxifying enzymes in response to the fungal entomopathogen *L. lecanii* in the spiraling whitefly *Aleurodicus dispersus* Russell (Hemiptera: Aleyrodidae) have recently been demonstrated (*Liu et al., 2013*). These changes in defensive enzymes are deserving further attention, due to its practical considerations.

Due to the lack of studies and the economic importance of *Bemisia tabaci*, the objective of this study was to determine the pathogenic effect of *L. lecanii* strain JMC-01 at the nymphal stages of *Bemisia tabaci* by evaluating the disruption of immune mechanisms.

## MATERIALS AND METHODS

### Entomopathogen strain and insect collection

*Lecanicillium lecanii* strain: the *L. lecanii* strain JMC-01 was isolated from *Bemisia tabaci* infected nymphs from a greenhouse in Yinchuan, Ningxia (38°33'N, 106°08'E), China in May 2017. The JMC-01 strain was deposited at the China Center for Type Culture Collection with the accession number [M2018303](#). The strain status was determined based on ITS sequence divergence to the reference strain (*Jiang, 2018*). The JMC-01 strain reference ITS nucleotide sequence was deposited in GenBank with the identification number [MH312006](#).

Insect: the whitefly *Bemisia tabaci* Q-biotype was collected from a tomato greenhouse in Yinchuan, Ningxia (38°33'N, 106°08'E) in July 2018. Biotype assignment was performed as previously described (*Gao, 2018*). The tomato cultivar Bijiao was planted in a greenhouse in Yinchuan, Ningxia (38°33'N, 106°08'E) and cultivated using drip irrigation technology. Tomato was used as the host plant for two generations, following which the synchronized nymphs were collected for experimentation.

### Preparation of the *L. lecanii* JMC-01 conidial suspension

The *L. lecanii* JMC-01 strain was inoculated on potato dextrose agar (PDA) plates, at 28 °C with a 12:12 (L:D, light:dark) photoperiod for 7 days (MJ-250 Mould Incubator; Jiangsu Zhengji Instruments Co. Ltd., Jiangsu, China). Spore suspensions were prepared by recovering the conidia from the PDA plates with a 0.05% Tween-80 solution. The solution was filtered with sterile cheesecloth to eliminate the hyphae, following which the concentration was adjusted to  $1.0 \times 10^8$  conidia/mL with sterile water using a hemocytometer (Qiujing, Shanghai, China).

### *Bemisia tabaci* nymph mortality induced by *L. lecanii* JMC-01

Tomato leaves with 1st-, 2nd-, 3rd- or 4th-instar nymphs (only one leaf was selected for each instar nymph) were immersed in *L. lecanii* JMC-01 solution at  $1.0 \times 10^8$  conidia/mL for, 30 s or in a control solution of 0.05% Tween-80. After immersion, each leaf was sealed in a standard Petri dish, with its petiole wrapped in a moistened cotton ball. The plates were incubated in an artificial climate chamber (RQX-250; Shanghai Yuejin

Medical Devices Co., Ltd., Shanghai, China) at  $28 \pm 2$  °C,  $70\% \pm 10\%$  RH, and 12:12 (L:D) photoperiod. There were three replicates per treatment. Deaths were recorded daily, and the cumulative corrected mortality was calculated as follows:

$$\text{Accumulative corrected mortality (\%)} = \frac{\text{Infection mortality} - \text{Control mortality}}{1 - \text{Control mortality}} \times 100\%$$

### Susceptibility of 3rd-instar *Bemisia tabaci* nymphs to different JMC-01 concentrations

The *L. lecanii* JMC-01 suspensions were prepared as described above at different conidial concentrations:  $1 \times 10^8$ ,  $1 \times 10^7$ ,  $1 \times 10^6$ ,  $1 \times 10^5$ , and  $1 \times 10^4$  conidia/mL.

Three tomato leaves with 3rd-instar *Bemisia tabaci* nymphs were immersed for, 30 s at each JMC-01 test concentration, and the leaves were incubated as described above. Deaths were recorded on a daily basis, and were used to determine the cumulative corrected mortality for each conidial concentration.

### Protective and detoxifying enzyme activity determination

Tomato leaves with 3rd-instar *Bemisia tabaci* nymphs were infected with *L. lecanii* JMC-01 at  $1 \times 10^8$  conidia/mL, using the immersion procedure described above. Treated and control (0.05% Tween-80) leaves were immersed in *L. lecanii* JMC-01 solution at  $1.0 \times 10^8$  conidia/mL for 30 s. After immersion, each leaf was sealed in a standard Petri dish, with its petiole wrapped in a moistened cotton ball. The plates were incubated in an artificial climate chamber (RQX-250; Shanghai Yuejin Medical Devices Co., Ltd., Shanghai, China) at  $28 \pm 2$  °C,  $70\% \pm 10\%$  RH, and 12:12 (L:D) photoperiod.

Sample processing: the animal tissue, was weighted and nine-times the volume of normal saline by weight was added (weight (g):volume (mL) = 1:9), the samples were then ground with liquid nitrogen to make a 10% tissue homogenate, which was then centrifuged at 2,500 rpm for 10 min (Sigma D-37520; Sigma-Aldrich; Nanjing Beiden Medical Co., Ltd. Nanjing, China). The supernatant was then diluted to 1% tissue homogenate with normal saline for experimentation.

### Protein content determination

The 563 µg/mL standard solution, working fluid, stop application solution and normal saline were purchased from the Jian Cheng Bioengineering Institute (Nanjing, China).

After combining the solutions, they were placed at room temperature for 5 min, and measured colorimetrically at 562 nm (L5S UV Spectrophotometer; Shanghai Yidian Analytical Instrument Co., Ltd., Shanghai, China) (Table 1). Double-distilled water served as the blank control.

The protein concentration was determined as follows:

$$\text{Protein}(\mu\text{gprot/mL}) = \frac{\text{Measure OD} - \text{Blank OD}}{\text{Standard OD} - \text{Blank OD}} \times \text{Standard solution (563}\mu\text{g/mL)} \\ \times \text{Sample dilution before determination}$$

**Table 1** The steps of protein content determination.

	Blank tube	Standard tube	Measuring tube
Double distilled water (μL)	20		
563 μg/mL standard solution (μL)		20	
Sample (μL)			20
Working fluid (μL)	250	250	250
Mix, set at 37 °C water bath for 30 min (digital thermostat water bath)			
Stop application solution (μL)	750	750	750

**Table 2** The steps of SOD activity determination.

Reagent	Measuring tube	Control tube
Reagent one application solution (mL)	1.0	1.0
Sample (mL)	0.1	
Double distilled water (mL)		0.1
Reagent two solution (mL)	0.1	0.1
Reagent three solution (mL)	0.1	0.1
Reagent four application solution (mL)	0.1	0.1
Mix, set at 37 °C water bath for 40 min (digital thermostat water bath)		
Chromogen solution (mL)	2	2

### SOD activity determination

Reagent one application solution, reagent two solution, reagent three solution, reagent four application solution, chromogen solution, and normal saline were purchased from the Jian Cheng Bioengineering Institute, Nanjing, China.

After combining the solutions, they were placed at room temperature for 10 min, and measured colorimetrically at 550 nm (Table 2). Double-distilled water served as the blank control.

Superoxide dismutase activity was determined as follows:

$$\text{SOD(U/mg prot)} = \frac{\text{Control OD} - \text{Measure OD}}{\text{Control OD}} \div 50\% \times \frac{\text{Total volume of reaction solution}}{\text{Sample size (mL)}} \div \text{Protein concentration of the sample to be tested (mg prot/mL)}$$

### POD activity determination

Reagent one solution, reagent two application solution, reagent three application solution, reagent four solution, and normal saline were purchased from the Jian Cheng Bioengineering Institute, Nanjing, China.

The solutions were combined and centrifuged at 3,500 rpm for 10 min (Sigma D-37520; Sigma-Aldrich, Germany), following which the supernatant was measured colorimetrically at 420 nm (Table 3). Double-distilled water served as the blank control.

**Table 3** The steps of POD activity determination.

	Blank tube	Measuring tube
Reagent one solution (mL)	2.4	2.4
Reagent two application solution (mL)	0.3	0.3
Reagent three application solution (mL)	0.2	0.2
Double distilled water (mL)	0.1	
Sample (mL)		0.1
Set at 37 °C water bath for 30 min (digital thermostat water bath)		
Reagent four (mL)	1.0	1.0

**Table 4** The steps of CAT activity determination.

	Control tube	Measuring tube
Sample (mL)		0.05
Reagent one solution (37 °C preheat) (mL)	1.0	1.0
Reagent two solution (37 °C preheat) (mL)	0.1	0.1
Mix, set at 37 °C water bath for 1 min (digital thermostat water bath)		
Reagent three solution (mL)	1.0	1.0
Reagent four solution (mL)	0.1	0.1
Sample (mL)	0.05	

Peroxidase activity was determined as follows:

$$\text{POD(U/mg prot)} = \frac{\text{Measure OD} - \text{Blank OD}}{12 \times 1} \times \frac{\text{Total volume of reaction solution}}{\text{Sample size (mL)}} \\ \div \text{Reaction time (30 min)} \\ \div \text{Protein concentration of the sample to be tested (mg prot/mL)} \\ \times 1,000$$

### CAT activity determination

Reagent one solution, reagent two solution, reagent three solution, reagent four solution, and normal saline were purchased from the Jian Cheng Bioengineering Institute, Nanjing, China.

After combining the solutions, they were measured colorimetrically at 405 nm (Table 4). Double-distilled water served as the blank control.

Catalase activity was determined as follows:

$$\text{CAT(U/mg prot)} = (\text{Control OD} - \text{Measure OD}) \times 271 \times \frac{1}{60 \times 0.05} \\ \div \text{Protein concentration of the sample to be tested (mg prot/mL)}$$

### CarE activity determination

The working fluid and normal saline were purchased from the Jian Cheng Bioengineering Institute, Nanjing, China.

Sample processing: the sample processing was as described in the protein content determination step above, except that the tissue homogenate was centrifuged at 12,000 rpm for 4 min.

**Table 5** The steps of AchE activity determination.

	Measuring tube	Control tube	Standard tube	Blank tube
Sample (mL)	0.1			
One $\mu\text{mol/mL}$ standard application solution (mL)			0.1	
Double distilled water (mL)				0.1
Substrate buffer (mL)	0.5	0.5	0.5	0.5
Chromogen application solution (mL)	0.5	0.5	0.5	0.5
Mix, set at 37 °C water bath for 6 min (digital thermostat water bath)				
Inhibitor solution (mL)	0.03	0.03	0.03	0.03
Transparent solution (mL)	0.1	0.1	0.1	0.1
Sample (mL)		0.1		

**Note:**

The same sample was added to the control tube and the measuring tube, but the order was different. The blank tube was not sampled and distilled water was used instead of the sample.

The steps were as follows:

1. The spectrophotometer was preheated for at least 30 min and the wavelength was adjusted to 450 nm. The machine was blanked with double-distilled water.
2. The working fluid was preheated at 37 °C for at least 30 min.
3. Blank tube: Five  $\mu\text{L}$  of distilled water was added to a blank glass cuvette, to which 1,000  $\mu\text{L}$  of preheated working solution was sequentially added to a one mL glass cuvette. The solution was rapidly mixed, and light absorption  $A_1$  and  $A_2$  was measured at 450 nm 10 and 190 s,  $\Delta A_{\text{Blank tube}} = A_2 - A_1$ .
4. Measuring tube: Five  $\mu\text{L}$  of supernatant was sequentially added to a one ml glass cuvette, 1,000  $\mu\text{L}$  of preheated working solution, and rapidly mixed, and light absorption of  $A_3$  and  $A_4$  were measured at 450 nm,  $\Delta A_{\text{Measuring tube}} = A_4 - A_3$ .

$$\text{CarE(U/mg prot)} = (\Delta A_{\text{Measuring tube}} - \Delta A_{\text{Blank tube}}) \times V \div (\text{Cpr} \times V_{\text{Sample}}) \div T$$

$V$ : total volume of the reaction solution, 1.005 mL;

$\text{Cpr}$ : protein concentration of the sample to be tested (mg prot/mL);

$V_{\text{Sample}}$ : adding of supernatant volume to the reaction system (mL), 0.005 mL;

$T$ : catalytic reaction time (min), 3 min.

### AchE activity determination

One  $\mu\text{mol/mL}$  standard application solution, substrate buffer, chromogen application solution, inhibitor solution, transparent solution, and normal saline were purchased from the Jian Cheng Bioengineering Institute, Nanjing, China.

After combining the solutions, they were placed at room temperature for 15 min and measured colorimetrically at 412 nm (Table 5). Double-distilled water served as the blank control.

**Table 6** The steps of enzymatic reaction.

	Measuring tube	Control tube
Matrix fluid (mL)	0.3	0.3
Sample (mL)	0.1	
Mix, set at 37 °C water bath for 10 min (digital thermostat water bath)		
Reagent two application solution (mL)	1	1
Anhydrous alcohol (mL)	1	1
Sample (mL)		0.1

**Table 7** The steps of chromogen reaction.

	Blank tube	Standard tube	Measuring tube	Control tube
GSH standard application solution (mL)	2			
20 µmol/mL GSH standard solution (mL)		2		
Supernatant (mL)			2	2
Reagent three application solution (mL)	2	2	2	2
Reagent four application solution (mL)	0.5	0.5	0.5	0.5

**Note:**

The same sample was added to the control tube and the measuring tube, but the order was different. The blank tube was not sampled and distilled water was used instead of the sample.

AchE activity was determined as follows:

$$\text{AchE(U/mg prot)} = \frac{\text{Measure OD} - \text{Control OD}}{\text{Standard OD} - \text{Blank OD}} \times \text{Standard concentration (1 } \mu\text{mol/mL)} \div \text{Protein concentration of the sample to be tested (mg prot/mL)}$$

**GST activity determination**

Matrix fluid, reagent two application solution, anhydrous alcohol, GSH standard application solution, 20 µmol/mL GSH standard solution, reagent three application solution, reagent four application solution, and normal saline were purchased from the Jian Cheng Bioengineering Institute, Nanjing, China.

Enzymatic reaction:

The solutions were combined and centrifuged at 3,500 rpm for 10 min (Sigma D-37520; Sigma-Aldrich, Germany) (Table 6). The supernatant was then used in the chromogen reaction.

Chromogen reaction:

The solutions were combined and placed at room temperature for 15 min, following which they were measured colorimetrically at 412 nm (Table 7). Double-distilled water served as the blank control.

Glutathione-S transferase activity was determined as follows:

$$\text{GST(U/mg prot)} = \frac{\text{Control OD} - \text{Measure OD}}{\text{Standard OD} - \text{Blank OD}} \times \text{Standard concentration (20 } \mu\text{mol/mL)} \times \text{Reaction system dilution factor (6 times)} \div \text{Reaction time (10 min)} \div [\text{Sample volume (0.1 mL)}] \times \text{Protein concentration of the sample to be tested (mg prot/mL)}$$



## Determination of weight, and water and fat content of the *Bemisia tabaci* nymphs after infestation with *L. lecanii* JMC-01

Tomato leaves with 3rd-instar *Bemisia tabaci* nymphs exposed to  $1.0 \times 10^8$  conidia/mL or the control treatment (0.05% Tween-80). The treated and control leaves were placed in similar Petri dishes. The same Petri dish method as above was then used.

The treatment and control group were selected one hundred 3rd-instar nymphs for experimentation after 0, 12, 24, 36, 48, 60, and 72 h, respectively. First determining the total fresh weight of 100 nymphs prior to infection (Mettler Toledo LE204E/02 electronic balance), the nymphs were dried by placing each batch at 60 °C for 48 h in an electrothermal blowing dry box (Shanghai Yiheng Technology Co., Ltd., Shanghai, China), and weighed in a similar method as for the determination of dry weight (dry mass, DM).

Water content (WC) was determined using the formula  $WC = (FW - DM)/FW \times 100\%$ , where DW is the dry mass determined as explained above, and FW is the fresh weight determined as above.

Lipid extraction was performed with the dried nymphs. The dried nymphs were grinded under liquid nitrogen in a centrifuge tube. One mL of chloroform isoamyl alcohol (24:1) and 0.5 mL of methanol (99.99%) was added to each tube, mixed, and then centrifuged at 4,500 rpm for 10 min. The supernatant was discarded. The precipitate was extracted again with one mL of chloroform isoamyl alcohol (24:1) and 0.5 mL of methanol (99.99%) by centrifugation at 4,500 rpm for 10 min. The final remaining precipitate was dried in an oven at 60 °C for 48 h to determine the constant dry mass (LDM).

Fat content (FC) was determined using the formula  $FC = (DM - LDM)/DM \times 100\%$ , where DM is the dry mass determined as explained above, and LDM is the constant dry mass determined after lipid extraction.

There were three replicates per treatment and time point, and 100 nymphs per replicate.

## Data analysis

Excel 2010 (Microsoft Corporation, Albuquerque, NM, USA) was used to process all the data. All results are expressed as the mean  $\pm$  standard deviation. Statistical analysis of the data was performed using one-way analysis of variance with SPSS version 21.0 (SPSS; IBM Corp., Armonk, NY, USA). Multiple comparisons of the means were performed using Duncan's (*D*) tests at a significance level of  $P = 0.05$ . All figures were produced using Origin 8.0.


## RESULTS

### Morphological characteristics of the *Bemisia tabaci* nymphs

Figure 1 shows the morphological characteristics of *Bemisia tabaci* under *L. lecanii* JMC-01 infection as observed under a microscope (Leica Microsystems Wetzlar GmbH, Wetzlar, Germany). The surface is covered with hyphae.


### Mortality of the *Bemisia tabaci* nymphs

Figure 2 indicates the cumulative mortality induced by *L. lecanii* JMC-01 to each *Bemisia tabaci* immature stage. The cumulative corrected mortality of the nymph instars was as follows (from high to low): 2nd instar > 3rd instar > 1st instar > 4th instar > egg.




**Figure 1** Morphological characteristics of the *B. tabaci* nymph induced by *L. lecanii* JMC-01.

Full-size DOI: 10.7717/peerj.7690/fig-1



**Figure 2** Cumulative corrected mortality of *L. lecanii* JMC-01 infestation on *B. tabaci* nymphs. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point.

Full-size DOI: 10.7717/peerj.7690/fig-2



**Figure 3** Cumulative corrected mortality of the 3rd-instar *B. tabaci* nymphs following exposure to different concentrations of *L. lecanii* JMC-01. Each data point indicates the corrected cumulative mortality for each time period. [Full-size !\[\]\(ba1b80118482ccef74a5d718ca4d7242\_img.jpg\) DOI: 10.7717/peerj.7690/fig-3](https://doi.org/10.7717/peerj.7690/fig-3)

The 2nd- and 3rd-instar nymphs were most affected, with corrected cumulative mortality percentages of 82.22% and 75.55%, respectively.

### The initial dose of *L. lecanii* JMC-01 affects the 3rd-instar *Bemisia tabaci* nymphs

As indicated in Fig. 3, increasing doses of *L. lecanii* JMC-01 (from  $1 \times 10^4$  to  $1 \times 10^8$  conidia/mL) also increased the corrected cumulative mortality of the 3rd-instar nymphs, reaching a maximum of 75.55% at  $1 \times 10^8$  conidia/mL after 6 days.


### Protective and detoxifying enzyme activity determination

The highest activity of SOD (43 U/mg prot) was detected on the 2nd day, reaching 1.1-fold that of the control (Fig. 4). The highest activities of POD and CAT were 20 and 6.3 U/mg prot on the 3rd day, respectively, and reached 2.4- and 1.4-fold that of the control level (Figs. 5 and 6). Following this, the activities of protective enzymes decreased. The lowest activities of SOD, POD, and CAT were 30, 8.5, and 1.3 U/mg prot on the 5th day, respectively (Figs. 4–6).


The highest activities of CarE and AchE were 10.5 and 0.32 U/mg prot. These levels were observed on the 3rd day and were 2.2- and 4.3-fold that of the control level, respectively (Figs. 7 and 8). The highest GST activity was 64 U/mg prot on the 2nd day and was 2.7-fold that of the control level (Fig. 9). After the 3rd day, the activities of detoxifying enzymes decreased, and the lowest activities of CarE, AchE, and GST, respectively, reached 3.5, 15, and 0.05 U/mg prot on the 5th day (Figs. 7–9).

### Determination of the weight and water and fat contents of the of *Bemisia tabaci* nymphs

The lowest changes in weight were observed at 24–36 h. At 72 h, the weight of the infected group was 0.78-fold that of the control (Fig. 10).




**Figure 4** Effects of SOD activities of the 3rd-instar *B. tabaci* nymphs infested with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size](#) DOI: 10.7717/peerj.7690/fig-4




**Figure 5** Effects of POD activities of the 3rd-instar *B. tabaci* nymphs infested with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size](#) DOI: 10.7717/peerj.7690/fig-5

The WC of *Bemisia tabaci* continuously decreased after infection with *L. lecanii*. At 72 h, the WCs of the infected and control groups were lowest reaching 56% and 66%, respectively (Fig. 11).




**Figure 6** Effects of CAT activities of the 3rd-instar *B. tabaci* nymphs infested with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac\_img.jpg\) DOI: 10.7717/peerj.7690/fig-6](https://doi.org/10.7717/peerj.7690/fig-6)




**Figure 7** Effects of CarE activities of the 3rd instar nymph of *B. tabaci* infested with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size !\[\]\(0c0f8cc6eca4f663c17a652926046967\_img.jpg\) DOI: 10.7717/peerj.7690/fig-7](https://doi.org/10.7717/peerj.7690/fig-7)

Until 36 h after infection, the changes in FC were not significantly different from the control level. At 72 h, the FC of the infected and control groups was the lowest, reaching 13% and 20.5%, respectively (Fig. 12).




**Figure 8** Effects of AchE activities of the 3rd instar nymph of *B. tabaci* infested with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90\_img.jpg\) DOI: 10.7717/peerj.7690/fig-8](https://doi.org/10.7717/peerj.7690/fig-8)




**Figure 9** Effects of GST activities of the 3rd instar nymph of *B. tabaci* infested with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size !\[\]\(b5d7dedcc48d5bfd2c56b334ed39e34f\_img.jpg\) DOI: 10.7717/peerj.7690/fig-9](https://doi.org/10.7717/peerj.7690/fig-9)

## DISCUSSION

The fungus penetrated the insect epidermis via the germ tubes and appressoria, following which the conidia invaded the nymphs and began to enter the hemocoel. Ultimately,




**Figure 10** Changes in weight of the 3rd instar *B. tabaci* nymphs infected with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size !\[\]\(fd7fe780e8fd8eece60268c87d0c3e04\_img.jpg\) DOI: 10.7717/peerj.7690/fig-10](https://doi.org/10.7717/peerj.7690/fig-10)



**Figure 11** Changes in water content of the 3rd instar *B. tabaci* nymphs infected with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size !\[\]\(86257f54800c9844bc7e863bea396fba\_img.jpg\) DOI: 10.7717/peerj.7690/fig-11](https://doi.org/10.7717/peerj.7690/fig-11)

the hyphae covered the host surface and had colonized the body cavity (Zhou *et al.*, 2017). Previously, *L. lecanii* caused over 90% mortality of vegetable pest, such as aphids, *Plutella xylostella* (Ravindran *et al.*, 2018; Saruhan, 2018; Sugimoto *et al.*, 2003). In this



**Figure 12** Changes in and fat content of the 3rd instar *B. tabaci* nymphs infected with *L. lecanii* JMC-01. Times marked with different uppercase letters on the same line are significantly different ( $P < 0.05$ ). The different lowercase letters indicate significant differences between the treatment and control groups ( $P < 0.05$ ) at the same time point. [Full-size !\[\]\(5fd6ef84f97f42d7f8b34275f1b65312\_img.jpg\) DOI: 10.7717/peerj.7690/fig-12](https://doi.org/10.7717/peerj.7690/fig-12)

study, mortality increased greatly during the first 5 days of infection, with the maximum mortality is 82.22% being reached on the 6th day. Accordingly, the activities of detoxification and protective enzymes were lowest on the 5th day, indicating that as the infected nymphs of *Bemisia tabaci* neared death on the 5th day, their enzyme activity was reduced.

Insects are protected from the stresses of adverse conditions by various physical barriers, including a cuticular exoskeleton, peritrophic membrane, and an immune system that reduces pathogen infection (Chen & Lu, 2017). These fungi stimulate the stress responses of the insect detoxification system and the protective enzyme system under adverse conditions by changing the function of ion channels (Zhang et al., 2017). The major components of the antioxidant defense system of insects include the antioxidant enzymes SOD, CAT, and POD (Li et al., 2016b). When insects are stimulated by exogenous compounds, SOD converts the superoxide radical  $O_2^-$  into  $H_2O_2$ . Then, POD and CAT convert the  $H_2O_2$  into  $H_2O$ . The imbalance between oxidative stress and antioxidant responses contributes to disease and the death of insect hosts (Felton & Summers, 1995).

Our study showed that, after infection of *Bemisia tabaci* by *L. lecanii*, the activities of SOD, CAT, and POD initially increased but then decreased thereafter, and the maximum activities protective enzymes were observed on the 2nd day or 3rd day. Previous studies (Yang et al., 2015; Ye et al., 2018; Zhou et al., 2017) indicated agricultural insects infested by entomogenous fungus, the activities of SOD, CAT, and POD initially increased but then decreased. The increased activity of SOD, CAT, and POD effectively preventing the formation of more toxic substances such as hydroxyl radicals and helped



increase the resistance of *Bemisia tabaci* (Ding, Zhang & Chi, 2015). Under *L. lecanii* infection, ROS and other toxic substances stimulated an immune system response in *Bemisia tabaci* (Li et al., 2016a). To resist the adverse environmental influence and maintain normal physiological activities, the enzyme activities sharply increased. However, the internal spread of the pathogen led to the destruction of the internal tissue structure of the insect and subsequent collapse of the immune system. In addition, the ROS scavenging system might not have been able to remove the excessive quantity of free radicals, leading to reduced enzyme activity and the death of the insect (Li et al., 2016a). So, the activities of SOD, CAT, and POD were decreased on the 5th day. GSTs participate in detoxification metabolism and catalyze a combination of toxic substances with glutathione and also promote the excretion of toxic chemicals and pathogenic substances (Mathews et al., 2002; Schama et al., 2016). CarEs can catalyze the hydrolysis of ester bonds, and their major physiological functions include lipid metabolism, detoxification metabolism of exogenous compounds and biochemical regulatory functions (Guo et al., 2015). AchE is a target for organophosphorus and carbamate insecticides (Ding et al., 2001). Some exogenous compounds, such as pesticides and pathogenic fungi, can be altered by insect detoxification enzymes. This suggests that *L. lecanii* can promote the detoxification metabolism of *Bemisia tabaci*, which is beneficial for the discharge of exogenous toxicants. With the increase in the level of *Bemisia tabaci* infection with *L. lecanii* exposure time, the exogenous toxicants overpowered the detoxification metabolism, resulting in the eventual reduction in enzyme activities, and ultimately, insect death. We found that the activities of CarEs, AchE, and GST initially increased but then decreased, and the maximum activities of CarEs, AchE, and GST were observed on the 2nd day or 3rd day after infection. Effects of *Isaria fumosorosea* infection on different enzyme activities in the adult in vivo of *Bemisia tabaci* indicated that the maximum activities of GSTs and CarE were observed on the 48–60 h (Tian, Diao & Ma, 2016a). Besides, these findings are similar to previous study (Liu et al., 2013; Zhang et al., 2015). Insects infested with entomopathogenic fungi initially exhibit elevated enzyme activities that decline as the fungal infection continues (Tian, Diao & Ma, 2016a). The entomopathogenic fungus *L. lecanii* can be used to control *Bemisia tabaci* nymphs, but the prevention and control effect is slower than with chemical insecticides. However, the use of entomopathogenic fungi in combination with insecticides to control pests could increase their speed and efficacy (Purwar & Sachan, 2006).

A study of the pathogenicity and control potential of *Beauveria bassiana* on the onion fly showed that the weight increment was smallest after 48 h. The WC and FC continued to decrease, and the WC and FC of the infection level were 0.81- and 0.69-fold that of the control level, respectively, at 72 h (Zhang, 2017). Screening of the strains of the highly pathogenic *Beauveria bassiana* on soybean pod borers and the assessment biophysical and biochemical effects on their hosts indicated reductions in weight, WC, and FC (Tian, 2014). These studies corroborate our findings. In the present study, *Bemisia tabaci* nymphs infested with *L. lecanii* JMC-01 gradually lost vitality until death. This process causes many physiological changes in the insects. Thus, *L. lecanii* could constitute a useful alternative biopesticide for *Bemisia tabaci* population management.

Biocontrol can reduce insecticide resistance and increase environmental and non-target organism safety.

## CONCLUSIONS

We observed that *L. lecanii* JMC-01 affected the viability of the *B. tabaci* Q-biotype, by inducing mortality, affecting the activities of protective and detoxifying enzymes, and by significantly reducing the weight, and water and FC. Thus, *L. lecanii* impacted the physiological functioning of *B. tabaci* by directly acting on molecular targets and by indirectly acting on detoxification and protective enzymes (Bantz *et al.*, 2018). These results indicate that this fungal strain could constitute an effective biological control for *B. tabaci* in agriculture.

## ACKNOWLEDGEMENTS

We are grateful to Master Kai Gao for providing the biotype of *B. tabaci* and we to Master Hui Wang for helping with the formatting of this paper. We are grateful to two anonymous reviewers and Jasmine Janes for their comments on an early version of the manuscript. We thank LetPub for its linguistic assistance during the preparation of this manuscript.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This research was funded by the Ningxia 13th Five-Year Plan for Key Research and Development Program (No. 2016BZ09-03 and No. 2018BBF02021-02) and the Natural Science Foundation of Ningxia Province (No. 2018AAC03038). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Ningxia 13th Five-Year Plan for Key Research and Development Program: 2016BZ09-03 and 2018BBF02021-02.

Natural Science Foundation of Ningxia Province: 2018AAC03038.

### Competing Interests

The authors declare that they have no competing interests.

The strain *Lecanicillium lecanii* JMC-01 is in the process of patenting (Patent Number: 201811310451.X), but the patent is currently in the trial stage.

### Author Contributions

- Ting Xie conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Ling Jiang performed the experiments, analyzed the data, prepared figures and/or tables.

- Jianshe Li performed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Bo Hong performed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Xinpu Wang performed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Yanxia Jia conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Shuangyu Zhang granted access to his private greenhouse for the duration of the experiments.

### Patent Disclosures

The following patent dependencies were disclosed by the authors:

The strain *Lecanicillium lecanii* JMC-01 has been deposited in China Center for Type Culture Collection on 2018.06.05 and is being patented (Patent Number: 201811310451.X), but the patent is in the trial stage.

### DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The JMC-01 strain reference ITS nucleotide sequence is available at GenBank: [MH312006](#).

### Data Availability

The following information was supplied regarding data availability:

The raw measurements for [Figs. 2–12](#) are available as a [Supplemental File](#).

The strain *Lecanicillium lecanii* JMC-01 is available in the China Center for Type Culture Collection (No: M2018303).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.7690#supplemental-information>.

## REFERENCES

- Askary H, Benhamou N, Brodeur J. 1999.** Ultrastructural and cytochemical characterization of aphid invasion by the hyphomycete *Verticillium lecanii*. *Journal of Invertebrate Pathology* **74**(1):1–13 DOI [10.1006/jipa.1999.4857](#).
- Bantz A, Camon J, Froger JA, Goven D, Raymond V. 2018.** Exposure to sublethal doses of insecticide and their effects on insects at cellular and physiological levels. *Current Opinion in Insect Science* **30**:73–78 DOI [10.1016/j.cois.2018.09.008](#).
- Chen K, Lu Z. 2017.** Immune responses to bacterial and fungal infections in the silkworm, *Bombyx mori*. *Developmental Comparative Immunology* **83**:3–11 DOI [10.1016/j.dci.2017.12.024](#).

- De Barro PJ. 2011.** *Bemisia tabaci*, the capacity to invade. *Life Sciences* 181–204.
- Ding SY, Li HY, Li XF, Zhang ZY. 2001.** Effects of Bt transgenic poplar on detoxification enzyme and AChE in American white moth larvae. *Journal of Northeast Forestry University* 29:28–29 DOI 10.13759/j.cnki.dlxb.2001.03.007.
- Ding JN, Zhang HH, Chi DF. 2015.** Effects of a pathogenic *Beauveria bassiana* (Hypocreales: Cordycipitaceae) strain on detoxifying and protective enzyme activities in *Xylotrechus rusticus* (Coleoptera: Cerambycidae) larvae. *Florida Entomologist* 98(4):1148–1156 DOI 10.1653/024.098.0419.
- Duan Y, Wu H, Ma Z, Yang L, Ma D. 2017.** Scanning electron microscopy and histopathological observations of *Beauveria bassiana* infection of Colorado potato beetle larvae. *Microbial Pathogenesis* 111:435–439 DOI 10.1016/j.micpath.2017.09.025.
- Esmail SM, Omara RI, Abdelaal KAA, Hafez YM. 2018.** Histological and biochemical aspects of compatible and incompatible wheat-*Puccinia striiformis* interactions. *Physiological and Molecular Plant Pathology* 106:120–128 DOI 10.1016/j.pmpp.2018.11.004.
- Felton GW, Summers CB. 1995.** Antioxidant systems in insects. *Archives of Insect Biochemistry and Physiology* 29(2):187–197 DOI 10.1002/arch.940290208.
- Foyer CH, Noctor G. 2013.** Redox signaling in plants. *Antioxidants & Redox Signaling* 18(16):2087–2090 DOI 10.1089/ars.2013.5278.
- Gao K. 2018.** *Biotype identification and control of Bemisia tabaci in Ningxia*. Ningxia University.
- Ghaffari S, Karimi J, Kamali S, Moghadam EM. 2017.** Biocontrol of *Planococcus citri* (Hemiptera: Pseudococcidae) by *Lecanicillium longisporum* and *Lecanicillium lecanii* under laboratory and greenhouse conditions. *Journal of Asia-Pacific Entomology* 20(2):605–612 DOI 10.1016/j.aspen.2017.03.019.
- Guo LT, Xie W, Wang S, Wu Q, Li R, Yang N, Yang X, Pan H, Zhang Y. 2015.** Detoxification enzymes of *Bemisia tabaci* B and Q: biochemical characteristics and gene expression profiles. *Pest Management Science* 70(10):1588–1594 DOI 10.1002/ps.3751.
- Jiang L. 2018.** *Biological characteristics of Lecanicillium lecanii and synergistic with insecticides Bemisia tabaci*. Ningxia University.
- Lacey LA, Frutos R, Kaya HK, Vail P. 2001.** Insect pathogens as biological control agents: do they have a future? *Biological Control* 21(3):230–248 DOI 10.1006/bcon.2001.0938.
- Li SG, Dou TT, Fu XL, Liu ZQ, Chao H. 2016a.** Changes of protective enzyme activities in *Pieris rapae* infected by *Metarhizium anisopliae*. *Plant Protection* 42:133–136 DOI 10.3969/j.issn.0529-1542.2016.03.022.
- Li XY, Liu QZ, Lewis EE, Tarasco E. 2016b.** Activity changes of antioxidant and detoxifying enzymes in *Tenebrio molitor* (Coleoptera: Tenebrionidae) larvae infected by the entomopathogenic nematode *Heterorhabditis beicherriana* (Rhabditida: Heterorhabditidae). *Parasitology Research* 115(1):1–10 DOI 10.1007/s00436-015-4785-4.
- Liu SA, Fu YG, Huang WR, Fu BL, Zeng DQ. 2013.** Variation in the activities of protective and detoxification enzymes in *Aleurodicus dispersus* infected by *Verticillium lecanii*. *Plant Protection* 39:7–11 DOI 10.3969/j.issn.0529-1542.2013.03.002.
- Liu W, Xie Y, Xue J, Gao Y, Zhang Y, Zhang X, Tan J. 2009.** Histopathological changes of *Ceroplastes japonicus* infected by *Lecanicillium lecanii*. *Journal of Invertebrate Pathology* 101(2):96–105 DOI 10.1016/j.jip.2009.03.002.
- Mathews PM, Guerra CB, Jiang Y, Grbovic OM, Kao BH, Schmidt SD, Dinakar R, Mercken M, Hille-Rehfeld A, Rohrer J, Mehta P, Cataldo AM, Nixon RA. 2002.** Alzheimer's disease-related overexpression of the cation-dependent mannose 6-phosphate receptor increases A $\beta$  secretion. *Journal of Biological Chemistry* 277(7):5299–5307 DOI 10.1074/jbc.M108161200.

- Purwar JP, Sachan GC. 2006.** Synergistic effect of entomogenous fungi on some insecticides against Bihar hairy caterpillar *Spilarctia obliqua* (Lepidoptera: Arctiidae). *Microbiological Research* **161**(1):38–42 DOI [10.1016/j.micres.2005.04.006](https://doi.org/10.1016/j.micres.2005.04.006).
- Ravindran K, Sivaramakrishnan S, Hussain M, Dash CK, Bamisile BS, Qasim M, Liande W. 2018.** Investigation and molecular docking studies of Bassianolide from *Lecanicillium lecanii* against *Plutella xylostella* (Lepidoptera: Plutellidae). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* **206–207**:65–72 DOI [10.1016/j.cbpc.2018.03.004](https://doi.org/10.1016/j.cbpc.2018.03.004).
- Saruhan I. 2018.** Efficacy of some entomopathogenic fungi against *Aphis fabae Scopoli* (Hemiptera: Aphididae). *Egyptian Journal of Biological Pest Control* **28**(1):61 DOI [10.1186/s41938-018-0096-2](https://doi.org/10.1186/s41938-018-0096-2).
- Schama R, Pedrini N, Juarez MP, Nelson DR, Torres AQ, Valle D, Mesquita RD. 2016.** *Rhodnius prolixus* supergene families of enzymes potentially associated with insecticide resistance. *Insect Biochemistry and Molecular Biology* **69**:91–104 DOI [10.1016/j.ibmb.2015.06.005](https://doi.org/10.1016/j.ibmb.2015.06.005).
- Shinya R, Aiuchi D, Kushida A, Tani M, Kuramochi K, Koike M. 2008.** Effects of fungal culture filtrates of *Verticillium lecanii* (*Lecanicillium* spp.) hybrid strains on *Heterodera glycines* eggs and juveniles. *Journal of Invertebrate Pathology* **97**(3):291–297 DOI [10.1016/j.jip.2007.11.005](https://doi.org/10.1016/j.jip.2007.11.005).
- Sugimoto M, Koike M, Hiyama N, Nagao H. 2003.** Genetic, morphological, and virulence characterization of the entomopathogenic fungus *Verticillium lecanii*. *Journal of Invertebrate Pathology* **82**(3):176–187 DOI [10.1016/S0022-2011\(03\)00014-4](https://doi.org/10.1016/S0022-2011(03)00014-4).
- Tang XT, Cai L, Shen Y, Du YZ. 2018.** Diversity and evolution of the endosymbionts of *Bemisia tabaci* in China. *PeerJ* **6**(1):e5516 DOI [10.7717/peerj.5516](https://doi.org/10.7717/peerj.5516).
- Tian YZ. 2014.** *The screening of the strains of the high pathological Beauveria bassiana on soybean pod borers and the biophysical and biochemical effects on their hosts*. Jilin Agricultural University.
- Tian J, Diao HL, Ma RY. 2016a.** Effects of *Isaria fumosorosea* infection on different enzyme activities in the adult in vivo of *Bemisia tabaci*. *Journal of Shanxi Agricultural Sciences* **44**:1007–1010 DOI [10.3969/j.issn.1002-2481.2016.07.27](https://doi.org/10.3969/j.issn.1002-2481.2016.07.27).
- Tian S, Wang XB, Li P, Wang H, Ji HT, Xie JY, Qiu QL, Shen D, Dong HS. 2016b.** Plant aquaporin AtPIP1;4 links apoplastic H<sub>2</sub>O<sub>2</sub> induction to disease immunity pathways. *Plant Physiology* **171**:1635–1650 DOI [10.1104/pp.15.01237](https://doi.org/10.1104/pp.15.01237).
- Wang Z, Zhao Z, Cheng X, Liu S, Wei Q, Scott IM. 2016.** Conifer flavonoid compounds inhibit detoxification enzymes and synergize insecticides. *Pesticide Biochemistry and Physiology* **127**:1–7 DOI [10.1016/j.pestbp.2015.09.003](https://doi.org/10.1016/j.pestbp.2015.09.003).
- Xie M, Zhang YJ, Peng DL, Zhou J, Zhang XL, Zhang ZR, Zhao JJ, Wu YH. 2015.** Persistence and viability of *Lecanicillium lecanii* in Chinese agricultural soil. *PLOS ONE* **10**(9):e0138337 DOI [10.1371/journal.pone.0138337](https://doi.org/10.1371/journal.pone.0138337).
- Xu Q, Chai F, An X, Han S. 2014.** Comparison of detoxification enzymes of *Bemisia tabaci* (Hemiptera: Aleyrodidae) biotypes B and Q after various host shifts. *Florida Entomologist* **97**(2):715–723 DOI [10.1653/024.097.0253](https://doi.org/10.1653/024.097.0253).
- Xu Y, Wang Z, He K, Bai S. 2006.** Effects of transgenic Bt corn expressing Cry1Ab toxin on activities of some enzymes in larvae of the Asian corn borer, *Ostrinia furnacalis* (Guenée) (Lepidoptera: Pyralidae). *Acta Entomologica Sinica* **49**(4):562–567 DOI [10.16380/j.kcxb.2006.04.005](https://doi.org/10.16380/j.kcxb.2006.04.005).
- Yang Q, Wang SH, Zhang WH, Yang TT, Liu YJ. 2015.** Toxicity of commonly used insecticides and their influences on protective enzyme activity of multicolored Asian lady beetle *Harmonia axyridis* (Pallas). *Journal of Plant Protection* **42**:258–263 DOI [10.13802/j.cnki.zwbhxb.2015.02.017](https://doi.org/10.13802/j.cnki.zwbhxb.2015.02.017).

- Ye BH, Zhang YB, Shu JP, Wu H, Wang XJ. 2018.** Effects of three different *Metarhizium* strains on virulence and protective enzymes activities of *Melanotus cribricollis* larvae. *Scientia Silvae Sinicae* **54**:100–108 DOI [10.11707/j.1001-7488.20180612](https://doi.org/10.11707/j.1001-7488.20180612).
- Zare R, Gams W. 2001.** A revision of *Verticillium* sect. Prostrata. VI. The genus *Haptocillium*. *Nova Hedwigia* **73**:1–50.
- Zhang H. 2017.** Research on virulence and control effect of *Beauveria bassiana* against Onion Maggot (Diptera: Anthomyiidae). Dissertation. Chinese Academy of Agricultural Sciences.
- Zhang YF, Wang ZH, Nong XQ, Cao GC, Li Z, Wang GJ, Zhang ZH. 2015.** Effect of *Metarhizium anisopliae* on protective enzyme and detoxification enzyme in the Midgut of *Locusta migratoria manilensis*. *Chinese Journal of Biological Control* **31**:876–881 DOI [10.16409/j.cnki.2095-039x.2015.06.009](https://doi.org/10.16409/j.cnki.2095-039x.2015.06.009).
- Zhang H, Wu SY, Wang SY, Lei ZR. 2017.** Effect of *Beauveria bassiana* on the activity of defense enzymes and cellular defense response of adult of *Delia antiqua* (Meigen). *Chinese Journal of Biological Control* **33**:198–205 DOI [10.16409/j.cnki.2095-039x.2017.02.009](https://doi.org/10.16409/j.cnki.2095-039x.2017.02.009).
- Zhou JY, Liu MK, Xiao LN, Chen DX, Ren CY, Diao CQ, Zhang GC, Wang XQ, Li MY. 2017.** Screening of high virulent isolate of *Beauveria* spp. against *Helicoverpa assulta* and changes of protective enzymes activities in the larvae infected by fungi. *Journal of Anhui Agricultural University* **44**:1119–1123 DOI [10.13610/j.cnki.1672-352x.20171214.034](https://doi.org/10.13610/j.cnki.1672-352x.20171214.034).
- Zhu H, Kim JJ. 2011.** Susceptibility of the tobacco whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae) biotype Q to entomopathogenic fungi. *Biocontrol Science and Technology* **21**(12):1471–1483 DOI [10.1080/09583157.2011.636482](https://doi.org/10.1080/09583157.2011.636482).