

# Huge Ventral Cervicomedullary Neurenteric Cyst: A Rare Entity with Good Surgical Outcome and Appraisal

## Abstract

Neurenteric cysts are rare congenital lesions of benign nature that can be encountered at any level of the neuraxis, starting from the cranium down to coccyx. Rewarding outcome can be achieved with early diagnosis and complete removal of these benign lesions. Here, we report a case of a huge neurenteric cyst in an 11-year-old boy at the ventral craniocervical junction, a rarely reported entity with literature review. In this article, we focus on the clinical presentation, pathogenesis, radiological findings, surgery, and surgical outcome of this benign lesion, as we succeeded to have gratifying result following surgery in our instance.

**Keywords:** Cervicomedullary, magnetic resonance imaging, neurenteric cysts

**Moududul Haque,  
Asifur Rahman,  
Nazmin Ahmed,  
Shamsul Alam**

*Department of Neurosurgery,  
Bangabandhu Sheikh Mujib  
Medical University, Dhaka,  
Bangladesh*

## Introduction

Neurenteric cysts (NC) are rare congenital lesions arising from the misplaced lining epithelium of respiratory or gastrointestinal tract, commonly found in the posterior mediastinum and often associated with vertebral dysplasia.<sup>[1-3]</sup> Cervicothoracic junction is the most common location in the craniospinal axis, while intracranial and craniocervical locations are rarer, where they tend to grow ventrally.<sup>[4-6]</sup> Here, we present a case of an 11-year-old boy having a huge neurenteric cyst ventral to the brainstem, extending from pontomedullary junction down to C2. Appraisal of literature is also deliberated.

## Case Report

An 11-year-old boy presented with gradually progressive neck pain and quadriparesis for the past 1 year. Although conscious and oriented, on examination, he was found to have severe neck ache and quadriparesis of Medical Research Council (MRC) Grade 3, rendering him unable to stand without support. His cranial nerves and autonomic functions were intact.

Magnetic resonance imaging (MRI) of cervical spine including craniocervical junction revealed a well-circumscribed, ventrally placed intradural, extramedullary

cystic lesion extending from the pontomedullary junction down to C2 measuring 4 cm vertically and 2.5 cm anteroposteriorly. There was a significant mass effect on the cervicomedullary junction and upper cervical spinal cord. The lesion was iso- to hypointense on T1-weighted image (T1WI), hyperintense on T2-weighted image (T2WI) [Figure 1a and b] with a peripheral hypointense rim, displacing the basilar artery posteriorly. There was no appreciable enhancement on postgadolinium sequences or appreciable cord edema and/or syringomyelia.

Image wise, the most likely differential diagnosis of NCs include arachnoid cyst, epidermoid cyst, and dermoid cyst.

With a “hockey stick” incision, a suboccipital craniotomy with opening of the foramen magnum, removal of the C1 posterior arch, and mobilization of the vertebral artery were done to accomplish a far lateral approach. After opening the dura in a curvilinear fashion, a well-circumscribed, encapsulated cyst ventral to the ponto-medullo-cervical region was encountered. After aspiration of dirty white fluid, the extra-axial cyst with a good dissecting plane between the capsule and the surrounding structures, especially the medulla and the cervical cord, was excised en mass.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** Haque M, Rahman A, Ahmed N, Alam S. Huge ventral cervicomedullary neurenteric cyst: A rare entity with good surgical outcome and appraisal. *Asian J Neurosurg* 2020;15:1016-9.

**Submitted:** 16-Jul-2020

**Revised:** 19-Sep-2020

**Accepted:** 27-Sep-2020

**Published:** 21-Dec-2020

## Address for correspondence:

*Dr. Moududul Haque,  
Department of Neurosurgery,  
Bangabandhu Sheikh Mujib  
Medical University, Kazi  
Nazrul Islam Avenue, Shabag,  
Dhaka 1000 Bangladesh.  
E-mail: moududneurosurg@  
gmail.com*

## Access this article online

**Website:** www.asianjns.org

**DOI:** 10.4103/ajns.AJNS\_351\_20

## Quick Response Code:




The postoperative period was uneventful. The patient started to show signs of improvement from the 3<sup>rd</sup> postoperative day (POD) by walking with support. His neck pain subsided, and on 7<sup>th</sup> POD, he was able to stand on one foot. At discharge on 10<sup>th</sup> POD, he was neurologically intact with MRC Grade 5 power in all four limbs. His postoperative MRI revealed a complete removal of the cyst [Figure 2a and b].

At 2½ years after surgery, the patient was doing well. He was neurologically intact and was attending his school.

## Discussion

In 1928, Kubie and Fulton first described neurenteric cyst as “teratomatous cysts.”<sup>[7]</sup> Later on, this rare entity was named as neurenteric cyst by Holcomb and Matson in 1954.<sup>[8]</sup> These cysts are actually considered hamartomas in many previously reported literatures, which is formed as a result of the failure of the separation between the primitive gastrointestinal tract to that of neural crest cells. Due to the pattern of embryogenesis, these lesions are rare in the central nervous system, accounting for 0.7%–1.3% of all spinal cord tumors – usually in the cervical or thoracic regions and ventral to the spinal cord.<sup>[9–11]</sup> Among them, about 90% are intradural extramedullary in location.<sup>[12]</sup> Considering the intracranial neurenteric cyst, the most common location is in the ventral brain stem, followed by the cerebellopontine angle.<sup>[13]</sup> While the spinal NCs are frequently associated with underlying bone abnormalities, intracranial NCs possess a rare association, which is consistent with our reported case.<sup>[14]</sup>

Male-female ratio of NC is approximately 2:1, and they generally present in the second and third decades of life. Nonetheless, in the pediatric population, the mean age of presentation is 6.4 years. Among them, NCs in the craniocervical junction are a rare occurrence, while the dorsally located craniocervical NCs are extremely rare.<sup>[15–19]</sup> Although not as rare as the dorsal craniocervical NCs, our male patient of the ventral craniocervical NC presented at an age more than that is average for presentation in the pediatric group.




**Figure 1:** Preoperative sagittal magnetic resonance imaging showing the ventrally placed intradural, extramedullary cystic lesion extending from the pontomedullary junction down to C2 with significant mass effect on the cervicomedullary junction and upper cervical spinal cord. The lesion was iso- to hypointense in T1-weighted image (a) and hyperintense in T2-weighted image (b)

The majority of the patients present with progressive worsening of the neck or back pain, depending on the location of the cysts. The patient may present with features of myelopathy and/or radiculopathy which depends on the extent and level of compression. In case of ventral brain stem location, there may be additional features of multiple cranial nerve palsies. The clinical course of these cases is usually gradually progressive or may have intermittent relapsing and remitting course because of periodic leakage of fluid content secondary to osmotic and hemodynamic factors or changes in the rate of production and reabsorption by the columnar epithelium the cyst wall or in cases of hemorrhagic events.<sup>[20–23]</sup> This patient of ours presented with progressive quadriparesis without any remission or relapse, indicating that the cyst had no leakage or intracystic hemorrhage during the course of its growth, and he improved dramatically following surgery. The relapsing and remitting nature of the disease is frequently misdiagnosed as central nervous system demyelinating diseases.<sup>[24]</sup> Sometimes, traumatic event may initiate or exacerbate such clinical pictures.<sup>[25,26]</sup> In addition, pediatric patients may also present with signs and symptoms of aseptic meningitis, pyogenic meningitis, pyrexia, bowel bladder disturbance, and paraparesis or paraplegia.<sup>[20,27–28]</sup>

As in our case, most NCs are usually isointense on T1-weighted sequence and homogeneously hyperintense on T2-weighted sequence on MRI. However, variations in the protein content of the fluid influence the appearance on T2WI, causing them to appear hyperintense to cerebrospinal fluid (CSF).<sup>[29–31]</sup> The cyst was hyperintense than the CSF intensity in T1WI in our case also and that was found to match with the cyst content peroperatively. Contrast enhancement is mostly absent, and there is also no enhancing solid component usually.<sup>[31]</sup>

On gross appearance, NCs usually appear smooth, regular, contain straw-colored fluid surrounded by a thickened membrane. Peroperatively, we found the cyst to be a well-capsulated one that was well separable from the surrounding structures with a good cleavage plane.




**Figure 2:** Follow-up magnetic resonance images at 2½ years after surgery showing complete resolution and absence of recurrence in T1-weighted image (a) and T2-weighted image (b)

Moreover, the cyst could easily be excised en mass after aspiration of the dirty white fluid inside. Because of differences in the fluid content, the cyst fluid may have diverse appearances such as “milky,” “blackish,” “CSF-like,” or “clear jelly-like”. These varied appearances may result from different degrees of protein contents or from intracystic hemorrhage.<sup>[32-34]</sup>

On histopathological examination, the neurenteric cyst demonstrates columnar or cuboidal epithelium with or without cilia and intracellular mucous globules, simulating the lining epithelium of respiratory or gastrointestinal tract.<sup>[29,35]</sup> In 1976, based on histopathology, Wilson and Odom classified NCs into three types: Type A: composed of a single layer of pseudostratified columnar or cuboidal epithelium; Type B: composed of complex invaginations with glandular organization; and Type C: consisting of ependymal or glial tissue.<sup>[36]</sup> Histopathology of our particular case fell under the category of Type A having cyst wall lined by columnar and pseudostratified ciliated epithelium with the presence of occasional goblet cells [Figure 3].

Complete surgical resection remains the gold standard of treatment for NC, which almost always gives a favorable outcome. In addition to the location of the cyst, the prognosis depends on the size of the cyst as well. As partial resection poses the risk of cyst recurrence, total surgical excision should be the norm, whenever feasible.<sup>[27]</sup> Depending on location, three basic surgical approaches: posterior, anterior, and lateral can be chosen for cyst excision. In spite of the ventral location, favorable results can be achieved after partial or complete resection of cyst wall through a posterior approach, as reported in different literature.<sup>[17,32,37]</sup> Now, 2½ years after surgery, our patient is leading a normal healthy life without any recurrence seen in his follow-up MR images [Figure 2a and b]. Although the location was a bit difficult, because of relatively smaller size, good cleavage plane, and following a total excision of the cyst



**Figure 3: Photomicrograph showing cyst wall lined by columnar and pseudostratified ciliated epithelium with the presence of occasional goblet cells**

through a far lateral approach, the boy is expected to have a recurrence-free vigorous life. Cysts that are located ventral to the brain stem surgery may be challenging and may result in incomplete resection, which often results in recurrence. Partial resection remains the most important risk factors for the recurrence in NCs, and recurrence rate ranges from 0% to 37% in the literature.<sup>[38,39]</sup> Because of complete resection, we believe that the boy will not have any recurrence in the future as evident in his postoperative follow-up MR image.

#### Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

#### Financial support and sponsorship

Nil.

#### Conflicts of interest

There are no conflicts of interest.

#### References

1. Macdonald RL, Schwartz ML, Lewis AJ. Neurenteric cyst located dorsal to the cervical spine: Case report. *Neurosurgery* 1991;28:583-7.
2. Harris CP, Dias MS, Brockmeyer DL, Townsend JJ, Willis BK, Apfelbaum RI. Neurenteric cysts of the posterior fossa: Recognition, management, and embryogenesis. *Neurosurgery* 1991;29:893-7.
3. Gauden AJ, Khurana VG, Tsui AE, Kaye AH. Intracranial neurenteric cysts: A concise review including an illustrative patient. *J Clin Neurosci* 2012;19:352-9.
4. de Oliveira RS, Cinalli G, Roujeau T, Sainte-Rose C, Pierre-Kahn A, Zerah M. Neurenteric cysts in children: 16 consecutive cases and review of the literature. *J Neurosurg* 2005;103:512-23.
5. Filho FL, Tatagiba M, Carvalho GA, Weichhold W, Klekamp J, Samii M. Neurenteric cyst of the craniocervical junction. Report of three cases. *J Neurosurg* 2001;94:129-32.
6. Bejjani GK, Wright DC, Schessel D, Sekhar LN. Endodermal cysts of the posterior fossa. Report of three cases and review of the literature. *J Neurosurg* 1998;89:326-35.
7. Kubie LS, Fulton JFA. Clinical and pathological study of two teratomatous cysts of the spinal cord, containing mucus and ciliated cells. *Surg Gynecol Obstet* 1928;47:297-311.
8. Holcomb GW Jr., Matson DD. Thoracic neurenteric cyst. *Surgery* 1954;35:115-21.
9. Fernandes ET, Custer MD, Burton EM, Boulden TF, Wrenn EL Jr., Whittle AP, *et al.* Neurenteric cyst: Surgery and diagnostic imaging. *J Pediatr Surg* 1991;26:108-10.
10. Lippman CR, Argenteanu M, Purohit D, Naidich TP, Camins MB. Intramedullary neurenteric cysts of the spine. Case report and review of the literature. *J Neurosurg* 2001;94:305-9.
11. Palma L, Di Lorenzo N. Spinal endodermal cysts without

- associated vertebral or other congenital abnormalities. Report of four cases and review of the literature. *Acta Neurochir (Wien)* 1976;33:283-300.
12. Fortuna A, Mercuri S. Intradural spinal cysts. *Acta Neurochir (Wien)* 1983;68:289-314.
  13. Shenoy SN, Raja A. Spinal neurenteric cyst. Report of 4 cases and review of the literature. *Pediatr Neurosurg* 2004;40:284-92.
  14. Kumar R, Prakash M. Unusual split cord with neurenteric cyst and cerebellar heterotopia over spinal cord. *Childs Nerv Syst* 2007;23:243-7.
  15. Arai Y, Yamauchi Y, Tsuji T, Fukasaku S, Yokota R, Kudo T. Spinal neurenteric cyst. Report of two cases and review of forty-one cases reported in Japan. *Spine (Phila Pa 1976)* 1992;17:1421-4.
  16. Rao MB, Rout D, Misra BK, Radhakrishnan VV. Craniospinal and spinal enterogenous cysts—report of three cases. *Clin Neurol Neurosurg* 1996;98:32-6.
  17. Agnoli AL, Laun A, Schönmayr R. Enterogenous intraspinal cysts. *J Neurosurg* 1984;61:834-40.
  18. Tan GS, Hortobágyi T, Al-Sarraj S, Connor SE. Intracranial laterally based supratentorial neurenteric cyst. *Br J Radiol* 2004;77:963-5.
  19. Gu J, Yang T, Xing X, Kuang Y, Cheng G, Zhang J, *et al.* A dorsally located giant posterior fossa neurenteric cyst in a Chinese woman. *J Clin Neurosci* 2015;22:917-8.
  20. Rivierez M, Buisson G, Kujas M, Ridarch A, Mignon E, Jouannelle A, *et al.* Intramedullary neurenteric cyst without any associated malformation. *Acta Neurochir* 1997;139:887-90.
  21. Bae KJ, Kim IM. Anterior surgical approach in recurrent cervical neurenteric cyst. *J Kor Neurosurg Soc* 2000;29:1258-61.
  22. Garg N, Sampath S, Yasha TC, Chandramouli BA, Devi BI, Kovoor JM. Is total excision of spinal neurenteric cysts possible? *Br J Neurosurg* 2008;22:241-51.
  23. Osenbach RK, Godersky JC, Traynelis VC, Schelper RD. Intradural extramedullary cysts of the spinal canal: Clinical presentation, radiographic diagnosis, and surgical management. *Neurosurgery* 1992;30:35-42.
  24. Vinters HV, Gilbert JJ. Neurenteric cysts of the spinal cord mimicking multiple sclerosis. *Can J Neurol Sci* 1981;8:159-61.
  25. D’Almeida AC, Stewart DH. Neurenteric cyst. *Neurosurgery* 1981;8:596-9.
  26. Jae CB, Kim TS, Park JT, Lim YJ, Leem W, Kim GK. Intradural extramedullary enterogenous cyst in cervical spine. *J Kor Neurosurg Soc* 1998;27:1741-5.
  27. Menezes AH, Traynelis VC. Spinal neurenteric cysts in the magnetic resonance imaging era. *Neurosurgery* 2006;58:97-105.
  28. Silvernail WI Jr, Brown RB. Intramedullary enterogenous cyst. Case report. *J Neurosurg* 1972;36:235-8.
  29. Preece MT, Osborn AG, Chin SS, Smirniotopoulos JG. Intracranial neurenteric cysts: Imaging and pathology spectrum. *AJNR Am J Neuroradiol* 2006;27:1211-6.
  30. Weiss MA, Gebarski SS, McKeever PE. Foramen magnum neurenteric cyst causing mollaret meningitis: MR findings. *AJNR Am J Neuroradiol* 1996;17:386-8.
  31. Brooks BS, Duvall ER, el Gammal T, Garcia JH, Gupta KL, Kapila A. Neuroimaging features of neurenteric cysts: Analysis of nine cases and review of the literature. *AJNR Am J Neuroradiol* 1993;14:735-46.
  32. Kim CY, Wang KC, Choe G, Kim HJ, Jung HW, Kim IO, *et al.* Neurenteric cyst: Its various presentations. *Childs Nerv Syst* 1999;15:333-41.
  33. Knight G, Griffiths T, Williams I. Gastrocystoma of the spinal cord. *Br J Surg* 1955;42:635-8.
  34. Mizuno J, Fiandaca MS, Nishio S, O’Brien MS. Recurrent intramedullary enterogenous cyst of the cervical spinal cord. *Childs Nerv Syst* 1988;4:47-9.
  35. Savage JJ, Casey JN, McNeill IT, Sherman JH. Neurenteric cysts of the spine. *J Craniovertebr Junction Spine* 2010;1:58-63.
  36. Wilkins R, Odom GL. Spinal intradural cysts. *Tumors Spinal Cord* 1976;20:55-102.
  37. Singhal BS, Parekh HN, Ursekar M, Deopujari CE, Manghani DK. Intramedullary neurenteric cyst in midthoracic spine in an adult. *Neurol India* 2001;49:302-4.
  38. Cai C, Shen C, Yang W, Zhang Q, Hu X. Intraspinal neurenteric cysts in children. *Can J Neurol Sci* 2008;35:609-15.
  39. Holmes GL, Trader S, Ignatiadis P. Intraspinal enterogenous cysts. A case report and review of pediatric cases in the literature. *Am J Dis Child* 1978;132:906-8.