

POSTER PRESENTATION

Open Access

Fucoidan extract enhances the anti-cancer activity of chemotherapeutic agents in breast cancer cells

Sanetaka Shirahata^{1,2*}, Zhonguan Zhang¹, Toshihiro Yoshida¹, Hiroshi Eto³, Kiichiro Teruya¹

From 23rd European Society for Animal Cell Technology (ESACT) Meeting: Better Cells for Better Health Lille, France. 23-26 June 2013

Background

Fucoidan, a fucose-rich polysaccharide isolated from brown alga, is currently under investigation as a new anti-cancer compound [1-4]. In the present study, fucoidan extract (FE) from *Cladosiphon navae-caledoniae* Kylin was prepared by enzymatic digestion. We investigated whether a combination of FE with chemotherapeutic agents had the potential to improve the therapeutic efficacy of cancer treatment.

Materials and methods

Estrogen receptor (ER)-positive MCF-7 and ER-negative MDA-MB-231 breast cancer cells were cultured in DME medium supplemented with 10% fetal bovine serum in a humidified atmosphere of 5% CO₂ at 37 °C. The abalone glycosidase-digested fucoidan extract (FE) was obtained from Daiichi Sangyo Corporation (Osaka, Japan). The cells were treated with FE and chemotherapeutic agents like cisplatin, tamoxifen or paclitaxel. The cell growth was determined by MTT assay. Apoptosis was evaluated using annexin V binding assay and flow cytometry analysis. Signaling proteins were analyzed by western blot. Intracellular reactive oxygen species (ROS) were determined using DCFH-DA and determined using IN Cell Analyzer 1000. The reduced glutathione (GSH) concentration was measured by the GSH assay kit.

Results

The co-treatments significantly induced cell growth inhibition, apoptosis, as well as cell cycle modifications in MDA-MB-231 and MCF-7 cells. FE enhanced apoptosis

in cancer cells that responded to treatment with cisplatin, tamoxifen, or paclitaxel after 48 h of treatment (Figure 1). FE enhanced the downregulation of the anti-apoptotic proteins Bcl-xL and Mcl-1 by these chemotherapeutic drugs. The combination treatments led to an obvious decrease in the phosphorylation of ERK and Akt in MDA-MB-231 cells, but increased the phosphorylation of ERK in MCF-7 cells. In addition, we observed that combination treatments enhanced intracellular ROS levels and reduced glutathione (GSH) levels in breast cancer cells, suggesting that induction of oxidative stress was an important event in the cell death induced by the combination treatments.

FE protected normal human fibroblast TIG-1 cells from apoptosis by cisplatin and tamoxifen, suggesting its favorable characteristic for application to cancer therapy.


Conclusions

- Combination of FE and three chemotherapeutic agents exhibit highly synergistic inhibitory effects on the growth of breast cancer cells.
- Combination treatments induced modifications in cell cycle distribution.
- Combination treatments modified the Bcl-2 expression, and ERK and Akt phosphorylation induced by FE, demonstrating different effects on apoptotic pathways in MDA-MB-231 cells and MCF-7 cells.
- Generation of intracellular ROS and depletion of GSH are related to the cell death in combination treated -breast cancer cells.

* Correspondence: sirahata@grt.kyushu-u.ac.jp

¹Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University, Fukuoka 812-8581, Japan

Full list of author information is available at the end of the article



Authors' details

¹Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University, Fukuoka 812-8581, Japan. ²Yosida Clinic, Osaka 532-0002, Japan.
³Daiichi Sangyo Co. Ltd., Osaka 530-0037, Japan.

Published: 4 December 2013

References

1. Ye J, Li Y, Teruya K, Katakura Y, Ichikawa A, Eto H, Hosoi M, Hosoi M, Nishimoto S, Shirahata S: Enzyme-digested fucoidan extracts derived from seaweed *Mozuku* of *Cladosiphon novae-caledoniae kylin* inhibit invasion and angiogenesis of tumor cells. *Cytotechnology* 2005, 47:117-126.
2. Zhang Z, Teruya K, Eto H, Shirahata S: Fucoidan extract induces apoptosis in MCF-7 Cells via a mechanism involving the ROS-dependent JNK activation and mitochondria-mediated pathways. *PLoS ONE* 2012, 6: e27441.
3. Zhang Z, Teruya K, Eto H, Shirahata S: Induction of apoptosis by low-molecular weight fucoidan through calcium- and caspase-dependent mitochondrial pathways in MDA-MB-231 breast cancer cells. *Biosci Biotechnol Biochem* 2012, 77:235-242.
4. Zhang Z, Teruya K, Yoshida T, Eto H, Shirahata S: Fucoidan extract enhances the anti-cancer activity of chemotherapeutic agents in MDA-MB-231 and MCF-7 breast cancer cells. *Marine Drugs* 2013, 11:81-98.

doi:10.1186/1753-6561-7-S6-P70

Cite this article as: Shirahata et al.: Fucoidan extract enhances the anti-cancer activity of chemotherapeutic agents in breast cancer cells. *BMC Proceedings* 2013 7(Suppl 6):P70.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

