

Review Article

The effects of marine eukaryote evolution on phosphorus, carbon and oxygen cycling across the Proterozoic–Phanerozoic transition

Timothy M. Lenton and Stuart J. Daines

Earth System Science Group, College of Life and Environmental Sciences, University of Exeter, Exeter, U.K.

Correspondence: Timothy M. Lenton (t.m.lenton@exeter.ac.uk)



A ‘Neoproterozoic oxygenation event’ is widely invoked as a causal factor in animal evolution, and often attributed to abiotic causes such as post-glacial pulses of phosphorus weathering. However, recent evidence suggests a series of transient ocean oxygenation events ~660–520 Ma, which do not fit the simple model of a monotonic rise in atmospheric oxygen (pO_2). Hence, we consider mechanisms by which the evolution of marine eukaryotes, coupled with biogeochemical and ecological feedbacks, potentially between alternate stable states, could have caused changes in ocean carbon cycling and redox state, phosphorus cycling and atmospheric pO_2 . We argue that the late Tonian ocean ~750 Ma was dominated by rapid microbial cycling of dissolved organic matter (DOM) with elevated nutrient (P) levels due to inefficient removal of organic matter to sediments. We suggest the abrupt onset of the eukaryotic algal biomarker record ~660–640 Ma was linked to an escalation of protozoan predation, which created a ‘biological pump’ of sinking particulate organic matter (POM). The resultant transfer of organic carbon (C_{org}) and phosphorus to sediments was strengthened by subsequent eukaryotic innovations, including the advent of sessile benthic animals and mobile burrowing animals. Thus, each phase of eukaryote evolution tended to lower P levels and oxygenate the ocean on $\sim 10^4$ year timescales, but by decreasing C_{org}/P burial ratios, tended to lower atmospheric pO_2 and deoxygenate the ocean again on $\sim 10^6$ year timescales. This can help explain the transient nature and $\sim 10^6$ year duration of oceanic oxygenation events through the Cryogenian–Ediacaran–Cambrian.

Introduction

Marine eukaryotes are important ecosystem engineers and their evolution over ~850–500 Ma (Figure 1a,b) surely had an impact on carbon, phosphorus and oxygen cycling [1]. While a monotonic rise in atmospheric oxygen (pO_2) in a ‘Neoproterozoic oxygenation event’ [2–5] is still regularly invoked to explain evidence of deeper ocean oxygenation in the Neoproterozoic Era, ocean oxygenation could equally have been caused by declining oxygen demand in deeper waters [1,6]. Here, ‘deeper’ could mean anything below the well-mixed surface layer of stratified shelf seas or the open ocean, but sampling of deep time is largely restricted to continental shelf-slope ocean margins (so does not extend to the deep open ocean) [6]. In deeper waters, oxygenation state is governed by the balance of oxygen supply and respiratory demand from organic matter input. Declining oxygen demand could occur due to: (i) a redistribution of organic matter (and its remineralisation) away from these depths, and/or (ii) a global decline in ocean nutrient levels and organic matter primary production. We focus on phosphorus as the ultimate limiting nutrient, because the ocean available nitrogen inventory — whether as NO_3^- , NH_4^+ or a mix of both [6] — tends to track changes in the phosphorus inventory [7], even though models predict nitrogen would have been much further below the ‘Redfield ratio’ to phosphorus (required by organisms) in a more anoxic ocean [6,8,9].

Received: 13 February 2018

Revised: 27 April 2018

Accepted: 1 May 2018

Version of Record published:

29 June 2018




Figure 1. Timeline of biological and environmental changes through the Tonian, Cryogenian, Ediacaran and Cambrian periods (850–500 Ma).

(a) Biomarker evidence [34]; sterane/hopane ratio indicates balance of eukaryotes/bacteria, followed by occurrence of eukaryotic steranes (with probable sources [34,35]) including cholestane (heterotrophic eukaryotes), stigmastane (chlorophytes), ergostane (general), cryostane (unknown), 24-ipc = 24-isopropylcholestane (demosponges), 24-npc = 24-*n*-propylcholestane (rhizarians). (b) Fossil evidence, updated from [1] based on studies cited in main text and [33]. (c) Ocean redox state from redox-sensitive elements (pink; global signature) [11,13] and iron speciation data at different depths (green, yellow, orange; local signature) [13,109,110]. (d) The carbon isotopic composition of marine carbonates.

Available data show that ocean redox conditions were temporally (as well as spatially) variable in the late Neoproterozoic-early Paleozoic (Figure 1c), while the carbon cycle underwent major fluctuations (Figure 1d). A series of transient ocean oxygenation events are inferred, beginning in the middle of the Cryogenian period (720–635 Ma) [10,11], and getting more frequent through the Ediacaran (635–540 Ma) [12] and early-mid Cambrian (540–500 Ma) [13]. If these transient events were due purely to changes in atmospheric pO_2 , then it must have peaked and declined repeatedly. They could equally have been caused by rapid reorganizations of nutrient and carbon cycling within the ocean, potentially between alternative stable states (anoxic high P recycling, oxic efficient P removal) [14]. Indeed, the onset of oxygenation appears sufficiently rapid in some cases [12] that $\sim 10^4$ year declines in ocean nutrient inventories look a more plausible explanation than $\sim 10^6$ year

increases in atmospheric pO_2 . These are not mutually exclusive scenarios as rapid biogeochemical reorganisations would ultimately be countered by a slower adjustment of pO_2 [14].

Here, we explore how eukaryote evolution could have changed the nutrient and redox state of their (submarine) world. We start by introducing key controls on ocean redox state, marine phosphorus cycling and oxygen cycling. Then, we consider possible causes of the rise to ecological prominence of eukaryotic algae, their biogeochemical effects, and those of early sessile animals, and later mobile burrowing animals.

Processes and timescales

We distinguish three timescales over which eukaryotic evolutionary or ecologically driven changes are coupled to biogeochemical cycles (Figure 2):

- (1) A $\sim 10^3$ year ocean circulation timescale [15] of effects on the nature and distribution of organic matter and corresponding oxygen demand in the water column (and between it and the sediments; Figure 2a).
- (2) A $\sim 10^4$ year timescale [16] of changes in P burial efficiency altering the ocean P inventory, which in turn controls the amount of organic matter produced and corresponding oxygen demand (Figure 2b). This modern P timescale could have differed in the Neoproterozoic.
- (3) A $\sim 10^6$ year timescale of changes in the oxygen source from organic carbon (C_{org}) burial altering atmospheric pO_2 (Figure 2c), which in turn controls oxygen supply to the water column. This timescale is estimated for the Neoproterozoic based on $pO_2 \sim 0.1$ PAL (present atmospheric level) supported by $\sim 2.5 \times 10^{12}$ mol year⁻¹ net O_2 source [17].

When looking at the low temporal resolution of the geological record, we expect that the effects of eukaryote evolution on organic matter cycling (1) cannot be resolved from their effects on phosphorus cycling (2), whereas the slower timescale of changes in atmospheric pO_2 (3) is resolvable from the others.

Today, a relatively efficient ‘biological pump’ of sinking particulate organic matter (POM) creates oxygen demand across a range of water depths, and transfers C_{org} and P to sediments.

On geologic timescales $>10^4$ years (i.e. longer than the residence time of P in the ocean), global phosphorus output to sediments must equal global input from rivers (ultimately derived from weathering). This means that changes in P input or the efficiency of P removal (the focus here) cause the ocean P inventory and corresponding output flux to adjust until output again matches input. Today, despite an efficient biological pump, $\sim 70\%$ of global phosphorus removal occurs in shelf sea settings and only $\sim 30\%$ on the deep ocean floor [16,18].




Figure 2. Timescales and processes of eukaryotic effects on biogeochemical cycling.

(a) Ocean circulation ($\sim 10^3$ year) timescale (black arrows): circulation redistributes O_2 supply from atmosphere and O_2 demand from DOM (green dotted background) and sinking POM (green downward wobble arrow) within the water column. (b) Phosphorus cycling ($\sim 10^4$ year) timescale (dark blue arrows), inset of shelf sea: the ocean P inventory adjusts to maintain balance between P input (via weathering) and P burial (primarily on shelves). P sequestration in sediments is enhanced by biological pump (downward wobble arrow) and by sessile benthic animals (downward arrow). P is preferentially recycled from sediments especially under euxinic conditions (upward arrow). (c) Oxygen cycling ($\sim 10^6$ year) timescale (red arrows): Organic carbon burial (C_{org} ; green downward arrow) provides O_2 source, governed by P input and C_{org} :P burial ratio which is redox-sensitive. Oxidative weathering of ancient C_{org} in sedimentary rocks (green upward arrow) provides O_2 sink.

Hence, here we ignore P burial in deep ocean sediments and focus on shelf seas, because P burial there largely controls the global P reservoir.

The burial of photosynthetically derived C_{org} also occurs mostly in shelf seas [19] and represents the major net source of oxygen to the atmosphere: $\text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{CH}_2\text{O}(\downarrow\text{buried}) + \text{O}_2(\uparrow)$. The C_{org} burial flux ($F_{C_{\text{org}}}$) is controlled by the phosphorus input/burial flux (F_{P} ; which limits production) and the burial ratio of C_{org} to total phosphorus in marine sediments [$(C_{\text{org}}/P)_{\text{burial}}$; which is redox-sensitive]: $F_{\text{P}} \times (C_{\text{org}}/P)_{\text{burial}} = F_{C_{\text{org}}}$. To first order, we do not expect evolutionary or ecologically driven changes in organic matter cycling within the ocean to have impacted atmospheric $p\text{O}_2$, because the ocean P inventory adjusts to changes in P burial efficiency such that the P burial flux again matches the (unchanged) P input flux — and the corresponding oxygen source from C_{org} burial is left unchanged.

In reality, the ratio of C_{org} to total phosphorus buried in new sedimentary rocks, $(C_{\text{org}}/P)_{\text{burial}}$, can change (thus affecting atmospheric $p\text{O}_2$), because it is sensitive to the redox state of ocean bottom waters and hence to ocean nutrient levels and (ultimately) atmospheric $p\text{O}_2$. Under anoxic and euxinic (sulfate-reducing) bottom-water conditions, phosphorus recycling to the water column is enhanced, increasing $(C_{\text{org}}/P)_{\text{burial}}$ [20,21], and oxygenation reduces P recycling from sediments [22] decreasing $(C_{\text{org}}/P)_{\text{burial}}$. This acts as a positive feedback amplifying changes in phosphorus levels, productivity and redox state on $\sim 10^4$ year timescales [23]. However, on $\sim 10^6$ year timescales, changes in atmospheric $p\text{O}_2$ provide negative feedback [14,24]. Less clear is what happens to phosphorus cycling under widespread anoxic but ‘ferruginous’ (iron-reducing) conditions [25]. While the formation of Fe(III) oxides and associated trapping of P in sediments will be suppressed, there is evidence that vivianite and mixed Fe(II)–Fe(III) minerals (‘green rust’) can provide a potentially large P sink [26,27]. This could change the sign of the feedback, such that ferruginous conditions enhance P removal suppressing their own spread, but amplify declining $p\text{O}_2$ [28,29].

Dissolved organic matter world

Picture the ocean in the mid to late Tonian period (~ 850 – 720 Ma). Despite the evolution of algae ~ 1.7 – 1.4 Ga [30], including multi-cellular red algae by ~ 1.1 Ga [31], and the radiation of red and green algae into the marine environment [32,33], they left no biomarker record [34] (Figure 1a). The low sterane/hopane ratio of biomarkers [34] indicates that bacteria dominated preserved lipids, with cholestane (probably from heterotrophic eukaryotes) and cryostane (of unknown origin [35]) the only eukaryotic steranes present. This requires that unicellular algae were either an ecologically insignificant contributor to marine productivity [34], or that eukaryotic biomass was efficiently recycled either in the water column [36] or in microbial mats [37]. Efficient water column recycling of primary production by prokaryotes (Figure 3a) and possibly small protists (Figure 3c) would have closed a ‘microbial loop’ ensuring dissolved organic matter (DOM; operationally defined as <0.22 or $<0.7 \mu\text{m}$) [38,39] dominated the water column. This DOM pool was not a massive, ancient one [40], rather it turned over on $<10^4$ year timescales consistent with modern observations [38,39].

A microbially dominated DOM recycling ecosystem would have had interesting implications for water column redox state [6]. Unlike the biological pump in the modern ocean (due to sinking POM), oxygen demand in the water column would be strongly influenced by advective and diffusive transport of DOM. This would still be able to drive anoxia in the open-ocean thermocline (see Figure 5f of [6]) and in shelf seas (Figure 4a), although less efficiently for a given nutrient level than particulate export. Shelf-sea environments would be expected to show strong latitudinal patterns in redox state, with permanent anoxia likely in permanently stratified regions at low-mid latitudes, and seasonal anoxia in seasonally stratifying high-latitude shelves.

With all oxygen demand used up at intermediate depths, deeper waters of the open ocean would be less prone to anoxia (Figure 5f of [6]), having their $[\text{O}_2]$ governed by the balance of supply from the atmosphere via high-latitude deep convection and (a smaller) demand from inorganic reductant input at mid-ocean ridges [6]. Any additional POM flux into deep waters could readily have driven them anoxic (Figure 5d of [6] and [41]). However, what little data we have from the truly deep Proterozoic ocean suggest that some oxygen was present [42,43], consistent with a DOM-dominated system without a significant biological pump [6].

Maintaining an oxygenated Proterozoic atmosphere (given tectonic reductant input) required greater than $\sim 25\%$ of present-day C_{org} burial [17]. A key puzzle is; how did organic matter — including P and C — get out of the bottom of this DOM-cycling system in solid, sedimentary form? Much of it must have been derived from seafloor microbial mats. It seems unlikely that heterotrophic bacterial mats could extract significant DOM from the water column, relative to much larger water column bacterial populations. Instead, occasional aggregation of DOM into POM [44] and of bacterial cells into larger POM, which sunk as ‘marine snow’, seems a




Figure 3. Community size structure in an idealized steady-state microbial food chain model [52,111], for assumed evolutionary steps. Green = autotrophs, brown = protist heterotrophs, blue = dissolved nitrogen.

Insets show assumed community composition (dots) in size classes (increasing in size upwards), with lines indicating trophic relationships. Graphs show the fraction of total nitrogen ('frac N') in dissolved form (blue) and in each component of the size structured population, as total nitrogen ('tot N') increases. Heterotrophs are stacked on top of autotrophs and for each, where different size classes coexist, the smallest size class is at the bottom with progressively larger size classes stacked on top: (a) cyanobacteria only — nutrients are drawn down to limiting concentration and cyanobacterial population increases (until light limitation, not shown); (b) including phagotrophy from heterotrophic nanoflagellates limits cyanobacterial population size, allowing nutrient levels to rise; (c) this allows autotrophic nanoflagellates to coexist with cyanobacteria; (d) eukaryophagy limits population size of autotrophic nanoflagellates and allows larger size classes of eukaryotic phytoplankton (at high total nitrogen levels).

more plausible food source for heterotrophic mats. Additionally, autotrophic shallow water mats would have produced POM.

In this 'DOM world', balancing the P cycle would have demanded a larger ocean P inventory (Figure 4a) to drive the same P output, consistent with geochemical inferences of high P levels by the Cryogenian [29,45]. As ocean nutrient inventory determines the total amount of organic matter production in the ocean, a high P ocean would have been prone to anoxia at the depths where DOM was respired, particularly under lower early Neoproterozoic atmospheric pO_2 [6].

Eukaryotic algae and eukaryophagy

Fossils indicate a diversification of heterotrophic eukaryotes in the late Tonian [46–48], including testate amoebae and organic or siliceous scales, followed by the advent of foraminifera and ciliates in the Cryogenian [49] (Figure 1b). Recent (re)analysis of the biomarker record [34] indicates that steranes from eukaryotic algae, demosponges and rhizarians (a group which includes foraminifera and radiolarians) all first appear in the Cryogenian period (720–635 Ma), between the Sturtian and Marinoan 'snowball Earth' glaciations ~660–640 Ma [50], and the sterane/hopane ratio increases, indicating the prevalence of eukaryotic over bacterial lipids [34] (Figure 1a).

The absence of a eukaryotic algal biomarker record prior to the Cryogenian has been attributed to nutrient scarcity [34,51]. However, in the modern ocean phagotrophy limits cyanobacterial population size allowing eukaryotic algae to coexist [52], where even in the most oligotrophic regions mixotrophic eukaryotes that




Figure 4. Scenarios for different shelf sea biogeochemical regimes through time, showing redox state after adjustment of P cycle ($\sim 10^4$ year) but before any adjustment of O_2 cycle.

Physical setting is a stratified shelf sea ~ 100 m deep with surface waters separated from deeper waters by a sharp thermocline. DOM, dissolved organic matter (pale green). POM, particulate organic matter (dark green). Arrow width roughly represents magnitude of organic matter flux. We assume constant P input flux throughout hence P burial flux into sediments is identical throughout, but this is achieved at different [P] and through a different balance of processes over time. (a) Tonian ‘DOM world’ either dominated by cyanobacterial productivity (small pale green dots) or including small green algae (dark green circles) and small phagotrophic eukaryotes (brown ‘Pac-Men’), overlying benthic mats (pale green layer). Dashed arrows indicate uncertainty surrounding POM pathway(s) to sediments from sinking ‘marine snow’ and/or photoautotrophic mats. (b) Cryogenian world of eukaryotic algae (larger dark green dots) and eukaryophagy (larger brown ‘Pac-Men’) with a biological pump transferring POM to sediments. (c) Late Ediacaran world of sessile animals, including rangeomorph fronds (brown diamonds on stalks) and filter-feeding sponges (brown clouds) transferring POM to sediments. Their location on top of benthic mats creates a sharp redox boundary supporting phosphorite and authigenic carbonate deposition. (d) Cambrian world of mobile animals (brown splodges) bioturbating (and thus oxygenating) upper sediments, which lowers the C_{org}/P burial ratio enabling a smaller sedimentary POM flux to maintain the required P output flux.

consume bacteria compete effectively with prokaryotes [53]. Also, during Phanerozoic ocean anoxic events that deplete nitrogen, eukaryotes comprise a significant fraction of productivity [54]. Furthermore, any boost in phosphorus levels during [45] or after the Sturtian glaciation would be transient [1] hence cannot explain the irreversible biomarker transition.

Instead, we suggest two scenarios linking the biomarker record of eukaryotic algae [34] with an apparent escalation in protistan predation [55,56] (Figure 3). In the first scenario, eukaryotic bacterivory was, for some reason, ineffective prior to the Cryogenian (despite being plesiomorphic among eukaryotes), allowing prokaryotic autotrophs to competitively exclude larger eukaryotic algae, because smaller cells are more efficient at diffusion-limited nutrient uptake [52] (Figure 3a). Then, an increase in the effectiveness of eukaryotic bacterivory limited cyanobacterial population size (Figure 3b, as in the modern ocean), allowing surface nutrient levels to rise and creating a niche for autotrophic eukaryotes (Figure 3c), followed by a rapid escalation of eukaryophagy to exploit this new resource [57] (Figure 3d). Alternatively, the advent of eukaryophagy put a selection pressure for predation resistance on an earlier cryptic population of small heterotrophic, autotrophic and mixotrophic eukaryotes (Figure 3c), driving increases in size (Figure 3d), armour, and therefore sinking speed and preservation potential — leading to the formation of the algal biomarker record. By filtering out smaller cyanobacterial cells, the advent of sponges [58–60] could also have provided a strong selection pressure for larger eukaryotic algal cells in shelf seas [1].

Thanks to their larger size and faster sinking rate, eukaryotes created a biological pump of POM [1]. The short-term effect would have been to increase oxygen demand in the deeper waters of shelf seas and the open

ocean [6,61]. However, by creating an efficient particulate P removal flux to shelf sediments, it would have lowered ocean P content and corresponding O₂ demand over ~10⁴ years, thus tending to oxygenate deeper waters of shelf seas (Figure 4b) and the open ocean [6]. This prediction is supported by redox proxy evidence of ocean oxygenation after the Sturtian [10,11,62]. The contribution of redox-sensitive P cycling feedback to this oxygenation is uncertain, but the partial oxygenation of ‘ferruginous’ background ocean waters (Figure 1c) apparently did not cause a major drop in the efficiency of P removal with Fe minerals [27–29], as that would have rapidly counteracted further oxygenation. The oxygenation event is inferred to have reversed before the Marinoan [11], perhaps because partial shelf seafloor oxygenation produced a lower C_{org} to total phosphorus burial ratio [21], thus triggering a decline in atmospheric pO₂.

Sessile animals and benthic mats

The nature and timing (Figure 1a,b) of the first animals remains contested. Recent molecular phylogenetic studies support the common sense view that sponges (Porifera) are the sister group to all other animals [63,64]. Relaxed molecular clocks put the origin of crown-group demosponges 872–657 Ma (across studies) [65] consistent with biomarkers ~660–640 Ma [58–60], but put silicification and spicule production later at 648–616 Ma [66]. Hence, the gap to the first widely accepted fossil sponges in the early Cambrian ~535 Ma [67–69] could be partly due to poor preservation potential. One ~600 Ma fossil [70] could also close this gap, if supported. A counter-view takes the fossil record at face value (rejecting the molecular clock and biomarker evidence) and argues sponges are not the basal animals and originated in the latest Ediacaran–Cambrian [68,69].

The first complex macroscopic body fossils of the Ediacara biota ~570–540 Ma are interpreted as a mix of stem- and crown-group animals [71,72]. Early ‘rangeomorph’ fronds that stuck up from the deep, dark seafloor could have fed by osmotrophy [73] — enhanced in deep, low-flow regimes by the establishment of a ‘canopy flow’ regime [74] — facilitating uptake of dissolved nutrients. However, if we accept that the Ediacara biota were eukaryotic, then they would have had other feeding modes — phagocytosis — and their own source of motility — flagella — with which to create advection that can significantly enhance nutrient uptake [75]. Either way, fronds could have provided a spatially concentrated source of POM to sediments (e.g. on death), thus lowering DOM and P concentration in the water column [76].

Whenever actively water-pumping, filter-feeding sponges appeared, they would have altered the size structure of the water column ecosystem, ocean nutrient levels and redox state (Figure 4c). As well as efficiently filtering bacteria and POM, in modern (oligotrophic) coral reef settings, the ‘sponge loop’ [77] converts DOM to POM — forming an effective nutrient concentration and recycling system and a mechanism for transferring POM to sediments [78]. Sponge symbionts also sequester polyphosphate, which can trigger apatite formation, thus removing P to sediments [79]. Thus, when sponges arose they could have provided a significant direct pathway of C and P to sediments, lowering overall oxygen demand in the water column and thus tending to oxygenate it. *Thectardis* ~565–555 Ma has been interpreted as a possible sponge [80], although others disagree [68,69]. Subsequently, there is geochemical evidence for the progressive expansion of siliceous sponges on the Yangtze Platform ~550–525 Ma, and associated decrease in the DOC pool, enhanced C_{org} and phosphorus burial, and water column oxygenation [81].

The combination of an oxygenated shelf water column overlying a sediment surface still covered by mats (and sessile animals) produced a shelf-sea ecosystem structure unique to the Neoproterozoic, with a sharp sediment-surface redox gradient that would favour the formation of authigenic carbonate [82] and phosphorite [83–85]. In modern environments, sulphide-oxidising bacterial mats (*Thiomargarita*, *Beggiatoa*) that bridge the water-sediment interface accumulate polyphosphate from oxic waters and utilise it under anoxic conditions, triggering apatite precipitation [86,87]. Neoproterozoic phosphorites are typically associated with stromatolites [88,89] and some contain filamentous microfossils that resemble modern sulphide-oxidising bacteria [90]. Phosphorites are associated with nearshore oxygen oases ~610 Ma [91], then shift to greater depths and areal extent ~570 Ma onwards (e.g. the Doushantuo Formation) [83,91].

Mobile animals

The first mobile trace-makers that scratched across mat surfaces appeared ~565 Ma [92], but did not significantly disrupt the ‘mat seal’ on the sediments, until the first burrowing animals evolved. Fine meiofaunal traces from 555 to 542 Ma have recently been described [93], and widespread burrows capable of significant sediment mixing slightly predate the Precambrian/Cambrian boundary [94].

Bioturbating animals are well-known ‘ecosystem engineers’ [95] that bring oxygen into contact with sediments, increase the turnover rate of C_{org} and O₂ [96], suppressing sulfate reduction [97] and trapping

phosphorus in iron oxides [98], while also increasing water exchange fluxes that release nitrogen and phosphorus to the water column [97]. It is hypothesised that by oxidising upper sediment layers, the evolution of bioturbators increased the sulfate inventory of the ocean [99], and by lowering $(C_{\text{org}}/P)_{\text{burial}}$, initially removed phosphorus and oxygenated the ocean ($\sim 10^4$ years) (Figure 4d) and then lowered atmospheric $p\text{O}_2$ and deoxygenated the ocean ($\sim 10^7$ years) restoring higher P levels [100]. What is unresolved is when these predicted effects became globally significant.

One view is that bioturbation increased in depth and intensity in the Cambrian ‘agronomic revolution’ in Stages 2–4 (~ 530 – 510 Ma) [101,102], causing initial ocean oxygenation and then declining atmospheric $p\text{O}_2$ and ocean deoxygenation over the next ~ 20 Myr [100]. A counter view is that the development of bioturbation only became globally significant from the late Silurian ~ 420 Ma onwards [103,104]. The argument depends on whether sediment P cycling responds linearly or non-linearly to increasing burrowing depth. Diagenetic modelling suggests the effects of burrowing animals are nonlinear and even shallow bioturbation significantly sequesters P [105].

These alternative hypotheses make distinct, testable predictions. Evidence that mid-depth waters of the Yangtze Platform oxygenated from Cambrian stages 2–4 has been used to question the early bioturbation model [106], but is actually consistent with the original predictions [100], which show that ocean oxygenation should accompany the initial onset of deep bioturbation (Stages 2–4), followed by a much slower deoxygenation (governed by the slow timescale of atmospheric $p\text{O}_2$ decline). Wider evidence shows an ocean oxygenation event ~ 520 Ma (broadly coincident with the ‘Cambrian explosion’) followed by deoxygenation [13,107,108].

The Cambrian evolution of large zooplankton would also have increased the efficiency of the biological pump [36], transferring organic matter to sediments, lowering the ocean P inventory and tending to oxygenate the ocean [1,107].

Conclusion

The Neoproterozoic–Cambrian transition was not unidirectional or driven solely by either rising atmospheric $p\text{O}_2$ or evolutionary innovations. We describe a series of eukaryotic innovations that created and strengthened the biological pump of POM from the ocean to the sediments, with major consequences for the phosphorus, carbon and oxygen cycles. We suggest that each phase of eukaryote evolution tended to lower P levels and oxygenate the ocean on $\sim 10^4$ year timescales, but by decreasing C_{org}/P burial ratios tended to lower atmospheric $p\text{O}_2$ and deoxygenate the ocean again on $\sim 10^6$ year timescales. Coupled with tectonic drivers, and biogeochemical and ecological feedback, potentially between alternate stable states [14], this could help explain the transient nature and $\sim 10^6$ year duration of oceanic oxygenation events through the Cryogenian–Ediacaran–Cambrian.

Summary

- The late Tonian ocean ~ 750 Ma was dominated by rapid microbial cycling of DOM with elevated nutrient (P) levels due to inefficient organic matter removal to sediments.
- We hypothesise that abrupt onset of the eukaryotic algal biomarker record in the Cryogenian ~ 660 – 640 Ma was linked to an escalation of protozoan predation (eukaryophagy).
- This ecological regime shift created a ‘biological pump’ of sinking POM, which transferred C_{org} and P to sediments.
- The Late Ediacaran advent of sessile benthic animals on top of microbial mats increased the efficiency of C_{org} and P transfer to sediments, contributing to the deposition of phosphorites.
- The Cambrian explosion of mobile burrowing animals broke the ‘mat seal’ on the upper sediments but by oxygenating them it enabled alternative means of efficient P retention.

- Each phase of eukaryote evolution tended to lower P levels and oxygenate the ocean on $\sim 10^4$ year timescales, but by decreasing C_{org}/P burial ratios, tended to lower atmospheric pO_2 and deoxygenate the ocean again on $\sim 10^6$ year timescales.
- This can help explain the transient nature and $\sim 10^6$ year duration of oceanic oxygenation events through the Cryogenian–Ediacaran–Cambrian periods.

Abbreviations

C_{org} , organic carbon; DOM, dissolved organic matter; POM, particulate organic matter.

Funding

This work was supported by the NERC-NSFC programme ‘Biosphere Evolution, Transitions and Resilience’ through grant NE/P013651/1.

Competing Interests

The Authors declare that there are no competing interests associated with the manuscript.

References

- 1 Lenton, T.M., Boyle, R.A., Poulton, S.W., Shields, G.A. and Butterfield, N.J. (2014) Co-evolution of eukaryotes and ocean oxygenation in the Neoproterozoic era. *Nat. Geosci.* **7**, 257–265 <https://doi.org/10.1038/ngeo2108>
- 2 Shields-Zhou, G. and Och, L. (2011) The case for a Neoproterozoic Oxygenation Event: geochemical evidence and biological consequences. *GSA Today* **21**, 4–11 <https://doi.org/10.1130/GSATG102A.1>
- 3 Och, L.M. and Shields-Zhou, G.A. (2012) The Neoproterozoic oxygenation event: environmental perturbations and biogeochemical cycling. *Earth-Sci. Rev.* **110**, 26–57 <https://doi.org/10.1016/j.earscirev.2011.09.004>
- 4 Lyons, T.W., Reinhard, C.T. and Planavsky, N.J. (2014) The rise of oxygen in Earth’s early ocean and atmosphere. *Nature* **506**, 307–315 <https://doi.org/10.1038/nature13068>
- 5 Planavsky, N.J., Reinhard, C.T., Wang, X., Thomson, D., McGoldrick, P., Rainbird, R.H. et al. (2014) Low Mid-Proterozoic atmospheric oxygen levels and the delayed rise of animals. *Science* **346**, 635–638 <https://doi.org/10.1126/science.1258410>
- 6 Lenton, T.M. and Daines, S.J. (2017) Biogeochemical transformations in the history of the ocean. *Ann. Rev. Mar. Sci.* **9**, 31–58 <https://doi.org/10.1146/annurev-marine-010816-060521>
- 7 Lenton, T.M. and Watson, A.J. (2000) Redfield revisited: 1. Regulation of nitrate, phosphate, and oxygen in the ocean. *Global Biogeochem. Cycles* **14**, 225–248 <https://doi.org/10.1029/1999GB900065>
- 8 Monteiro, F.M., Pancost, R.D., Ridgwell, A. and Donnadieu, Y. (2012) Nutrients as the dominant control on the spread of anoxia and euxinia across the Cenomanian-Turonian oceanic anoxic event (OAE2): model-data comparison. *Paleoceanography* **27**, PA4209 <https://doi.org/10.1029/2012PA002351>
- 9 Lenton, T.M., Daines, S.J. and Mills, B.J.W. (2018) COPSE reloaded: an improved model of biogeochemical cycling over Phanerozoic time. *Earth Sci. Rev.* **178**, 1–28 <https://doi.org/10.1016/j.earscirev.2017.12.004>
- 10 Zhang, F., Zhu, X., Yan, B., Kendall, B., Peng, X., Li, J. et al. (2015) Oxygenation of a Cryogenian ocean (Nanhua Basin, South China) revealed by pyrite Fe isotope compositions. *Earth Planet. Sci. Lett.* **429**, 11–19 <https://doi.org/10.1016/j.epsl.2015.07.021>
- 11 Lau, K.V., Macdonald, F.A., Maher, K. and Payne, J.L. (2017) Uranium isotope evidence for temporary ocean oxygenation in the aftermath of the sturtian snowball earth. *Earth Planet. Sci. Lett.* **458**, 282–292 <https://doi.org/10.1016/j.epsl.2016.10.043>
- 12 Hardisty, D.S., Lu, Z., Bekker, A., Diamond, C.W., Gill, B.C., Jiang, G. et al. (2017) Perspectives on Proterozoic surface ocean redox from iodine contents in ancient and recent carbonate. *Earth Planet. Sci. Lett.* **463**, 159–170 <https://doi.org/10.1016/j.epsl.2017.01.032>
- 13 Sahoo, S.K., Planavsky, N.J., Jiang, G., Kendall, B., Owens, J.D., Wang, X. et al. (2016) Oceanic oxygenation events in the anoxic Ediacaran ocean. *Geobiology* **14**, 457–468 <https://doi.org/10.1111/gbi.12182>
- 14 Handoh, I.C. and Lenton, T.M. (2003) Periodic mid-Cretaceous Oceanic Anoxic Events linked by oscillations of the phosphorus and oxygen biogeochemical cycles. *Global Biogeochem. Cycles* **17**, 1092 <https://doi.org/10.1029/2003GB002039>
- 15 England, M.H. (1995) The age of water and ventilation timescales in a global ocean model. *J. Phys. Oceanogr.* **25**, 2756–2777 [https://doi.org/10.1175/1520-0485\(1995\)025<2756:TAOWAV>2.0.CO;2](https://doi.org/10.1175/1520-0485(1995)025<2756:TAOWAV>2.0.CO;2)
- 16 Wallmann, K. (2010) Phosphorus imbalance in the global ocean? *Global Biogeochem. Cycles* **24**, GB4030 <https://doi.org/10.1029/2009GB003643>
- 17 Daines, S.J., Mills, B.J.W. and Lenton, T.M. (2017) Atmospheric oxygen regulation at low Proterozoic levels by incomplete oxidative weathering of sedimentary organic carbon. *Nat. Commun.* **8**, 14379 <https://doi.org/10.1038/ncomms14379>
- 18 Slomp, C.P. and Van Cappellen, P. (2007) The global marine phosphorus cycle: sensitivity to oceanic circulation. *Biogeosciences* **4**, 155–171 <https://doi.org/10.5194/bg-4-155-2007>
- 19 Burdige, D.J. (2007) Preservation of organic matter in marine sediments: controls, mechanisms, and an imbalance in sediment organic carbon budgets? *Chem. Rev.* **107**, 467–485 <https://doi.org/10.1021/cr050347q>
- 20 Ingall, E. and Jahnke, R. (1994) Evidence for enhanced phosphorus regeneration from marine sediments overlain by oxygen depleted waters. *Geochim. Cosmochim. Acta* **58**, 2571–2575 [https://doi.org/10.1016/0016-7037\(94\)90033-7](https://doi.org/10.1016/0016-7037(94)90033-7)

- 21 Ingall, E.D., Bustin, R.M. and Van Cappellen, P. (1993) Influence of water column anoxia on the burial and preservation of carbon and phosphorus in marine shales. *Geochim. Cosmochim. Acta* **57**, 303–316 [https://doi.org/10.1016/0016-7037\(93\)90433-W](https://doi.org/10.1016/0016-7037(93)90433-W)
- 22 Sommer, S., Clemens, D., Yücel, M., Pfannkuche, O., Hall, P.O.J., Almroth-Rosell, E. et al. (2017) Major bottom water ventilation events do not significantly reduce basin-wide benthic N and P release in the eastern Gotland basin (Baltic Sea). *Front. Mar. Sci.* **4**, 18 <https://doi.org/10.3389/fmars.2017.00018>
- 23 Van Cappellen, P. and Ingall, E.D. (1994) Benthic phosphorus regeneration, net primary production, and ocean anoxia: a model of the coupled marine biogeochemical cycles of carbon and phosphorus. *Paleoceanography* **9**, 677–692 <https://doi.org/10.1029/94PA01455>
- 24 Van Cappellen, P. and Ingall, E.D. (1996) Redox stabilization of the atmosphere and oceans by phosphorus-limited marine productivity. *Science* **271**, 493–496 <https://doi.org/10.1126/science.271.5248.493>
- 25 Poulton, S.W. and Canfield, D.E. (2011) Ferruginous conditions: a dominant feature of the ocean through Earth's history. *Elements* **7**, 107–112 <https://doi.org/10.2113/gselements.7.2.107>
- 26 Zegeye, A., Bonneville, S., Benning, L.G., Sturm, A., Fowle, D.A., Jones, C.A. et al. (2012) Green rust formation controls nutrient availability in a ferruginous water column. *Geology* **40**, 599–602 <https://doi.org/10.1130/G32959.1>
- 27 Derry, L.A. (2015) Causes and consequences of mid-Proterozoic anoxia. *Geophys. Res. Lett.* **42**, 8538–8546 <https://doi.org/10.1002/2015GL065333>
- 28 Laakso, T.A. and Schrag, D.P. (2014) Regulation of atmospheric oxygen during the Proterozoic. *Earth Planet. Sci. Lett.* **388**, 81–91 <https://doi.org/10.1016/j.epsl.2013.11.049>
- 29 Reinhard, C.T., Planavsky, N.J., Gill, B.C., Ozaki, K., Robbins, L.J., Lyons, T.W. et al. (2017) Evolution of the global phosphorus cycle. *Nature* **541**, 386–389 <https://doi.org/10.1038/nature20772>
- 30 Parfrey, L.W., Lahr, D.J.G., Knoll, A.H. and Katz, L.A. (2011) Estimating the timing of early eukaryotic diversification with multigene molecular clocks. *Proc. Natl Acad. Sci. U.S.A.* **108**, 13624–13629 <https://doi.org/10.1073/pnas.1110633108>
- 31 Butterfield, N.J. (2000) *Bangiomorpha pubescens* n. gen., n. sp.: implications for the evolution of sex, multicellularity, and the Mesoproterozoic/Neoproterozoic radiation of eukaryotes. *Paleobiology* **26**, 386–404 [https://doi.org/10.1666/0094-8373\(2000\)026<0386:BPNGNS>2.0.CO;2](https://doi.org/10.1666/0094-8373(2000)026<0386:BPNGNS>2.0.CO;2)
- 32 Sánchez-Baracaldo, P., Raven, J.A., Pisani, D. and Knoll, A.H. (2017) Early photosynthetic eukaryotes inhabited low-salinity habitats. *Proc. Natl Acad. Sci. U.S.A.* **114**, E7737–E7745 <https://doi.org/10.1073/pnas.1620089114>
- 33 Butterfield, N.J. (2015) Proterozoic photosynthesis - a critical review. *Palaeontology* **58**, 953–972 <https://doi.org/10.1111/pala.12211>
- 34 Brocks, J.J., Jarrett, A.J.M., Sirantoine, E., Hallmann, C., Hoshino, Y. and Liyanage, T. (2017) The rise of algae in Cryogenian oceans and the emergence of animals. *Nature* **548**, 578–581 <https://doi.org/10.1038/nature23457>
- 35 Brocks, J.J., Jarrett, A.J.M., Sirantoine, E., Kenig, F., Moczyłowska, M., Porter, S. et al. (2016) Early sponges and toxic protists: possible sources of cryostane, an age diagnostic biomarker antedating Sturtian Snowball Earth. *Geobiology* **14**, 129–149 <https://doi.org/10.1111/gbi.12165>
- 36 Logan, G.A., Hayes, J.M., Hieshima, G.B. and Summons, R.E. (1995) Terminal Proterozoic reorganization of biogeochemical cycles. *Nature* **376**, 53–56 <https://doi.org/10.1038/376053a0>
- 37 Pawłowska, M.M., Butterfield, N.J. and Brocks, J.J. (2013) Lipid taphonomy in the Proterozoic and the effect of microbial mats on biomarker preservation. *Geology* **41**, 103–106 <https://doi.org/10.1130/G33525.1>
- 38 Hansell, D.A., Carlson, C.A., Repeta, D.J. and Schlitzer, R. (2009) Dissolved organic matter in the ocean: a controversy stimulates new insights. *Oceanography* **22**, 202–211 <https://doi.org/10.5670/oceanog.2009.109>
- 39 Jiao, N., Herndl, G.J., Hansell, D.A., Benner, R., Kattner, G., Wilhelm, S.W. et al. (2010) Microbial production of recalcitrant dissolved organic matter: long-term carbon storage in the global ocean. *Nat. Rev. Microbiol.* **8**, 593–599 <https://doi.org/10.1038/nrmicro2386>
- 40 Rothman, D.H., Hayes, J.M. and Summons, R.E. (2003) Dynamics of the Neoproterozoic carbon cycle. *Proc. Natl Acad. Sci. U.S.A.* **100**, 8124–8129 <https://doi.org/10.1073/pnas.0832439100>
- 41 Reinhard, C.T., Planavsky, N.J., Olson, S.L., Lyons, T.W. and Erwin, D.H. (2016) Earth's oxygen cycle and the evolution of animal life. *Proc. Natl Acad. Sci. U.S.A.* **113**, 8933–8938 <https://doi.org/10.1073/pnas.1521544113>
- 42 Slack, J.F., Grenne, T. and Bekker, A. (2009) Seafloor-hydrothermal Si-Fe-Mn exhalites in the Pecos greenstone belt, New Mexico, and the redox state of ca. 1720 Ma deep seawater. *Geosphere* **5**, 302–314 <https://doi.org/10.1130/GES00220.1>
- 43 Slack, J.F., Grenne, T., Bekker, A., Rouxel, O.J. and Lindberg, P.A. (2007) Suboxic deep seawater in the late Paleoproterozoic: evidence from hematitic chert and iron formation related to seafloor-hydrothermal sulfide deposits, central Arizona, USA. *Earth Planet. Sci. Lett.* **255**, 243–256 <https://doi.org/10.1016/j.epsl.2006.12.018>
- 44 Engel, A., Thoms, S., Riebesell, U., Rochelle-Newall, E. and Zondervan, I. (2004) Polysaccharide aggregation as a potential sink of marine dissolved organic carbon. *Nature* **428**, 929–932 <https://doi.org/10.1038/nature02453>
- 45 Planavsky, N.J., Rouxel, O.J., Bekker, A., Lalonde, S.V., Konhauser, K.O., Reinhard, C.T. et al. (2010) The evolution of the marine phosphate reservoir. *Nature* **467**, 1088–1090 <https://doi.org/10.1038/nature09485>
- 46 Porter, S.M. and Knoll, A.H. (2000) Testate amoebae in the Neoproterozoic Era: evidence from vase-shaped microfossils in the Chuar Group, Grand Canyon. *Paleobiology* **26**, 360–385 [https://doi.org/10.1666/0094-8373\(2000\)026<0360:TAITNE>2.0.CO;2](https://doi.org/10.1666/0094-8373(2000)026<0360:TAITNE>2.0.CO;2)
- 47 Cohen, P.A., Schopf, J.W., Butterfield, N.J., Kudryavtsev, A.B. and Macdonald, F.A. (2011) Phosphate biomineralization in mid-Neoproterozoic protists. *Geology* **39**, 539–542 <https://doi.org/10.1130/G31833.1>
- 48 Javaux, E. (2011) Early eukaryotes in Precambrian oceans. In *Origins and Evolution of Life: An Astrobiological Perspective* (Gargaud, M., Lopez-Garcia, P. and Martin, H., eds), pp. 414–449, Cambridge University Press, Cambridge
- 49 Cohen, P.A. and Macdonald, F.A. (2015) The proterozoic record of eukaryotes. *Paleobiology* **41**, 610–632 <https://doi.org/10.1017/pab.2015.25>
- 50 Rooney, A.D., Macdonald, F.A., Strauss, J.V., Dudás, F.Ö., Hallmann, C. and Selby, D. (2014) Re-Os geochronology and coupled Os-Sr isotope constraints on the Sturtian snowball Earth. *Proc. Natl Acad. Sci. U.S.A.* **111**, 51–56 <https://doi.org/10.1073/pnas.1317266110>
- 51 Knoll, A.H. (2017) Food for early animal evolution. *Nature* **548**, 528–530 <https://doi.org/10.1038/nature23539>
- 52 Armstrong, R.A. (1994) Grazing limitation and nutrient limitation in marine ecosystems: steady state solutions of an ecosystem model with multiple food chains. *Limnol. Oceanogr.* **39**, 597–608 <https://doi.org/10.4319/lo.1994.39.3.0597>
- 53 Hartmann, M., Grob, C., Tarran, G.A., Martin, A.P., Burkill, P.H., Scanlan, D.J. et al. (2012) Mixotrophic basis of Atlantic oligotrophic ecosystems. *Proc. Natl Acad. Sci. U.S.A.* **109**, 5756–5760 <https://doi.org/10.1073/pnas.1118179109>

- 54 Higgins, M.B., Robinson, R.S., Husson, J.M., Carter, S.J. and Pearson, A. (2012) Dominant eukaryotic export production during ocean anoxic events reflects the importance of recycled NH_4^+ . *Proc. Natl Acad. Sci. U.S.A.* **109**, 2269–2274 <https://doi.org/10.1073/pnas.1104313109>
- 55 Knoll, A.H. (2014) Paleobiological perspectives on early eukaryotic evolution. *Cold Spring Harbor Perspect. Biol.* **6**, a016121 <https://doi.org/10.1101/cshperspect.a016121>
- 56 Porter, S. (2011) The rise of predators. *Geology* **39**, 607–608 <https://doi.org/10.1130/focus062011.1>
- 57 Knoll, A.H. and Follows, M.J. (2016) A bottom-up perspective on ecosystem change in Mesozoic oceans. *Proc. Royal Soc. B: Biol. Sci.* **283**, 20161755 <https://doi.org/10.1098/rspb.2016.1755>
- 58 Love, G.D., Grosjean, E., Stalvies, C., Fike, D.A., Grotzinger, J.P., Bradley, A.S. et al. (2009) Fossil steroids record the appearance of Demospongiae during the Cryogenian period. *Nature* **457**, 718–721 <https://doi.org/10.1038/nature07673>
- 59 Gold, D.A., Grabenstatter, J., de Mendoza, A., Riesgo, A., Ruiz-Trillo, I. and Summons, R.E. (2016) Sterol and genomic analyses validate the sponge biomarker hypothesis. *Proc. Natl Acad. Sci. U.S.A.* **113**, 2684–2689 <https://doi.org/10.1073/pnas.1512614113>
- 60 Love, G.D. and Summons, R.E. (2015) The molecular record of Cryogenian sponges - a response to Antcliffe (2013). *Palaeontology* **58**, 1131–1136 <https://doi.org/10.1111/pala.12196>
- 61 Meyer, K.M., Ridgwell, A. and Payne, J.L. (2016) The influence of the biological pump on ocean chemistry: implications for long-term trends in marine redox chemistry, the global carbon cycle, and marine animal ecosystems. *Geobiology* **14**, 207–219 <https://doi.org/10.1111/gbi.12176>
- 62 Kunzmann, M., Gibson, T.M., Halverson, G.P., Hodgskiss, M.S.W., Bui, T.H., Carozza, D.A. et al. (2017) Iron isotope biogeochemistry of Neoproterozoic marine shales. *Geochim. Cosmochim. Acta* **209**, 85–105 <https://doi.org/10.1016/j.gca.2017.04.003>
- 63 Simion, P., Philippe, H., Baurain, D., Jager, M., Richter, D.J., Di Franco, A. et al. (2017) A large and consistent phylogenomic dataset supports sponges as the sister group to all other animals. *Curr. Biol.* **27**, 958–967 <https://doi.org/10.1016/j.cub.2017.02.031>
- 64 Feuda, R., Dohrmann, M., Pett, W., Philippe, H., Rota-Stabelli, O., Lartillot, N. et al. (2017) Improved modeling of compositional heterogeneity supports sponges as sister to all other animals. *Curr. Biol.* **27**, 3864–3870.e4 <https://doi.org/10.1016/j.cub.2017.11.008>
- 65 Schuster, A., Vargas, S., Knapp, I., Pomponi, S.A., Toonen, R.J., Erpenbeck, D. et al. (2017) Divergence times in demosponges (Porifera): first insights from new mitogenomes and the inclusion of fossils in a birth-death clock model. *bioRxiv* <https://doi.org/10.1101/159806>
- 66 Ma, J.-Y. and Yang, Q. (2016) Early divergence dates of demosponges based on mitogenomics and evaluated fossil calibrations. *Palaeoworld* **25**, 292–302 <https://doi.org/10.1016/j.palwor.2015.03.004>
- 67 Botting, J.P., Cárdenas, P. and Peel, J.S. (2015) A crown-group demosponge from the early Cambrian Sirius Passet Biota, North Greenland. *Palaeontology* **58**, 35–43 <https://doi.org/10.1111/pala.12133>
- 68 Antcliffe, J.B., Callow, R.H.T. and Brasier, M.D. (2014) Giving the early fossil record of sponges a squeeze. *Biol. Rev.* **89**, 972–1004 <https://doi.org/10.1111/brv.12090>
- 69 Botting, J.P. and Muir, L.A. (2018) Early sponge evolution: a review and phylogenetic framework. *Palaeoworld* **27**, 1–29 <https://doi.org/10.1016/j.palwor.2017.07.001>
- 70 Yin, Z., Zhu, M., Davidson, E.H., Bottjer, D.J., Zhao, F. and Tafforeau, P. (2015) Sponge grade body fossil with cellular resolution dating 60 Myr before the Cambrian. *Proc. Natl Acad. Sci. U.S.A.* **112**, E1453–E1460 <https://doi.org/10.1073/pnas.1414577112>
- 71 Droser, M.L., Tarhan, L.G. and Gehling, J.G. (2017) The rise of animals in a changing environment: global ecological innovation in the late Ediacaran. *Annu. Rev. Earth Planet. Sci.* **45**, 593–617 <https://doi.org/10.1146/annurev-earth-063016-015645>
- 72 Muscente, A.D., Boag, T.H., Bykova, N. and Schiffbauer, J.D. (2018) Environmental disturbance, resource availability, and biologic turnover at the dawn of animal life. *Earth-Science Reviews*. **177**, 248–264 <https://doi.org/10.1016/j.earscirev.2017.11.019>
- 73 Laflamme, M., Xiao, S. and Kowalewski, M. (2009) Osmotrophy in modular Ediacara organisms. *Proc. Natl Acad. Sci. U.S.A.* **106**, 14438–14443 <https://doi.org/10.1073/pnas.0904836106>
- 74 Ghisalberti, M., Gold, D.A., Laflamme, M., Clapham, M.E., Narbonne, G.M., Summons, R.E. et al. (2014) Canopy flow analysis reveals the advantage of size in the oldest communities of multicellular eukaryotes. *Curr. Biol.* **24**, 305–309 <https://doi.org/10.1016/j.cub.2013.12.017>
- 75 Solari, C.A., Ganguly, S., Kessler, J.O., Michod, R.E. and Goldstein, R.E. (2006) Multicellularity and the functional interdependence of motility and molecular transport. *Proc. Natl Acad. Sci. U.S.A.* **103**, 1353–1358 <https://doi.org/10.1073/pnas.0503810103>
- 76 Budd, G.E. and Jensen, S. (2017) The origin of the animals and a ‘Savannah’ hypothesis for early bilaterian evolution. *Biol. Rev.* **92**, 446–473 <https://doi.org/10.1111/brv.12239>
- 77 de Goeij, J.M., van Oevelen, D., Vermeij, M.J.A., Osinga, R., Middelburg, J.J., de Goeij, A.F.P.M. et al. (2013) Surviving in a marine desert: the sponge loop retains resources within coral reefs. *Science* **342**, 108–110 <https://doi.org/10.1126/science.1241981>
- 78 Middelburg, J.J. (2018) Reviews and syntheses: to the bottom of carbon processing at the seafloor. *Biogeosciences* **15**, 413–427 <https://doi.org/10.5194/bg-15-413-2018>
- 79 Zhang, F., Blasiak, L.C., Karolin, J.O., Powell, R.J., Geddes, C.D. and Hill, R.T. (2015) Phosphorus sequestration in the form of polyphosphate by microbial symbionts in marine sponges. *Proc. Natl Acad. Sci. U.S.A.* **112**, 4381–4386 <https://doi.org/10.1073/pnas.1423768112>
- 80 Sperling, E.A., Peterson, K.J. and Laflamme, M. (2011) Rangeomorphs, Thectardis (Porifera?) and dissolved organic carbon in the Ediacaran oceans. *Geobiology* **9**, 24–33 <https://doi.org/10.1111/j.1472-4669.2010.00259.x>
- 81 Tatzel, M., von Blanckenburg, F., Oelze, M., Bouchez, J. and Hippler, D. (2017) Late Neoproterozoic seawater oxygenation by siliceous sponges. *Nat. Commun.* **8**, 621 <https://doi.org/10.1038/s41467-017-00586-5>
- 82 Cui, H., Kaufman, A.J., Xiao, S., Zhou, C. and Liu, X.-M. (2017) Was the Ediacaran Shuram Excursion a globally synchronized early diagenetic event? Insights from methane-derived authigenic carbonates in the uppermost Doushantuo Formation, South China. *Chem. Geol.* **450**, 59–80 <https://doi.org/10.1016/j.chemgeo.2016.12.010>
- 83 Cui, H., Xiao, S., Zhou, C., Peng, Y., Kaufman, A.J. and Plummer, R.E. (2016) Phosphogenesis associated with the Shuram Excursion: petrographic and geochemical observations from the Ediacaran Doushantuo Formation of South China. *Sediment. Geol.* **341**, 134–146 <https://doi.org/10.1016/j.sedgeo.2016.05.008>
- 84 Crosby, C.H. and Bailey, J.V. (2012) The role of microbes in the formation of modern and ancient phosphatic mineral deposits. *Front. Microbiol.* **3**, 241 <https://doi.org/10.3389/fmicb.2012.00241>

- 85 Omelon, S., Ariganello, M., Bonucci, E., Grynpsas, M. and Nanci, A. (2013) A review of phosphate mineral nucleation in biology and geobiology. *Calif. Tissue Int.* **93**, 382–396 <https://doi.org/10.1007/s00223-013-9784-9>
- 86 Goldhammer, T., Brüchert, V., Ferdelman, T.G. and Zabel, M. (2010) Microbial sequestration of phosphorus in anoxic upwelling sediments. *Nat. Geosci.* **3**, 557–561 <https://doi.org/10.1038/ngeo913>
- 87 Schulz, H.N. and Schulz, H.D. (2005) Large sulfur bacteria and the formation of phosphorite. *Science* **307**, 416–418 <https://doi.org/10.1126/science.1103096>
- 88 Papineau, D. (2010) Global biogeochemical changes at both ends of the proterozoic: insights from phosphorites. *Astrobiology* **10**, 165–181 <https://doi.org/10.1089/ast.2009.0360>
- 89 Gómez-Peral, L.E., Kaufman, A.J. and Poiré, D.G. (2014) Paleoenvironmental implications of two phosphogenic events in Neoproterozoic sedimentary successions of the Tandilia System, Argentina. *Precambrian Res.* **252**, 88–106 <https://doi.org/10.1016/j.precamres.2014.07.009>
- 90 Bailey, J.V., Corsetti, F.A., Greene, S.E., Crosby, C.H., Liu, P. and Orphan, V.J. (2013) Filamentous sulfur bacteria preserved in modern and ancient phosphatic sediments: implications for the role of oxygen and bacteria in phosphogenesis. *Geobiology* **11**, 397–405 <https://doi.org/10.1111/gbi.12046>
- 91 Drummond, J.B.R., Pufahl, P.K., Porto, C.G. and Carvalho, M. (2015) Neoproterozoic peritidal phosphorite from the Sete Lagoas Formation (Brazil) and the Precambrian phosphorus cycle. *Sedimentology* **62**, 1978–2008 <https://doi.org/10.1111/sed.12214>
- 92 Liu, A.G., Mclroy, D. and Brasier, M.D. (2010) First evidence for locomotion in the Ediacara biota from the 565 Ma Mistaken Point Formation, Newfoundland. *Geology* **38**, 123–126 <https://doi.org/10.1130/G30368.1>
- 93 Parry, L.A., Boggiani, P.C., Condon, D.J., Garwood, R.J., de Leme, J.M., Mclroy, D. et al. (2017) Ichological evidence for meiofaunal bilaterians from the terminal Ediacaran and earliest Cambrian of Brazil. *Nat. Ecol. Evol.* **1**, 1455–1464 <https://doi.org/10.1038/s41559-017-0301-9>
- 94 Jensen, S., Saylor, B.Z., Gehling, J.G. and Germs, G.J.B. (2000) Complex trace fossils from the terminal Proterozoic of Namibia. *Geology* **28**, 143–146 [https://doi.org/10.1130/0091-7613\(2000\)28<143:CTFFTT>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<143:CTFFTT>2.0.CO;2)
- 95 Mermillod-Blondin, F. and Rosenberg, R. (2006) Ecosystem engineering: the impact of bioturbation on biogeochemical processes in marine and freshwater benthic habitats. *Aquat. Sci.* **68**, 434–442 <https://doi.org/10.1007/s00027-006-0858-x>
- 96 Kristensen, E. (2001) Impact of polychaetes (*Nereis* spp. and *Arenicola marina*) on carbon biogeochemistry in coastal marine sediments. *Geochem. Trans.* **2**, 92 <https://doi.org/10.1186/1467-4866-2-92>
- 97 Mermillod-Blondin, F., Rosenberg, R., François-Carcaillet, F., Norling, K. and Mauclair, L. (2004) Influence of bioturbation by three benthic infaunal species on microbial communities and biogeochemical processes in marine sediment. *Aquat. Microb. Ecol.* **36**, 271–284 <https://doi.org/10.3354/ame036271>
- 98 Zhang, L., Gu, X., Fan, C., Shang, J., Shen, Q., Wang, Z. et al. (2010) Impact of different benthic animals on phosphorus dynamics across the sediment-water interface. *J. Environ. Sci.* **22**, 1674–1682 [https://doi.org/10.1016/S1001-0742\(09\)60305-3](https://doi.org/10.1016/S1001-0742(09)60305-3)
- 99 Canfield, D.E. and Farquhar, J. (2009) Animal evolution, bioturbation, and the sulfate concentration of the oceans. *Proc. Natl Acad. Sci. U.S.A.* **106**, 8123–8127 <https://doi.org/10.1073/pnas.0902037106>
- 100 Boyle, R.A., Dahl, T.W., Dale, A.W., Shields-Zhou, G.A., Zhu, M., Brasier, M.D. et al. (2014) Stabilization of the coupled oxygen and phosphorus cycles by the evolution of bioturbation. *Nat. Geosci.* **7**, 671–676 <https://doi.org/10.1038/ngeo2213>
- 101 Mángano, M.G. and Buatois, L.A. (2017) The Cambrian revolutions: trace-fossil record, timing, links and geological impact. *Earth-Sci. Rev.* **173**, 96–108 <https://doi.org/10.1016/j.earscirev.2017.08.009>
- 102 Zhang, L.-J., Qi, Y.-A., Buatois, L.A., Mángano, M.G., Meng, Y. and Li, D. (2017) The impact of deep-tier burrow systems in sediment mixing and ecosystem engineering in early Cambrian carbonate settings. *Sci. Rep.* **7**, 45773 <https://doi.org/10.1038/srep45773>
- 103 Tarhan, L.G. and Droser, M.L. (2014) Widespread delayed mixing in early to middle Cambrian marine shelfal settings. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **399**, 310–322 <https://doi.org/10.1016/j.palaeo.2014.01.024>
- 104 Tarhan, L.G., Droser, M.L., Planavsky, N.J. and Johnston, D.T. (2015) Protracted development of bioturbation through the early Palaeozoic Era. *Nat. Geosci.* **8**, 865–869 <https://doi.org/10.1038/ngeo2537>
- 105 Dale, A.W., Boyle, R.A., Lenton, T.M., Ingall, E.D. and Wallmann, K. (2016) A model for microbial phosphorus cycling in bioturbated marine sediments: significance for phosphorus burial in the early Paleozoic. *Geochim. Cosmochim. Acta* **189**, 251–268 <https://doi.org/10.1016/j.gca.2016.05.046>
- 106 Li, C., Jin, C., Planavsky, N.J., Algeo, T.J., Cheng, M., Yang, X. et al. (2017) Coupled oceanic oxygenation and metazoan diversification during the early–middle Cambrian? *Geology* **45**, 743–746 <https://doi.org/10.1130/G39208.1>
- 107 Dahl, T.W., Connelly, J.N., Kouchinsky, A., Gill, B.C., Månsson, S.F. and Bizzarro, M. (2017) Reorganisation of Earth's biogeochemical cycles briefly oxygenated the oceans 520 Myr ago. *Geochem. Perspect. Lett.* **3**, 210–220 <https://doi.org/10.7185/geochemlet.1724>
- 108 Chen, X., Ling, H.-F., Vance, D., Shields-Zhou, G.A., Zhu, M., Poulton, S.W. et al. (2015) Rise to modern levels of ocean oxygenation coincided with the Cambrian radiation of animals. *Nat. Commun.* **6**, 7142 <https://doi.org/10.1038/ncomms8142>
- 109 Canfield, D.E., Poulton, S.W., Knoll, A.H., Narbonne, G.M., Ross, G., Goldberg, T. et al. (2008) Ferruginous conditions dominated later neoproterozoic deep-water chemistry. *Science* **321**, 949–952 <https://doi.org/10.1126/science.1154499>
- 110 Canfield, D.E., Poulton, S.W. and Narbonne, G.M. (2007) Late-Neoproterozoic deep-ocean oxygenation and the rise of animal life. *Science* **315**, 92–95 <https://doi.org/10.1126/science.1135013>
- 111 Ward, B.A., Dutkiewicz, S. and Follows, M.J. (2014) Modelling spatial and temporal patterns in size-structured marine plankton communities: top–down and bottom–up controls. *J. Plankton Res.* **36**, 31–47 <https://doi.org/10.1093/plankt/ftb097>