



Reply to Ben-Dor and Barkai: A low Zn isotope ratio is not equal to a low Zn content

Klervia Jaouen^{a,1}, Thomas Tütken^b, Nicolas Bourgon^{b,c}, Tina Lüdecke^d, Geoff M. Smith^e, Domingo C. Salazar-García^{f,g}, Jean-Jacques Hublin^{c,h}, Vanessa Villalba-Mouco^{ij}, and Pauline Méjean^a

We are grateful to Miki Ben-Dor and Ran Barkai for their interest in our study (1). We acknowledge that a high-fat diet should have been discussed in our hypothesis as an additional dietary option.

However, we are afraid that there was a misunderstanding in the interpretation of our data. Ben-Dor and Barkai (2) wrote that “One explanation presented in the [our] paper was that Neandertals may have consumed body parts and animals that were relatively poor in zinc, like the liver or deer.” This sentence is not reflecting our conclusions: We discussed Zn isotope ratios (i.e., ⁶⁶Zn/⁶⁴Zn ratio expressed as a $\delta^{66}\text{Zn}$ value) and not Zn concentrations. To explain low $\delta^{66}\text{Zn}$ values of the Neandertal tooth, we propose that Neandertals ate food items depleted in heavy Zn isotopes (and therefore enriched in light Zn isotopes), but those foods can have variable Zn contents, independent of the Zn isotope composition as such. A large fraction of Zn with low $\delta^{66}\text{Zn}$ needs to be ingested with the diet to shift Neandertal enamel mineralizing to low $\delta^{66}\text{Zn}$ values. Thus, the proposition of Ben-Dor and Barkai (2) that “a high animal fat consumption with zero zinc content as a probable explanation for the extremely low $\delta^{66}\text{Zn}$ in the Gabasa Neandertal” does not work as the zinc content of fat is too low to contribute to the enamel $\delta^{66}\text{Zn}$. Furthermore, fat and bone marrow do not even have low $\delta^{66}\text{Zn}$ values (3, 4).

To support this statement, we here give an example based on the numbers suggested by Ben-Dor and Barkai (2). Assuming 1) a concentration of Zn in fat of 0.06 mg/100 g and a concentration of Zn in meat (0% fat) of 4 mg/100 g (5), 2) a consumption of 1 kg meat with 20% of protein and 15% of fat per day, 3) Zn isotope compositions of meat ($\delta^{66}\text{Zn} = 0.15\text{‰}$) and of fat ($\delta^{66}\text{Zn} = 0.4\text{‰}$) (3), 4) and that proteins contribute to 4 kcal/g and fat to 9 kcal/g in the diet (6), we obtain a $\delta^{66}\text{Zn}$ of 0.1506‰ (Table 1). Thus, compared with a fat-free meat diet, the $\delta^{66}\text{Zn}$ value is only increasing by 0.0006‰, which is two orders of magnitude lower than the analytical error of $\delta^{66}\text{Zn}$ measurement (<0.05‰, 1 SD) and therefore not significant.

The consumption of bone grease has been reported in Pyrenean Mousterian sites as described by the work of Costamagno (7, 8) in the nearby location of the Noisetier Cave and Blasco (9) through the exploitation of bone marrow for the very site of Gabasa. Hence, the consumption of fat

Table 1. Contribution of fat to the Zn isotope composition of 1 kg meat (see text for references)

	Mass in g	Contribution in kcal	% in the diet	Amount of Zn (mg)	$\delta^{66}\text{Zn}$ (‰)
Proteins	200	800	37%		
Lipids	150	1,350	63%	0.09	0.4
Meat	850	800	37%	34	0.15
without fat					
Total amount of meat eaten	1,000	2,150	100%	34.09	0.1506

could potentially be substantial at these sites, but at this stage, zinc isotope ratios do not allow us to quantify it. As shown by our mass balance, a high-fat diet cannot explain the low $\delta^{66}\text{Zn}$ values that we observe in the Neandertal tooth of Gabasa as it would have a very small impact on the $\delta^{66}\text{Zn}$ bulk value of the diet.

Author affiliations: ^aGéosciences Environnement Toulouse/CNRS, Observatoire Midi Pyrénées, Toulouse 31400, France; ^bApplied and Analytical Palaeontology, Institute of Geosciences, Johannes Gutenberg University, Mainz 55122, Germany; ^cDepartment of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Leipzig 04103, Germany; ^dEmmy Noether Group for Hominin Meat Consumption, Max Planck Institute for Chemistry, Mainz 55128, Germany; ^eSchool of Anthropology and Conservation, University of Kent, Canterbury CT2 7NZ, United Kingdom; ^fDepartament de Prehistòria, Arqueologia i Història Antiga, Universitat de València, València 46101, Spain; ^gDepartment of Geological Sciences, University of Cape Town, Cape Town 7700, South Africa; ^hChaire internationale de Paléontologie, Collège de France, Paris 75005, France; ⁱInstitute of Evolutionary Biology, Consejo Superior de Investigaciones Científicas-Universitat Pompeu Fabra, Barcelona 08003, Spain; and ^jResearch Group “Primeros Pobladores y Patrimonio Arqueológico del Valle del Ebro” Department of Sciences of Antiquity-Prehistory, University of Zaragoza, Zaragoza 50009, Spain

Author contributions: K.J., T.T., N.B., D.C.S.-G., T.L., and J.-J.H. designed research; K.J., T.T., N.B., T.L., G.M.S., D.C.S.-G., V.V.-M., and P.M. performed research; K.J., T.T., N.B., T.L., G.M.S., D.C.S.-G., J.-J.H., V.V.-M., and P.M. analyzed data; and K.J., T.T., N.B., T.L., G.M.S., D.C.S.-G., V.V.-M., and P.M. wrote the paper.

The authors declare no competing interest.

Copyright © 2023 the Author(s). Published by PNAS. This article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

¹To whom correspondence may be addressed. Email: klervia.jaouen@get.omp.eu.

Published January 31, 2023.

1. K. Jaouen *et al.*, A Neandertal dietary conundrum: New insights provided by tooth enamel Zn isotopes from Gabasa, Spain. *Proc. Natl. Acad. Sci. U.S.A.* **119**, 43 (2022).
2. M. Ben-Dor, R. Birkai, A limited protein high-fat diet may explain the low $\delta^{64}\text{Zn}$ conundrum in the Neandertal from Gabasa. *Proc. Natl. Acad. Sci. U.S.A.* **11**, 301 (2022).
3. F. Moynier, T. Fujii, A. S. Shaw, M. Le Borgne, Heterogeneous distribution of natural zinc isotopes in mice. *Metallomics* **5**, 693–699 (2013).
4. V. Balter *et al.*, Contrasting Cu, Fe, and Zn isotopic patterns in organs and body fluids of mice and sheep, with emphasis on cellular fractionation. *Metallomics* **5**, 1470–1482 (2013).
5. FoodData Central, <https://fdc.nal.usda.gov/>. Accessed 2 November 2022
6. J. D. Speth, K. A. Spielmann, Energy source, protein metabolism, and hunter-gatherer subsistence strategies. *J. Anthropol. Archaeol.* **2**, 1–31 (1983).
7. S. Costamagno, "Bone grease rendering in mousterian contexts: The case of Noisetier cave (Fréchet-Aure, Hautes-Pyrénées, France)" in *Zooarchaeology and Modern Human Origins: Human Hunting Behavior during the Later Pleistocene, Vertebrate Paleobiology, Paleoanthropology*, J. L. Clark, J. D. Speth, Eds. (Springer, Netherlands, 2013), pp. 209–225.
8. S. Costamagno, J.-P. Rigaud, L. Laud, "Ligaud de la graisse au Pal au Pa" in *138 e Congrès National Des Sociétés Historiques et Scientifiques, Histoire de l'alimentation humaine: entre choix et contraintes*, S. Costamagno, Ed. (CTHS, 2013), pp. 134–152, Édition électronique du CTHS (Actes des congrès des sociétés historiques et scientifiques).
9. M. F. Blasco, In the pursuit of game: The Mousterian cave site of Gabasa 1 in the Spanish Pyrenees. *J. Anthropol. Res.* **53**, 177–217 (1997).