




# How Can We Best Diagnose Severity Levels of Dry Eye Disease: Current Perspectives

Thomas Chester<sup>1</sup>, Sumit (Sam) Garg<sup>2</sup>, Josh Johnston<sup>3,4</sup>, Brandon Ayers<sup>5</sup>, Preeya Gupta<sup>6,7</sup>

<sup>1</sup>Cleveland Eye Clinic, Brecksville, OH, USA; <sup>2</sup>Department of Ophthalmology, Gavin Herbert Eye Institute, University of California-Irvine, Irvine, CA, USA; <sup>3</sup>Georgia Eye Partners, Atlanta, GA, USA; <sup>4</sup>Southern College of Optometry, Memphis, TN, USA; <sup>5</sup>Ophthalmic Partners PC, Cornea Service, Wills Eye Hospital, Philadelphia, PA, USA; <sup>6</sup>Triangle Eye Consultants, Raleigh, NC, USA; <sup>7</sup>Department of Ophthalmology, Tulane University, New Orleans, LA, USA

Correspondence: Thomas Chester, 7001 S. Edgerton Rd, Brecksville, OH, 44141, USA, Tel +1 440 526 1974, Fax +1 440 526 2352, Email [drchester@clevelandeyeclinic.com](mailto:drchester@clevelandeyeclinic.com)

**Abstract:** Dry eye disease (DED) is a common ocular condition, but the diagnosis relative to other ocular conditions and the evaluation of severity of the condition has often been difficult. This challenge can be due to clinical signs and symptoms not always correlating with each other. An understanding of the various components which create the condition, as well as the diagnostic measures used to evaluate these components, is useful to the clinician working with DED patients. This review paper will discuss traditional diagnostic options, diagnostic imaging, and Advanced Point of Care testing capabilities to determine the severity level of dry eye disease more adequately.

**Keywords:** dry eye disease, diagnostic imaging, POC testing, biomarkers

## Introduction

Dry eye disease (DED) and ocular allergy (OA) are the most common inflammatory disorders of the ocular surface.<sup>1</sup> The prevalence of DED globally is 5–50%, and more than 16 million Americans have been diagnosed with DED while others carry the condition.<sup>2</sup> Approximately one-third of the global population is affected by allergic disease and OA symptoms occur in 40% to 80% of the affected individuals.<sup>3</sup> DED and OA often mimic each other and frequently coexist.<sup>4</sup> An irregular correlation between reported symptoms and observed signs is potentially attributed to the challenge of diagnosing and evaluating the severity of DED.<sup>5</sup>

DED is most often diagnosed in adults over 50 years old but can occur at any age. Risk factors include age, female gender, use of contact lenses, refractive surgeries, smoking, poor health, use of antidepressants or oral steroids, systemic drug effects, autoimmune diseases, thyroid disease, and environmental factors.<sup>6</sup> Up to 46% DED and OA patients impact on their quality of life activities of daily living.<sup>7–10</sup> Significant psychological impact has been reported with DED and even a willingness to shorten their lifespan to be free of DED.<sup>11</sup> Similarly, the most prevalent forms of ocular allergy, seasonal allergic conjunctivitis (SAC) and perennial allergic conjunctivitis (PAC), can cause extreme discomfort and have a significant impact on quality of life and often predispose or exacerbate DED.<sup>7</sup>

Symptoms by themselves are often inadequate for the differential diagnosis of ocular surface disease (OSD), because a wide variety of ocular surface and tear film disorders such as DED, OA, blepharitis, meibomian gland dysfunction (MGD) and bacterial or viral infections can produce the same symptoms.<sup>12,13</sup> Signs and symptoms of DED do not often correlate.<sup>14</sup> Only about two-thirds of patients with dry eye symptoms test positive for DED with traditional confirmatory tests.<sup>15,16</sup> Despite some patients having minimal objective signs such as punctate corneal fluorescein staining, they complain of severe ocular irritation, whereas others present with severe staining and have minimal symptomatology.<sup>16–18</sup> The ability to practically examine tear biomarkers reveals that specific biomarkers correlate to the signs and symptoms in all forms of dry eye.<sup>19</sup>

Typical ocular surface-related symptoms include eye irritation, burning, tearing, stinging, a gritty or foreign body sensation, photophobia, fluctuating vision, redness, intermittent sharp pain, and transient blurred vision. Environments that generate desiccating stress including high temperature, low humidity, air conditioning, and activities that lead to reduced blinking such as driving, computer use, reading, and watching television can worsen dry eye symptoms.<sup>20–22</sup> Ocular itching, frequently

referenced as the hallmark OA symptom, is not just an allergic issue, just as tear film instability is not just related to DED.<sup>1,23</sup> DED symptoms have features of neuropathic ocular pain, including spontaneous pain, hyperalgesia and dysesthesias.<sup>24,25</sup> Initial ocular surface damage or inflammation may cause central and/or peripheral sensitization, producing hypersensitivity and persistent DED symptoms despite the source of inflammation and/or damage being removed.<sup>26</sup>

The ocular surface is composed of several components working in tandem to create and maintain a healthy refractive surface on the cornea including the conjunctiva, glandular epithelium, nerves that control the blink reflex, main and accessory lacrimal glands, eyelashes and associated glands, meibomian glands, and the nasolacrimal duct.<sup>27,28</sup> A disruption to any component of the ocular surface can lead to an unstable surface, which in turn, can lead to tear film abnormalities and disrupt the refractive capabilities creating visual disturbances and OSD symptoms.

The normal tear film contains a complex mixture of proteins, electrolytes, mucins, cytokines, lysozymes, immunoglobulins, lactoferrin, and growth factors.<sup>29</sup> The tear film also consists of three layers: the mucin layer, the aqueous layer, and the lipid layer. These layers play an important role in maintaining the ocular surface stability. Normal tear film dynamics include distribution, turnover (and drainage), evaporation, and absorption of the tears.<sup>30</sup> DED is a multifactorial disorder of the tear film caused by decreased tear production and/or increased evaporation that can lead to a loss of homeostasis, tear film instability, inflammation, hyperosmolarity, and neurosensory dysfunction.<sup>6</sup> OA has been shown to be associated with changes in the composition of the tear film that leads to tear film dysfunction.<sup>31</sup> Even though DED and OA are different ocular conditions, there is significant clinical overlap between OA and DED and are often thought of as facilitating, or in the least predisposing, each other, or co-existing conditions.<sup>4,32</sup> It is essential to consider the implications of both DED and OA together in the effective management of OSD.

Changes in the quantity and quality of tears will lead to the development of OSD and these parameters are important for making a correct diagnosis. DED is separated into two primary underlying mechanisms: 1) aqueous-deficient dry eye disease (ADDE) and 2) evaporative dry eye disease (EDE). The aqueous layer accounts for the majority of the tear film thickness and is produced primarily by the lacrimal glands. The lipid layer is made up of meibomian glands, and its function is to stabilize the tear film and protect it from evaporation.<sup>33</sup> ADDE reflects a reduction of lacrimal secretion while EDE has normal lacrimal secretion but is due to excessive fluid loss from the exposed ocular surface, both of which lead to tear film instability and increased tear film osmolarity. However, both ADDE and EDE are often present at the same time and may be associated with nerve abnormalities.<sup>34</sup>

Tears in patients with DED have more proinflammatory markers, do not contain as many protective proteins when compared with patients without DED,<sup>35,36</sup> and due to inadequate production of mucins, both soluble and membrane-bound, lack an even distribution of the tear film.<sup>37</sup> A cycle of tear film instability and ocular surface damage result from desiccant stress and a proinflammatory tear film.<sup>38</sup>

Beyond quality-of-life impairment, undiagnosed and untreated OSD prior to ocular surgery leads to less accurate pre-surgical measurements,<sup>39</sup> mild or asymptomatic DED to become severe or chronic DED and leads to worse visual acuity outcomes.<sup>40</sup> Similarly, OA is a significant risk factor for myopic regression and haze after photorefractive keratectomy.

Having standardized methods for diagnosing DED as well as establishing modalities for monitoring the efficacy of different treatments for DED are necessary.<sup>41</sup> Since OSD is multifactorial, one test has been insufficient to diagnose DED,<sup>42</sup> and a panel or battery of tests have been recommended to improve the effectiveness of diagnosis.<sup>12</sup> Moreover, some patients with no significant clinical signs have considerable discomfort, while those with severe dry eye and vision-threatening ocular complications may suffer from only mild symptoms.<sup>43</sup> An ideal OSD diagnostic test should be rapid (less than 15 minutes), objective, quantitative, specific, reproducible and should directly impact clinical management and/or therapeutic decisions. Testing should be simple to allow implementation by ancillary staff and efficient workflow. Advanced tear-based point-of-care quantitative results support early diagnosis and intervention as well as therapeutic monitoring to confirm adequate disease control or progression. The relationship between DED and OA is well described, and any technology that can rapidly quantify biomarkers like MMP-9, immunoglobulin E and lactoferrin in tears will allow for differentiation between DED and OA and more specific interventions.<sup>44</sup>

## Traditional Diagnostics

Historically, DED was determined by a combination of a history and physical examination of the lid and meibomian morphology and expression, and questionnaires followed by conventional diagnostic tests for initial assessment that includes corneal staining, tear film assessment, conjunctival staining, tear break-up time (TBUT), and the Schirmer test.<sup>45,46</sup>

## Questionnaires

Questionnaires assess and quantify patient's subjective symptoms of DED to create an objective score so that it can be used to better follow patients over time. The most used questionnaires include the Ocular Surface Disease Index (OSDI) and National Eye Institute Visual Function Questionnaire-25 (NEIVFQ-25) due to their multi-dimensionality and the inclusion of patient's quality of life changes.<sup>47-49</sup> Other questionnaires include the Dry Eye Questionnaire 5, McMonnies, and the Standard Patient Evaluation of Eye Dryness (SPEED).<sup>50</sup>

The OSDI is a 12-question assessment developed to grade the symptoms of ocular irritation in DED and vision-related function over the prior week.<sup>51</sup> The OSDI measures symptoms of dry eye symptoms, light sensitivity, grittiness, and pain, but not other common symptoms like foreign body sensation and tearing. The OSDI is reported to have a specificity of (83%) and a sensitivity of (60%) when diagnosing patients with DED.<sup>51</sup> The OSDI was not shown to correlate with Schirmer tear testing<sup>19</sup> (Lam 2013) but correlates with tear break up time (TBUT) in 35% and increases to 42% when the OSDI  $\geq 35$ .<sup>52</sup> OSDI and TBUT show no correlation to any elevation in cytokines or MMP-9.<sup>53</sup>

The 25-item NEIVFQ-25 is a questionnaire derived from a validated 51-question survey. It examines different health-related quality of life aspects including vision, health overall, difficulty with distance and near vision, limitations in social functioning and limitations of role due to vision, dependency on others and mental health symptoms impacted by vision, expectations for vision in the future, difficulties with driving, and discomfort and/or pain around the eyes. This questionnaire may be less sensitive at capturing symptoms associated with milder dry eye disease. Subscales show correlations ranging from 39% to 69%<sup>48</sup> for DED. The NEI-VFQ is only moderately associated with Sjogren's patients when the ocular surface parameters (tear film break-up time, Schirmer score with and without anesthesia, Oxford score, van Bijsterveld score) exist.<sup>49</sup>

The Dry Eye Questionnaire (DEQ) was developed with the intention of diagnosing dry eye disease and quantifying its severity level for the prior week. It quantifies several symptoms using four measurements: level of irritation, frequency of irritation, intensity of the irritation in the morning, and intensity of the irritation late in the day.<sup>54</sup> There is no correlation between DEQ and clinical parameters but instead can differentiate between patient groups such as 1) dry eye vs non-dry eye, 2) Sjogren's syndrome [SS] with keratoconjunctivitis sicca [KCS] vs non-SS-KCS, and 3) control vs SS-KCS and non-SS-KCS.<sup>55</sup> The DEQ shows a correlation coefficient of  $r = 0.76$ .<sup>56</sup>

Standard Patient Evaluation of Eye Dryness (SPEED) survey is a validated DED symptom survey based on eight questions that assess both the frequency and severity of symptoms over a 3-month period. The patient grades the frequency of their symptoms as well as the severity. The severity is graded on a scale of zero to four with zero being no symptoms and four being intolerable symptoms. The SPEED questionnaire can differentiate symptomatic from asymptomatic patients and is associated with a sensitivity of 90% and specificity of 80%.<sup>50</sup>

## Physical Examination

A slit-lamp examination is very informative in the overall assessment of OSD. It is important to look for abnormalities of the lids, conjunctiva, cornea, and tear film. Misalignment of the lids may lead to trichiasis or corneal exposure and can initiate or exacerbate tear film dysfunction.<sup>57,58</sup> Meibomian gland dysfunction (MGD) and blepharitis are common ocular disorders that are frequently associated with DED and with OA, especially with chronic and severe forms (eg AKC).<sup>59</sup> Moreover, DED impacts corneal sensitivity which may be elevated during the early stages and reduced in chronically progressed stages of the disease.<sup>60</sup> Decreased corneal sensation can potentially disrupt the corneal innervation, which can disrupt the homeostasis of the ocular surface, potentially leading to epithelial disruption and cell loss.<sup>61</sup>

## Tear Break-Up Time

The quality of meibum may result in tear film instability, creating a desiccating stress on the ocular surface.<sup>62,63</sup> Tear break-up time (TBUT) is an assessment of tear film instability. TBUT is measured after introducing fluorescein dye

instillation into the eye. Once fluorescein is instilled into the eye and under cobalt blue light, the patient is asked to stare without blinking and the cornea/tear film is observed. The time from a blink, to the time a portion of the cornea becomes exposed from under fluorescein-stained tears is measured. A TBUT >8–10 seconds is generally considered normal.<sup>64</sup> Faster TBUT values less than 5–10 seconds are observed in patients with tear instability and indicate mild-to-moderate dry eye disease.<sup>57</sup> TBUT shows a sensitivity of 75% and specificity of 60%.<sup>65</sup>

TBUT is not accurate or reproducible if performed once, but if the average score of 2–3 separate TBUT measurements are used, it increases its utility.<sup>15,28</sup> TBUT measurement does not provide direct information on tear evaporation and the amount of fluorescein that is instilled can potentially alter the TBUT measurements.<sup>66</sup> Patients with MGD frequently present with a rapid TBUT (<10 sec); however, tear film instability and a rapid TBUT can also be observed in patients with mucin deficiency and/or goblet cell loss.<sup>67</sup> The pattern of tear break-up may suggest where the abnormality lies such as surface epithelial layer/membrane-associated mucins or aqueous tear layer/secretory mucins.<sup>68</sup>

## Schirmer Tear Test

The Schirmer tear test (STT) is used to quantitatively measure tears produced by the lacrimal gland during fixed time period in patients suspected of having DED.<sup>10</sup> The STT I test measures total tear production, including both the basal and reflex tears.<sup>69</sup> The test is performed by inserting Schirmer tear strips into the lower conjunctival sac without anesthesia.<sup>70</sup> The Schirmer tear test II test (performed with topical anesthesia) measures tear production of lacrimal gland by stimulating the lacrimal reflex for 5 minutes with a wetting result of <10mm considered abnormal.<sup>70</sup>

The amount of tear production correlates with the length of wetting of the strips with values between 10 and 33 mm being normal.<sup>71</sup> Patients identified with aqueous deficiency have a reduced STT score, and scores of less than 5 mm frequently are seen in moderate-to-severe aqueous deficiency.<sup>12</sup>

STT I test demonstrates a sensitivity 42% and specificity 76% for identifying DED in Sjogren's patients.<sup>72</sup> Several factors that may reduce the accuracy of STT results include manipulation of the lid causing reflex tearing,<sup>66</sup> nasal versus temporal lid position during the STT, disruptive tear drainage, and inferior gaze. All these confounders produce a falsely higher result.<sup>73</sup> Finally, a study that compared STT I and STT II determined that results using anesthesia was found to be more reliable and objective in diagnosing DED.<sup>74</sup> In general, in more advanced cases of DED STT is more reproducible.<sup>15</sup>

Similar to the STT, a phenol red thread test is used to measure tear volume by inserting a thread that has the dye implanted with pH sensitive phenol into the lower eyelid. The thread color changes from red to yellow once in contact with tears measured after 15 seconds. Studies showed that phenol red thread tests were equally sensitive in detecting dry eye as Schirmer testing.<sup>75</sup>

## Tear Clearance

The fluorescein clearance test analyzes reflex and basal tears and utilizes Schirmer paper, proparacaine and 5  $\mu$ L of 0.25% fluorescein with 0.4% benoxinate hydrochloride. The disappearance of the dye and wetting of the strip are both measured in 10-minute increments. If both a value of 3 mm or greater dye is seen in the first 10-minute time-frame and if the dye is not detected after 20 minutes, the tear clearance is normal.<sup>76</sup> The clearance test indirectly measures dry eye induced inflammation and may prove valuable in identifying patient's response to anti-inflammatory therapy.<sup>77</sup>

## Tear Function Index

Like the STT, the tear function index also evaluates tear production. A 10  $\mu$ L drop of fluorescein is instilled and after 5 minutes, Schirmer tear strip wetted amount is quantified, and the color intensity is compared to a color standard. The rate at which the color of the fluorescein dye fades is the tear clearance rate, while the tear function index is a calculation achieved by dividing the value of STT1 by the rate of tear clearance. A tear function index score less than 96 is consistent with DED and less than 34 suggests Sjogren's syndrome.<sup>78</sup> Compared to the tear clearance alone or STT1, the function index improves sensitivity and specificity in diagnosing DED.<sup>78</sup> The tear function index is subjective, time-consuming and intraday variations in tear film dynamics limit the clinical value of this test.<sup>79</sup> A higher tear function index numerical value indicates a healthier the ocular surface, whereas values below 96 suggest DED.<sup>68</sup>

## Tear Meniscus

The tear meniscus, also known as the tear lake, is the height of tears at the intersection of the lower eyelid margin and the bulbar conjunctiva used to determine whether or not DED is present.<sup>80</sup> A normal tear meniscus height is generally 0.2–0.5 mm while a patient with aqueous-deficient dry eye usually will present with a height less than 0.2 mm as measured by OCT.<sup>79</sup>

## Staining

Vital dyes such as Lissamine Green (LG), Rose Bengal (RB), and fluorescein are used to diagnose ocular surface disease, tear film quality, and dryness severity.<sup>81</sup> Healthy epithelium is protected by tight junctions and does not take up dye. Damage to the cells will cause the dye to absorb and cause staining.<sup>82</sup> Unfortunately, no clear relationship exists between ocular surface damage and DED specific staining and a patient's symptoms.<sup>83</sup>

Vital dyes do not detect early DED or differentiate DED from other conditions causing staining,<sup>84</sup> and therefore, vital dyes lack the power to discriminate between mild-to-moderate cases of dry eye.<sup>85</sup> Studies show poor reliability of inferior corneal staining absence or presence in dry eye patients,<sup>17</sup> and in some patients with moderate-to-severe symptoms, no surface staining was present.<sup>86</sup>

Both RB and LG stain dead, devitalized cells and potentially stain healthy cells. RB can be toxic to the cells on the corneal epithelium.<sup>87</sup> and staining occurs in areas that lack membrane-associated mucins on the cornea or conjunctiva.<sup>88</sup> Thus, LG is preferable to RB for DED evaluations as it is associated with less discomfort and corneal toxicity despite being less sensitive at detecting defects.<sup>87,89</sup> Fluorescein stains degenerating or dead cells seen in epithelial erosions and preferentially stains the cornea more than the conjunctiva.<sup>87</sup> If there is an intact epithelium, any pooled fluorescein on the surface will be lifted by a simple cotton wisp applied to the area revealing an unstained area underneath. Mild cases of DED are detected using RB more than fluorescein stain and the cornea shows less staining than the conjunctiva.<sup>90</sup>

Staining tests have been found to have poor repeatability<sup>15</sup> although the level of staining will typically match the level of severity of aqueous deficiency.<sup>91</sup> While mild cases of DED exhibit normal tear production without fluorescein staining, an increase in inflammatory cell mediators may be present in the tears and serve as more accurate measures of DED than just tear production or staining.

## Meibomian Gland Expression and Secretion

The diagnosis of MGD can be made by several diagnostic and physical examination tests looking for altered morphology features such as gland atrophy, terminal duct obstruction and gland drop out. Physiologic changes can be examined with diagnostic tests that grade the qualitative and quantitative changes in meibum.<sup>92</sup> Gland expression diagnostic tests are useful to evaluate these physiologic changes in meibum, grading meibomian gland expression and meibomian gland secretion. Meibum quality, quantity, and expressibility are generally thought to represent meibomian gland function.<sup>93</sup> Applying digital pressure to the central lower lid provides gland expression which when followed by assessing the ocular surface damage was recommended by the Diagnostic Subcommittee at the International Workshop on Meibomian Gland Dysfunction.<sup>92</sup>

The expressibility of meibum is commonly evaluated by applying the digital pressure to the glands along the entire length of the eyelid.<sup>94–96</sup> Healthy eyelids secrete clear meibum that is easily expressed with gentle pressure. In patients with MGD, the meibum can become cloudy and opaque with a viscosity like toothpaste and difficult to express.<sup>93</sup> The various qualities of meibum, as well as the ease of expression are rated in different grading schemes. These grading schemes can include counting the number and location of expressible glands, as well as the response of the glands to different levels of digitally applied pressure. Grading and scoring expression provides information related directly to meibomian gland condition.<sup>12,97</sup> However, the value of meibomian gland expressibility and meibomian gland duct appearance has not been fully established in diagnosing DED.<sup>93</sup>

## Impression Cytology

Impression cytology is a minimally invasive, rapid, and sensitive method of collecting conjunctival epithelial, goblet, and inflammatory cells<sup>98</sup> that serves as an improved alternative to biopsy of the ocular surface. Ocular surface changes such as goblet cell loss and keratinization are evaluated by examining the collected superficial layers microscopically. It requires



appropriate staining technique and enhanced evaluation microscopically.<sup>98</sup> A recognizable deficiency in goblet cells is the primary characteristic of goblet cell loss. Pathological alterations on the ocular surface can also be used to identify the keratinization of the superficial epithelial layers revealing an unhealthy ocular surface.<sup>8</sup>

## Diagnostic Imaging

Digital imaging can be used to support the differential diagnosis of blepharitis and MGD as well as to evaluate tear break-up time and assess tear film stability using corneal topography, aberrometry, functional visual acuity assessment, interferometry, and confocal microscopy.<sup>99</sup> MGD is characterized by posterior lid margin alterations, including prominent blood vessel development, turbid or thickened meibomian gland secretions, and gland orifice plugging which alter the tear film lipid composition and overall tear quality. Specifically, the Keratograph 5M (OCULUS Optikgeraete GmbH, Wetzlar, Germany) and LipiView (Johnson & Johnson Vision, Jacksonville, FL, USA) interferometer devices and LipiScan Imager (Johnson & Johnson Vision, Jacksonville, FL, USA) allow for imaging of the Meibomian glands.<sup>100</sup> Tear protein and lipid analysis may help distinguish dry eye with or without meibomian gland disease.<sup>19,101</sup>

## Functional Visual Acuity

Functional visual acuity is assessed monocularly with a device that measures visual over a 30-second blink-free period. In dry eye patients, functional visual acuity was significantly lower than with controls.<sup>22</sup> This may serve as an indicator of tear film instability and underlying ocular surface disease.

## Topography

Corneal topography and high-speed videokeratoscopy evaluate indices like surface asymmetry index (SAI), surface regularity index (SRI), and topographic pattern to assess tear film stability and corneal surface regularity. These predict consequent effects on image quality.<sup>102</sup> Alternatively, tear film stability may be assessed automatically using either a grid pattern or Placido disk image by evaluating the distortion change of the reflected ocular surface image.<sup>32,65</sup>

Tear film stability analysis system (TSAS) for the TMS-2N corneal topography instrument (Tomey Technology, Nagoya, Japan) is a sensitive videokeratographer.<sup>103</sup> TSAS uses videokeratography to capture 10 corneal surface topography images at 1-second intervals and analyzes any changes in tear film stability over 10 seconds by quantitatively measuring any areas of corneal irregularity. The more irregularities that are detected, the more severe the underlying tear film instability/DED.

## Interferometry

Interferometry is based on interference between light that is reflected from the surface of the lipid layer and light reflected from the interface between the lipid layer and the aqueous layer of the tear film. Colored images of the superficial lipid layer can be created and evaluated over the entire ocular surface. Interference patterns may be used to assess the thickness and quality of the lipid layer.<sup>104</sup>

The lipid layer thickness, blink rate, and images of the meibomian gland structures can be measured using the LipiView (Johnson & Johnson Vision, Jacksonville, FL, USA).<sup>105</sup> It may be used as a non-invasive tear break-up time assessment to determine the interval between the last blink and the appearance of the first lipid layer discontinuity.<sup>106</sup> Thin lipid layers, those less than 60 nm, are correlated with DED while reduced dry eye symptoms are often associated with thick lipid layers, those more than 75 nm.<sup>107</sup> However, since the lipid thickness layer is only measured at one point in time, this may present as a limitation and may not necessarily correlate with TBUT.<sup>108</sup>

## Meibography

Meibomian glands make the lipid layer of the tear film. Meibography is a method of evaluating the meibomian gland structure and enables objectivity.<sup>92</sup> Meibography can be used to examine meibomian gland changes such as dilation, truncation and dropout.<sup>109,110</sup>

## Wavefront Aberrometry

DED patients show increased optical aberrations relative to normal control eyes because of irregularities to the tear film.<sup>102</sup> Wavefront aberrometry allows for the non-invasive evaluation of higher order aberrations present from tear film instability. Alterations and irregularities in the tear film thickness associated with tear film instability can lead to reduced retinal image quality from higher order aberrations. Moreover, changes in tear volume and quality can create changes in higher order aberrations consistent with DED.<sup>111,112</sup>

## OCT

Anterior segment optical coherence tomography (OCT) is a noninvasive diagnostic test that uses emission and reflection of light waves to create a high-resolution, three-dimension, cross-sectional image of biologic structures.<sup>113</sup> OCT devices may also quantify aspects of the tear film including tear volume, tear meniscus height, tear film thickness and cross-sectional.<sup>114</sup> A reduced tear meniscus measurement correlates with symptoms of DED and the Schirmer test.<sup>115</sup>

## Confocal Microscopy

Confocal microscopy is another technology providing high-resolution evaluation non-invasively. Confocal microscopy can capture images of all layers of the cornea, including sub-basal corneal nerve density, morphology, and the presence of and inflammatory cells, as well as goblet cells and inflammation of the conjunctiva.<sup>116,117</sup> Confocal microscopy requires interpretation by a skilled clinician.

## Advanced Point-of-Care (POC) Testing

### Biomarkers

The tear film plays a critical role in supporting the ocular surface health and serves as a reservoir for potential biomarkers. The most desirable biomarkers are rapidly quantifiable, and their assay results directly influence treatment decisions. Easy access to a patient's tears allows for rapid advanced tear-based point-of-care diagnostics to objectively assess these biomarkers. These tests are typically performed by a technician during the patient workup, and since the results are readily available within 10–15 minutes, test results can be used to manage patients during the same office visit. Early detection of objective biomarkers such as tear osmolarity, lactoferrin, total IgE, and matrix metalloproteinase 9 (MMP-9) may not only lead to faster intervention but can also allow for a more targeted approach to therapeutic interventions and disease monitoring over time.

### Osmolarity

Increased tear osmolarity is thought to be a hallmark sign of DED and does not differentiate aqueous deficiency from increased evaporation.<sup>6</sup> Hyperosmolarity of the tear film can occur through increased evaporation and instability of the tears and/or decreased flow of the aqueous layer produced from the lacrimal gland. Increased osmolarity of the tear film causes the release of inflammatory cytokines, promotes cell destruction, and produces a decrease in goblet cell number.<sup>30,118</sup> Although the gold standard for tear osmolarity is freezing point depression (FPD), it may be confirmed through vapor pressure, electrical conductivity, or impedance.<sup>119</sup>

TearLab (Trukera, San Diego, CA, USA) osmometer measures tear osmolarity using impedance after collection of a 50 nL tear sample in less than 20 seconds (Table 1). This test must be done before instilling ocular anesthesia drops and other diagnostic tests that create reflex tearing may lead to variable results.<sup>120</sup> This test should be performed bilaterally with the highest number recorded as the determinant value. It is accepted that an osmolarity reading of 308 mOsm/L or higher in one eye or a difference in measurements between the two eyes of greater than 8 mOsm/L indicates an abnormal tear osmolarity.<sup>121</sup>

Hyper osmolarity >308 mOsm/L indicates mild DED and readings above 312 mOsm/L indicate moderate-to-severe DED.<sup>121</sup> Despite a higher tear osmolarity being an indicator of DED, the higher reading may not correlate with other clinical signs or with a patient's symptoms since there is often poor correlation between signs and symptoms in DED patients.<sup>122</sup> There is a modest correlation between staining and symptoms, but none was found between tear osmolarity and symptoms or between tear osmolarity and corneal fluorescein staining.<sup>123</sup>

**Table 1** Advanced Point-of-Care Diagnostic Tear Tests

Clinical Condition	Analyte	Result	Sample Size	Time to Results	Variability from Reflex Tearing	Guides Therapeutic Decisions
Dry eye	Osmolarity	Quantitative	50 nanoliters	Less than 30 seconds	Yes	No
Ocular allergy	IgE	Quantitative	1 microliter	Less than 10 minutes	No	Yes
Aqueous deficient dry eye	Lactoferrin	Quantitative	1 microliter	Less than 10 minutes	No	Yes
Inflammatory dry eye	Metalloproteinase-9 (MMP-9)	Qualitative	10 microliters	Less than 15 minutes	No	Yes

Clinical studies have created concern on the utility of tear osmolarity to distinguish healthy controls from DED given the wide variability in tear osmolarity seen in healthy controls and the impact reflex tearing leading to a false lowering of results.<sup>124–128</sup> Tear osmolarity measurements could vary in dry eye patients during the day on average by 21.9 mOsmol/L.<sup>129</sup> A meta-analysis reveals a sensitivity of 67% and specificity 46% for DED diagnosis using a cutoff of 294 mOsmol/L and a sensitivity of 40% and specificity of 100% using a cutoff of 310 mOsmol/L in Sjogren's syndrome patients.<sup>127</sup> Since osmolarity has challenges with clinical reproducibility and does not differentiate between aqueous sufficient dry eye and evaporative dry eye, it cannot be used solely as an indicator of dry eye or to guide clinical management and therapeutic decisions.<sup>130</sup> If other tear-based testing will be performed such as Schirmer testing, lactoferrin, IgE and MMP-9 Inflammatory testing, tear osmolarity must be performed first to reduce the impact of reflex testing.

## Lactoferrin

The lacrimal gland secretions that produce most of the tear film aqueous layer contain proteins including lactoferrin and lysozyme enzymes, immunoglobulins, and electrolytes that are involved in maintaining ocular surface health. The lacrimal gland acinar cells produce lactoferrin.<sup>131</sup> A tear film sample used to measure the proteins in lacrimal secretions, such as lactoferrin, has shown discordant levels in patients with DED. This allows lactoferrin to serve as a novel biomarker for DED.<sup>35,119,132–136</sup> Lysozyme, another lacrimal gland protein often associated with lactoferrin has also been found to be reduced in Sjogren's and/or medicated glaucoma patients. Also, despite some studies showing a lack of differentiation between Sjogren's syndrome, non-Sjogren's DED and non-DED patients, lysozyme may still be useful in monitoring damage from beta-adrenergic receptor blocking medications.<sup>137,138</sup>

Lactoferrin is a glycoprotein created by the acinar cells of the main and accessory lacrimal glands<sup>139</sup> and binds iron, providing both an antibacterial and anti-inflammatory effect on the ocular surface.<sup>140</sup> Lactoferrin is a key predictor of the volume and stability of the tear film. The lactoferrin concentration is reduced in the setting of aqueous-deficient dry eye and is shown to be dramatically reduced in tears of both non-Sjogren's and Sjogren's syndrome patients with DED<sup>35</sup> as well as patients with chronic medication induced DED.<sup>141</sup> Tear volumes from the lacrimal gland are noted to positively correlate with the concentration of lactoferrin. Lactoferrin levels have an inverse correlation with corneal staining scores and OSDI scores.<sup>142</sup> Patients with lower tear volume tend to have reduced lactoferrin concentration.<sup>143,144</sup> Further, a direct correlation between lactoferrin concentration and TBUT time<sup>35</sup> as well as a significant negative correlation between Rose Bengal staining and the level of tear lactoferrin in DED with Sjogren's syndrome as well as DED without a diagnosis of Sjogren's syndrome.<sup>140,143,145</sup> In comparison with traditional diagnostics such as the STT, the vital dyes and the tear film break-up-time, the lactoferrin test is reported to be the most reliable, single marker in the diagnosis of KCS.<sup>42</sup> Lactoferrin levels are normal, and not reduced, in the setting of meibomitis related rosacea<sup>146</sup> and OA.<sup>147</sup> A reduction in lactoferrin has been often reported as an indication of contact lens intolerance.<sup>148</sup> Improvement in lactoferrin levels is seen with dry eye disease treatment.<sup>148</sup>



A quantitative point-of-care test to measure tear film lactoferrin in patients suspected of DED is available (Table 1). Lactoferrin can be detected with the use of the T-POC Lactoferrin (Verséa Ophthalmics, Tampa, FL, USA) tear-based testing platform. A glass capillary tube is used to directly collect 1 $\mu$ L of tears from each eye and transferred to an immunoassay test cassette containing a reagent. The test cassette is then inserted into the T-POC digital lateral flow reader which yields a quantitative numeric result within 6–8 minutes. Lactoferrin is an abundant tear protein and has normal tear concentration of 2.2 mg/mL.<sup>140</sup> Since lactoferrin is produced by the accessory and lacrimal gland, it has been seen in similar concentrations in both basal tears and in reflex tears.<sup>149</sup> A cutoff for lactoferrin below 0.9–1.1 mg/mL provides model accuracy for the diagnosis of DED<sup>150</sup> and confirms the presence of ADDE. Lactoferrin has been proposed for the diagnosis of primary Sjogren's syndrome, since the test had a sensitivity of 72% and specificity of 95% while STT I had a sensitivity of 64% and specificity of 85%.<sup>151</sup> Whereas in patients with non-Sjogren's Syndrome DED, lactoferrin showed a sensitivity 79.4% and specificity of 78.3%.<sup>152</sup> A recent meta-analysis confirms lactoferrin concentrations are significantly decreased in tears of DED patients.<sup>153</sup> Lower levels of lactoferrin also correlate with severity of DED,<sup>132</sup> and DED progression is associated with a reduction of lactoferrin.<sup>154</sup>

Lactoferrin concentrations respond to treatment and allow for therapeutic monitoring. Treatment of DED with punctal occlusion<sup>155</sup> and cyclosporine 0.2%<sup>156</sup> was associated with increased tear lactoferrin levels.

## MMP-9

Matrix metalloproteinase-9 (MMP-9) is part of a multidomain calcium and zinc ion-dependent enzyme family, produced by epithelial cells, and is involved in wound healing, tissue remodeling, and inflammation.<sup>157</sup> Increased MMP-9 disrupts corneal epithelial barrier function, increases corneal epithelial desquamation, and leads to irregularities of the corneal surface.<sup>157</sup> Corneal cell disruption leads to primary ocular symptoms of dry eye including pain, fluctuating vision, and tearing.<sup>157</sup>

Increased levels of tear MMP-9 are common in patients with ocular surface disease. MMP-9 is involved with rebuilding of the extracellular matrix after a wound to the corneal surface. MMP-9s are linked to sterile corneal ulcers, active pterygia, fungal keratitis, ocular allergy, advanced keratoconus with irregular surface, ocular burns, blepharitis, conjunctivochalasis, and DED.<sup>158–160</sup> Tear MMP-9 is considered a biomarker of damage to the ocular surface and inflammation<sup>27</sup> and is a more sensitive marker of ocular surface disease than other clinical signs.<sup>157</sup>

MMP-9 levels have been observed to be elevated with DED<sup>6,27</sup> and other OSD. A disposable single-use, qualitative immunoassay point-of-care test, InflammDry® (Quidel Corp) is made to identify MMP-9 levels in tears of patients suspected of DED (Table 1).<sup>161</sup> The acceptable level of tear MMP-9's in healthy patients is between 3 and 40 ng/mL.<sup>16</sup> In a diseased state, MMP-9's can exceed 40 ng/mL in patients with moderate-to-severe DED.<sup>161</sup> The InflammDry test is performed by dabbing 6–8 times against the lower conjunctiva for a goal of collecting 10 $\mu$ L of tear fluid. The test is activated with a running buffer and results are ready for interpretation after 10–15 minutes.<sup>16,162</sup> Sample volume affects the detection limit. Even if the MMP-9 concentration in the tear film is elevated, if the tear volume is extremely reduced then there is a chance the result will be negative or weak positive. The test is subject to variability because it is qualitative and highly dependent on adequate saturation of the sampling fleece which is influenced by practitioner sampling proficiency. Insufficient tear samples (less than 10  $\mu$ L) may produce false-negative results, whereas excessive tear samples (more than 20  $\mu$ L) may produce false-positive results.<sup>163</sup> InflammDry shows a sensitivity 80–85% and 94–97% specificity in diagnosing dry eye.<sup>16,162</sup>

Inflammatory biomarkers are not unique to DED since many have been identified with conditions such as infection, contact lens use, and ocular allergy.<sup>164</sup>

While inflammation is known to be involved with DED, it is not always recognized in symptomatic patients as even high dose topical steroids or topical cyclosporine only relieve symptoms in 40–80% of the patients.<sup>165–167</sup> Artificial tear eye drops alone failed to demonstrate any reduction in MMP-9 levels.<sup>158</sup> However, confirmation of the presence of clinically significant inflammation can impact clinical decisions. By avoiding punctal occlusion in the presence of high levels of inflammatory mediators and predicting a favorable therapeutic response for immunomodulator or anti-inflammatory treatments such as topical cyclosporine, lifitegrast or steroids treatment failures and unnecessary office visits can be reduced.<sup>168,169</sup>

## IgE

OA is an conjunctival inflammatory response to a variety of allergens.<sup>170</sup> Allergens come into contact with the conjunctiva and produce local IgE, which then assimilates into the tears through the blood tear barrier, resulting in higher levels of IgE in the tears.<sup>171</sup> Many ocular surface diseases that contribute to tear film dysfunction, infections, blepharitis, and mechanical and toxic conjunctivitis may mimic the clinical pictures of OA<sup>172</sup> as well as long-term antihistamine usage that has anticholinergic properties due to their high muscarinic receptor binding.<sup>4</sup> Chronic OA is known to create dry eye, which is classified as either reduced tear production or evaporative.<sup>172</sup> Some authors have shown a correlation of total tear IgE to serum IgE,<sup>173–175</sup> but the primary contributor of seasonal allergic conjunctivitis (SAC) severity is the locally produced IgE.<sup>176</sup> When decrease in TBUT is present without the other signs of DED, it may represent a feature of OA.<sup>177</sup>

Patients with forms of OA that are chronic, such as patients with atopic keratoconjunctivitis (AKC) and vernal conjunctivitis (VKC), have reduced tear film break-up times,<sup>178</sup> reduced conjunctival mucins in the tear fluid layer<sup>179</sup> and reduced density of the conjunctival goblet cells.<sup>177</sup>

OA adversely affects tear film stability, which facilitates the ocular allergic process to persist and recur.<sup>177,180,181</sup> Activation of Eosinophils and simultaneous release of inflammatory mediators could damage the conjunctival epithelium and goblet cells. As a result, the deficiency of the mucin layer creates tear film instability. Significantly thicker lipid layer and lipid layer changes that are typical of DED have been found in SAC patients without corneal fluorescein staining pointing to the instability of the tear film caused by changes to the lipid layer.<sup>147</sup> A point-of-care test available commercially measures the IgE level in tears. Patients with IgE levels at least 80 ng/mL point to an allergic conjunctivitis diagnosis, with the tear film level of IgE correlated.<sup>173</sup> Interferometry shows increased lipid layer thickness relative to healthy patients and is negatively correlated with TBUT.<sup>130</sup> Further, DED may create the OA or initiate it. The decreased tear clearance in DED is somewhat responsible for the reduced clearance of allergens and inflammatory mediators, facilitating an antigen that may be insufficient to cause a reaction systemically but may initiate a more limited allergic reaction locally.<sup>44</sup>

Mucins, secreted by conjunctival epithelium goblet cells, serve several functions: they enable the tear film to be spread evenly over the ocular surface, they assist with tear film stability, and they provide a protective layer to the corneal epithelium.<sup>182</sup> Decreased mucins are found in both DED and OA patients and show correlation with Schirmer test, TBUT, and the density of the goblet cells. It has a negative correlation with corneal staining. Dry eye symptoms are greater with lower levels of mucin.<sup>183,184</sup>

The main laboratory tests for allergic conjunctivitis (AC) include skin prick tests, cytological examination, and IgE antibody detection.<sup>185</sup> The tear samples need to be collected from each eye because of potential asymmetrical conditions preferentially collected with a capillary tube as cells and mediators could bind to the strips or sponges. Eosinophil presence with cytology is highly indicative of OA, whereas their lack of presence does not necessarily exclude it.<sup>186</sup> Skin prick tests are a relatively invasive procedure and use many needle pricks containing small volumes of different allergens. Based on the large ocular surface reaction created with the conjunctival provocation test, there is limited clinical use for this test.<sup>187</sup> The identification of conjunctival eosinophils is a difficult and complex process and yielding a low number (approximately 50%) in the conjunctival scrapings of patients with mild allergic conditions.<sup>188</sup> Serum total IgE tests do not differentiate the pathogenesis and severity of the disease and are not diagnostically recommended.<sup>189</sup> There has also been conflicted data between total IgE in tears and serum. For example, one study was unable to demonstrate a correlation between serum total IgE and tear total IgE levels in patients with SAC, while another study found a high correlation in patients with moderate-to-severe SAC.<sup>174,190</sup>

Most tear IgE is produced through the local create of IgE antibodies in the eye.<sup>175,191,192</sup> In OA, IgE may be quantified in tears of patients with perennial allergic conjunctivitis (PAC), SAC, AKC and VKC.<sup>173,174,189</sup> SAC is dominated by local production and symptoms, compared to serum total IgE levels generally revealing the immune response of the whole body with only a modest correlation to serum IgE.<sup>193</sup> Tear IgE varies from 159 IU/mL to less than 1 IU/mL compared with controls of 8 IU/mL to less than 1 IU/mL.<sup>194</sup> Nomura et al sampled tears from patients with AKC, epidemic keratoconjunctivitis (EKC), bacterial conjunctivitis (BC) and normal controls, and determined that their


tear IgE concentrations were significantly increased in allergic conjunctivitis when compared to healthy patients and there was no significant difference found between EKC, BC and the healthy controls.

Following treatment of OA, the tear concentration of total IgE and symptoms improved.<sup>193</sup> The decrease between the measured amount of total tear IgE concentrations at the initial and the primary follow-up visit showed positive correlation with the total clinical scores recorded during both visits. Likewise, at every 2-week follow-up, there was a significant correlation between the change in the clinical score and total tear IgE concentration. Patients at risk for recurrent disease show total IgE concentration significantly higher than those without recurrent disease. Higher initial presenting tear total IgE concentration was also associated with an improvement delay following treatment and may help to gauge the course and efficacy of treatment by observing changes that occur in the total IgE concentration of the tears.<sup>193</sup>

## Conclusion

DED is a complex multifactorial condition that frequently predisposes or co-exists with OA. Both conditions lead to tear instability and ocular surface inflammation that can lead to chronic symptoms, morbidity, and negatively impact surgical outcomes.

Traditional diagnostics are less reliable than newer advanced tear point-of-care diagnostics that allow improved diagnostic accuracy and therapeutic monitoring (Figures 1A–D). Digital imaging can also serve as a diagnostic adjunct to



**Figure 1** (A and B): Identification and management of aqueous-deficient dry eye. (C and D): Identification and management of evaporative dry eye.

tear-based testing by providing visual images which can be used to support a diagnosis and garner patient acceptance and compliance with treatment plans.

## Disclosure

Dr Tom Chester reports the following relationships: Allergan – consultant, research; Dompe – consultant; Glaukos – consultant, speaker; Novartis – consultant; Oyster Point – consultant; Sightscience- consultant, speaker, research; Sun – consultant; Tarsus – consultant; Versea – consultant.

Dr Sumit (Sam) Garg reports personal fees from Alderya, personal fees from Dompe, personal fees from JJV, personal fees from Novartis, personal fees from Oyster Point, personal fees from Sight Sciences, personal fees from Tarsus, personal fees from Versea, outside the submitted work.

Dr Josh Johnston reports the following relationships: Allergan – consultant, speaker, research; Aldeyra – consultant; Alcon – consultant; Aescula Tech – consultant; Avellino – consultant; Azura – consultant; BioTissue – consultant, speaker; Bruder – consultant; Dompe – consultant; E-Swinn – speaker; Glaukos – consultant, speaker; Horizon Therapeutics – consultant; Johnson & Johnson – spouse works there; LacriSciences – share holder; Sight Sciences – consultant, speaker, research; Sun – consultant, speaker; Tarsus – consultant, researcher; Thea – consultant; Visus – consultant; Quidel – consultant; Zeiss – consultant; Ocuterra – consultant; Oyster Point – speaker, consultant; Orasis – consultant; SeaGen – consultant; Versea Biologics – consultant.

Brandon Ayers MD reports the following relationships: Alcon, Allergan, Bausch and Lomb, CorneaGen, Expert Opinion, WL Gore, Glaukos, Rayner, Sight Sciences, Sun Pharma, Tarsus, Trukera, and Carl Zeiss Meditech.

Dr Preeya Gupta reports the following relationships: Azura – consultant, stock options; Alcon- consultant, Aldeyra – consultant; Allergan – consultant; Expert Opinion – consultant, stock options; HanAll Biopharma – consultant; J&J Vision – consultant; Kala – consultant; New World Medical – consultant; Novartis – consultant; Ocular Science – consultant; Ocular Therapeutix – consultant; Orasis – consultant, stock options; Oyster Point – consultant; Santen – consultant; Sight Sciences – consultant; Spyglass – consultant, stock options; Surface Ophthalmics – consultant, stock options, Chief Medical Advisor; Sun Pharmaceuticals – consultant; Tarsus – consultant, stock options; Tear Lab – consultant; Tear Clear – consultant, stock options; Tissue Tech, Inc – consultant; Visant – stock options, Visionology – consultant, stock options; Zeiss – consultant.

The authors report no other conflicts of interest in this work.

## References

1. Hom MM, Nguyen AL, Bielory L. Allergic conjunctivitis and dry eye syndrome. *Ann Allergy Asthma Immunol.* 2012;108:163–166.
2. Farrand KF, Fridman M, Stillman IO, Schaumberg DA. Prevalence of diagnosed dry eye disease in the United States among adults aged 18 years and older. *Am J Ophthalmol.* 2017;182:90–98. doi:10.1016/j.ajo.2017.06.033
3. Leonardi A, Castegnaro A, Valerio AL, Lazzarini D. Epidemiology of allergic conjunctivitis: clinical appearance and treatment patterns in a population-based study. *Curr Opin Allergy Clin Immunol.* 2015;15:482–488. doi:10.1097/ACI.0000000000000204
4. Leonardi A, Luigi R, Salami M, Salami E. Allergy and dry eye disease. *Ocular Immunol Inflamm.* 2021;29:1168–1176.
5. Clayton JA. Dry eye. *N Engl J Med.* 2018;378(23):2212–2223. doi:10.1056/NEJMra1407936
6. Craig JP, Nichols KK, Akpek EK, et al. TFOS DEWS II definition and classification report. *Ocul Surf.* 2017;15:276–283.
7. Palmares J, Delgado L, Cidade M, Quadrado MJ, Filipe HP. Allergic conjunctivitis: a national cross-sectional study of clinical characteristics and quality of life. *Eur J Ophthalmol.* 2010;20(2):257–264. doi:10.1177/112067211002000201
8. Uchino M, Uchino Y, Dogru M, et al. Dry eye disease and work productivity loss in visual display users: the Osaka study. *Am J Ophthalmol.* 2014;157(2):294–300. doi:10.1016/j.ajo.2013.10.014
9. Grubbs JR, Tolleson-Rinehart S, Huynh K, Davis RM. A review of quality-of-life measures in dry eye questionnaires. *Cornea.* 2014;33(2):215–218. doi:10.1097/ICO.0000000000000038
10. Paulsen AJ, Cruickshanks KJ, Fischer ME, et al. Dry eye in the beaver dam offspring study: prevalence, risk factors, and health-related quality of life. *Am J Ophthalmol.* 2014;157:799–806. doi:10.1016/j.ajo.2013.12.023
11. Schiffman RM, Walt JG, Jacobsen F, et al. Utility assessment among patients with dry eye disease. *Ophthalmology.* 2003;110:1412–1419. doi:10.1016/S0161-6420(03)00462-7
12. Pflugfelder SC, Tseng SC, Sanabria O, et al. Evaluation of subjective assessments and objective diagnostic tests for diagnosing tear-film disorders known to cause ocular irritation. *Cornea.* 1998;17:38–56. doi:10.1097/00003226-199801000-00007
13. Suárez-Cortés T, Merino-Inda N, Benitez-Del-Castillo JM. Tear and ocular surface disease biomarkers: a diagnostic and clinical perspective for ocular allergies and dry eye disease. *Exp Eye Res.* 2022;221:109121. doi:10.1016/j.exer.2022.109121
14. Bartlett JD, Keith MS, Sudharshan L, Snedecor SJ. Associations between signs and symptoms of dry eye disease: a systematic review. *Clin Ophthalmol.* 2015;9:1719–1730. doi:10.2147/OPTH.S89700



15. Nichols KK, Mitchell GL, Zadnik K. The repeatability of clinical measurements of dry eye. *Cornea*. 2004;23(3):272–285. doi:10.1097/00003226-200404000-00010
16. Sambursky R, Davitt WF, Friedberg M, Tauber S. Prospective, multicenter, clinical evaluation of point-of-care matrix metalloproteinase-9 test for confirming dry eye disease. *Cornea*. 2014;33:812–818. doi:10.1097/ICO.0000000000000175
17. Nichols KK, Nichols JJ, Mitchell GL, Mitchell GL. The lack of association between signs and symptoms in patients with dry eye disease. *Cornea*. 2004;23(8):762–770.5. doi:10.1097/01.ico.0000133997.07144.9e
18. Pflugfelder SC, Maskin SL, Anderson B, et al. A randomized, double-masked, placebo-controlled, multicenter comparison of loteprednol etabonate ophthalmic suspension, 0.5%, and placebo for treatment of keratoconjunctivitis sicca in patients with delayed tear clearance. *Am J Ophthalmol*. 2004;138:444–457. doi:10.1016/j.ajo.2004.04.052
19. Lam H, Bleiden L, de Paiva CS, et al. Tear cytokine profiles in dysfunctional tear syndrome. *Am J Ophthalmol*. 2009;47(2):198–205, e1. doi:10.1016/j.ajo.2008.08.032
20. Tsubota K, Nakamori K. Dry eyes and video display terminals. *N Engl J Med*. 1993;328:584. doi:10.1056/NEJM199302253280817
21. Abusharha AA, Pearce EI. The effect of low humidity on the human tear film. *Cornea*. 2013;32(4):429–434. doi:10.1097/ICO.0b013e31826671ab
22. Ishida R, Kojima T, Dogru M, et al. The application of a new continuous functional visual acuity measurement system in dry eye syndromes. *Am J Ophthalmol*. 2005;139:253–258. doi:10.1016/j.ajo.2004.08.075
23. Fauquert JL, Jedrzejczak-Czechowicz M, Rondon C, et al. Conjunctival allergen provocation test: guidelines for daily practice. *Allergy Eur J Allergy Clin Immunol*. 2017;72(1):43–54.
24. Galor A, Levitt RC, Felix ER, et al. Neuropathic ocular pain: an important yet under evaluated feature of dry eye. *Eye*. 2015;29:301–312. doi:10.1038/eye.2014.263
25. Galor A, Zlotcavitch L, Walter SD, et al. Eye symptom severity and persistence are associated with symptoms of neuropathic pain. *Br J Ophthalmol*. 2015;99:665–668. doi:10.1136/bjophthalmol-2014-306057
26. Rosenthal P, Borsook D. The corneal pain system. Part I: the missing piece of the dry eye puzzle. *Ocul Surf*. 2012;10:2–14. doi:10.1016/j.jtos.2012.01.002
27. Bron AJ, de Paiva CS, Chauhan SK, et al. TFOS DEWS II pathophysiology report. *Ocul Surf*. 2017;15:438–510. doi:10.1016/j.jtos.2017.05.011
28. Willcox MDP, Argueso P, Georgiev GA, et al. TFOS DEWS II tear film report. *Ocul Surf*. 2017;15:366–403. doi:10.1016/j.jtos.2017.03.006
29. Tiffany JM. The normal tear film. *Dev Ophthalmol*. 2008;41:1–20.
30. Tomlinson A, Khanal S, Ramaesh K, et al. Tear film osmolarity: determination of a referent for dry eye diagnosis. *Invest Ophthalmol Vis Sci*. 2006;47:4309–4315. doi:10.1167/iovs.05-1504
31. Kari O, Määttä M, Tervahartiala T, et al. Tear fluid concentration of MMP-8 is elevated in non-allergic eosinophilic conjunctivitis and correlates with conjunctival inflammatory cell infiltration. *Graefes Arch Clin Exp Ophthalmol*. 2009;247:681–686. doi:10.1007/s00417-009-1042-5
32. Gumus K, Cavanagh DH. The role of inflammation and anti-inflammation therapies in keratoconjunctivitis sicca. *Clin Ophthalmol*. 2009;3:57–67.
33. Paranjpe V, Tan J, Nguyen J, et al. Clinical signs of meibomian gland dysfunction (MGD) are associated with changes in meibum sphingolipid composition. *Ocul Surf*. 2019;2:318–326. doi:10.1016/j.jtos.2018.12.006
34. Galor A, Moein HR, Lee C, et al. Neuropathic pain and dry eye. *Ocul Surf*. 2018;16:31–44. doi:10.1016/j.jtos.2017.10.001
35. Ohashi Y, Ishida R, Kojima T, et al. Abnormal protein profiles in tears with dry eye syndrome. *Am J Ophthalmol*. 2003;136:291–299. doi:10.1016/S0002-9394(03)00203-4
36. Solomon A, Dursun D, Liu Z, Xie Y, Macri A, Pflugfelder SC. Pro- and anti-inflammatory forms of interleukin-1 in the tear fluid and conjunctiva of patients with dry-eye disease. *Invest Ophthalmol Vis Sci*. 2001;42:2283–2292.
37. Tei M, Spurr-Michaud SJ, Tisdale AS, Gipson IK. Vitamin A deficiency alters the expression of mucin genes by the rat ocular surface epithelium. *Invest Ophthalmol Vis Sci*. 2000;41:82–88.
38. Baudouin C, Messmer EM, Aragona P, et al. Revisiting the vicious circle of dry eye disease: a focus on the pathophysiology of meibomian gland dysfunction. *Br J Ophthalmol*. 2016;100:300–306. doi:10.1136/bjophthalmol-2015-307415
39. Trattler WB, Majmudar PA, Donnenfeld ED, McDonald MB, Stonecipher KG, Goldberg D. The Prospective Health Assessment of Cataract Patients' Ocular Surface (PHACO) study: the effect of dry eye. *Clin Ophthalmol*. 2017;11:1423–1430. doi:10.2147/OPHT.S120159
40. Ambrósio R Jr, Tervo T, Wilson SE. LASIK-associated dry eye and neurotrophic epitheliopathy: pathophysiology and strategies for prevention and treatment. *J Refract Surg*. 2008;24(2):396–407. doi:10.3928/1081597X-20080401-14
41. Pinto-Fraga J, Enriquez-de-salamanca A, Calonge M, et al. Severity, therapeutic, and activity tear biomarkers in dry eye disease: an analysis from a Phase III clinical trial. *Ocul Surf*. 2018;16:368–376. doi:10.1016/j.jtos.2018.05.001
42. Goren MB, Goren SB. Diagnostic tests in patients with symptoms of keratoconjunctivitis sicca. *Am J Ophthalmol*. 1988;106:570–574. doi:10.1016/0002-9394(88)90587-9
43. Sullivan BD, Crews LA, Messmer EM, et al. Correlations between commonly used objective signs and symptoms for the diagnosis of dry eye disease: clinical implications. *Acta Ophthalmol*. 2014;9:161–166. doi:10.1111/aos.12012
44. Fujishima H, Toda I, Shimazaki J, Tsubota K. Allergic conjunctivitis and dry eye. *Br J Ophthalmol*. 1996;80(11):994–997. doi:10.1136/bjo.80.11.994
45. Serin D, Karsloglu S, Kyan A, Alagoz G. A simple approach to the repeatability of the Schirmer test without anesthesia: eyes open or closed? *Cornea*. 2007;26:903–906. doi:10.1097/ICO.0b013e3180950083
46. Yamaguchi T. Inflammatory response in dry eye. *Invest Ophthalmol Vis Sci*. 2018;59:192–199. doi:10.1167/iovs.17-23651
47. Nichols KK, Mitchell GL, Zadnik K. Performance and Repeatability of the NEI-VFQ-25 in patients with dry eye. *Cornea*. 2002;21(6):578–583. doi:10.1097/00003226-200208000-00009
48. Mangione CM. Development of the 25-list-item national eye institute visual function questionnaire. *Arch Ophthalmol*. 2001;119(7):1050–1058. doi:10.1001/archophth.119.7.1050
49. Vitale S, Goodman LA, Reed GF, Smith JA. Comparison of the NEI-VFQ and OSDI questionnaires in patients with Sjogren's syndrome-related dry eye. *Health Qual Life Outcomes*. 2004;2:44. doi:10.1186/1477-7525-2-44



50. Ngo W, Situ P, Keir N, et al. Psychometric properties and validation of the standard patient evaluation of eye dryness questionnaire. *Cornea*. 2013;32(9):1204–1210. doi:10.1097/ICO.0b013e318294b0c0
51. Schiffman RM. Reliability and validity of the ocular surface disease index. *Arch Ophthalmol*. 2000;118:615–621. doi:10.1001/archophth.118.5.615
52. Unlü C, Güney E, Akçay Bİ, et al. Comparison of ocular-surface disease index questionnaire, tear film break-up time, and Schirmer tests for the evaluation of the tear film in computer users with and without dry-eye symptomatology. *Clin Ophthalmol*. 2012;6:1303–1306. doi:10.2147/OPHT.S33588
53. Van Der Meid KR, Su SP, Ward KW, Zhang JG. Correlation of tear inflammatory cytokines and matrix metalloproteinases with four dry eye diagnostic tests. *Invest Ophthalmol Vis Sci*. 2012;53:1512–1518. doi:10.1167/iovs.11-7627
54. Chalmers RL, Begley CG, Caffery B. Validation of the 5-Item Dry Eye Questionnaire (DEQ-5): discrimination across self-assessed severity and aqueous tear deficient dry eye diagnoses. *Cont Lens Anterior Eye*. 2010;33(2):55–60. doi:10.1016/j.clae.2009.12.010
55. Begley CG, Caffery B, Chalmers RL, et al. Use of the dry eye questionnaire to measure symptoms of ocular irritation in patients with aqueous tear deficient dry eye. *Cornea*. 2002;21:664–670. doi:10.1097/00003226-200210000-00007
56. Simpson TL, Situ P, Jones LW, Fonn D. Dry eye symptoms assessed by four questionnaires. *Optom Vis Sci*. 2008;85:E692–E699. doi:10.1097/OPX.0b013e318181ae36
57. Abelson MB, Ousler GW, Nally LA, et al. Alternative reference values for tear film breakup time in normal and dry eye populations. *Adv Exp Med Biol*. 2002;506:1121–1125.
58. Tsubota K, Nakamori K. Effects of ocular surface area and blink rate on tear dynamics. *Arch Ophthalmol*. 1995;113:155–158. doi:10.1001/archophth.1995.01100020037025
59. Schaumberg DA, Nichols JJ, Papas EB, Tong L, Uchino M, Nichols KK. The international workshop on meibomian gland dysfunction: report of the subcommittee on the epidemiology of, and associated risk factors for, MGD. *Investig Ophthalmol Vis Sci*. 2011;52:1994–2005. doi:10.1167/iovs.10-6997e
60. Bourcier T, Acosta MC, Borderie V, et al. Decreased corneal sensitivity in patients with dry eye. *Invest Ophthalmol Vis Sci*. 2005;46:2341–2345.8. doi:10.1167/iovs.04-1426
61. Belmonte C, Nichols JJ, Cox SM, et al. TFOS DEWS II pain and sensation report. *Ocul Surf*. 2017;15:404–437. doi:10.1016/j.jtos.2017.05.002
62. Shine WE, McCulley JP. The role of cholesterol in chronic blepharitis. *Invest Ophthalmol Vis Sci*. 1991;32:2272–2280.
63. Shine WE, Silvany R, McCulley JP. Relation of cholesterol-stimulated Staphylococcus aureus growth to chronic blepharitis. *Invest Ophthalmol Vis Sci*. 1993;34:2291–2296.
64. Cox SM, Nichols KK, Nichols JJ. Agreement between automated and traditional measures of tear film breakup. *Optom Vis Sci*. 2015;92:e257–e63. doi:10.1097/OPX.0000000000000648
65. Goto T, Zheng X, Okamoto S, Ohashi Y. Tear film stability analysis system: introducing a new application for video keratography. *Cornea*. 2004;23:S65e70. doi:10.1097/01.ico.0000136685.88489.70
66. Cedarstaff TH, Tomlinson A. Human tear volume, quality, and evaporation: a comparison of Schirmer, tear break-up time and resistance hygrometry techniques. *Ophthalmic Physiol Opt*. 1983;3:239–245. doi:10.1111/j.1475-1313.1983.tb00609.x
67. Pflugfelder SC, Tseng SC, Yoshino K, et al. Correlation of goblet cell density and mucosal epithelial membrane mucin expression with Rose Bengal staining in patients with ocular irritation. *Ophthalmology*. 1997;104:223–235. doi:10.1016/S0161-6420(97)30330-3
68. Yokoi N, Georgiev GA, Kato H, et al. Classification of fluorescein breakup patterns: a novel method of differential diagnosis for dry eye. *Am J Ophthalmol*. 2017;180:72–85. doi:10.1016/j.ajo.2017.05.022
69. Javadi MA, Feizi S. Dry eye syndrome. *J Ophthalmic Vis Res*. 2011;6(3):192–198.
70. Tsubota K, Kaido M, Yagi Y, Fujihara T, Shimmura S. Diseases associated with ocular surface abnormalities: the importance of reflex tearing. *Br J Ophthalmol*. 1999;83(1):89–91. doi:10.1136/bjo.83.1.89
71. Jordan A, Baum J. Basic tear flow. Does it exist? *Ophthalmology*. 1980;87:920–930. doi:10.1016/S0161-6420(80)35143-9
72. Versura P, Frigato M, Cellini M, et al. Diagnostic performance of tear function tests in Sjogren's syndrome patients. *Eye*. 2007;21:229–237.
73. Bitton E, Wittich W. Influence of eye position on the Schirmer tear test. *Cont Lens Anterior Eye*. 2014;37(4):257–261.
74. Li N, Deng XG, He MF. Comparison of the Schirmer I test with and without topical anesthesia for diagnosing dry eye. *Int J Ophthalmol*. 2012;5:478–481. doi:10.3980/j.issn.2222-3959.2012.04.14
75. Vashisht S, Singh S. Evaluation of phenol red thread test versus Schirmer test in dry eyes: a comparative study. *Int J Appl Basic Med Res*. 2011;1(1):40–42. doi:10.4103/2229-516X.81979
76. Prabhasawat P, Tseng SC. Frequent association of delayed tear clearance in ocular irritation. *Br J Ophthalmol*. 1998;82:666–675. doi:10.1136/bjo.82.6.666
77. de Paive CS, Pflugfelder SC. Tear clearance implication for ocular surface health. *Exp Eye Res*. 2004;78(3):395–397. doi:10.1016/j.exer.2003.08.003
78. Xu KP, Yagi Y, Toda I, Tsubota K. Tear function index: a new measure of dry eye. *Arch Ophthalmol*. 1995;113:84–88. doi:10.1001/archophth.1995.01100010086025
79. Savini G, Barboni P, Zanini M. Tear meniscus evaluation by optical coherence tomography. *Ophthalmic Surg Lasers Imaging*. 2006;37(37):112–118. doi:10.3928/1542-8877-20060301-06
80. Mainstone JC, Bruce AS, Golding TR. Tear meniscus measurement in the diagnosis of dry eye. *Curr Eye Res*. 1996;15(6):653–661. doi:10.3109/02713689609008906
81. Bron AJ, Evans VE, Smith JA. Grading of corneal and conjunctival staining in the context of other dry eye tests. *Cornea*. 2003;22:640–650. doi:10.1097/00003226-200310000-00008
82. Bron AJ, Argueso P, Irkeç M, Bright FV. Clinical staining of the ocular surface: mechanisms and interpretations. *Prog Retin Eye Res*. 2015;44:36–61. doi:10.1016/j.preteyeres.2014.10.001
83. Khan-Lim D, Berry M. Still confused about Rose Bengal? *Curr Eye Res*. 2004;29:311–317. doi:10.1080/02713680490516864
84. Caffery BE, Josephson JE. Corneal staining after sequential instillations of fluorescein over 30 days. *Optom Vis Sci*. 1991;68:467–469. doi:10.1097/00006324-199106000-00011

85. Sullivan BD, Whitmer D, Nichols KK, et al. An objective approach to dry eye disease severity. *Invest Ophthalmol Vis Sci.* 2010;51:6125–6130. doi:10.1167/iovs.10-5390
86. Begley CG, Chalmers RL, Abetz L, et al. The relationship between habitual patient-reported symptoms and clinical signs among patients with dry eye of varying severity. *Invest Ophthalmol Vis Sci.* 2003;44(11):4753–4761. doi:10.1167/iovs.03-0270
87. Kim J, Foulks GN. Evaluation of the effect of lissamine green and Rose Bengal on human corneal epithelial cells. *Cornea.* 1999;18(3):328–332. doi:10.1097/00003226-199905000-00015
88. Feenstra RP, Tseng SC. What is actually stained by Rose Bengal? *Arch Ophthalmol.* 1992;110:984–993. doi:10.1001/archophth.1992.01080190090035
89. Machado LM, Castro RS, Fontes BM. Staining patterns in dry eye syndrome: Rose Bengal versus lissamine green. *Cornea.* 2009;28(7):732–734. doi:10.1097/ICO.0b013e3181930c03
90. Watanabe S, Tanaka M. Rose Bengal staining and expression of mucin-like glycoprotein in cornea epithelium. *Invest Ophthalmol Vis Sci.* 1996;37:S357.
91. Savini G, Prabhawat P, Kojima T, et al. The challenge of dry eye diagnosis. *Clin Ophthalmol.* 2008;2:31–55. doi:10.2147/OPHT.S1496
92. Tomlinson A, Bron AJ, Korb DR, et al. The international workshop on meibomian gland dysfunction: report of the diagnosis subcommittee. *Invest Ophthalmol Vis Sci.* 2011;52:2006–2049. doi:10.1167/iovs.10-6997f
93. Wolffsohn JS, Arita R, Chalmers R, et al. TFOS DEWS II diagnostic methodology report. *Ocul Surf.* 2017;15(3):539–574. doi:10.1016/j.jtos.2017.05.001
94. Mathers WD, Shields WJ, Sachdev MS, Petroll WM, Jester JV. Meibomian gland dysfunction in chronic blepharitis. *Cornea.* 1991;10(4):277–285. doi:10.1097/00003226-199107000-00001
95. Bron AJ, Benjamin L, Snibson GR. Meibomian gland disease. Classification and grading of lid changes. *Eye.* 1991;5(Pt 4):395–411. doi:10.1038/eye.1991.65
96. Mathers WD, Shields WJ, Sachdev MS, Petroll WM, Jester JV. Meibomian gland morphology and tear osmolarity: changes with accutane therapy. *Cornea.* 1991;10(4):286–290. doi:10.1097/00003226-199107000-00002
97. Blackie CA, Korb DR. The diurnal secretory characteristics of individual meibomian glands. *Cornea.* 2010;29(1):34–38. doi:10.1097/ICO.0b013e3181ac9fd0
98. Calonge M, Diebold Y, Sáez V, et al. Impression cytology of the ocular surface: a review. *Exp Eye Res.* 2004;78:457–472. doi:10.1016/j.exer.2003.09.009
99. Smith J, Nichols KK, Baldwin EK. Current patterns in the use of diagnostic tests in dry eye evaluation. *Cornea.* 2008;27:656–662. doi:10.1097/01.ico.0000611384.81547.8d
100. Zang S, Cui Y, Cui Y, Fei W. Meibomian gland dropout in Sjogren's syndrome and non-Sjogren's dry eye patients. *Eye.* 2018;32:1681–1687. doi:10.1038/s41433-018-0149-5
101. Enriquez-de-salamanca A, Castellanos E, Stern ME, et al. Tear cytokine and chemokine analysis and clinical correlations in evaporative-type dry eye disease. *Mol Vis.* 2010;16:862–873.
102. Montes-Mico R, Caliz A, Alio JL. Wavefront analysis of higher order aberrations in dry eye patients. *J Refract Surg.* 2004;20(3):243–247. doi:10.3928/1081-597X-20040501-08
103. Goto T, Zheng X, Klyce SD, et al. A new method for tear film stability analysis using videokeratography. *Am J Ophthalmol.* 2003;135(5):607–612. doi:10.1016/S0002-9394(02)02221-3
104. King-Smith PE, Kimball SH, Nichols JJ. Tear film interferometry and corneal surface roughness. *Invest Ophthalmol Vis Sci.* 2014;55:2614–2618. doi:10.1167/iovs.14-14076
105. Doane MG. An instrument for in vivo tear film interferometry. *Optom Vis Sci.* 1989;66(6):383–388. doi:10.1097/00006324-198906000-00008
106. Finis D, Pischel N, Schrader S, Geerling G. Evaluation of lipid layer thickness measurement of the tear film as a diagnostic tool for meibomian gland dysfunction. *Cornea.* 2013;32:1549–1553. doi:10.1097/ICO.0b013e3182a7f3e1
107. Blackie CA, Solomon JD, Scaffidi RC, Greiner JV, Lemp MA, Korb DR. The relationship between dry eye symptoms and lipid layer thickness. *Cornea.* 2009;28(7):789–794. doi:10.1097/ICO.0b013e318191b870
108. Jung JW, Park SY, Kim JS, Kim EK, Seo KY, Kim TI. Analysis of factors associated with the tear film lipid layer thickness in normal eyes and patients with dry eye syndrome. *Invest Ophthalmol Vis Sci.* 2016;57:4076–4083. doi:10.1167/iovs.16-19251
109. Jackson MA. A systematic approach to dry eye using lipiflow treatment. *US Ophthalmic Rev.* 2014;7:104–108. doi:10.17925/USOR.2014.07.02.104
110. Ngo W, Srinivasan S, Schulze M, Jones L. Repeatability of grading meibomian gland dropout using two infrared systems. *Optom Vis Sci.* 2014;91:658–667. doi:10.1097/OPX.0000000000000279
111. Koh S, Maeda N, Hirohara Y, et al. Serial measurements of higher-order aberrations after blinking in normal subjects. *Invest Ophthalmol Vis Sci.* 2006;47:3318. doi:10.1167/iovs.06-0018
112. Lin YY, Carrel H, Wang IJ, et al. Effect of tear film break-up on higher order aberrations of the anterior cornea in normal, dry, and post-LASIK eyes. *J Refract Surg.* 2005;21(5):S525–9. doi:10.3928/1081-597X-20050901-21
113. Liang Q, Pan Z, Zhou M, et al. Evaluation of optical coherence tomography meibography in patients with obstructive meibomian gland dysfunction. *Cornea.* 2015;34:1193–1199. doi:10.1097/ICO.0000000000000563
114. Tung CI, Perin AF, Gumus K, Pflugfelder SC. Tear meniscus dimensions in tear dysfunction and their correlation with clinical parameters. *Am J Ophthalmol.* 2014;157:301.e1. doi:10.1016/j.ajo.2013.09.024
115. Nguyen P, Huang D, Li Y, et al. Correlation between optical coherence tomography-derived assessments of lower tear meniscus parameters and clinical features of dry eye disease. *Cornea.* 2012;31(6):680–685. doi:10.1097/ICO.0b013e3182261577
116. Chan TCY, Wan KH, Shih KC, Jhanji V. Advances in dry eye imaging: the present and beyond. *Br J Ophthalmol.* 2018;102(3):295–301. doi:10.1136/bjophthalmol-2017-310759
117. Villani E, Baudouin C, Efron N, et al. In vivo confocal microscopy of the ocular surface: from bench to bedside. *Curr Eye Res.* 2014;39(39):213–231. doi:10.3109/02713683.2013.842592
118. Bron AJ, Tomlinson A, Foulks GN, et al. Rethinking dry eye disease: a perspective on clinical implications. *Ocul Surf.* 2014;12:S1–S31. doi:10.1016/j.jtos.2014.02.002

119. Versura P, Nanni P, Bavelloni A, et al. Tear proteomics in evaporative dry eye disease. *Eye*. 2010;24:1396–1402. doi:10.1038/eye.2010.7
120. Mathews PM, Karakus S, Agrawal D, Hindman HB, Ramulu PY, Akpek EK. Tear osmolarity and correlation with ocular surface parameters in patients with dry eye. *Cornea*. 2017;36:1352–1357. doi:10.1097/ICO.0000000000001364
121. Lemp MA, Bron AJ, Baudouin C, et al. Tear osmolarity in the diagnosis and management of dry eye disease. *Am J Ophthalmol*. 2011;151:792–8 e1. doi:10.1016/j.ajo.2010.10.032
122. Keech A, Senchyna M, Jones L. Impact of time between collection and collection method on human tear fluid osmolarity. *Curr Eye Res*. 2013;38(38):428–436. doi:10.3109/02713683.2013.763987
123. Amparo F, Jin Y, Hamrah P, Schaumberg DA, Dana R. What is the value of incorporating tear osmolarity measurement in assessing patient response to therapy in dry eye disease? *Am J Ophthalmol*. 2014;157(1):69–77.e2. doi:10.1016/j.ajo.2013.07.019
124. Szalai E, Berta A, Szekanez Z, Szűcs G, L M Jr. Evaluation of tear osmolarity in non-Sjögren and Sjögren syndrome dry eye patients with the TearLab system. *Cornea*. 2012;31:867–871. doi:10.1097/ICO.0b013e3182532047
125. Tashbayev B, Utheim TP, Utheim ØA, et al. Utility of tear osmolarity measurement in diagnosis of dry eye disease. *Sci Rep*. 2020;10:5542. doi:10.1038/s41598-020-62583-x
126. Bunya VY, Fuerst NM, Pistilli M, et al. Variability of tear osmolarity in patients with dry eye. *JAMA Ophthalmol*. 2015;133(6):662–667. doi:10.1001/jamaophthalmol.2015.0429
127. Baenninger PB, Voegeli S, Bachmann LM, et al. Variability of tear osmolarity measurements with a point-of-care system in healthy subjects-systematic review. *Cornea*. 2018;37(7):938–945. doi:10.1097/ICO.0000000000001562
128. Messmer EM, Bulgen M, Kampik A. Hyperosmolarity of the tear film in dry eye syndrome. *Dev Ophthalmol*. 2010;45:129–138.
129. Li M, Du C, Zhu D, Shen M, Cui L, Wang J. Daytime variations of tear osmolarity and tear meniscus volume. *Eye Contact Lens*. 2012;38(5):282–287. doi:10.1097/ICL.0b013e31825fed57
130. Suzuki S, Goto E, Dogru M, et al. Tear film lipid layer alterations in allergic conjunctivitis. *Cornea*. 2006;25:277–280. doi:10.1097/01.ico.0000178277.29010.bc
131. Sánchez L, Calvo M, Brock JH. Biological role of lactoferrin. *Arch Dis Child*. 1992;67(5):657–661. doi:10.1136/adc.67.5.657
132. Zhou L, Beuerman RW, Chan CM, et al. Identification of tear fluid biomarkers in dry eye syndrome using iTRAQ quantitative proteomics. *J Proteome Res*. 2009;8(8):4889–4905. doi:10.1021/pr900686s
133. Narayanan S, Redfern RL, Miller WL, Nichols KK, McDermott AM. Dry eye disease and microbial keratitis: is there a connection? *Ocul Surf*. 2013;11:75–92. doi:10.1016/j.jtos.2012.12.002
134. Seal DV, McGill JI, Mackie IA, Liakos GM, Jacobs P, Goulding NJ. Bacteriology and tear protein profiles of the dry eye. *Br J Ophthalmol*. 1986;70:122–125. doi:10.1136/bjo.70.2.122
135. Boukes RJ, Boonstra A, Breebaart AC, et al. Analysis of human tear protein profiles using high performance liquid chromatography (HPLC). *Doc Ophthalmol*. 1987;67:105–113. doi:10.1007/BF00142704
136. Grus FH, Podust VN, Bruns K, et al. SELDI-TOF-MS protein chip array profiling of tears from patients with dry eye. *Investig Ophthalmol Vis Sci*. 2005;46:863. doi:10.1167/iovs.04-0448
137. D'Souza S, Tong L. Practical issues concerning tear protein assays in dry eye. *Eye Vis*. 2014;1:6. doi:10.1186/s40662-014-0006-y
138. Mackie IA, Seal DV. Confirmatory tests for the dry eye of Sjogren's syndrome. *Scand J Rheumatol Suppl*. 1986;61:220–223.
139. Gillette TE, Allansmith MR. Lactoferrin in human ocular tissues. *Am J Ophthalmol*. 1980;90:30–37. doi:10.1016/S0002-9394(14)75074-3
140. Kijlstra A, Jeurissen SH, Koning KM. Lactoferrin levels in normal human tears. *Br J Ophthalmol*. 1983;67(3):199–202. doi:10.1136/bjo.67.3.199
141. Ono M, Fujita H, Goto E, Naka H, Walcott B, Tsubota K. Decreased tear lactoferrin secretion by cellular dysfunction. *Invest Ophthalmol Vis Sci*. 1998;39:S1100.
142. Sonobe H, Ogawa Y, Yamada K, et al. A novel and innovative paper-based analytical device for assessing tear lactoferrin of dry eye patients. *Ocul Surf*. 2019;17:160–166. doi:10.1016/j.jtos.2018.11.001
143. Danjo Y, Lee M, Horimoto K, et al. Ocular surface damage and tear lactoferrin in dry eye syndrome. *Acta Ophthalmol*. 1994;72:433–437. doi:10.1111/j.1755-3768.1994.tb02791.x
144. Glasson MJ, Stapleton F, Keay L, et al. Differences in clinical parameters and tear film of tolerant and intolerant contact lens wearers. *Investig Ophthalmol Vis Sci*. 2003;44:5116–5124. doi:10.1167/iovs.03-0685
145. Yamada K, Takaki S, Komuro N, Suzuki K, Citterio D. An antibody-free microfluidic paper-based analytical device for the determination of tear fluid lactoferrin by fluorescence sensitization of Tb<sup>3+</sup>. *Analyst*. 2014;139:1637–1643. doi:10.1039/c3an01926h
146. Kiratli H, Irkeç M, Orhan M. Tear lactoferrin levels in chronic meibomitis associated with acne rosacea. *Eur J Ophthalmol*. 2000;10:11–14.
147. Li K, Liu X, Chen Z, Huang Q, Wu K. Quantification of tear proteins and sPLA2-IIa alteration in patients with allergic conjunctivitis. *Mol Vis*. 2010;16:2084–2091.
148. Chiva A. Electrophoresis of tear proteins as a new diagnostic tool for two high risk groups for dry eye: computer users and contact lens wearers. *J Med Life*. 2011;154:228–233.
149. Fullard RJ, Snyder C. Protein levels in nonstimulated and stimulated tears of normal human subjects. *Invest Ophthalmol Vis Sci*. 1990;31:1119–1126.
150. Fukuda M, Wang HF. Dry eye and closed eye tears. *Cornea*. 2000;19:S44–8. doi:10.1097/00003226-200005001-00009
151. Da Dalt S, Moncada A, Priori R, Valesini G, Pivetti-Pezzi P. The lactoferrin tear test in the diagnosis of Sjogren's syndrome. *Eur J Ophthalmol*. 1996;6:284–286. doi:10.1177/112067219600600311
152. Wang HF, Fukuda M, Shimomura Y. Diagnosis of dry eye. *Semin Ophthalmol*. 2005;20:53–62. doi:10.1080/08820530590931115
153. Ponzini E, Scotti L, Grandori R, Tavazzi S, Zambon A. Lactoferrin concentration in human tears and ocular diseases: a meta-analysis. *Invest Ophthalmol Vis Sci*. 2020;61(12):9. doi:10.1167/iovs.61.12.9
154. Flanagan JL, Willcox MD. Role of lactoferrin in the tear film. *Biochimie*. 2009;91(1):35–43. doi:10.1016/j.biochi.2008.07.007
155. Reese V, Youngbar PR. The effect of punctal occlusion on tear lactoferrin in aqueous deficient dry eye patients. *Adv Exp Med Biol*. 2002;506:1269–1271.
156. Williams DL. A comparative approach to topical cyclosporine therapy. *Eye*. 1997;11(11):453–464. doi:10.1038/eye.1997.126

157. Chotikavanich S, de Paiva CS, Li de Q, et al. Production and activity of matrix metalloproteinase-9 on the ocular surface increase in dysfunctional tear syndrome. *Invest Ophthalmol Vis Sci.* 2009;50:3203–3209. doi:10.1167/iovs.08-2476
158. Aragona P, Aguennouz M, Rania L, et al. Matrix metalloproteinase 9 and transglutaminase 2 expression at the ocular surface in patients with different forms of dry eye disease. *Ophthalmology.* 2015;122:62–71. doi:10.1016/j.ophtha.2014.07.048
159. Sakimoto T, Ohnishi T, Ishimori A. Simultaneous study of matrix metalloproteinases, proinflammatory cytokines, and soluble cytokine receptors in the tears of noninfectious corneal ulcer patients. *Graefes Arch Clin Exp Ophthalmol.* 2014;252:1451–1456. doi:10.1007/s00417-014-2708-1
160. Acera A, Vecino E, Duran JA. Tear MMP-9 levels as a marker of ocular surface inflammation in conjunctivochalasis. *Invest Ophthalmol Vis Sci.* 2013;54(13):8285–8291. doi:10.1167/iovs.13-12235
161. Kaufman HE. The practical detection of mmp-9 diagnoses ocular surface disease and may help prevent its complications. *Cornea.* 2013;32:211–216. doi:10.1097/ICO.0b013e3182541e9a
162. Sambursky R, Davitt WF 3rd, Latkany R, et al. Sensitivity and specificity of a point-of-care matrix metalloproteinase 9 immunoassay for diagnosing inflammation related to dry eye. *JAMA Ophthalmol.* 2013;131:24–28. doi:10.1001/jamaophthol.2013.561
163. Bang SP, Son MJ, Kim H, Lee YH, Jun JH. In vitro validation of the tear matrix metalloproteinase 9 in-situ immunoassay. *Sci Rep.* 2020;10(1):15126. doi:10.1038/s41598-020-71977-w
164. Lanza NL, Valenzuela F, Perez VL, Galor A. The matrix metalloproteinase 9 point-of-care test in dry eye. *Ocul Surf.* 2016;14:189–195. doi:10.1016/j.jtos.2015.10.004
165. McCabe E, Narayanan S. Advancements in anti-inflammatory therapy for dry eye syndrome. *Optometry.* 2009;80(10):555–566. doi:10.1016/j.optm.2009.02.010
166. Pflugfelder SC. Anti-inflammatory therapy for dry eye. *Am J Ophthalmol.* 2004;137(2):337–342. doi:10.1016/j.ajo.2003.10.036
167. Sall K, Stevenson OD, Mundorf TK, Reis BL. Two multicenter, randomized studies of the efficacy and safety of cyclosporine ophthalmic emulsion in moderate to severe dry eye disease. CsA Phase 3 Study Group. *Ophthalmology.* 2000;107:631–639. doi:10.1016/S0161-6420(99)00176-1
168. Park JY, Kim BG, Kim JS, Hwang JH. Matrix metalloproteinase 9 point-of-care immunoassay result predicts response to topical cyclosporine treatment in dry eye disease. *Transl Vis Sci Technol.* 2018;7:31. doi:10.1167/tvst.7.5.31
169. Sambursky R. Presence or absence of ocular surface inflammation directs clinical and therapeutic management of dry eye. *Clin Ophthalmol.* 2016;10:2337–2343. doi:10.2147/OPHTH.S121256
170. Dupuis P, Prokopich CL, Hynes A, Kim H. A contemporary look at allergic conjunctivitis. *Allergy Asthma Clin Immunol.* 2020;6:1–18.
171. Shoji J. Ocular allergy test and biomarkers on the ocular surface: clinical test for evaluating the ocular surface condition in allergic conjunctival diseases. *Allergol Int.* 2020;69:496–504. doi:10.1016/j.alit.2020.05.003
172. Leonardi A, Lazzarini D, Bortolotti M, Piliago F, Midena E, Fregona I. Corneal confocal microscopy in patients with vernal keratoconjunctivitis. *Ophthalmology.* 2012;119(3):509–515. doi:10.1016/j.ophtha.2011.09.018
173. Nomura K, Takamura E. Tear IgE concentrations in allergic conjunctivitis. *Eye.* 1998;12(12):296–298. doi:10.1038/eye.1998.68
174. Mimura T, Usui T, Mori M, et al. Relationship between total tear and serum IgE in allergic conjunctivitis. *Int Arch Allergy Immunol.* 2011;154:349–352. doi:10.1159/000321828
175. Inada N, Shoji J, Kato H, Kiely S, Mulyanto SM, Sawa M. Clinical evaluation of total IgE in tears of patients with allergic conjunctivitis disease using a novel application of the immunochromatography method. *Allergol Int.* 2009;58(4):585–589. doi:10.2332/allergolint.09-OA-0101
176. Aalders-Deenstra V, Kok PT, Bruynzeel PL. Measurement of total IgE antibody levels in lacrimal fluid of patients suffering from atopic and non-atopic eye disorders. Evidence for local IgE production in atopic eye disorders? *Br J Ophthalmol.* 1985;69(5):380–384. doi:10.1136/bjo.69.5.380
177. Toda I, Shimazaki J, Tsubota K. Dry eye with only decreased tear break-up time is sometimes associated with allergic conjunctivitis. *Ophthalmology.* 1995;102:302–309. doi:10.1016/S0161-6420(95)31024-X
178. Onguchi T, Dogru M, Okada N, et al. The impact of the onset time of atopic keratoconjunctivitis on the tear function and ocular surface findings. *Am J Ophthalmol.* 2006;141:569–571. doi:10.1016/j.ajo.2005.09.032
179. Dogru M, Asano-Kato N, Tanaka M, et al. Ocular surface and MUC5AC alterations in atopic patients with corneal shield ulcers. *Curr Eye Res.* 2005;30:897–908. doi:10.1080/02713680500196715
180. Lobefalo L, D'Antonio E, Colangelo L, et al. Dry eye in allergic conjunctivitis: role of inflammatory infiltrate. *Int J Immunopathol Pharmacol.* 1999;12:133–137. doi:10.1177/205873929901200304
181. Berdy GJ, Hedqvist B. Ocular allergic disorders and dry eye disease: associations, diagnostic dilemmas, and management. *Acta Ophthalmol Scand.* 2000;230:32–37. doi:10.1034/j.1600-0420.2000.078s230032.x
182. Gipson IK, Argüeso P. Role of mucins in the function of the corneal and conjunctival epithelia. *Int Rev Cytol.* 2003;231:1–49.
183. Jumblatt MM, McKenzie RW, Jumblatt JE. MUC5AC mucin is a component of the human precorneal tear film. *Investig Ophthalmol Vis Sci.* 1999;40:43–49.
184. Zhang J, Yan X, Li H. Analysis of the correlations of mucins, inflammatory markers, and clinical tests in dry eye. *Cornea.* 2013;32:928–932. doi:10.1097/ICO.0b013e3182801622
185. Aydin E, Gokhale M, Azizoglu S, Suphioglu C. To see or not to see: a systematic review of the importance of human ocular surface cytokine biosignatures in ocular allergy. *Cells.* 2019;8:620. doi:10.3390/cells8060620
186. Leonardi A, Doan S, Fauquert JL, et al. Diagnostic tools in ocular allergy. *Allergy.* 2017;72:1485–1498. doi:10.1111/all.13178
187. Bielory L, Delgado L, Katelaris CH, Leonardi A, Rosario N, Vichyanoud P. ICON: diagnosis and management of allergic conjunctivitis. *Ann Allergy Asthma Immunol.* 2020;124:118–134. doi:10.1016/j.anai.2019.11.014
188. Tsubota K, Takamura E, Hasegawa T, Kobayashi T. Detection by brush cytology of mast cells and eosinophils in allergic and vernal conjunctivitis. *Cornea.* 1991;10:525–531. doi:10.1097/00003226-199110000-00011
189. Wakamatsu TH, Satake Y, Igarashi A, et al. IgE and eosinophil cationic protein (ECP) as markers of severity in the diagnosis of atopic keratoconjunctivitis. *Br J Ophthalmol.* 2012;96:581–586. doi:10.1136/bjophthalmol-2011-300143
190. Kari O, Salo OP, Björkstén F, Backman A. Allergic conjunctivitis, total and specific IgE in the tear fluid. *Acta Ophthalmol.* 1985;63:97–99. doi:10.1111/j.1755-3768.1985.tb05224.x

191. Mimura T, Usui T, Yamagami S, Miyai T, Amano S. Relationship between total tear IgE and severity of acute seasonal allergic conjunctivitis. *Curr Eye Res*. 2012;37(10):864–870. doi:10.3109/02713683.2012.689069
192. Mimura T, Usui T, Yamagami S, Miyai T, Amano S. Relationship between total tear IgE and specific serum IgE in autumnal allergic conjunctivitis. *Cornea*. 2013;32(1):14–19. doi:10.1097/ICO.0b013e318253214b
193. Bao J, Tian L, Meng Y, et al. Clinical and translational allergy. *Clin Transl Allergy*. 2022;12:e12139. doi:10.1002/ctt2.12139
194. Insler MS, Lim JM, Queng JT, et al. Tear and serum IgE concentrations by Tandem-R IgE immunoradiometric assay in allergic patients. *Ophthalmology*. 1987;94(8):945–948. doi:10.1016/S0161-6420(87)33355-X

## Clinical Ophthalmology

Dovepress

### Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-ophthalmology-journal>