



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Encephalomyocarditis virus is potentially derived from eastern bent-wing bats living in East Asian countries

Karla Cristine C. Doysabas<sup>a,1</sup>, Mami Oba<sup>b,1</sup>, Masaya Furuta<sup>a,1</sup>, Keisuke Iida<sup>a,1</sup>, Tsutomu Omatsu<sup>b</sup>, Tetsuya Furuya<sup>b</sup>, Takashi Okada<sup>c</sup>, Kripitch Sutummaporn<sup>b</sup>, Hiroshi Shimoda<sup>d</sup>, Min-Liang Wong<sup>e</sup>, Chung-Hsin Wu<sup>f</sup>, Yasushige Ohmori<sup>a</sup>, Ryosuke Kobayashi<sup>a</sup>, Yupadee Hengjan<sup>a</sup>, Kenzo Yonemitsu<sup>d</sup>, Ryusei Kuwata<sup>d</sup>, Yoo-Kyung Kim<sup>g</sup>, Sang-Hyun Han<sup>g</sup>, Joon-Hyuk Sohn<sup>h</sup>, Sang-Hoon Han<sup>i</sup>, Kazuo Suzuki<sup>j</sup>, Junpei Kimura<sup>h</sup>, Ken Maeda<sup>d</sup>, Hong-Shik Oh<sup>g</sup>, Daiji Endoh<sup>k</sup>, Tetsuya Mizutani<sup>b</sup>, Eiichi Hondo<sup>a,\*</sup>

<sup>a</sup> Laboratory of Animal Morphology, Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya, Japan

<sup>b</sup> Laboratory of Veterinary Microbiology, Cooperative Department of Veterinary Medicine, Tokyo University of Agriculture and Technology, Japan

<sup>c</sup> Department of Pediatric Infectious Diseases, Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan

<sup>d</sup> Yamaguchi University, Yamaguchi, Japan

<sup>e</sup> National Chung Hsing University, Taiwan

<sup>f</sup> National Taiwan Normal University, Taiwan

<sup>g</sup> Institute of Science Education, Jeju National University, Jeju, South Korea

<sup>h</sup> Laboratory of Veterinary Anatomy and Cell Biology and Research Institute for Veterinary Science, College of Veterinary Medicine, Seoul National University, Seoul, South Korea

<sup>i</sup> Natural Institute of Biological Resources, South Korea

<sup>j</sup> Hikiwa Park Center, Tanabe, Wakayama, Japan

<sup>k</sup> Department of Veterinary Radiology, School of Veterinary Medicine, Rakuno Gakuen University, 582 Midorimachi, Bunkyo-dai, Ebetsu-shi 069-8501, Japan

### ARTICLE INFO

#### Keywords:

Encephalomyocarditis virus (EMCV)  
*Miniopterus fuliginosus*

### ABSTRACT

Bats are reservoir hosts of many zoonotic viruses and identification of viruses that they carry is important. This study aimed to use high throughput screening to identify the viruses in fecal guano of Taiwanese insectivorous bats caves in order to obtain more information on bat-derived pathogenic viruses in East Asia. Guano samples were collected from two caves in Taiwan, pooled, and then subjected to Multiplex PCR-based next generation sequencing for viral identification. Subsequently, encephalomyocarditis virus (EMCV) sequence was detected and confirmed by reverse transcription PCR. EMCV is considered as rodent virus and thus, animal species identification through cytochrome oxidase I (COI) barcoding was further done to identify the viral source. Finally, determination of distribution and verification of the presence of EMCV in guano obtained from Japanese and South Korean caves was also done. We concluded that the guano collected was not contaminated with the excrement of rodents which were reported and presumed to live in Taiwan. Also, EMCV genome fragments were found in guanof of Japanese and South Korean caves. It is possible that the eastern bent-wing bat (*Miniopterus fuliginosus*) is one of the natural hosts of EMCV in East Asia.

### 1. Introduction

Bats are considered natural reservoirs for several zoonotic viruses including lyssaviruses (Davis et al., 2013a, b; Albas et al., 2011; Almeida et al., 2011, 2005), henipaviruses (Roche et al., 2015; Breed et al., 2011; Rahman et al., 2010), coronaviruses (Anthony et al., 2017;

Chen et al., 2016; Anindita et al., 2015; Annan et al., 2013; Anthony et al., 2013), and/or picornaviruses (Wu et al., 2012; Lau et al., 2011). Most viruses, which originate from bats and are pathogenic in humans and other mammals, appear to cause no clinical signs of disease in bats under natural or experimental infection (Calisher et al., 2006). Bats are also considered the origin of the Middle East respiratory syndrome

\* Corresponding author at: Laboratory of Animal Morphology, Department of Animal Science, Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-8601 Japan.

E-mail address: [ehondo@agr.nagoya-u.ac.jp](mailto:ehondo@agr.nagoya-u.ac.jp) (E. Hondo).

<sup>1</sup> This manuscript is equally contributed by Karla Cristine C. Doysabas, Mami Oba, Masaya Furuta, and Keisuke Iida.

<https://doi.org/10.1016/j.virusres.2018.10.020>

Received 20 September 2018; Received in revised form 19 October 2018; Accepted 30 October 2018

Available online 01 November 2018

0168-1702/ © 2018 Elsevier B.V. All rights reserved.

coronavirus (MERS-CoV) that was first reported in Saudi Arabia (Wang et al., 2014), and severe acute respiratory syndrome coronavirus (SARS-CoV) that caused outbreaks in China in 2003 (Ge et al., 2016, 2013; Balboni et al., 2012; Wang et al., 2006). A novel coronavirus, which is closely related to SARS-CoV and uses Angiotensin Converting Enzyme 2 (ACE-2) receptor from humans, civets, and Chinese horseshoe bats for cell entry, was isolated from Chinese horseshoe bats suggesting these bats might be natural reservoirs of SARS-CoV (Ge et al., 2013).

In 2012, an outbreak of rabies occurred among ferret badgers in Taiwan (Chiou et al., 2016; Tsai et al., 2016; Chiou et al., 2014) after the country had been considered rabies-free since the last reported human case in 1959 and non-human animal in 1961 (Huang et al., 2015; Chiou et al., 2014). During 2016–2017, a putative new lyssavirus was found in the Japanese pipistrelle bat in the same country (Hu et al., 2018). SARS-related coronavirus and *Alphacoronavirus* were also detected in the country's bat population (Chen et al., 2016). To obtain more information on bat-derived pathogenic viruses in East Asia, this study aimed to determine the viruses present in fecal guano of Taiwanese insectivorous bats caves.

## 2. Materials and methods

### 2.1. Sampling

Sampling was done in the two caves of Taiwan located in Taoyuan City, Dasi district: 24°50'35"N, 121°17'58"E (Taiwan 1) and New Taipei City, Ruifang district: 25°07'34"N, 121°49'59"E (Taiwan 2) wherein fecal guanios were collected randomly at several points in the cave and then pooled. During the process of viral identification, the encephalomyocarditis virus (EMCV) genome was found. To determine the distribution of the virus and confirm the presence of EMCV in fecal guanios, sampling was also performed in other countries including the main island of South Korea (36°39'33"N, 128°11'34"E); Jeju Island, South Korea (33°26'08"N, 126°50'15"E); Nishi-Yoshino, Nara, Japan (34°18'58"N, 135°46'28"E); Kawakami, Nara, Japan (34°17'02"N, 136°00'09"E); Sabae-shi, Fukui, Japan (35°56'54"N, 136°08'08"E); and Wakasa, Fukui, Japan (35°31'52"N, 135°50'22"E).

### 2.2. Primer design (Multiplex PCR)

The viral genera used for primer design (Supplementary Table 1) were selected in reference to the ICTV Master Species List (King et al., 2018). Thirty-seven genera including viruses possibly infectious to mammals and birds were selected. The data sets for primer design of each genus were compiled using viral strains specified by RefSeq on GenBank. The CoCoMo primer algorithms (Jimba et al., 2010) and CLC Genomics Workbench 6.5.1 (CLC bio) were used for the design. If there was only one RefSeq sequence per genus, another sequence of the same genus was added to the data set. The product size of the PCR was designed at a maximum of 1000 bp. All primers used for the Multiplex PCR are presented in Supplementary Table 1. Previously reported universal primers for Herpesvirus, Adenovirus, Orthoreovirus, and Bornavirus (Weissenböck et al., 2009; Wellehan et al., 2004; Spinner and Di Giovanni, 2001; Mizutani et al., 1999; VanDevanter et al., 1996) were also used.

### 2.3. Animal species identification

Identification of animal species was done from the guano through DNA barcoding based on cytochrome c oxidase I (COI) sequence using the specific primers (5'-GGTCAACAAATCATAAAGATATTGG, 5'-TAAACCTTCAGGGTGACCAAAAAATCA).

### 2.4. Data sets for mapping

For viral identification, the sequences used for primer design were

used as the data set for mapping reads from next generation sequencing (NGS). For animal species identification, sequences from a total of 75 species of mammals (Supplementary Table 2) that had been recorded and/or presumed to be present in Taiwan, and/or distributed worldwide were used as reference (Vincenot, 2017; Heaney and Molur, 2016; Ruedas et al., 2016; Challender et al., 2014; Lin and Motokawa, 2014; Bates et al., 2008; Hai Yin and Richardson, 2008; Hutson et al., 2008; Smith and Johnston, 2008; Wilson and Reeder, 2005).

### 2.5. Total nucleic acid extraction and cDNA synthesis

Total viral nucleic acid from pooled guano from each cave was extracted using the High Pure Viral Nucleic Acid Kit (Roche, Basel, Switzerland); half of the samples were used for RNA extraction by purification with the TURBO DNA-free™ Kit (Life Technologies, CA, USA) according to the manufacturer's instruction. Complementary DNA (cDNA) was generated using random hexamers and the SuperScript™ IV First-Strand Synthesis System (Thermo Fisher Scientific, California, USA) by incubating for 10 min at 23°C, 10 min at 50°C, and 10 min at 80°C.

### 2.6. Multiplex PCR

A primer mix was prepared for each well so that multiplex PCR using approximately 700 primers could be performed on one 96-well plate. Primers were mixed for each viral genus making the maximum number of primers in one well as 10, occupying 94 wells. The cDNA library was amplified using KOD-Multi & Epi-® (TOYOBO, Osaka, Japan) with initial denaturation at 94°C for 10 min., followed by 40 cycles with denaturation at 98°C for 10 s, annealing at 45°C for 30 s, and extension at 68°C for 30 s, and a final extension at 72°C for 10 min. PCR products were then collected from all wells (94 wells), mixed, and purified with Agencourt AMPure XP (Beckman Coulter, Brea, Calif., USA, USA) using 80% ethanol to remove primer dimers of less than 100 bp.

### 2.7. Next generation sequencing (NGS) and data analysis

A NGS library was first constructed according to the manufacturer's protocol using the NEBNext® Ultra™ RNA Library Prep Kit for Illumina® (New England Bio Labs, Massachusetts, USA). Sequencing of paired ends was performed with the MiSeq Reagent Kit v2 (300 cycles) (Illumina®, San Diego, CA, USA) wherein each read was output as a FASTQ format file from the MiSeq reporter and analyzed using CLC genomics Workbench 6.5.1 (Filgen, Nagoya, Japan). Commands for trimming sequences with low quality and commands for removing 20 bases corresponding to primers from the 5'-end of each read were executed. Trimmed reads were mapped to each sequence of the prepared data sets for viral identification (Supplementary Table 1) with a 50% length fraction and 80% similarity fraction. Reads with a minimum consensus length of 200 and a minimum mapping read number of 100 were selected, and the reference sequence that was mapped at the forward and reverse end of the predicted fragment between the primers' binding site was selected and judged as positive. For animal species identification, NGS data that showed less than 50% similarity to the COI reference in consecutive full-length sequence of 5' and 3' ends of NGS data (less than 151 bp each sequence) were eliminated. Since only one COI genomic sequence per species was included in the reference despite the intraspecies variation, the maximum variation in COI barcoding was computed with the formula:  $\text{similarity} + [(100 - \text{similarity}) \times 0.11]$ . The maximum similarity was used as the threshold to determine the species.

### 2.8. Amplification of EMCV genomic fragments

When the EMCV genome was detected using NGS, reverse-

**Table 1**  
Primers for EMCV genome.

EMCV genes	Forward Primer (5'-3')	Reverse Primer (5'-3')	Amplicon size (bp)
Promoter (1st set)	TGAATGTCGTGAAGGAAGCAGT	ACCTCGACTAAACACATGT	315
Promoter (2nd set)	TGGAAGCTTCTTGAAGACA	AGATCAGATCCCACATAAT	240
1A	ATTCCACCTCCTCAGACAAGA	AGCTAGCAATGGAAGCATAT	206
2B (1st set)	TTCATGTTTAGACCAAGGAAACA	AGAGATTGTGGGAAACCGT	432
2B (2nd set)	AGACCAAGGAGCGGCAGTGT	TGTCTTGAACITAGCTGCTAT	361
3A/B (1st set)	TCCAGTAGACGAGGTCACT	TGAAATGTCCAACAACCTGCA	321
3A/B (2nd set)	TCCAGCAGCTTAAAGCAAGACA	AACTCTCGCCGTCTCAATTGTA	256


transcription and PCR amplification of EMCV genes was done using SuperScript™ IV First-Strand Synthesis System (Thermo Fisher Scientific, California, USA) first by incubating for 10 min at 23°C, 10 min at 50°C, and 10 min at 80°C. Amplification was carried out using the specific primers for the EMCV genome (Table 1), which were designed based on the highly conserved regions of EMCV (Liu et al., 2016; Duke et al., 1992), and KOD-Plus-Neo (Toyobo, Japan) at the following conditions: 2 min at 94°C; 35 cycles of 10 ss at 98°C, 30 ss at 58°C, and 30 ss at 68°C, with a final extension at 72°C for 5 min.

### 3. Results

Guanos from two caves in Taiwan were subjected to NGS using the designed primers and the reads were mapped to the viral sequences used for primer design. In two caves (Taiwan 1 and Taiwan 2), sequences matched to the reference were homologous to EMCV, enterovirus F strain, rabbit vesivirus, and Spanish goat encephalitis virus (Supplementary Table 3). To confirm the presence of these viral genomes, conventional RT-PCR followed by Sanger sequencing was performed using specific primers for EMCV (5'CACCATATTGCCGCTTT, 5'CCACAACCTATCCAACCTCAC), enterovirus F strain (5'CCACGTGGCGC TAGTACC, 5'ACGGGGTACCGAAAGTAG), rabbit vesivirus (5'TCAAAC CCAATTGGCATCTTC, 5'GCTTTGATGTAGTAGAATTGCCCA), and Spanish goat encephalitis virus (5'TGCGTGTACAACATGATGGGA, 5' CACGCAGTCATCACCCTGA). However, only the EMCV genomic sequences were amplified. To confirm the presence of EMCV, RT-PCR was performed using the specific primers for the EMCV genome (Table 1) wherein only the promoter region was amplified (Fig. 1A). To eliminate the possibility that the EMCV genome is incorporated in the genome of rodents, *Rhinolophinae*, and *Miniopterinae*, the obtained sequence by RT-

PCR was then subjected to BLAST wherein the result showed the absence of EMCV in the genome of these animals.

Rodents have been known to be the natural reservoir of EMCV. To determine if the guanos collected also contained the excreta of some rodents, identification of animal species was done through DNA bar-coding based on the 5'-end and 3'-end regions of the mitochondrial gene, COI sequence. Using the next generation sequencer with CLC genomics software, the number of reads in Taiwan 1 and Taiwan 2 caves were 5'-end: 3256, 3'-end: 3526, and 5'-end: 3198, 3'-end: 3198, respectively. After elimination of reads whose similarity was less than 50% to the COI reference, the number of mapped reads was reduced to 5'-end: 2167, 3'-end: 1936 (Taiwan 1), and 5'-end: 880, 3'-end: 518 (Taiwan 2). Among these read sequences, a total of 189 kinds of consensus sequences were found. The 189 consensus sequences were mapped to 48 out of 76 reference species. Of the 48 reference species, 11 were from the order Chiroptera (*Miniopterus fuliginosus*, *Myotis adversus*, *Myotis formosus*, *Nyctalus noctula*, *Murina puta*, *Tadarida teniotis*, *Rhinolophus Monoceros*, *Hipposideros armiger*, *Eptesicus serotinus*, *Rhinolophus Monoceros*, *Rhinolophus ferrumequinum*), 19 from Rodentia (*Rattus exulans*, *Rattus rattus*, *Rattus tanezumi*, *Rattus tiomanicus*, *Rhizomys pruinosus*, *Euchoreutes naso*, *Mesocricetus auratus*, *Eospalax baileyi*, *Petaurista alborufus*, *Callosciurus erythraeus*, *Tamiops maritimus*, *Apodemus agrarius*, *Mus caroli*, *Mus musculus*, *Eothenomys chinensis*, *Micromys minutus*, *Rattus losea*, *Demonys pernyi*), three from Eulipotyphla (*Condylura cristata*, *Mogera robusta*, *Hylomys suillus*), five from Soricomorpha (*Crocidura attenuata*, *Crocidura suaveolens*, *Suncus murinus*, *Chimarrogale himalayica*, *Soriculus fumidus*), seven from Carnivora (*Prionailurus bengalensis*, *Paguma larvata*, *Viverricula indica*, *Melogale moschata*, *Canis familiaris*, *Ursus thibetanus*, *Mustela nivalis*), and three from Artiodactyla (*Cervus nippon taiouanus*, *Cervus unicolor*



**Fig. 1.** Electrophoresis result of RT-PCR of EMCV genes. **A**, RT-PCR result of 4 EMCV genomic fragments using the primers for EMCV genome (Table 1): 1, promoter; 2, 1A; 3, 2B; 4, 3A/B; in samples collected from Taiwan 1 (Taoyuan city, Dasi district) and Taiwan 2 (New Taipei city) showing positive PCR-fragment for EMCV promoter after sequencing (arrows). **B**, RT-PCR result of EMCV promoter for samples collected in Korea (Main Island, South Korea), Jeju Is. (Jeju Island, South Korea), Nara 1 (Nishi-Yoshino, Nara, Japan), Nara 2 (Kawakami, Nara, Japan), Fukui 1 (Sabae-shi, Fukui, Japan), and Fukui 2 (Wakasa, Fukui, Japan); showing positive PCR-fragment for EMCV promoter after sequencing (arrow) in all the samples.



*swinhoe*, and *Capricornis swinhoei*) (Supplementary Table 3). Since the maximal intraspecific distances of COI is between 0.00–11.00% (Yassin et al., 2010), we considered 89.00% similarity, with the computed maximum variation, as the maximum threshold for identifying the species. Exactly 10/189 consensus sequences were above the threshold similarity to the reference species. Four of these, total of six reads in 5' and 3' ends, were 96.44%–100% matched to the reference animal species *Miniopterus fuliginosus*. These sequences were 97.60%–100% similar to the same species when subjected to BLAST (Supplementary Table 4). One (1 read) was 91.63% similar to *Petaurista alborfus*, two (1 read/consensus) were 96.71% similar to *Prionailurus bengalensis*, one (1 read) was 93.68% similar to *Melogale moschata*, and two (1 and 7 reads) were 91.28% similar to *Mogera robusta*. When subjected to BLAST for confirmation, all of them were 100% homologous to species of insects (*Cochyllis hospes*, *Fosterina rustica*, *Alysiinae sp.*) and a spider (*Argyroides zonatus*). The remaining consensus sequences (150/189) that were below 89.00% similarity, were also subjected to BLAST to determine the species, where 120 of 150 sequences showed  $\geq 89.00\%$  similarity to various species of insects and other species such as scorpions, fungi, gastropods, and even a nematode (Supplementary Table 5). Moreover, in 23 of the 150 sequences consensus was below 89.00%, with regard to both the reference species and to other species when BLASTed while 7 of the 150 species had no matching species when BLASTed and thus, these sequences could not be identified. All of the highest homologies of 23 sequences did not belong to vertebrates.

To further confirm the presence of the EMCV genome in the feces of insectivorous bats, guano samples were also collected in the bat caves in Japan and Korea. The samples from all of the caves examined contained EMCV promoter region (Fig. 1B), which was confirmed by sequencing. This region was reported to be 100% homologous to Chinese, Korean, and Belgian swine isolates; and other strains from chimpanzee and mice (Liu et al., 2016).

#### 4. Discussion

This study showed that the EMCV genome was widely found in fecal guanos of East Asian countries, and the possibility of interfusion of rodents' excretion to the guanos was very low. It is possible that *Miniopterus fuliginosus* is one of the natural reservoirs of EMCV.

EMCV, under the family *Picornaviridae*, genus *Cardiovirus*, has a single-stranded, positive sense RNA and the virus has worldwide distribution (Billinis et al., 2004; King et al., 2000). Although it is considered a rodent virus (Liu et al., 2017, 2016; Spyrou et al., 2004), it has wide-spectrum of hosts including monkeys and porcupines (Cardeti et al., 2016), elephants (Lamglait et al., 2015), wild boar (Liu et al., 2017, 2016), dogs (Luo et al., 2017), tapir (Vercammen et al., 2017), with pigs being the most sensitive (Liu et al., 2017; Maurice et al., 2016; Vansteenkiste et al., 2016). It causes myocarditis, encephalitis, neurological diseases, reproductive disorders, and diabetes (Luo et al., 2017; Carocci and Bakkali-Kassimi, 2012). As a potentially zoonotic virus, its discovery in these caves could pose a threat to animals and humans living nearby because this virus is resistant to a hostile environment and thus can remain infectious for days (Carocci and Bakkali-Kassimi, 2012).

Although the sample areas were mainly inhabited by *Miniopterus fuliginosus* and *Rhinolophus ferrumequinum*, detection of EMCV in fecal guanos could not conclude that the source of the viral genome was indeed these bats since the samples were not collected directly from their bodies. Guanos might be composed of excreta from rodents, the natural reservoir of EMCV, aside from bats. To determine the species that contributed most to the guano formation collected, animal species identification was done through NGS using the 1545-bp, at the 5'-end of the mitochondrial gene *COI* as the DNA barcode region. This *COI* gene locus, being easily amplified using universal primers, shows low intraspecific and high interspecific divergence (Dentinger et al., 2011), and possesses a higher range of phylogenetic signal than any other

mitochondrial genes (Hebert et al., 2003a), and has been proven to be highly effective in identifying the species of various organisms (Andujar et al., 2018; Bingpeng et al., 2018; Changbunjong et al., 2018; Yassin et al., 2010; Hubert et al., 2008; Ward et al., 2005; Hebert et al., 2004, 2003b).

In both sample areas, a total of 189 consensus sequences were mapped to 48 out of 75 reference sequences wherein 10/189 were  $\geq 89.00\%$  similar to the reference species (Supplementary table 3). Four of these mapped sequences, containing two and four reads in Taiwan 1 and Taiwan 2, respectively, were 96.44%–100% matched to the species *Miniopterus fuliginosus*, and were 98.00%–100% similar to the same species when subjected to BLAST (Table 3). The majority of other sequences belonged to the order Diptera followed by Lepidoptera, Hymenoptera, Coleoptera, Trichoptera, Orthoptera, and Thysanoptera, which are the main diet of the insectivorous bats including *Miniopterus fuliginosus* (Hu et al., 2011; Lumsden and Bennett, 2005; Funakoshi and Takeda, 1998; Bernard and D'avidson, 1996). Also, it was unlikely that other unidentified sequences were derived from vertebrates. That is, the possibility that the guanos contained excreta of vertebrate other than *Miniopterus fuliginosus* was very low. The relative frequency of the reads from *Miniopterus fuliginosus* was very low compared to that from insects. Insectivorous bats can consume 3.60–8.00 g/day, 4.20–7.60 insects/minute in a foraging time of 20.30–462.70 min (Encarnação and Dietz, 2006). Thus, the proportion of the DNA of this bat, from its intestinal epithelium, in fecal guano could be low in comparison to the insects that had been ingested and excreted and/or insects that directly jumped into the guano and finally died. These results suggest that the main source of the guano is the *Miniopterus fuliginosus* and that the possibility of interfusion of rodents' excreta to the guano was very low.

Apart from *Miniopterus fuliginosus*, a few numbers of *Rhinolophus ferrumequinum* were also seen in the caves; however, species identification though COI barcoding did not confirm this bat's genome. This does not eliminate the possibility of this species being a host of EMCV. Since the population of this species was overwhelmed by *Miniopterus fuliginosus*, it is possible that the feces from *Rhinolophus ferrumequinum* were not sampled in the representative collection points in the cave. Even in the same cave, the group of *Miniopterus fuliginosus* and *Rhinolophus ferrumequinum* were segregated because bats usually inhabit separate rooms of the roosting site (Hengjan et al., 2017; Schmidly and Bradley, 2016). Bats are known to have a high viral sharing network connection, suggesting that viruses may pass easily among bat species (Luis et al., 2015). Thus, it is important to also consider *Rhinolophus ferrumequinum* as a possible host of EMCV.

One of the many traits of bats that makes them suited to hosting and transmitting viruses is their ability to fly long distances during seasonal migrations in search for food (Calisher et al., 2006). Microchiropterans can travel from 10 km up to 80 km per night (Mildenstein and De Jong, 2011; Wong et al., 2007). This ability is a major factor of viral transmission to various locations. For instance, rabies virus associated with some species of bats have been identified throughout the geographic range and migratory routes of these bats (Calisher et al., 2006). Aside from flight ability, accidental and intentional translocation of bats has also been reported both intra- and intercontinentally. In a review by Constantine (2003), translocation of bats happens accidentally through ships and shipping containers where they land and/or roost, through aircrafts such as the eastern pipistrelle bat found in a plane that travelled from Mexico to Texas, or intentionally when these bats are used for captivity or for release after being rescued. All of these translocation incidents are modes of viral expansion. This is supported in this study as the EMCV genome was also found in samples from Korea and Japan. Moreover, migration of the bats between these countries was previously reported according to genetic analysis (Iida et al., 2017).

This study suggests the presence of EMCV sequences in *Miniopterus fuliginosus* in Taiwan, Korea, and Japan. It is possible that *Miniopterus fuliginosus* is one of the natural hosts of EMCV in East Asia.

## Acknowledgements

We thank the e-Asia Joint Research Program (Japan Agency for Medical Research and Development) for supporting this study.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.virusres.2018.10.020>.

## References

- Albas, A., Campos, A.C., Araujo, D.B., Rodrigues, C.S., Sodre, M.M., Durigon, E.L., Favoretto, S.R., 2011. Molecular characterization of rabies virus isolated from non-haematophagous bats in Brazil. *Rev. Soc. Bras. Med. Trop.* 44, 678–683.
- Almeida, M.F., Martorelli, L.F.A., Aires, C.C., Sallum, P.C., Durigon, E.L., Massad, E., 2005. Experimental rabies infection in haematophagous bats *Desmodus rotundus*. *Epidemiol. Infect.* 133, 523–527. <https://doi.org/10.1017/S0950268804003656>.
- Almeida, M.F., Favoretto, S.R., Martorelli, L.F., Trezza-Netto, J., Campos, A.C., Ozahata, C.H., Sodre, M.M., Kataoka, A.P., Sacramento, D.R., Durigon, E.L., 2011. Characterization of rabies virus isolated from a colony of *Eptesicus fuscus* bats in Brazil. *Rev. Inst. Med. Trop. Sao Paulo* 53, 31–37.
- Andujar, C., Arribas, P., Yu, D.W., Vogler, A.P., Emerson, B.C., 2018. Why the COI barcode should be the community DNA metabarcode for the metazoa. *Mol. Ecol.* 27 (20), 3968–3975. <https://doi.org/10.1111/mec.14844>.
- Anindita, P.D., Sasaki, M., Setiyono, A., Handharyani, E., Orba, Y., Kobayashi, S., Rahmadani, I., Taha, S., Adiani, S., Subangkit, M., Nakamura, I., Sawa, H., Kimura, T., 2015. Detection of coronavirus genomes in moluccan naked-backed fruit bats in Indonesia. *Arch. Virol.* 160, 1113–1118. <https://doi.org/10.1007/s00705-015-2342-1>.
- Annan, A., Baldwin, H.J., Corman, V.M., Klose, S.M., Owusu, M., Nkrumah, E.E., Badu, E.K., Anti, P., Agbenyega, O., Meyer, B., Oppong, S., Sarkodie, Y.A., Kalko, E.K., Lina, P.H., Godlevska, E.V., Reusken, C., Seebens, A., Gloza-Rausch, F., Vallo, P., Tschapka, M., Drosten, C., Drexler, J.F., 2013. Human betacoronavirus 2c emc/2012-related viruses in bats, Ghana and Europe. *Emerg. Infect. Dis.* 19, 456–459. <https://doi.org/10.3201/eid1903.121503>.
- Anthony, S.J., Ojeda-Flores, R., Rico-Chavez, O., Navarrete-Macias, I., Zambrana-Torrel, C.M., Rostal, M.K., Epstein, J.H., Tipples, T., Liang, E., Sanchez-Leon, M., Sotomayor-Bonilla, J., Aguirre, A.A., Avila-Flores, R., Medel, R.A., Goldstein, T., Suzan, G., Daszak, P., Lipkin, W.I., 2013. Coronaviruses in bats from Mexico. *J. Gen. Virol.* 94, 1028–1038. <https://doi.org/10.1099/vir.0.049759-0>.
- Anthony, S.J., Gilardi, K., Menachery, V.D., Goldstein, T., Ssebide, B., Mbabazi, R., Navarrete-Macias, I., Liang, E., Wells, H., Hicks, A., Petrosov, A., Byarugaba, D.K., Debbink, K., Dinno, K.H., Scobey, T., Randell, S.H., Yount, B.L., Cranfield, M., Johnson, C.K., Baric, R.S., Lipkin, W.I., Mazet, J.A., 2017. Further evidence for bats as the evolutionary source of Middle East respiratory syndrome coronavirus. *MBio* 8. <https://doi.org/10.1128/mBio.00373-17>.
- Balboni, A., Battilani, M., Prosperi, S., 2012. The sars-like coronaviruses: the role of bats and evolutionary relationships with sars coronavirus. *New Microbiol.* 35, 1–16.
- Bates, P., Kingston, T., Francis, C., Rosell-Ambal, G., Heaney, L., Gozales, J.-C., Molur, S., Srinivasulu, C., 2008. *Scotophilus kuhlii*. The IUCN Red List of Threatened Species 2008: e.T20068A9142479. <https://doi.org/10.2305/IUCN.UK.2008.RLTS.T20068A9142479.en>. August 29, 2018.
- Bernard, R.T.F., D'avidson, A.A.G., 1996. Does calcium constrain reproductive activity in insectivorous bats? Some empirical evidence for Schreiber's long-fingered bat (*Miniopterus schreibersii*). *Afr. Zool.* 31, 218–220.
- Billinis, C., Leontides, L., Psychas, V., Spyrou, V., Kostoulas, P., Koenen, F., Papadopoulos, O., 2004. Effect of challenge dose and age in experimental infection of pigs with encephalomyocarditis virus. *Vet. Microbiol.* 99, 187–195. <https://doi.org/10.1016/j.vetmic.2004.01.002>.
- Bingpeng, X., Heshan, L., Zhilan, Z., Chunguang, W., Yanguo, W., Jianjun, W., 2018. DNA barcoding for identification of fish species in the Taiwan Strait. *PLoS One* 13, e0198109. <https://doi.org/10.1371/journal.pone.0198109>.
- Breed, A.C., Breed, M.F., Meers, J., Field, H.E., 2011. Evidence of endemic hendra virus infection in flying-foxes (*Pteropus conspicillatus*)—implications for disease risk management. *PLoS One* 6, e28816. <https://doi.org/10.1371/journal.pone.0028816>.
- Calisher, C.H., Childs, J.E., Field, H.E., Holmes, K.V., Schountz, T., 2006. Bats: important reservoir hosts of emerging viruses. *Clin. Microbiol. Rev.* 19, 531–545. <https://doi.org/10.1128/CMR.00017-06>.
- Cardeti, G., Mariano, V., Eleni, C., Aloisi, M., Grifoni, G., Sittinieri, S., Dante, G., Antognetti, V., Foglia, E.A., Cersini, A., Nardi, A., 2016. Encephalomyocarditis virus infection in *Macaca sylvanus* and *hystrix cristata* from an Italian rescue centre for wild and exotic animals. *Virol. J.* 13, 193. <https://doi.org/10.1186/s12985-016-0653-9>.
- Carocci, M., Bakkali-Kassimi, L., 2012. The encephalomyocarditis virus. *Virulence* 3, 351–367.
- Challender, D., Baillie, J., Ades, G., Kaspal, P., Chan, B., Khatiwada, A., Xu, L., Chin, S., KC, R., Nash, H., Hsieh, H., 2014. *Manis pentadactyla*. The IUCN Red List of Threatened Species 2014: e.T12764A45222544. <https://doi.org/10.2305/IUCN.UK.2014-2.RLTS.T12764A45222544.en>. August 29, 2018.
- Changbunjong, T., Bhusri, B., Sedwisai, P., Weluwanarak, T., Nitiyamatawat, E., Chareonviriyaphap, T., Ruangsittichai, J., 2018. Species identification of horse flies (Diptera: Tabanidae) in Thailand using DNA barcoding. *Vet. Parasitol.* 259, 35–43. <https://doi.org/10.1016/j.vetpar.2018.07.002>.
- Chen, Y.N., Phuong, V.N., Chen, H.C., Chou, C.H., Cheng, H.C., Wu, C.H., 2016. Detection of the severe acute respiratory syndrome-related coronavirus and alphacoronavirus in the bat population of Taiwan. *Zoonoses Public Health* 63, 608–615. <https://doi.org/10.1111/zph.12271>.
- Chiou, H.Y., Hsieh, C.H., Jeng, C.R., Chan, F.T., Wang, H.Y., Pang, V.F., 2014. Molecular characterization of cryptically circulating rabies virus from ferret badgers, Taiwan. *Emerg. Infect. Dis.* 20, 790–798. <https://doi.org/10.3201/eid2005.131389>.
- Chiou, H.Y., Jeng, C.R., Wang, H.Y., Inoue, S., Chan, F.T., Liao, J.W., Chiou, M.T., Pang, V.F., 2016. Pathology and molecular detection of rabies virus in ferret badgers associated with a rabies outbreak in Taiwan. *J. Wildl. Dis.* 52, 57–69. <https://doi.org/10.7589/2015-01-007>.
- Constantine, D.G., 2003. Geographic translocation of bats: known and potential problems. *Emerg. Infect. Dis.* 9, 17–21. <https://doi.org/10.3201/eid0901.020104>.
- Davis, A.D., Gordy, P.A., Bowen, R.A., 2013a. Unique characteristics of bat rabies viruses in big brown bats (*Eptesicus fuscus*). *Arch. Virol.* 158, 809–820. <https://doi.org/10.1007/s00705-012-1551-0>.
- Davis, A.D., Jarvis, J.A., Pouliott, C., Rudd, R.J., 2013b. Rabies virus infection in *Eptesicus fuscus* bats born in captivity (naive bats). *PLoS One* 8, e64808. <https://doi.org/10.1371/journal.pone.0064808>.
- Dentinger, B.T., Didukh, M.Y., Moncalvo, J.M., 2011. Comparing COI and ITS as DNA barcode markers for mushrooms and allies (Agaricomycotina). *PLoS One* 6, e25081. <https://doi.org/10.1371/journal.pone.0025081>.
- Duke, G.M., Hoffman, M.A., Palmenberg, A.C., 1992. Sequence and structural elements that contribute to efficient encephalomyocarditis virus translation. *J. Virol.* 66, 1602–1609.
- Encarnação, J.A., Dietz, M., 2006. Estimation of food intake and ingested energy in daubenton's bats (*Myotis daubentonii*) during pregnancy and spermatogenesis. *Eur. J. Wildl. Res.* 52, 221–227. <https://doi.org/10.1007/s10344-006-0046-2>.
- Funakoshi, K., Takeda, Y., 1998. Food habits of sympatric insectivorous bats in southern Kyushu, Japan. *Mammal Study* 23, 49–62. <https://doi.org/10.3106/mammalstudy.23.49>.
- Ge, X.-Y., Li, J.-L., Yang, X.-L., Chmura, A.A., Zhu, G., Epstein, J.H., Mazet, J.K., Hu, B., Zhang, W., Peng, C., Zhang, Y.-J., Luo, C.-M., Tan, B., Wang, N., Zhu, Y., Cramer, G., Zhang, S.-Y., Wang, L.-F., Daszak, P., Shi, Z.-L., 2013. Isolation and characterization of a bat sars-like coronavirus that uses the ACE2 receptor. *Nature* 503, 535–538. <https://doi.org/10.1038/nature12711>.
- Ge, X.Y., Wang, N., Zhang, W., Hu, B., Li, B., Zhang, Y.Z., Zhou, J.H., Luo, C.M., Yang, X.L., Wu, L.J., Wang, B., Zhang, Y., Li, Z.X., Shi, Z.L., 2016. Coexistence of multiple coronaviruses in several bat colonies in an abandoned mineshaft. *Virol. Sin.* 31, 31–40. <https://doi.org/10.1007/s12250-016-3713-9>.
- Hai Yin, W., Richardson, M., 2008. *Maca cyclopis*. The IUCN Red List of Threatened Species 2008: e.T12550A3355290. <https://doi.org/10.2305/IUCN.UK.2008.RLTS.T12550A3355290.en>. August 29, 2018.
- Heaney, L., Molur, S., 2016. *Rattus tanezumii*. The IUCN Red List of Threatened Species 2016: e.T19366A115149780. <https://doi.org/10.2305/IUCN.UK.2016-3.RLTS.T19366A22445589.en>. August 22, 2018.
- Hebert, P.D., Cywinska, A., Ball, S.L., deWaard, J.R., 2003a. Biological identifications through DNA barcodes. *Proc. Biol. Sci.* 270, 313–321. <https://doi.org/10.1098/rspb.2002.2218>.
- Hebert, P.D., Ratnasingham, S., deWaard, J.R., 2003b. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proc. Biol. Sci.* 270 (Suppl. 1), S96–9. <https://doi.org/10.1098/rspb.2003.0025>.
- Hebert, P.D., Stoeckle, M.Y., Zemplak, T.S., Francis, C.M., 2004. Identification of birds through DNA barcodes. *PLoS Biol.* 2, e312. <https://doi.org/10.1371/journal.pbio.0020312>.
- Hengjan, Y., Iida, K., Doysabas, K.C.C., Pichitrasilp, T., Ohmori, Y., Hondo, E., 2017. Diurnal behavior and activity budget of the golden-crowned flying fox (*Acerodon jubatus*) in the Subic Bay forest reserve area, the Philippines. *J. Vet. Med. Sci.* 79, 1667–1674. <https://doi.org/10.1292/jvms.17-0329>.
- Hu, K.L., Wei, L., Zhu, T.T., Wang, X.Z., Zhang, L.B., 2011. Dietary composition, echolocation pulses and morphological measurements of the long-fingered bat *Miniopterus fuliginosus* (Chiroptera: Vespertilioninae). *Zool. Res.* 32, 163–167.
- Hu, S.C., Hsu, C.L., Lee, M.S., Tu, Y.C., Chang, J.C., Wu, C.H., Lee, S.H., Ting, L.J., Tsai, K.R., Cheng, M.C., Tu, W.J., Hsu, W.C., 2018. Lyssavirus in Japanese pipistrelle, Taiwan. *Emerg. Infect. Dis.* 24, 782–785. <https://doi.org/10.3201/eid2404.171696>.
- Huang, A.S., Chen, W.C., Huang, W.T., Huang, S.T., Lo, Y.C., Wei, S.H., Kuo, H.W., Chan, P.C., Hung, M.N., Liu, Y.L., Mu, J.J., Yang, J.Y., Liu, D.P., Chou, J.H., Chuang, J.H., Chang, F.Y., 2015. Public health responses to reemergence of animal rabies, Taiwan, July 16–December 28, 2013. *PLoS One* 10, e0132160. <https://doi.org/10.1371/journal.pone.0132160>.
- Hubert, N., Hanner, R., Holm, E., Mandrak, N.E., Taylor, E., Burridge, M., Watkinson, D., Dumont, P., Curry, A., Bentzen, P., Zhang, J., April, J., Bernatchez, L., 2008. Identifying Canadian freshwater fishes through DNA barcodes. *PLoS One* 3. <https://doi.org/10.1371/journal.pone.0002490>.
- Hutson, A.M., Kingston, T., Francis, C., 2008. *Myotis adversus*. The IUCN Red List of Threatened Species 2008: e.T14138A4407658. <https://doi.org/10.2305/IUCN.UK.2008.RLTS.T14138A4407658.en>. August 29, 2018.
- Iida, K., Kobayashi, R., Hengjan, Y., Nagata, N., Yonemitsu, K., Nunome, M., Kuwata, R., Suzuki, K., Ichiyana, K., Maeda, K., Ohmori, Y., Hondo, E., 2017. The genetic diversity of d-loop sequences in eastern bent-winged bats (*Miniopterus fuliginosus*) living in Wakayama prefecture, Japan. *J. Vet. Med. Sci.* 79, 1142–1145. <https://doi.org/10.1292/jvms.17-0152>.
- Jimba, M., Takeshima, S., Matoba, K., Endoh, D., Aida, Y., 2010. Blv-cocomo-qPCR: quantitation of bovine leukemia virus proviral load using the cocomo algorithm.

- Retrovirology 7, 91.
- King, A.M.Q., Brown, F., Christian, P., Hovi, T., Hyypiä, T., Knowles, N.J., Lemon, S.M., Minor, P.D., Palmenberg, A.C., Skern, T., Stanway, G., 2000. Picornaviridae. In: Van Regenmortel, M.H.V., Fauquet, C.M., Bishop, D.H.L., Calisher, C.H., Carsten, E.B., Estes, M.K., Lemon, S.M., Maniloff, J., Mayo, M.A., McGeoch, D.J., Pringle, C.R., Wickner, R.B. (Eds.), *Virus Taxonomy. Seventh Report of the International Committee for the Taxonomy of Viruses*. Academic Press, New York, pp. 657–673.
- King, A.M.Q., Lefkowitz, E.J., Mushegian, A.R., Adams, M.J., Dutilh, B.E., Gorbalenya, A.E., Harrach, B., Harrison, R.L., Junglen, S., Knowles, N.J., Kropinski, A.M., Krupovic, M., Kuhn, J.H., Nibert, M.L., Rubino, L., Sabanadzovic, S., Sanfacon, H., Siddell, S.G., Simmonds, P., Varsani, A., Zerbini, F.M., Davison, A.J., 2018. Changes to taxonomy and the international code of virus classification and nomenclature ratified by the international committee on taxonomy of viruses. *Arch. Virol.* 2018. <https://doi.org/10.1007/s00705-018-3847-1>.
- Lamglait, B., Joris, A., Romey, A., Bakkali-Kassimi, L., Lemberger, K., 2015. Fatal encephalomyocarditis virus infection in an african savanna elephant (*loxodonta africana*) in a french zoo. *J. Zoo Wildl. Med.* 46, 393–396. <https://doi.org/10.1638/2014-0132R1.1>.
- Lau, S.K., Woo, P.C., Lai, K.K., Huang, Y., Yip, C.C., Shek, C.T., Lee, P., Lam, C.S., Chan, K.H., Yuen, K.Y., 2011. Complete genome analysis of three novel picornaviruses from diverse bat species. *J. Virol.* 85, 8819–8828. <https://doi.org/10.1128/JVI.02364-10>.
- Lin, L.K., Motokawa, M., 2014. *Soricomorpha. Mammals of Taiwan Vol. 1. Center for Tropical Ecology and Biodiversity, Tunghai University, Taichung*, pp. 89.
- Liu, H., He, X., Song, X., Xu, L., Zhang, Y., Zhou, G., Zhu, W., Chang, C., Yin, Z., Shi, Y., Wang, C., Chang, H., 2016. Isolation and molecular and phylogenetic analyses of encephalomyocarditis virus from wild boar in central china. *Infect. Genet. Evol.* 40, 67–72. <https://doi.org/10.1016/j.meegid.2016.02.025>.
- Liu, H., Li, Y., Zhang, G., Sang, S., Wang, C., Chang, H., 2017. Complete genome sequences and phylogenetic analysis of encephalomyocarditis virus strains isolated from pigs and rats origin. *Infect. Genet. Evol.* 55, 277–280. <https://doi.org/10.1016/j.meegid.2017.09.032>.
- Luis, A.D., O’Shea, T.J., Hayman, D.T.S., Wood, J.L.N., Cunningham, A.A., Gilbert, A.T., Mills, J.N., Webb, C.T., 2015. Network analysis of host-virus communities in bats and rodents reveals determinants of cross-species transmission. *Ecol. Lett.* 18, 1153–1162. <https://doi.org/10.1111/ele.12491>.
- Lumsden, L.F., Bennett, A.F., 2005. Scattered trees in rural landscapes: foraging habitat for insectivorous bats in south-eastern australia. *Biol. Conserv.* 122, 205–222. <https://doi.org/10.1016/j.biocon.2004.07.006>.
- Luo, Y.K., Liang, L., Tang, Q.H., Zhou, L., Shi, L.J., Cong, Y.Y., Lin, W.C., Cui, S.J., 2017. Isolation and characterization of encephalomyocarditis virus from dogs in china. *Sci. Rep.* 7, 438. <https://doi.org/10.1038/s41598-017-00435-x>.
- Maurice, H., Thulke, H.H., Schmid, J.S., Stegeman, A., Nielen, M., 2016. The impact of compartmentalised housing on direct encephalomyocarditis virus (emcv) transmission among pigs; insight from a model. *Prev. Vet. Med.* 127, 105–112. <https://doi.org/10.1016/j.prevetmed.2016.03.006>.
- Mildenstein, T., De Jong, C., 2011. Natural history, ecological and socio-economic value of bats. In: Newman, S.H., Field, H., Epstein, J.H., De Jong, C.E. (Eds.), *Investigating the Role of Bats in Emerging Zoonoses 20* FAO Animal Production and Health, Rome.
- Mizutani, T., Nishino, Y.H.K., Takashima, I., 1999. Reverse transcription-nested polymerase chain reaction for detecting p40 rna of borna disease virus, without risk of plasmid contamination. *J. Vet. Med. Sci.* 61, 77–80.
- Rahman, S.A., Hassan, S.S., Olival, K.J., Mohamed, M., Chang, L.Y., Hassan, L., Saad, N.M., Shohaimi, S.A., Mamat, Z.C., Naim, M.S., Epstein, J.H., Suri, A.S., Field, H.E., Daszak, P., Ecology, Henipavirus, Research, G., 2010. Characterization of nipah virus from naturally infected pteropus vampyrus bats, Malaysia. *Emerg. Infect. Dis.* 16, 1990–1993. <https://doi.org/10.3201/eid1612.091790>.
- Roche, S.E., Costard, S., Meers, J., Field, H.E., Breed, A.C., 2015. Assessing the risk of nipah virus establishment in australian flying-foxes. *Epidemiol. Infect.* 143, 2213–2226. <https://doi.org/10.1017/S0950268813003336>.
- Ruedas, L., Heaney, L., Molur, S., 2016. *Rattus exulans*. The IUCN Red List of Threatened Species 2016. <https://doi.org/10.2305/IUCN.UK.2016-3.RLTS.T19330A22440938.en>. August 22, 2018.
- Schmidly, J.S., Bradley, R.D., 2016. *The Mammals of Texas*. University of Texas Press, United States of America.
- Smith, A.T., Johnston, C.H., 2008. *Lepus sinensis*. The IUCN Red List of Threatened Species 2008: e.T41286A10433753. <https://doi.org/10.2305/IUCN.UK.2008.RLTS.T41286A10433753.en>. August 29, 2018.
- Spinner, M.L., Di Giovanni, G.D., 2001. Detection and identification of mammalian reoviruses in surface water by combined cell culture and reverse transcription-pcr. *Appl. Environ. Microbiol.* 67, 3016–3020. <https://doi.org/10.1128/AEM.67.7.3016-3020.2001>.
- Spyrou, V., Maurice, H., Billinis, C., Papanastassopoulou, M., Psalla, D., Nielen, M., Koenen, F., Papadopoulos, O., 2004. Transmission and pathogenicity of encephalomyocarditis virus (emcv) among rats. *Vet. Res.* 35, 113–122. <https://doi.org/10.1051/vetres:2003044>.
- Tsai, K.J., Hsu, W.C., Chuang, W.C., Chang, J.C., Tu, Y.C., Tsai, H.J., Liu, H.F., Wang, F.I., Lee, S.H., 2016. Emergence of a sylvatic enzootic formosan ferret badger-associated rabies in taiwan and the geographical separation of two phylogenetic groups of rabies viruses. *Vet. Microbiol.* 182, 28–34. <https://doi.org/10.1016/j.vetmic.2015.10.030>.
- VanDevanter, D.R., Warren, P., Bennett, L., Schultz, E.R., Coulter, S., Garber, R.L., Rose, T.M., 1996. Detection and analysis of diverse herpesviral species by consensus primer pcr. *J. Clin. Microbiol.* 34, 1666–1671.
- Vansteenkiste, K., Van Limbergen, T., Decaluwe, R., Tignon, M., Cay, B., Maes, D., 2016. Clinical problems due to encephalomyocarditis virus infections in two pig herds. *Porcine Health Manage.* 2, 19. <https://doi.org/10.1186/s40813-016-0036-z>.
- Vercammen, F., Bosseler, L., Tignon, M., Cay, A.B., 2017. Encephalomyocarditis virus in a captive malayan tapir (*tapirus indicus*). *Open Vet. J.* 7, 100–103. <https://doi.org/10.4314/ovj.v7i2.4>.
- Vincenot, C., 2017. *Pteropus dasymallus*. The IUCN Red List of Threatened Species 2017: e.T18722A22080614. <https://doi.org/10.2305/IUCN.UK.2017-2.RLTS.T18722A22080614.en>. August 29, 2018.
- Wang, L.F., Shi, Z., Zhang, S., Field, H., Daszak, P., Eaton, B.T., 2006. Review of bats and sars. *Emerg Infect Dis* 12, 1834–1840. <https://doi.org/10.3201/eid1212.060401>.
- Wang, Q., Qi, J., Yuan, Y., Xuan, Y., Han, P., Wan, Y., Ji, W., Li, Y., Wu, Y., Wang, J., Iwamoto, A., Woo, P.C., Yuen, K.Y., Yan, J., Lu, G., Gao, G.F., 2014. Bat origins of mers-cov supported by bat coronavirus hku4 usage of human receptor cd26. *Cell Host Microbe* 16, 328–337. <https://doi.org/10.1016/j.chom.2014.08.009>.
- Ward, R.D., Zemlak, T.S., Innes, B.H., Last, P.R., Hebert, P.D., 2005. DNA barcoding australia’s fish species. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 360, 1847–1857. <https://doi.org/10.1098/rstb.2005.1716>.
- Weissenböck, H., Bakonyi, T., Sekulin, K., Ehrensperger, F., Doneley, R.J., Durrwald, R., Hoop, R., Erdelyi, K., Gal, J., Kolodziejek, J., Nowotny, N., 2009. Avian bornaviruses in psittacine birds from europe and australia with proventricular dilatation disease. *Emerg Infect Dis* 15, 1453–1459. <https://doi.org/10.3201/eid1509.090353>.
- Wellehan, J.F., Johnson, A.J., Harrach, B., Benkö, M., Pessier, A.P., Johnson, C.M., Garner, M.M., Childress, A., Jacobson, E.R., 2004. Detection and analysis of six lizard adenoviruses by consensus primer pcr provides further evidence of a reptilian origin for the atadenoviruses. *J. Virol.* 78, 13366–13369. <https://doi.org/10.1128/JVI.78.23.13366-13369.2004>.
- Wilson, D.E., Reeder, D.M., 2005. *Mammal Species of the World*. Johns Hopkins University Press.
- Wong, S., Lau, S., Woo, P., Yuen, K.Y., 2007. Bats as a continuing source of emerging infections in humans. *Rev. Med. Virol.* 17, 67–91. <https://doi.org/10.1002/rmv.520>.
- Wu, Z., Ren, X., Yang, L., Hu, Y., Yang, J., He, G., Zhang, J., Dong, J., Sun, L., Du, J., Liu, L., Xue, Y., Wang, J., Yang, F., Zhang, S., Jin, Q., 2012. Virome analysis for identification of novel mammalian viruses in bat species from chinese provinces. *J. Virol.* 86, 10999–11012. <https://doi.org/10.1128/JVI.01394-12>.
- Yassin, A., Markow, T.A., Narechania, A., O’Grady, P.M., DeSalle, R., 2010. The genus drosophila as a model for testing tree- and character-based methods of species identification using DNA barcoding. *Mol. Phylogenet. Evol.* 57, 509–517. <https://doi.org/10.1016/j.ympev.2010.08.020>.