

Gene expression

# VeTra: a tool for trajectory inference based on RNA velocity

Guangzheng Weng <sup>1,2,3</sup>, Junil Kim<sup>2,3,4,\*</sup> and Kyoung Jae Won<sup>2,3,\*</sup>

<sup>1</sup>Department of Biology, The Bioinformatics Centre, University of Copenhagen, 2200 Copenhagen N, Denmark, <sup>2</sup>Biotech Research and Innovation Centre (BRIC), University of Copenhagen, 2200 Copenhagen N, Denmark, <sup>3</sup>Novo Nordisk Foundation Center for Stem Cell Biology, DanStem, Faculty of Health and Medical Sciences, University of Copenhagen, 2200 Copenhagen N, Denmark and <sup>4</sup>Department of Bioinformatics, School of Systems Biomedical Science, Soongsil University, 06978 Seoul, South Korea

\*To whom correspondence should be addressed.

Associate Editor: Inanc Biro

Received on February 16, 2021; revised on April 11, 2021; editorial decision on April 29, 2021; accepted on May 10, 2021

## Abstract

**Motivation:** Trajectory inference (TI) for single cell RNA sequencing (scRNAseq) data is a powerful approach to interpret dynamic cellular processes such as cell cycle and development. Still, however, accurate inference of trajectory is challenging. Recent development of RNA velocity provides an approach to visualize cell state transition without relying on prior knowledge.

**Results:** To perform TI and group cells based on RNA velocity we developed VeTra. By applying cosine similarity and merging weakly connected components, VeTra identifies cell groups from the direction of cell transition. Besides, VeTra suggests key regulators from the inferred trajectory. VeTra is a useful tool for TI and subsequent analysis.

**Availability and implementation:** The VeTra is available at <https://github.com/wgzgithub/VeTra>.

**Contact:** [kyoung.won@bric.ku.dk](mailto:kyoung.won@bric.ku.dk) or [junil.kim@bric.ku.dk](mailto:junil.kim@bric.ku.dk)

**Supplementary information:** [Supplementary data](#) are available at *Bioinformatics* online.

## 1 Introduction

Trajectory analysis using single cell transcriptomics is useful to understand temporal transition of cell states. By showing transcriptional changes along the obtained trajectory (or pseudo-time), Trajectory inference (TI) has been applied to studying various biological processes including cell cycle, cancer development and cell differentiation (Tran and Bader, 2020).

Besides pseudo-temporal analysis, TI has been used to assist identifying gene regulatory rules. Visual inspection of gene expression changes along the inferred trajectory suggested potential regulators for the biological processes (Trapnell *et al.*, 2014). Systemic approaches have been developed to reconstruct gene regulatory networks (GRNs) using the gene expression changes along the inferred trajectory (Kim *et al.*, 2021; Matsumoto *et al.*, 2017). Well inferred trajectory helped improve the performance of GRN reconstruction (Qiu *et al.*, 2020).

At present, more than 70 methods have been published to infer trajectory from scRNAseq data (Saelens *et al.*, 2019). Majority of TI approaches have been developed based on the transcriptomic similarity among cells. Pseudo-time is obtained from the ordered cells based on the transcriptomic similarities (Saelens *et al.*, 2019). However, transcriptomic similarity cannot specify the initial and the terminal points of the trajectory. When using TI algorithms based on the transcriptomic similarity, a user has to provide the direction

of a cellular process, roots/terminals using the marker genes or any prior knowledge about experiments to obtain pseudo-time (Haghverdi *et al.*, 2016; Setty *et al.*, 2016; Trapnell *et al.*, 2014).

It is more challenging to infer trajectory when cell dynamics are complex. It is difficult to assign an accurate fate (lineage) to a cell around branching regions. To improve robustness in detecting TI, there have been attempts to confine the inferred trajectory to a set of fixed topological structures, e.g. linear, bifurcation and cycle (Saelens *et al.*, 2019). However, the use of pre-defined structures can restrict the opportunity to identify new structures.

Recently, RNA velocity has been suggested to analyze scRNAseq data by incorporating mRNA dynamics (La Manno *et al.*, 2018). Notably, RNA velocity can provide the direction and the speed of movement of individual cells and visualize the dynamic cell transitions without any prior knowledge. The direction information from RNA velocity provides a potential solution to determine the developmental trajectory of a cell around branching points. Therefore, RNA velocity can be useful in developing precise TI tools.

To obtain precise TI and subsequent analysis without prior information, we developed VeTra. VeTra identifies cell groups belonging to the same lineage and suggests potential regulators in the identified groups. Compared with previous undirected graph based methods such as minimum spanning tree (Street *et al.*, 2018) and graph-partitioning algorithm (Wolf *et al.*, 2019), VeTra builds directed graph


from RNA velocity. VeTra identifies isolated cell transition paths by searching for weakly connected components (WCC) which represent the coarse-grained structure of connected community (An *et al.*, 2004). The isolated paths are further grouped together using the hierarchical clustering algorithm to finally form potential development lineages. As the results, VeTra identifies cell transition trajectory and determines the memberships of cells to each trajectory even around branching points. VeTra further suggests key regulators along each trajectory by integrating the engine of TENET (Kim *et al.*, 2021). VeTra is a user-friendly tool for TI and subsequent analysis.

## 2 Materials and methods

### 2.1 VeTra groups the cells belonging to the same stream of trajectory

VeTra is an RNA velocity based TI tool that enables accurate lineage tracing and subsequent analysis for gene regulation. VeTra performs lineage tracing from the root to the terminal states by grouping cells based on the similarity in direction of cell transition. This enables VeTra to perform TI without prior knowledge or pre-defined lineage topology.

VeTra reconstructs the pseudo-temporal order of cells based on the coordinates and the velocity vector of cells in the low-dimensional embedding. The velocity vectors are estimated by extrapolating the spliced/unspliced read ratio to the local neighboring cells (La Manno *et al.*, 2018). Given velocity vectors (Fig. 1A), VeTra reconstructs multiple directed graphs. To link cells based on transition,  $k$  nearest neighbors of a cell with similar direction are selected using cosine similarity ( $\cos_1$ ) (Fig. 1B, Section 2). Among them, the nearby cell located upstream with the highest cosine similarity ( $\cos_2$ ) is selected (Fig. 1B, Section 2). Once all cells were investigated for their next transition, multiple directed graphs are obtained (Fig. 1C). To find a coarse-grained structure of the directed graph, VeTra identifies WCCs where every cell is reachable from every other cell regardless of the direction of relationships (Steven, 1990) (Fig. 1D). The WCCs are grouped together when they are similar and close each other (Section 2, Fig. 1E). Finally, we obtained the pseudo-time ordering of cell groups (Fig. 1F) by projecting the member cells onto the principal curve (Hastie and Stuetzle, 1989).



**Fig. 1.** VeTra reconstructs single-cell trajectories for multiple cell lineages. (A) A 2D embedding plot using the scRNAseq for pancreatic development. (B) Cosine similarity to search for the neighboring cells with similar direction.  $\cos_1$  finds the vectors with similar direction and  $\cos_2$  identifies the cell to transit from a cell. (C) The directed graph obtained by applying cosine similarity. (D) The WCCs obtained using all possible paths. (E) The grouped WCCs using a hierarchical clustering algorithm. (F) The pseudo-time for each lineage identified by VeTra

### 2.2 VeTra's trajectory matches well with the known lineage or biological process

We applied VeTra to infer the trajectory for various scRNAseq datasets with known cell dynamics structures for pancreatic development (Bastidas-Ponce *et al.*, 2019; Bergen *et al.*, 2020), chromaffin cell differentiation (Furlan *et al.*, 2017; La Manno *et al.*, 2018), neural lineages in the hippocampus (Hochgerner *et al.*, 2018; La Manno *et al.*, 2018) and cell cycle (Xia *et al.*, 2019). For benchmarking we used Slingshot (Street *et al.*, 2018), FateID (Herman and Grün, 2018) and PAGA (Wolf *et al.*, 2019) since they showed best performance in recent benchmarking tests (Saelens *et al.*, 2019). We also include CellRank (Bergen *et al.*, 2020) and CellPath (Zhang and Zhang, 2021a,b) as they are developed based on RNA velocity. We used Diffusion pseudo-time (DPT) to provide pseudo-time ordering to CellRank. We provided known roots and terminals cell to run Slingshot and FateID.


#### 2.2.1 Pancreatic development (a topology with crossing branches)

During the pancreatic development endocrine progenitors differentiate into alpha or beta cells (Bastidas-Ponce *et al.*, 2019). Ductal cells are originated from common progenitors, which are fated to endocrine and exocrine lineages (Reichert and Rustgi, 2011). From the transcriptome of pancreatic development (E15.5), VeTra identified three major trajectory groups; (i) alpha cell differentiation from endocrine progenitors (EP), (ii) beta/epsilon differentiation from EP and (iii) ductal development (Fig. 2A). The first two clusters were commonly originated from EPs and bifurcated into different lineages.

CellRank identified similar three trajectories with a slightly different starting point close to ductal cells for alpha and beta cell trajectories (Fig. 2A). FateID identified the trajectory to alpha and beta cells but failed in identifying ductal lineage. Slingshot did not find delta/beta cell lineage as well as ductal cell lineage (Fig. 2A). PAGA inferred coarse-grained trajectories from EP to alpha, beta, delta and epsilon cells but it also found trajectories between unrelated cell types (e.g. between alpha and beta cells) (Supplementary Fig. S1). CellPath did not find the path to ductal cells (Supplementary Fig. S2B).

#### 2.2.2 Chromaffin cell development (a topology with diverging branches)

During the development, the chromaffin cells and sympathoblasts cells are generated from Schwann cell precursors (SCPs) (Furlan *et al.*, 2017). VeTra, FateID and Slingshot successfully captured the



**Fig. 2.** Performance assessment of the TI tools using the scRNAseq datasets with various topologies. The trajectories inferred by VeTra, CellRank, FateID and Slingshot using the scRNAseq data for (A) pancreatic development, (B) chromaffin and sympathoblasts development, (C) cell cycle and (D) hippocampus development

branches toward sympathoblasts and chromaffin cells (Fig. 2B). CellRank identified SCPs as initial states but failed to identify proper terminal states (Fig. 2B). PAGA connected bridging cells presumably falsely as their fates are already committed to each branch (based on RNA velocity) (Supplementary Fig. S1). CellPath only identified the path to the chromaffin cells (Supplementary Fig. S2C).

### 2.2.3 Cell cycle (a cycling topology)

We applied TI tools on the scRNAseq dataset from U-2 OS cells (Xia *et al.*, 2019) and investigate the topology for cell cycle. VeTra identified a single circular trajectory, starting at G1 phase toward G1/S, G2/M and M phase, and finally going back to G1 phase (Fig. 2C). Given the root cells in G1 phase, Slingshot identified similar trajectory. On the other hand, CellRank and CellPath estimated a linear-like trajectory (Fig. 2C, Supplementary Fig. S2D). PAGA connected cell populations with cycle, but inferred wrong trajectories between separated cell cycles (Supplementary Fig. S1).

### 2.2.4 Hippocampus development (complex diverging branches)

We further evaluated the performance using the scRNAseq data for hippocampus development, which has been identified to have five diverging branches: astrocytes, oligodendrocyte progenitor cells (OPC), dentate gyrus granule neurons and pyramidal neurons including CA1, CA2, CA3 and subiculum (La Manno *et al.*, 2018). Among these five main branches, astrocytes and OPCs are commonly branched from radial glial cells and the other neuronal branch cells are originated from immature neuroblast cells (La Manno *et al.*, 2018).

VeTra identified five lineages toward CA1-subiculum, CA2-3-4, granule, astrocytes, OPC (Fig. 2D). VeTra also predicted that CA2-3-4 cells and granules share the same neuroblastic origin (Nbl1) and astrocytes and oligodendrocyte progenitors share the same intermediate progenitor origin (nIPC), which is consistent with the original RNA velocity results (La Manno *et al.*, 2018). CellRank identified one correct (Nbl1) and two incorrect (Granule, and CA1-Sub) initial states (Fig. 2D). The five terminal states estimated by CellRank were CA, CA2-3-4, ImmAstro, GliaProg and nIPC, including three potential misclassified intermediate states (GliaProg, nIPC and CA). Due to potential incorrect initial and terminal state assignment, the trajectory inferred by CellRank failed in distinguishing the five separated trajectories. Slingshot detected five lineages (Fig. 2D) but determined wrong initial states (always from ImmAstro). FateID did not infer any meaningful trajectories (Fig. 2D). PAGA inferred clear trajectories from Nbl1 to CA1-sub, CA2-3-4 and Granule. However, PAGA falsely connected ImmAstro and GlialProg (Supplementary Fig. S1). CellPath showed severely truncated paths for this dataset (Supplementary Fig. S2A).

### 2.3 Performance evaluation using simulated datasets

The assessment using scRNAseq data demonstrated the outstanding performance of VeTra. However, the gold standard annotation for individual cells is usually not provided and it is hard to quantify the performance. For quantitative assessment, we simulated scRNAseq datasets using Dyngen (Cannoodt *et al.*, 2020) and generated three lineage structures: binary tree, trifurcation and converging. We additionally included a disconnected structure from VeloSim (Zhang and Zhang, 2021a,b). Beside visualization, we calculated the matching score by obtaining the average value of the normalized hamming distances ( $D$ ) between the predicted and the known lineages. We used  $(1 - D) * 100$  as the score.

VeTra successfully identified the identified lineages for four simulated datasets. Slingshot identified successfully except for the disconnected path. However, other approaches were not successful compared with VeTra or Slingshot (Supplementary Fig. S3). The overall comparison (Supplementary Table S1) demonstrates the robustness of VeTra in TI.

### 2.4 VeTra provides condition-specific master regulator

It has been studied that TI can influence the performance of GRN reconstruction. VeTra is equipped with a function to suggest key regulators by adopting the engine of TENET (Kim *et al.*, 2021) that we previously developed to reconstruct GRN. TENET has a function to identify key regulators from the inferred causal relationships with their target genes (Kim *et al.*, 2021). Incorporating TENET, VeTra has a function to identify most influential transcription factors (TFs) for the inferred trajectory.

Using the RNAseq dataset from pancreas development (Bastidas-Ponce *et al.*, 2019), VeTra found key regulators for the three predicted major branches: alpha, beta and ductal (Fig. 3A). The key predicted regulators that distinguishing alpha cell development against others were Pax6, and Arx. Pax6 and Arx have been known to have a role during alpha cell development (Ashery-Padan *et al.*, 2004; Gosmain *et al.*, 2011). Pax6 regulates genes related to glucagon production (Ashery-Padan *et al.*, 2004). Arx has been shown to play a similar role in alpha-cell development (Gosmain *et al.*, 2011). VeTra also identified Neurog3 an endocrine (alpha and beta) specific key regulator (Gradwohl *et al.*, 2000; Gu *et al.*, 2002; Schwitzgebel *et al.*, 2000).

In addition, VeTra revealed condition-specific regulators for the five lineages of hippocampus development (Fig. 3B). For instance, Nr4a3 (Nor-1) is predicted a CA2-3-4 specific regulator by VeTra. Nor-1 deficient mice appear to have abnormal pyramidal cell layer in CA1 to CA3 (Pönniö and Conneely, 2004).

## 3 Algorithm

To pick the most appropriate cell to which the vector of the cell  $i$  is pointing,  $k$  closest neighbor cells are collected from the head of the vector of a cell  $i$  in the low-dimensional space (Fig. 1B). Among the neighbor cells, cells with similar direction are selected using a cosine similarity criterion between the cell  $i$  and the neighbor cell  $j$  ( $\cos_{1ij} = \frac{v_i \cdot v_j}{\|v_i\| \|v_j\|} > 0.5$ ) (Fig. 1B, where  $v_i$  and  $v_j$  denote velocity vectors (2D coordinates) of cell  $i$  and cell  $j$ ). The cell with the highest similarity for  $\cos_{2ij} (= \frac{\delta_{ij}}{\|v_i\| \|v_j\|})$ , where  $\delta_{ij}$  is the vector from  $v_i$  to  $v_j$ ) located upstream of the cell  $i$  is finally selected to obtain a directed graph in the same stream of trajectory (Fig. 1B).




Fig. 3. Key regulators for each trajectory identified by VeTra. (A) The key regulators identified from the trajectories for ductal, alpha and beta cell development. (B) The key regulators identified from the five developmental trajectories in mouse hippocampus

The hierarchical clustering is applied to the WCCs for further grouping (Fig. 1E). The distance between the two sub-graphs is defined by the maximum distance of all the closest pairs of cells. To calculate the distance between cells, we calculated the Euclidean distance in the four-dimensional space (two dimensions using the gene expression and the other two using the velocity vector in the reduced dimensional space). To obtain full trajectory (from the root to the branch), we extended the memberships if a cell located nearby is similar ( $\cos_{1ij} > 0.7$ ). We obtained the pseudo-time ordering by projecting the member cells onto the principal curve (Hastie and Stuetzle, 1989).

## 4 Discussion

TI is a widely used approach to understand temporal dynamics of cells from scRNAseq data. A number of approaches have already been developed to infer the trajectories. As shown in the examples using the simulated as well as real scRNAseq data (Fig. 2 and Supplementary Figs S1–S3), however, there is still room for algorithmic improvement for accurate detection of trajectory. We have noticed that the results of current TI tools do not always match with the cell dynamics observed by RNA velocity (La Manno et al., 2018). RNA velocity, by providing the direction of cell transition, enabled visual inspection of cell trajectory. VeTra is designed to group cells in accordance with the cell transition drawn by RNA velocity. To fully use the vector space provided by RNA velocity, VeTra used cosine similarity and obtained cell groups. The use of cosine similarity makes VeTra rely on the vector space for TI.

Our benchmarking test on the scRNAseq datasets exemplifies that VeTra successfully identified the trajectory drawn by RNA velocity and produced reasonable results that matched with the known topological structures for cell development and biological processes. It is interesting that VeTra outperformed other RNA velocity based methods such as CellRank (Bergen et al., 2020) and CellPath (Zhang and Zhang, 2021a,b) in our tests. Instead of using RNA velocity directly, CellRank calculates the probabilities of each cell transition to all terminals from initials. Therefore, the terminal states are not deterministic for CellRank. This may have influenced the performance of CellRank. CellPath uses meta-cells to obtain smoothed RNA velocity and reduce the computing cost (Zhang and Zhang, 2021a,b). In our test, CellPath has failed to annotate large number of cells for each lineage. A meta-cell is a cluster of cells. We suspect that meta-cells may smooth cell direction too much when cells exhibit diverse direction. Compared with them, VeTra tried to follow the velocity vector using cosine similarity without further processing them.

As the results, VeTra relies on RNA velocity calculation. RNA velocity can heavily depend on the correct measurement of spliced and unspliced RNA abundance. A recent study showed that the choice of experimental setting can change the results of RNA velocity (Soneson et al., 2021). As a downstream analysis tool of RNA velocity, VeTra can use the output from Velocyto (La Manno et al., 2018) or scVelo (Bergen et al., 2020). A previous study shows that the results of Velocyto and scVelo can be different (Cannoodt et al., 2020), which can also cause different results for VeTra. We investigated VeTra's output after running Velocyto and scVelo. For the cell cycle and chromaffin development datasets, both Velocyto and scVelo showed similar results. However, we found different results for the hippocampus and the pancreas development datasets (Supplementary Fig. S4). For VeTra, CellRank and CellPath, we selected the RNA velocity approach that produces the better performance for each dataset.

VeTra is equipped with the tool to predict key regulators by adopting the gear of TENET, a GRN reconstructor based on TE. It was discussed previously that TENET is capable of detecting key regulators using the aligned scRNAseq (Kim et al., 2021). VeTra equipped with the TENET engine will be useful for downstream analysis of TI.

VeTra requires the 2D coordinates embedded from expression profile and the RNA velocity vectors for each cell. A user can specify the number of groups. The output files include (i) 2D embedding

figures colored by grouping of cells, (ii) 2D embedding figures colored by pseudo-time ordering, (iii) lists of selected cells for each group and (iv) the pseudo-time ordering information for each group.

## Funding

This work was supported by The Novo Nordisk Foundation Center for Stem Cell Biology [NNF17CC0027852]; Lundbeck Foundation's Ascending Investigator [R313-2019-421]; and the Independent Research Fund Denmark [0135-00243B].

*Conflict of Interest:* none declared.

## References

- An, Y. et al. (2004) Characterizing and mining the citation graph of the computer science literature. *Knowledge Inf. Syst.*, 6, 664–678.
- Ashery-Padan, R. et al. (2004) Conditional inactivation of Pax6 in the pancreas causes early onset of diabetes. *Dev. Biol.*, 269, 479–488.
- Bastidas-Ponce, A. et al. (2019) Comprehensive single cell mRNA profiling reveals a detailed roadmap for pancreatic endocrinogenesis. *Development*, 146, dev173849.
- Bergen, V. et al. (2020) Generalizing RNA velocity to transient cell states through dynamical modeling. *Nat. Biotechnol.*, 38, 1408–1414.
- Cannoodt, R. et al. (2020) dyngen: a multi-modal simulator for spearheading new single-cell omics analyses. *BioRxiv*. Doi:10.1101/2020.02.06.936971
- Furlan, A. et al. (2017) Multipotent peripheral glial cells generate neuroendocrine cells of the adrenal medulla. *Science*, 357, eaal3753.
- Gosmain, Y. et al. (2011) Glucagon gene expression in the endocrine pancreas: the role of the transcription factor Pax6 in  $\alpha$ -cell differentiation, glucagon biosynthesis and secretion. *Diabetes Obesity Metab.*, 13, 31–38.
- Gradwohl, G. et al. (2000) neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas. *Proc. Natl. Acad. Sci. USA*, 97, 1607–1611.
- Gu, G. et al. (2002) Direct evidence for the pancreatic lineage: NGN3+ cells are islet progenitors and are distinct from duct progenitors. *Development*, 129, 2447–2457.
- Haghverdi, L. et al. (2016) Diffusion pseudotime robustly reconstructs lineage branching. *Nat. Methods*, 13, 845–848.
- Hastie, T. and Stuetzle, W. (1989) Principal curves. *J. Am. Stat. Assoc.*, 84, 502–516.
- Herman, J.S. and Grün, D. (2018) FateID infers cell fate bias in multipotent progenitors from single-cell RNA-seq data. *Nat. Methods*, 15, 379.
- Hochgerner, H. et al. (2018) Conserved properties of dentate gyrus neurogenesis across postnatal development revealed by single-cell RNA sequencing. *Nat. Neurosci.*, 21, 290–299.
- Kim, J. et al. (2021) TENET: gene network reconstruction using transfer entropy reveals key regulatory factors from single cell transcriptomic data. *Nucleic Acids Res.*, 49, e1.
- La Manno, G. et al. (2018) RNA velocity of single cells. *Nature*, 560, 494–498.
- Matsumoto, H. et al. (2017) SCODE: an efficient regulatory network inference algorithm from single-cell RNA-Seq during differentiation. *Bioinformatics*, 33, 2314–2321.
- Pönniö, T. and Conneely, O.M. (2004) nor-1 regulates hippocampal axon guidance, pyramidal cell survival, and seizure susceptibility. *Mol. Cell. Biol.*, 24, 9070–9078.
- Qiu, X. et al. (2020) Inferring causal gene regulatory networks from coupled single-cell expression dynamics using Scribe. *Cell Syst.*, 10, 265–274.e211.
- Reichert, M. and Rustgi, A.K. (2011) Pancreatic ductal cells in development, regeneration, and neoplasia. *J. Clin. Investig.*, 121, 4572–4578.
- Saelens, W. et al. (2019) A comparison of single-cell trajectory inference methods. *Nat. Biotechnol.*, 37, 547–554.
- Schwitzgebel, V.M. et al. (2000) Expression of neurogenin3 reveals an islet cell precursor population in the pancreas. *Development*, 127, 3533–3542.
- Setty, M. et al. (2016) Wishbone identifies bifurcating developmental trajectories from single-cell data. *Nat. Biotechnol.*, 34, 637–645.
- Soneson, C. et al. (2021) Preprocessing choices affect RNA velocity results for droplet scRNA-seq data. *PLoS Comput. Biol.*, 17, e1008585.
- Street, K. et al. (2018) Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics*, 19, 1–16.
- Tran, T.N. and Bader, G.D. (2020) Tempora: cell trajectory inference using time-series single-cell RNA sequencing data. *PLoS Comput. Biol.*, 16, e1008205.

- Trapnell, C. *et al.* (2014) The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat. Biotechnol.*, **32**, 381–386.
- Steven, S. (1990) *Implementing Discrete Mathematics: Combinatorics and Graph Theory with Mathematica*. Addison-Wesley.
- Wolf, F.A. *et al.* (2019) PAGA: graph abstraction reconciles clustering with trajectory inference through a topology preserving map of single cells. *Genome Biol.*, **20**, 59–59.
- Xia, C. *et al.* (2019) Spatial transcriptome profiling by MERFISH reveals sub-cellular RNA compartmentalization and cell cycle-dependent gene expression. *Proc. Natl. Acad. Sci. USA*, **116**, 19490–19499.
- Zhang, Z. and Zhang, X. (2021a) Inference of high-resolution trajectories in single cell RNA-Seq data from RNA velocity. *bioRxiv*, doi:<https://doi.org/10.1101/2020.09.30.321125>
- Zhang, Z. and Zhang, X. (2021b) VeloSim: simulating single cell gene-expression and RNA velocity. *bioRxiv*. doi:<https://doi.org/10.1101/2021.01.11.426277>