

Tg2576 Cortical Neurons That Express Human Ab Are Susceptible to Extracellular A β -Induced, K⁺ Efflux Dependent Neurodegeneration

Shannon Ray¹, Claire Howells¹, Emma D. Eaton¹, Chris W. Butler¹, Lana Shabala¹, Paul A. Adlard², Adrian K. West¹, William R. Bennett¹, Gilles J. Guillemin³, Roger S. Chung^{1*}

1 NeuroRepair Group, Menzies Research Institute, University of Tasmania, Hobart, Tasmania, Australia, **2** Synaptic Neurobiology Lab, Mental Health Research Institute, Melbourne, Victoria, Australia, **3** Neuroinflammation Group, University of New South Wales, Sydney, New South Wales, Australia

Abstract

Background: One of the key pathological features of AD is the formation of insoluble amyloid plaques. The major constituent of these extracellular plaques is the beta-amyloid peptide (A β), although A β is also found to accumulate intraneuronally in AD. Due to the slowly progressive nature of the disease, it is likely that neurons are exposed to sublethal concentrations of both intracellular and extracellular A β for extended periods of time.

Results: In this study, we report that daily exposure to a sublethal concentration of A β ₁₋₄₀ (1 μ M) for six days induces substantial apoptosis of cortical neurons cultured from Tg2576 mice (which express substantial but sublethal levels of intracellular A β). Notably, untreated Tg2576 neurons of similar age did not display any signs of apoptosis, indicating that the level of intracellular A β present in these neurons was not the cause of toxicity. Furthermore, wildtype neurons did not become apoptotic under the same chronic A β ₁₋₄₀ treatment. We found that this apoptosis was linked to Tg2576 neurons being unable to maintain K⁺ homeostasis following A β treatment. Furthermore, blocking K⁺ efflux protected Tg2576 neurons from A β -induced neurotoxicity. Interestingly, chronic exposure to 1 μ M A β ₁₋₄₀ caused the generation of axonal swellings in Tg2576 neurons that contained dense concentrations of hyperphosphorylated tau. These were not observed in wildtype neurons under the same treatment conditions.

Conclusions: Our data suggest that when neurons are chronically exposed to sublethal levels of both intra- and extracellular A β , this causes a K⁺-dependent neurodegeneration that has pathological characteristics similar to AD.

Citation: Ray S, Howells C, Eaton ED, Butler CW, Shabala L, et al. (2011) Tg2576 Cortical Neurons That Express Human Ab Are Susceptible to Extracellular A β -Induced, K⁺ Efflux Dependent Neurodegeneration. PLoS ONE 6(4): e19026. doi:10.1371/journal.pone.0019026

Editor: Malu G. Tansey, Emory University, United States of America

Received: February 17, 2011; **Accepted:** March 13, 2011; **Published:** April 27, 2011

Copyright: © 2011 Ray et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by research grants from the Australian Research Council (ARC, ID# DP0556630; DP0984673; LP0774820), National Health and Medical Research Council (ID# 490025, 544913) and Jack & Ethel Goldin Foundation. RSC holds an ARC Research Fellowship. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: rschung@utas.edu.au

Introduction

Alzheimer's disease (AD) is characterised by profound synaptic loss and neuronal death, and the accumulation of a number of key pathological hallmarks; senile plaques, dystrophic neurites and neurofibrillary tangles [1]. The β -amyloid peptide (A β) is the principle component of plaques, and is thought to contribute significantly to the pathogenesis of the disease [2]. However, the precise mechanisms that underlie the role of A β in AD are not clearly understood.

The localisation of A β is likely to have an important role in governing its toxic actions upon neurons. In this regard, it is well known that acute extracellular administration of aggregated forms of A β (and in particular oligomers) to cultured neurons is neurotoxic [3]. This is in accordance with the presence of amyloid plaques in AD, of which extracellular, aggregated forms of A β are the major constituent [4]. However, a growing body of evidence suggests that intraneuronal localisation of A β may also play a significant role in AD. For example, A β accumulates in processes

and synapses prior to, and with the onset of extracellular A β plaque formation [5,6], and in transgenic mice that develop A β plaques [7]. There is also some evidence that cognitive impairment in AD patients does not always correlate to the level of A β plaque deposition [8]. Similarly A β immunisation studies in Tg2576 [9] or PDAPP [10] transgenic mice reversed memory loss, but had no impact upon amyloid plaque levels. These studies suggest that the intraneuronal accumulation of A β may be important in disease progression and symptom onset.

Indeed, intracellular A β appears to increase the susceptibility of neurons to neurodegeneration. For example, Abdul *et al* [11] reported that cortical neurons cultured from APP/PS1 transgenic mice were more vulnerable to oxidative stress, mitochondrial dysfunction and apoptosis. Yao and colleagues [12] have demonstrated that 3xTg-AD mice exhibit increased hydrogen peroxide production and lipid peroxidation. Furthermore, hippocampal neurons cultured from these mice exhibited significantly decreased mitochondrial respiration and increased glycolysis [12]. Notably, cultured neurons transfected with constructs expressing

APP that contains familial-linked AD mutations that substantially increase levels of A β are also susceptible to apoptosis-inducing treatments [13]. Others have linked intracellular A β directly to apoptosis, reporting that transfection of constructs expressing A β into neuroblastoma cells resulted in activation of a P53-dependent apoptotic pathway [14]. A similar outcome has been observed when synthetic A β peptides were microinjected into neurons, which induced cytotoxicity via a p53-Bax apoptotic pathway [15]. Notably, exogenously applied A β is rapidly internalised by cultured neurons, where it could subsequently act in a neurotoxic manner like endogenous A β . Uptake of A β by neuronal cells occurs via the low-density lipoprotein receptor LRP1 [16]. Treatment of PC12 cells [17] or primary neuron cultures [18] with exogenous A β leads to accumulation of reactive oxygen species, and a decrease in redox activity and ATP levels [17]. Treatment with alpha-tocopherol (Vitamin E) can block these A β -induced changes in neuronal cells (see reviews by [19,20]).

AD is a progressive disease, in which neurons are likely to be exposed to both intracellular and extracellular A β at sublethal concentrations for extended periods of time. To experimentally model this situation, we have exposed cultured neurons from Tg2576 mice (which accumulate substantial amounts of intracellular A β) to daily treatment with exogenous A β ₁₋₄₀ for 6 days. Chronic A β ₁₋₄₀ treatment induced substantial apoptosis of cortical Tg2576 neurons but not wildtype neurons, suggesting that both intra- and extra-cellular A β are required to induce apoptosis. Apoptosis was linked to an inability of Tg2576 neurons to maintain K⁺ homeostasis following acute treatment with extracellular A β ₁₋₄₀. Chronic exposure to 1 μ M A β ₁₋₄₀ also caused the generation of hyperphosphorylated tau-immunoreactive axonal swellings in Tg2576 but not wildtype neurons. Our data suggest that chronic exposure to both intra- and extra-cellular A β induces neurodegenerative changes that bear similarities to some of the pathological hallmarks of AD.

Results

Uptake of exogenous A β by cultured transgenic Tg2576 and wildtype cortical neurons

Mouse cortical neurons (wildtype and Tg2576) were maintained in culture for seven days *in vitro* (DIV), at which time they were treated with 10 μ M soluble monomeric A β ₁₋₄₀. After 24 hours, neurons were fixed and A β detected by immunostaining. In untreated Tg2576 cortical neurons, A β was distributed within the cytoplasm and processes, but generally not in the nucleus (Figure 1A). In untreated wildtype cortical neurons, there was no A β detected (results not shown). In wildtype neurons treated with A β , A β was detected in a punctate distribution within the cytoplasm and processes (Figure 1B). When A β was applied to Tg2576 neurons, the distribution of A β resembled both of these scenarios, with both punctate and non-punctate regions of A β immunoreactivity observed within the cytoplasm and nucleus (Figure 1C). When fluorescently tagged A β ₁₋₄₀ (10 μ M) was applied to either wildtype or Tg2576 cortical neurons, we did not observe any difference in neuronal uptake or distribution of A β (results not shown).

Intraneuronal A β increases the vulnerability of cortical neurons to neurotoxicity induced by extracellular A β

7DIV mouse cortical neurons (wildtype and Tg2576) were treated with soluble monomeric A β ₁₋₄₀ (1–10 μ M) daily for a period of six days. Measurement of neuronal viability by an Alamar Blue assay revealed that while daily treatment with 1 μ M A β ₁₋₄₀ was not toxic to wildtype cortical neurons at any point in

the six day timecourse, this treatment resulted in a significant reduction in intracellular metabolism of Tg2576 neuronal cultures by approximately 30% after six days of continual A β ₁₋₄₀ treatment (Figure 2A). 10 μ M A β ₁₋₄₀ was mildly neurotoxic to wildtype neurons, resulting in approximately 20% cell death after seven days of treatment (Figure 2B). However, Tg2576 neurons were far more susceptible to A β ₁₋₄₀ (Figure 2B). The neurotoxic actions of chronic exposure to A β ₁₋₄₀ upon Tg2576 cortical neurons were confirmed in propidium iodide uptake studies (Figure 2C), whereby only dying cells internalise and incorporate propidium iodide in the nucleus. These results suggest that the non-toxic accumulation of intraneuronal A β in Tg2576 cortical neurons increases their vulnerability to subsequent neurotoxicity induced by chronic exposure to normally sublethal (1–10 μ M) levels of extracellular A β . Notably, in all cases when either wildtype or Tg2576 neurons were treated with vehicle alone, no change in viability was observed (results not shown).

To investigate whether chronic exposure to both intra- and extracellular A β had initiated an apoptotic pathway of cell death, immunostaining for activated caspase-3 was performed. Almost no caspase-3 labelled cells were observed in 14 DIV Tg2576 neuronal cultures (Figure 3A). However, when 7 DIV Tg2576 neurons were treated with 1 μ M A β ₁₋₄₀ for six consecutive days, caspase-3 immunoreactivity correlated with condensed or fragmented nuclei in approximately 30% of cells (Figure 3B), and direct counting found that the number of caspase-3 labelled neurons was equivalent to the number of neurons that incorporated propidium iodide (results not shown), indicating that the combination of both intra- and extraneuronal A β triggered an apoptotic pathway of neuronal death.

Treatment with A β causes K⁺ flux-dependent neurotoxicity in Tg2576 neurons

There are a number of reports suggesting that extracellular A β triggers changes in ionic homeostasis of neurons, and that these changes contribute directly to neurotoxicity. To investigate whether intracellular A β alters the ability of neurons to maintain ionic homeostasis following extracellular A β treatment, we used a novel non-invasive microelectrode ion flux (MIFE) measuring technique. Using the MIFE approach, we directly observed that A β treatment triggered rapid efflux of K⁺ from wildtype neurons (Figure 4A), which returned to homeostasis within 10 minutes after A β ₁₋₄₀ treatment. However, K⁺ flux in Tg2576 neurons treated with A β ₁₋₄₀ did not return to homeostasis (Figure 4B). Instead, transgenic neurons exhibited a continual efflux of potassium for more than 120 minutes after A β ₁₋₄₀ treatment (Figure 5A). Measurement of total K⁺ flux over 25 minutes following A β ₁₋₄₀ treatment revealed that significantly more potassium was extruded from Tg2576 neurons than wildtype neurons (Figure 4C). Interestingly, continuous treatment of wildtype neurons for three days with 1 μ M A β ₁₋₄₀ did not alter their ability to maintain K⁺ homeostasis following A β treatment (results not shown).

Concurrently, we also measured H⁺ flux of neurons in response to A β ₁₋₄₀ treatment. Wildtype neurons treated with A β ₁₋₄₀ demonstrated a rapid influx of H⁺, which stabilised within 5 minutes and remained stable for the entire recording period (Figure 4D). Tg2576 neurons displayed a similar response to A β ₁₋₄₀ (Figure 4E). However, from about 10 minutes post-treatment, Tg2576 neurons underwent a slow gradual influx of H⁺ (Figure 4E). The latter was further increased during the next 120 min post-treatment with A β ₁₋₄₀ (Figure 5B). Measurement of total H⁺ flux over 25 minutes after A β treatment revealed that A β ₁₋₄₀ induced significantly greater influx of H⁺ into Tg2576 neurons in comparison to wildtype neurons (Figure 4F).




Figure 1. Uptake of soluble A β by wildtype and Tg2576 cortical neurons *in vitro*. Wildtype and Tg2576 cortical neurons were treated with 10 μ M of monomeric A β ₁₋₄₀, and immunostained for A β after 24 hours. In untreated Tg2576 neurons, A β was smoothly distributed throughout the cytoplasm and processes of all neurons (A). When A β ₁₋₄₀ was applied to wildtype neurons, it was internalised and distributed in a punctate manner within the cytoplasm and processes (B). Notably, not all wildtype neurons internalised A β ₁₋₄₀ (B). When A β ₁₋₄₀ was applied to Tg2576 neurons, both smooth and punctately distributed A β was detected within neurons (C). scale bar = 25 μ m. doi:10.1371/journal.pone.0019026.g001

It is generally accepted that excessive potassium efflux is a key early step in apoptosis. To confirm that prolonged extrusion of K⁺ is responsible for A β -induced neurotoxicity in Tg2576 neurons, cells were pre-treated with the classic K⁺ channel blocker 4-aminopyridine (4-AP) prior to A β treatment. This resulted in a significant reduction of A β -induced neuronal death in Tg2576 neurons (Table 1).

The presence of both intra- and extracellular A β causes the formation of axonal swellings in cultured neurons

Immunolabelling of the axonal cytoskeleton (tau) revealed that there was no discernible difference in axonal morphology between wildtype (Figure 6A) and Tg2576 cortical neurons (Figure 6B) over 7-14 DIV. Daily treatment of 7DIV wildtype neurons with 1 μ M A β ₁₋₄₀ over the six day experimental timecourse did not markedly alter axonal morphology (Figure 6C). However, substantial changes in tau-labelling were observed in A β ₁₋₄₀ treated Tg2576 neurons. Treatment of Tg2576 neurons with 1 μ M A β ₁₋₄₀ resulted in a substantial increase in intensity of tau immunostaining after 24 hours (Figure 6D). After two daily treatments of Tg2576 neurons with 1 μ M A β ₁₋₄₀, blebbing and axonal fragmentation was apparent (Figure 6D), which worsened after four continuous days of A β ₁₋₄₀ treatment (Figure 6D). Furthermore, after four

consecutive days of 1 μ M A β ₁₋₄₀ treatment, a number of axonal swellings were observed in Tg2576 neuron cultures (Figure 6E), but never in wildtype neurons treated with 1 μ M A β ₁₋₄₀ for the same period of time (results not shown). These intra-axonal swellings were highlighted by dense accumulations of the microtubule-associated protein tau, including dense accumulations of hyperphosphorylated (AT-8 immunoreactive) tau (Figure 6E).

Discussion

AD is a progressive neurodegenerative disease, in which neurons are likely to be exposed to sublethal concentrations of both intracellular and extracellular A β for extended periods of time. To experimentally model this situation, Tg2576 neurons (which accumulate substantial amounts of intracellular A β) and wildtype neurons received daily treatment with soluble, monomeric A β for 6 days. While this chronic exposure to A β did not kill wildtype neurons, it caused substantial apoptosis of Tg2576 neurons. Further studies revealed that Tg2576 neurons were unable to maintain K⁺ and H⁺ homeostasis following A β treatment, leading to prolonged extrusion of potassium and influx of protons into Tg2576 neurons. Furthermore, chronic exposure to 1 μ M A β for six days caused the generation of hyperphosphorylated tau-immunoreactive axonal swellings in Tg2576 but




Figure 2. Tg2576 cortical neurons are more vulnerable to soluble A β -induced neurotoxicity. Wildtype and Tg2576 cortical neurons were treated daily with 1 μ M (A) or 10 μ M (B) of monomeric A β ₁₋₄₀ for 6 days, and neuronal viability (intracellular metabolism as assessed by Alamar Blue assay) assessed every 24 hours. At 1 μ M concentrations, only Tg2576 neurons were vulnerable to A β ₁₋₄₀, resulting in approximately 30% cell death after 6 days (A). 10 μ M A β ₁₋₄₀ was mildly toxic to wildtype neurons over the experimental timecourse, but killed 40% of Tg2576 neurons after 6 days (B). The Alamar Blue neurotoxicity assay produced similar results to direct counting of dying cells via propidium iodide uptake following treatment with 10 μ M A β ₁₋₄₀ (C). * - $p < 0.05$, ANOVA. Error bars represent standard error values from at least three replicates per experimental condition. This graph is representative of the results observed from 4 different experiments. doi:10.1371/journal.pone.0019026.g002

not wildtype neurons. In summary, our data suggest that chronic exposure to sublethal levels of both intra- and extra-cellular A β induces neurodegenerative changes in cultured neurons that bear similarities to pathological hallmarks observed in AD. These changes appear to be driven by an inability of Tg2576 to maintain normal K⁺ homeostasis in response to continual exposure to extracellular A β .

The mechanism by which A β causes neurotoxicity or neuronal dysfunction remains to be fully resolved. Numerous studies have used cultured neurons to investigate the neurotoxic actions of A β , and can generally be classified into two paradigms; experiments whereby A β is applied acutely or chronically to cultured neurons (testing the effect of extracellular A β upon neurons), and experiments using neurons cultured from transgenic AD mice which express human A β (to test the effect of intracellular A β upon neurons). The former experiments have been particularly informative, revealing important information regarding the concentration and biochemical form of extracellular A β that exhibits toxicity upon cultured neurons; from such studies it is proposed that soluble oligomeric forms of A β at $>5 \mu$ M concentrations are the most toxic form of extracellular A β to neurons [3]. In a similar manner, experiments using neurons cultured from transgenic AD mice have reported that intraneuronal A β increases the vulnerability of neurons to stressful cellular environments such as excitotoxicity and oxidative stress [11]. In some cases, the intraneuronal expression of A β itself can trigger apoptosis of neurons via a p53-dependent mechanism [21,14]. It is worth noting that one simplistic manner in which exogenous A β may induce neurodegenerative changes in Tg2576 neurons is due to uptake of A β via LRP1 [16], which increases intracellular A β levels above a threshold level leading to decreased viability and alterations in tau distribution and phosphorylation. However, two pieces of evidence argue against this possibility. Firstly, the distribution of endogenous A β and internalised A β in Tg2576 is quite different, suggesting that these two pools of A β are not able to act in the same manner. And secondly, our MIFE studies

demonstrate rapid and direct changes in ionic homeostasis of Tg2576 neurons triggered by application of exogenous A β , indicating that exogenously applied A β is probably acting in a different manner to intracellular endogenous A β .

While the studies discussed above have provided important information regarding the effect of extracellular and intracellular A β upon cultured neurons, it is important to consider that AD is a progressive condition, in which neurons are likely to be continuously exposed to sublethal concentrations of both intracellular and extracellular A β . To more accurately model this situation, Tg2576 neurons (which accumulate A β intraneuronally) were cultured in the presence of extracellular A β . This combination of intra- and extracellular A β induced caspase-3 dependent apoptosis of Tg2576 but not wildtype neurons. To elucidate the mechanisms underlying this, we used a non-invasive MIFE approach to observe changes in net ion flux of K⁺ and H⁺ ions in response to A β . Using this approach, we were able to continuously observe net ion flux of K⁺ and H⁺ for more than 2 hours. We found that A β treatment of wildtype neurons caused an immediate efflux of K⁺, which gradually returned to homeostasis within 10 minutes. However, we found that A β -treated Tg2576 neurons were unable to maintain K⁺ homeostasis, leading to prolonged leakage of K⁺ out of neurons. K⁺ efflux from cells is a key early initiator of apoptosis, as a low potassium intracellular microenvironment assists apoptosome formation and the activation of caspases and endonucleases. This suggests that the prolonged extrusion of K⁺ from A β -treated Tg2576 neurons was the cause of apoptosis in this study. This was confirmed by undertaking the same experiments in the presence of the K⁺ channel blocker 4-AP, which provided significant protection against A β -induced toxicity in Tg2576 neurons.

A novel element to the MIFE approach in this application is that it allows measurement of total flux of K⁺ into- or out- of cells, rather than the flow of K⁺ through particular channels/transporters that is observed through patch-clamp recording (electrogenic transporters) and pharmacological inhibitor studies.




Figure 3. Soluble A β triggers caspase-3 expression in Tg2576 cortical neurons. Tg2576 neurons cultured for 14 days *in vitro* (DIV) showed no signs of caspase-3 activation (A). However, when 7 DIV Tg2576 neurons were treated with 1 μ M A β ₁₋₄₀ daily for 6 days, a substantial number of neurons were found to express caspase-3 (B). doi:10.1371/journal.pone.0019026.g003




Figure 4. Soluble A β induces rapid efflux of K⁺ in Tg2576 cortical neurons. Treatment with 1 μ M A β_{1-40} triggered rapid efflux of K⁺ from wildtype neurons (A), which returned to homeostasis within 10 minutes. However, Tg2576 neurons treated with A β_{1-40} displayed a continual efflux of K⁺ over the recording period (B). Measurement of total K⁺ flux over 25 minutes following A β_{1-40} treatment revealed that significantly more potassium was extruded from Tg2576 neurons than wildtype neurons (C). A β_{1-40} treatment caused a rapid influx of H⁺ in both wildtype (D) and Tg2576 (E) neurons, which stabilised within 5 minutes. However, from about 10 minutes post-treatment, Tg2576 neurons underwent a slow gradual influx of H⁺ (E). Measurement of total H⁺ flux over 25 minutes revealed that A β_{1-40} induced significantly greater influx of H⁺ into Tg2576 neurons in comparison to wildtype neurons (F). * - $p < 0.05$, t-test. Error bars represent standard error values. doi:10.1371/journal.pone.0019026.g004

Our observations are in accordance with previous studies demonstrating that the neurotoxicity elicited by soluble A β upon neurons involves elevated K⁺ efflux, mediated through multiple pathways including enhanced activity of voltage-gated potassium channels [22,23] and the Na⁺/K⁺ ATPase [24,25]. Furthermore, increasing the extracellular K⁺ level to prevent K⁺ loss is also able to block A β -induced neuronal apoptosis [26]. We now demonstrate the direct measurement of net K⁺ flux of neurons in response to A β , and report that simultaneous exposure to both intra- and extracellular A β significantly impairs the ability of neurons to regulate K⁺ homeostasis. Given that intracellular accumulation of A β within neurons in the AD brain is only observed later in life, our data provides a potential mechanism that could explain why neurons in the AD brain become vulnerable to apoptosis later in life despite being continuously exposed to extracellular A β for many years.

While the toxicity of A β has been extensively studied in cultured neurons, two considerable limitations of these approaches have been that they have often involved the use of relatively immature neuronal phenotypes (cultured for between 1–4 days *in vitro*), and a treatment period of up to 24 hours. In this study we have cultured neurons at relatively high density for seven days *in vitro* prior to experimentation. Under these conditions, neuronal cultures contained dense networks of processes more representative of mature neurons. Chronic exposure of wildtype neurons to sublethal concentrations of A β did not alter axonal morphology of cultured neurons. However, Tg2576 neurons continuously treated with A β displayed substantial axonal pathology, including increased intensity of tau immunolabelling, axonal fragmentation and degeneration and the formation of axonal swellings that were packed with hyperphosphorylated tau. While it is well described that A β treatment of cultured neurons can directly cause tau hyperphosphorylation within 24 hours [27], our study is the first

that we are aware of that demonstrates that A β can cause the generation of hyperphosphorylated tau-immunoreactive axonal swellings in mature neuronal cultures. Notably, these axonal swellings took four days to develop, suggesting that they represent a slowly evolving, secondary phase of A β -induced neurodegeneration. These dystrophic axonal manifestations resemble some of the key neuritic pathologies observed in the AD brain, suggesting that the prolonged effect of intra- and extracellular A β exposure upon neurons is a critical step in the neurodegenerative process underlying AD.

In summary, we propose that in the AD brain, intraneuronal accumulation of A β increases the vulnerability of neurons to subsequent chronic exposure to soluble A β . This combined exposure to intra- and extracellular A β leads to degenerative changes in neurons (such as axonal swelling and fragmentation) and apoptosis through a K⁺ efflux mediated mechanism.

Methods

Ethics Statement

All animal experimentation was performed under the guidelines stipulated by the University of Tasmania Animal Ethics Committee, which is in accordance with the Australian code of practice for the care and use of animals for scientific purposes.

Cortical neuron cultures from Tg2576 and wildtype mice

Hemizygous Tg2576 male mice on a hybrid B6SjL background (Taconic) were crossed with wildtype B6SjLFl1 females, and cortical tissue was removed from individual embryos from the pregnant wildtype mice (the embryos are either transgenic or wildtype), and cortical neurons isolated as we have described previously [28]. Cortical neurons from individual pups were maintained in Neurobasal medium (Gibco) containing 10% fetal




Figure 5. Soluble A β induces prolonged efflux of K⁺ in Tg2576 cortical neurons. Treatment with 1 μ M A β_{1-40} triggered rapid efflux of K⁺ from Tg2576 neurons (A), which continued over the 120 minutes of recording. A β_{1-40} treatment caused a rapid influx of H⁺ in Tg2576 neurons (B), which did not stabilise over the 120 minute recording period.
doi:10.1371/journal.pone.0019026.g005

calf serum, at 37°C in humidified air containing 5% CO₂. The culture medium was replaced with serum- and glutamic acid-free culture medium after 24 hours, followed by half media changes twice weekly. To test the affect of chronic A β treatment upon cortical neurons, 1 μ M or 10 μ M of soluble monomeric A β_{1-40}

was added exogenously to 7DIV cultured neurons daily for 6 days with assessment of cellular viability conducted each day. Lyophilised A β_{1-40} peptide was purchased from EZBiolab, and solubilised in sterile MQ water (which served as the vehicle control for all experiments). The A β_{1-40} peptide was used in this study because in our experience the A β_{1-42} peptide induces substantial acute neurotoxicity at 10 μ M concentrations, making this peptide unsuitable for long-term studies. Note that each day, the entire culture medium was replaced with fresh media to which A β_{1-40} was added. All experiments were performed without knowing the genotype identity (wildtype or transgenic) of the individual neuron cultures. We generally obtained cultures from 10 embryos per pregnant animal, and they were usually split evenly between transgenic and wildtype. At the conclusion of each experiment, immunocytochemical detection of A β (using the 6E10 antibody, see details below) was used to genotype the cultures and reveal their identity for data analysis and interpretation. Untreated neuronal cultures at 14 days *in vitro* were used for this purpose, and neurons could be easily distinguished as containing either high- or low- levels of A β .

Table 1. Tg2576 cortical neurons are sensitive to soluble A β in a K⁺-dependent manner.

Treatment	Viability (relative to vehicle)
vehicle	100 \pm 6.9
1 μ M A β	60.1 \pm 5.8*
1 μ M A β + 2 mM 4-AP	87.4 \pm 6.0*#

Tg2576 cortical neurons were treated daily with either vehicle (PBS) or 1 μ M of monomeric A β_{1-40} for 6 days, and neuronal viability assessed 24 hours after the final treatment. This resulted in a reduction in neuronal viability to approximately 60% of vehicle controls (* - $p < 0.01$, student's t-test). However, when Tg2576 neurons were pre-treated with 2 mM 4-AP for 30 minutes prior to each A β_{1-40} treatment, this significantly reduced the neurotoxic action of exogenous A β_{1-40} .

(# - $p < 0.01$, student's t-test).

doi:10.1371/journal.pone.0019026.t001

Alamar blue viability assay

Neuronal viability was measured by the degree of cellular metabolic reduction of Alamar Blue[®], as we have reported




Figure 6. Soluble A β causes disruptions in tau distribution in Tg2576 cortical neurons. Tau immunolabelling of the axonal cytoskeleton demonstrated that axonal morphology was similar between wildtype (A) and Tg2576 (B) cortical neurons over 7–14 DIV. Six daily 1 μ M A β _{1–40} treatments of 7DIV wildtype neurons had no discernible effect upon axonal morphology (C). However, substantial changes in tau-labelling were observed in A β _{1–40} treated Tg2576 neurons (D); including increased intensity of tau immunostaining after 24 hours, followed by blebbing and axonal fragmentation which worsened after four days of treatment (D). Furthermore, after four consecutive days of 1 μ M A β _{1–40} treatment a number of axonal swellings, with dense accumulations of hyperphosphorylated tau, were observed in Tg2576 neuron cultures (E). scale bars = 30 μ m (A–D), 15 μ m (E).

doi:10.1371/journal.pone.0019026.g006

previously [29,30]. Briefly, viability was determined by determination of the fluorescence of Alamar Blue in culture wells (excitation 535 nm, emission 595 nm), and was expressed as the percentage of the signal obtained from the vehicle-treated culture. Alamar Blue was applied at a 1:10 dilution in culture media for 30 minutes, after which time it was collected and fluorescence measured on a fluorescent plate reader (Tecan Genios). In experiments involving repeated measurements of viability over a consecutive series of days, following Alamar Blue collection fresh media was applied to the neurons and the viability assay procedure repeated each day. Alamar Blue is non-toxic, and we have not observed any decline in neuronal viability when using this technique on consecutive days for up to one week (results not shown).

Immunocytochemical labelling of neurons

At the completion of experiments, cells were fixed with 4% paraformaldehyde for 20 minutes and an antibody diluent containing 0.03% Triton-X detergent was applied. For immunocytochemistry, rabbit anti-tau (1:5000; DAKO), mouse anti-A β (6E10; 1:1000; DAKO) and mouse anti-hyperphosphorylated tau (AT-8; 1:1000; Chemicon) antibodies were applied, and detected

with appropriate Alexa-Fluor-488 or -594 conjugated secondary antibodies at a 1:1000 concentration (Molecular Probes).

Non-invasive microelectrode ion flux (MIFE) measurements in cultured neurons

The theory of MIFE measurements was reviewed recently [31] and the complete experimental procedure including ion-selective microelectrode fabrication [32] and neuronal culture preparation, immobilisation and recording were performed as described previously [30,33]. Briefly, ion selective microelectrodes were silanised and filled with commercially available ionophore cocktails (Fluke catalog no. 60031 and 95297 for for K⁺ and H⁺, respectively). Microelectrodes for MIFE measurements were prepared on a daily basis and calibrated before and after measurements in a range of K⁺ and H⁺ concentrations. Cortical neurons for the MIFE measurements were grown for six days at a density of 1×10^5 cells/well on poly-L-lysine-coated coverslips and chronically treated with soluble monomeric A β _{1–40} as described above. A cover slip with neural cells was washed in and adapted to the MIFE artificial CSF (aCSF) for one hour prior to experiments. The composition of the aCSF was: 150 mM NaCl, 0.5 mM KCl, 0.5 mM CaCl₂, 1.5 mM MgCl₂, 1.25 mM NaH₂PO₄, 5 mM

NaHCO₃, 25 mM glucose, pH 7.4. Electrodes were co-focused and positioned ~5 μ m above the neuronal monolayer and moved up and down by a computer-controlled stepper motor providing a travel range between 5 and 50 μ m from the cell surface at a frequency of 0.1 Hz. After recording steady state flux for ~5 min, neurons were treated with A β and data acquired at a rate of 15 samples/sec and later averaged over 8 second intervals. For all ion flux measurements, the sign convention is 'influx positive' for a cation. The data were analysed using MIFE software with ion fluxes expressed in nmol m⁻² s⁻¹. A total flux was calculated as area between flux curve over the indicated experimental time-frame (25 min) and the starting flux value.

Statistical analysis

For each experiment unless otherwise stated, a minimum of four wells from at least three separate cultures (derived from different

animals), were used for quantification. Statistical analysis was completed using SPSS 16.0 (SPSS). When data was unequally distributed, data was transformed so that the residuals were approximately normally distributed. Statistical significance was calculated using t-test when only two experimental samples were compared, and One-Way ANOVA with Tukey's Post Hoc Test when multiple samples were compared. All graphical data is presented as mean \pm SEM, significance p<0.05.

Author Contributions

Conceived and designed the experiments: SR CH EDE LS WRB RSC. Performed the experiments: SR CH EDE CWB LS RSC. Analyzed the data: SR CH LS RSC. Contributed reagents/materials/analysis tools: AKW RSC. Wrote the paper: PAA GJG RSC.

References

- Selkoe DJ, Podlisny MB (2002) Deciphering the genetic basis of Alzheimer's disease. *Annual Review of Genomics and Human Genetics* 3: 67–99.
- Glenner GG, Wong CW (1984) Alzheimer's Disease - Initial Report of the Purification and Characterization of a Novel Cerebrovascular Amyloid Protein. *Biochemical and Biophysical Research Communications* 120: 885–890.
- Hung LW, Ciccostoto GD, Giannakis E, Tew DJ, Perez K, et al. (2008) Amyloid-beta peptide (A β) neurotoxicity is modulated by the rate of peptide aggregation: A β dimers and trimers correlate with neurotoxicity. *J Neurosci* 28(46): 11950–8.
- Mattson MP (2004) Pathways towards and away from Alzheimer's disease. *Nature* 430: 631–639.
- Takahashi RH, Almeida CG, Kearney PF, Yu F, Lin MT, et al. (2004) Oligomerization of Alzheimer's beta-amyloid within processes and synapses of cultured neurons and brain. *J Neurosci* 24(14): 3592–9.
- Gouras GK, Tsai J, Naslund J, Vincent B, Edgar M, et al. (2000) Intraneuronal A β 42 accumulation in human brain. *Am J Pathol* 156(1): 15–20.
- Oddo S, Caccamo A, Shepherd JD, Murphy MP, Golde TE, et al. (2003) Triple-transgenic model of Alzheimer's disease with plaques and tangles: intracellular A β and synaptic dysfunction. *Neuron* 39(3): 409–21.
- Giannakopoulos P, Herrmann FR, Bussière T, Bouras C, Kovari E, et al. (2003) Tangle and neuron numbers, but not amyloid load, predict cognitive status in Alzheimer's disease. *Neurology* 60(9): 1495–500.
- Kotilinek LA, Bacskai B, Westerman M, Kawarabayashi T, Younkin L, et al. (2002) Reversible memory loss in a mouse transgenic model of Alzheimer's disease. *J Neurosci* 22(15): 6331–5.
- Dodart JC, Bales KR, Gannon KS, Greene SJ, DeMattos RB, et al. (2002) Immunization reverses memory deficits without reducing brain A β burden in Alzheimer's disease model. *Nat Neurosci* 5(5): 452–7.
- Sompol P, Ittarat W, Tangpong J, Chen Y, Doubinskaia I, et al. (2008) A neuronal model of Alzheimer's disease: an insight into the mechanisms of oxidative stress-mediated mitochondrial injury. *Neuroscience* 153(1): 120–30.
- Yao J, Irwin RW, Zhao L, Nilsen J, Hamilton RT, et al. (2009) Mitochondrial bioenergetic deficit precedes Alzheimer's pathology in female mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 106(34): 14670–5.
- Esposito L, Gan L, Yu GQ, Essrich C, Mucke L (2004) Intracellularly generated amyloid-beta peptide counteracts the antiapoptotic function of its precursor protein and primes proapoptotic pathways for activation by other insults in neuroblastoma cells. *J Neurochem* 91(6): 1260–74.
- Ohyagi Y, Asahara H, Chui DH, Tsuruta Y, Sakae N, et al. (2005) Intracellular A β 42 activates p53 promoter: a pathway to neurodegeneration in Alzheimer's disease. *FASEB J* 19(2): 255–7.
- Zhang Y, McLaughlin R, Goodyer C, LeBlanc A (2002) Selective cytotoxicity of intracellular amyloid beta peptide 1–42 through p53 and Bax in cultured primary human neurons. *J Cell Biol* 156(3): 519–29.
- Fuentealba RA, Liu Q, Zhang J, Kanekiyo T, Hu X, et al. (2010) Low-density lipoprotein receptor-related protein 1 (LRP1) mediates neuronal A β 42 uptake and lysosomal trafficking. *PLoS One* 5(7): e11884.
- Pereira C, Santos MS, Oliveira C (1999) Involvement of oxidative stress on the impairment of energy metabolism induced by A β peptides on PC12 cells: protection by antioxidants. *Neurobiol Dis* 6(3): 209–19.
- Yatin SM, Varadarajan S, Link CD, Butterfield DA (1999) In vitro and in vivo oxidative stress associated with Alzheimer's amyloid beta-peptide (1–42). *Neurobiol Aging* 20(3): 325–30.
- Butterfield DA, Castegna A, Drake J, Scapagnini G, Calabrese V (2002) Vitamin E and neurodegenerative disorders associated with oxidative stress. *Nutr Neurosci* 5(4): 229–39.
- Boothby LA, Doering PL (2005) Vitamin C and vitamin E for Alzheimer's disease. *Ann Pharmacother* 39(12): 2073–80.
- Ohyagi Y, Asahara H, Chui DH, Tsuruta Y, Sakae N, et al. (2005) Intracellular A β 42 activates p53 promoter: a pathway to neurodegeneration in Alzheimer's disease. *FASEB J* 19(2): 255–7.
- Ramsden M, Plant LD, Webster NJ, Vaughan PF, Henderson Z, et al. (2001) Differential effects of unaggregated and aggregated amyloid beta protein (1–40) on K(+) channel currents in primary cultures of rat cerebellar granule and cortical neurones. *J Neurochem* 79(3): 699–712.
- Kerrigan TL, Atkinson L, Peers C, Pearson HA (2008) Modulation of 'A'-type K+ current by rodent and human forms of amyloid beta protein. *Neuroreport* 19(8): 839–43.
- Bores GM, Smith CP, Wirtz-Brugger F, Giovanni A (1998) Amyloid beta-peptides inhibit Na+/K+-ATPase: tissue slices versus primary cultures. *Brain Res Bull* 46(5): 423–7.
- Dickey CA, Gordon MN, Wilcock DM, Herber DL, Freeman MJ, et al. (2005) Dysregulation of Na+/K+ ATPase by amyloid in APP+PS1 transgenic mice. *BMC Neurosci* 6: 7.
- Yu SP, Farhangrazi ZS, Ying HS, Yeh CH, Choi DW (1998) Enhancement of outward potassium current may participate in beta-amyloid peptide-induced cortical neuronal death. *Neurobiol Dis* 5(2): 81–8.
- De Felice FG, Wu D, Lambert MP, Fernandez SJ, Velasco PT, et al. (2008) Alzheimer's disease-type neuronal tau hyperphosphorylation induced by A β oligomers. *Neurobiol Aging* 29(9): 1334–47.
- King AE, Dickson TC, Blizzard CA, Woodhouse A, Foster SS, et al. (2009) Neuron-glia interactions underlie ALS-like axonal cytoskeletal pathology. *Neurobiol Aging* Epub May 6.
- Töugu V, Karafin A, Zovo K, Chung RS, Howells C, et al. (2009) Zn(II)- and Cu(II)-induced non-fibrillar aggregates of amyloid-beta (1–42) peptide are transformed to amyloid fibrils, both spontaneously and under the influence of metal chelators. *J Neurochem* 110(6): 1784–95.
- Chung RS, Howells C, Eaton ED, Shabala L, Zovo K, et al. (2010) The native copper- and zinc-binding protein metallothionein blocks copper-mediated A β aggregation and toxicity in rat cortical neurons. *PLoS One* 5(8): e12030.
- Shabala L, Ross T, McMeekin T, Shabala, S (2006) Non-invasive microelectrode ion flux measurements to study adaptive responses of microorganisms to the environment. *FEMS Microbiol Rev* 30: 472–486.
- Shabala L, Ross T, Newman I, McMeekin T, Shabala, S (2001) Measurements of net fluxes and extracellular changes of H⁺, Ca²⁺, K⁺, and NH₄⁺ in *Escherichia coli* using ion-selective microelectrodes. *J Microbiol Methods* 46(2): 119–29.
- Shabala L, Howells C, West AK, Chung RS (2010) Prolonged A β treatment leads to impairment in the ability of primary cortical neurons to maintain K⁺ and Ca²⁺ homeostasis. *Mol Neurodegener* 5: 30.