

RESEARCH ARTICLE

Maternal Micronutrient Supplementation and Long Term Health Impact in Children in Rural Bangladesh

Tania Mannan^{1,2}, Sultan Ahmed¹, Evana Akhtar¹, Anjan Kumar Roy¹, Md Ahsanul Haq¹, Adity Roy¹, Maria Kippler³, Eva-Charlotte Ekström⁴, Yukiko Wagatsuma⁵, Rubhana Raqib^{1*}

1 Immunobiology, Nutrition and Toxicology Laboratory, Infectious Diseases Division, icddr,b, Dhaka, Bangladesh, **2** Department of Immunology, Bangladesh University of Health Sciences, Mirpur, Dhaka, Bangladesh, **3** Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden, **4** International Maternal and Child Health (IMCH), Department of Women's and Children's Health, Uppsala University, Uppsala, Sweden, **5** Department of Clinical Epidemiology, Faculty of Medicine, University of Tsukuba, Tsukuba, Ibaraki, Japan

* rubhana@icddr.org



OPEN ACCESS

Citation: Mannan T, Ahmed S, Akhtar E, Roy AK, Haq MA, Roy A, et al. (2016) Maternal Micronutrient Supplementation and Long Term Health Impact in Children in Rural Bangladesh. PLoS ONE 11(8): e0161294. doi:10.1371/journal.pone.0161294

Editor: Jacobus van Wouwe, TNO, NETHERLANDS

Received: June 9, 2016

Accepted: August 3, 2016

Published: August 18, 2016

Copyright: © 2016 Mannan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data underlying the findings in our study are freely available in the paper and supplemental files. For additional information please refer to <http://www.icddr.org/> policies.

Funding: This work was supported by the Japan Society for the Promotion of Science (JSPS; Grant No. 18256005), Karolinska Institutet (Grant No. 00933), and icddr (Grant 384, SWE-2008-065). icddr,b also gratefully acknowledges the following donors who provide unrestricted support: Government of the People's Republic of Bangladesh; Global Affairs Canada (GAC); Swedish International Development Cooperation Agency (Sida) and the Department for International Development, (UKAid).

Abstract

Background

Limited data is available on the role of prenatal nutritional status on the health of school-age children. We aimed to determine the impact of maternal micronutrient supplementation on the health status of Bangladeshi children.

Methods

Children (8.6–9.6 years; n = 540) were enrolled from a longitudinal mother-child cohort, where mothers were supplemented daily with either 30mg iron and 400µg folic acid (Fe30F), or 60mg iron and 400µg folic acid (Fe60F), or Fe30F including 15 micronutrients (MM), in rural Matlab. Blood was collected from children to determine the concentration of hemoglobin (Hb) and several micronutrients. Anthropometric and Hb data from these children were also available at 4.5 years of age and mothers at gestational week (GW) 14 and 30.

Results

MM supplementation significantly improved ($p \leq 0.05$) body mass index-for-age z-score (BAZ), but not Hb levels, in 9 years old children compared to the Fe30F group. MM supplementation also reduced markers of inflammation ($p \leq 0.05$). About 28%, 35% and 23% of the women were found to be anemic at GW14, GW30 and both time points, respectively. The prevalence of anemia was 5% and 15% in 4.5 and 9 years old children, respectively. The adjusted odds of having anemia in 9 year old children was 3-fold higher if their mothers were anemic at both GW14 and GW30 [Odds Ratio (OR) = 3.05; 95% Confidence Interval

Competing Interests: The authors have declared that no competing interests exist.

(CI) 1.42, 6.14, $P = 0.002$] or even higher if they were also anemic at 4.5 years of age [OR = 5.92; 95% CI 2.64, 13.25; $P < 0.001$].

Conclusion

Maternal micronutrient supplementation imparted beneficial effects on child health. Anemia during pregnancy and early childhood are important risk factors for the occurrence of anemia in school-age children.

Introduction

A growing body of evidence indicates that the nutritional background of a woman during pregnancy and especially during early life is a critical determinant of her offspring's subsequent health outcomes such as sub-optimum growth and mortality, suggesting an intergenerational transfer of poor health from mother to child [1,2]. The underlying cause of about 60% of childhood death below 5 years of age in Bangladesh is malnutrition which involves both calorie and micronutrient deficiencies [3]. In this age-group children suffer from one or more forms of malnutrition including stunting (32%), underweight (30%) and anemia (33%) [4]. Iron deficiency and iron deficiency anemia is a major public health problem worldwide, mostly in pregnant women, infants, and young children in developing countries [5]. It adversely affects some vital aspects of human health, including poor cognitive development, decreased immune function and work productivity [5–7]. The most common causes of anemia are nutritional (vitamin and mineral deficiencies such as iron, folate and B vitamins, malnutrition in general) and non-nutritional factors (acute and chronic infections, genetic disorders, poverty, sociocultural and maternal factors) [8]. The majority of the studies reporting maternal anemia as a risk factor for childhood anemia have focused on under-5 children [9,10]. However, the long-term effect of maternal anemia during pregnancy in older school-age children remains unclear.

A large number of the studies emphasized the effects of maternal micronutrient supplementation on pregnancy outcomes, birth outcomes, neonatal/child survival, child growth and morbidity outcomes [11,12]. Data is scarce on the consequence of long-term effects of maternal nutrition supplementation on nutritional status in school going children. In this study, we aimed to evaluate whether maternal micronutrient supplementation during pregnancy influences nutritional and micronutrient status in school-age children (~9 years) in a longitudinal mother-child cohort in rural Bangladesh [13,14]. A secondary aim was to investigate whether anemia during pregnancy has a long-term impact on anemia in pre-adolescent school-age children and whether this is influenced by maternal micronutrient supplementation.

Methods

Study area and population

This study is a follow-up of the MINIMat (the Maternal and Infant Nutrition Interventions, Matlab) (ISRCTN16581394) mother-child cohort [13]. It was conducted between 2012 and 2013 in Matlab, a rural area located 53 km southeast of Dhaka, Bangladesh. The icddr,b, runs a health and demographic surveillance system in Matlab, as well as a central hospital and four connected sub-centers that provide health care to the resident population (about 220,000) in the area.

The MINIMat trial included 4436 women identified as pregnant between November 2001 and October 2003 [14]. The enrolled women were randomly assigned to one of three different micronutrient supplementations: (i) 30 mg iron and 400 µg folic acid (Fe30F), (ii) 60 mg iron and 400 µg folic acid (Fe60F), or (iii) the UNICEF preparation of 15 different micronutrients (MM) including 30 mg iron and 400 µg folic acid [14]. Baseline data for mothers were taken by trained health research workers at the first sub-center visit in pregnancy week 8. Hemoglobin (Hb) levels of mothers were measured by HemoCue photometer (HemoCue® AB, Ängelholm, Sweden) during GW14 and GW30. Socioeconomic status (SES) of the families was estimated via an asset score, generated through principal component analysis of household assets and grouped into tertiles [15].

A total of $n = 2735$ MINIMat children were studied at the age of 4.5 years. To reduce the burden of various types of investigations, the children were divided into two groups on the basis of the calendar year of birth. Group A children ($n = 1432$) were born between April 2002 to May 2003. Group B children ($n = 1303$) were born between June 2003 to June 2004 and were studied for asthma, allergy (total group B children, $n = 1303$, referred in Hawlader et al) [16,17] as well as immune function and bone growth studies ($n = 640$ group B children reported in Ahmed et al) [13,18]. These group B children were earlier involved in various studies from birth and at the age of 4.5 years [13,18–22]. Anthropometric data and family SES in 4.5 years old children were determined as described earlier [13]. Hb levels at 4.5 years of age were measured by HemoCue photometer. In the current study, field workers carried out a survey by visiting the household of children studied at 4.5 years of age ($n = 640$) for availability at 9 years of age [13]. Eighty nine children were not available for various reasons (Fig 1). A total of 551 children were enrolled in the current study, among them 7 children refused to give blood and 4 blood samples were clotted. Finally, 540 blood samples were available for analysis (Fig 1).

Data collection for children at 9 years of age was carried out in a similar way as done for earlier follow up at 4.5 years of age [13]. In brief, anthropometric measurements were taken at the sub-center clinic visit by trained nurses. Weight was measured with a digital scale (TANITA HD-318, Tanita Corporation, Japan), accurate to ± 10 g. Weight scales were standardized daily. Height was measured using a free standing stadiometer Leicester Height Measure with millimeter marks (Seca214, UK). The stadiometer was calibrated before the start of the study and in every 6 months. The measured weight and height were converted to weight-for-age (WAZ), height-for-age (HAZ) and body mass index (BMI)-for-age Z-scores (BAZ) (standard deviation (SD) scores), using the WHO Multicentre Growth Reference Study child growth standards [23,24]. Children with WAZ < -2 SD from WHO reference population were considered as underweight, with HAZ < -2 SD as stunted, and those with BAZ < -2 SD were considered as thin. Family SES data were updated as described above.

Assessment of plasma biomarkers

Five ml of fasting venous blood was collected in Lithium heparin treated tubes (Sarstedt Mon-evette®, Sweden) by trained paramedics in the subcenters in Matlab. A drop of blood was used to measure Hb by HemoCueHb 201+ (HemoCue AB, Ängelholm, Sweden). The blood samples were transported to the Matlab Laboratory for separation of plasma which was thereafter stored in -80°C freezer until transported to the Laboratory in icddr,b, Dhaka. Ferritin, vitamin B₁₂ and folate were analyzed in plasma by chemiluminescence method using Cobas e601 (Roche Diagnostics, Mannheim, Germany). C-reactive protein and sTfR were assessed in plasma using Hitachi 902 (Roche Diagnostics, Mannheim, Germany). Plasma hepcidin was measured using a commercial ELISA kit (DRG International, GmbH, Germany). Flame atomic




Fig 1. Flow chart describing the enrolment of the children in the current study. The group A children (n = 1432) were born between April 2002 to May 2003 and group B (n = 1303) were born between June 2003 to June 2004.

doi:10.1371/journal.pone.0161294.g001

absorption spectroscopy was used to analyze plasma zinc (Zn) concentrations (Shimadzu Corporation, Kyoto, Japan). Isocratic reverse-phase HPLC and UV detection were used to analyze concentrations of plasma retinol/vitamin A (Shimadzu Corporation, Kyoto, Japan). The mean inter-day coefficient of variation was <5% for sTfR, ferritin, folate, vitamin B₁₂, Zn, vitamin A, and CRP and <8% for hepcidin. For an independent assessment of the laboratory's analytical performance, it participates in external quality assurance programs such as VITAL EQA of Centers for Disease Control and Prevention (CDC) for the above parameters excepting Zn. The values are given in [S1 Table](#).

The complete blood count (CBC) including mean corpuscular volume (MCV), mean corpuscular Hb (MCH), mean corpuscular Hb concentration (MCHC) and other blood indices were determined using Hematology Analyzer (Sysmex XT-1800i, Kobe, Japan). Erythrocyte Sedimentation Rate (ESR) was measured by Wintrobe Method [\[25\]](#).

Hb and anemia

Anemia is defined as Hb<115g/l for 9years and <110g/l for <4.5years old children according to WHO guidelines [\[26\]](#) ([S2 Table](#)). Mild anemia is defined if Hb level is 110-114g/l, moderate is 80–109 g/l, and severe anemia is <80g/l [\[27\]](#). The cut-off to define anemia in pregnancy was set to Hb<110 g/l according to WHO [\[5\]](#). The cut-off values to define iron deficiency (ID), iron deficiency anemia (IDA) [\[26,28\]](#), and micronutrient deficiencies [\[29–31\]](#) have been described in [S2 Table](#).

Ethics

The institutional review board at icddr,b consists of two review committees, namely the Research Review Committee (RRC) and Ethical Review Committee (ERC). The study was approved by both committees. Written informed consent was obtained from the parents or guardian of each child prior to participation in the study.

Statistical analyses

Data analyses were done using the Statistical package for the Social Science (SPSS) for Windows (version 20; Armonk, NY: IBM SPSS corp.; 2011) and Stata/IC, version 13 (StataCorp, Texas, USA). Normality (data distribution patterns) and homogeneity of variances were formally checked. Independent sample t-test, Mann-Whitney U-test, ANOVA or Kruskal-Wallis test was used as appropriate for an initial assessment. Analysis of covariance with least significant difference (LSD) was used for multiple comparisons of child nutritional status and plasma biomarkers in relation to supplementation groups. Linear regression analyses were used to evaluate the influence of plasma biomarkers on Hb levels. Multivariable adjusted logistic regression analyses were used to evaluate the risk of having anemia in 9 years old children if they were anemic at 4.5 years of age or their mothers were anemic during pregnancy. Statistical models were adjusted for covariates that were associated with exposure and outcome, or biologically relevant or changed the effect estimate more than 5%. These were SES, mother's education, mother's occupation, child BMI, child sex, and plasma CRP. P values <0.05 were considered statistically significant.

Results

Anthropometric measurements

Basic characteristics of the children and their mothers were presented in [Table 1](#). At 9 years of age, about 22% children were stunted and 40% were underweight ([Table 1](#)). Among the stunted

Table 1. Basic characteristics of mothers and their children.

	All children (n = 540)	Fe30F (n = 188)	Fe60F (n = 185)	MM (n = 167)
Mothers				
^a Hb at GW14 (g/l)	116.98±12.70	117.08±11.87	116.69±13.68	117.21±12.50
^b Hb at GW30 (g/l)	113.98±11.55	113.24±12.45	114.76±11.48	113.92±10.55
Updated family SES (at 9 years)				
1 st tertile, n (%)	180 (33.46%)	61 (32.45%)	64 (34.59%)	55 (33.33%)
2 nd tertile, n (%)	179 (33.27%)	65 (34.57%)	54 (29.19%)	60 (36.36%)
3 rd tertile, n (%)	179 (33.27%)	62 (32.98%)	67 (36.22%)	50 (30.30%)
Education				
Illiterate, n (%)	59 (11.01%)	13 (6.99%)	24 (12.97%)	22 (13.33%)
Primary level, n (%)	190 (35.45%)	72 (38.71%)	56 (30.27%)	62 (37.58)
≥Secondary, n (%)	287 (53.54%)	101 (54.30)	105 (56.76)	81 (49.09%)
Parity, n(%)				
1–2	215 (39.81%)	78 (41.49%)	69 (37.30%)	68 (40.72%)
3	212 (39.26%)	70 (37.23%)	79 (42.70%)	63 (37.72%)
≥4	113 (20.93%)	40 (21.28%)	37 (20.00%)	36 (21.56%)
Children at 4.5 years				
Age (month)	55.93±1.38	55.88±1.38	56.03±1.44	55.86±1.32
Hb (g/l)	128.61±13.33	127.68±12.53	129.84±13.45	128.28±14.03
^d Stunted, n (%)	158 (29.80%)	53 (28.19%)	51 (27.56%)	54 (32.33%)
^d Underweight, n (%)	209 (39.40%)	75 (39.90%)	73 (39.46%)	61 (36.53%)
^d Thinness, n (%)	69 (12.78%)	23 (12.23%)	26 (14.05%)	20 (11.97%)
Children at 9 years				
Age (month)	106.43±1.50	106.45±1.69	106.42±1.51	106.40±1.26
Hb (g/l)	123.81±9.06	123.37±8.89	124.43±9.49	123.62±8.77
^d Stunted, n (%)	116 (21.50%)	36 (19.15%)	41 (22.16%)	39 (23.35%)
^d Underweight, n (%)	215 (39.80%)	77 (40.96%)	75 (40.54%)	63 (37.32%)
^d Thinness, n (%)	137 (25.50%)	60 (31.91%)	41 (22.16%)	36 (21.56%)

Abbreviation: Hb, hemoglobin; SES, socioeconomic status.: Fe30F, 30mg iron and 400µg folic acid; Fe60F, 60mg iron and 400µg folic acid; MM, Fe30F including 15 micronutrients.

Data are presented as mean ± standard deviation or numbers with percentage in parentheses.

^aHb at GW14 (n = 482)

^bHb at GW30 (n = 461)

^cEducation: Illiterate, never admitted to school; Primary level, completed 1–5 years of schooling; Secondary level & above secondary level, completed ≥6 years of schooling.

^dStunting, underweight, and thinness were defined as children with height for age, weight for age and body mass index (BMI) for age <-2 SD from the median value of height, weight and BMI for age of reference population, respectively.

doi:10.1371/journal.pone.0161294.t001

children, 10% were severely stunted (HAZ<-3SD) while among the underweight children, 26% were severely underweight (WAZ<-3SD). About 26% children were thin (Table 1); among them 21% were severely thin (BAZ<-3SD).

At 4.5 years of age, about 30% children were stunted, among them, 16% were severely stunted. About 39% children were underweight and among them about 19% were severely underweight. Again, about 13% children were thin, of which 10% were severely thin.

The demographic characteristics of the children in the current study (n = 540) were similar to those of the rest of the group B children (S3 Table) who were available (n = 500) (n = 163

not studied) for the background information from another ongoing study at that time (Tofail F, unpublished).

Status of nutritional biomarkers

The mean Hb concentration of 9 years old children was 123.81 ± 9.06 g/l (mean \pm SD) (Table 1). There was no difference in the Hb concentration between boys and girls. The prevalence of anemia was about 15% in 9 years and 5% in 4.5 years old children. Among the 9 years old anemic children, 55% had mild, and 45% had moderate anemia. Severe anemia was not present among the children.

Descriptive statistics of all nutritional biomarkers have been given in Table 2. Folate deficiency (<5.2 nmol/l) was not observed among the study participants; only 5 children (0.93%) had vitamin B₁₂ deficiency (<182 pmol/l) (S2 Table). About 13% children had folate (>29.5 nmol/l) and 3.1% had vitamin B₁₂ concentration (>867 pmol/l) above the reference range. Zinc and vitamin A deficiencies were present in 8% and 6.2% children respectively (S2 Table). All studied children were hepcidin deficient when 53.5 μ g/l [32] was considered as a cut-off, but only 5 children were deficient when the cut-off was <1 μ g/l [33]. Iron deficiency was found in 8.0% children when sTfR cut-off was used. However, it was only 0.2% when ferritin cut-off was used (S2 Table). About 2.5% children had IDA according to the defined sTfR cut-off and 0.2% (one child) when applying the ferritin cut-off.

Based on morphological classification of anemia, about 61% of anemic children ($n = 50$) and 92% of non-anemic children ($n = 431$) had normocytic normochromic RBC. About 34% of anemic children and 6.3% of non-anemic children had microcytic RBC while 4.9% of anemic and 1.3% of non-anemic children had normocytic hypochromic RBC.

Influence of maternal supplementation on child growth and inflammatory markers

Compared to the Fe30F group, children in the Fe60F as well as MM group had significantly higher BAZ scores at 4.5 and 9 years of age (Table 3). However, no significant differences were seen in HAZ or WAZ scores with respect to supplementation. Maternal supplementation did not affect the number of stunted or underweight children either at 4.5 or 9 years of age. However, 9 years old children belonging to the MM and Fe60F groups had fewer thin children ($n = 36$ and 41 , respectively) compared to those in Fe30F group ($n = 60$, χ^2 $P = 0.02$, and 0.04 , respectively) (Data not shown).

When markers of inflammation or infection were considered, children in MM group had significantly lower levels of CRP and ESR compared to the Fe30F group (Table 3). Children in the MM group had significantly lower concentration of plasma folate compared to the Fe30F group.

When considering RBC indices, children in Fe60F group had higher MCH concentration compared to Fe30F group. RDW values were significantly lower in the Fe60F group compared to Fe30F group (Table 3).

Association of Hb with plasma biomarkers and effect of maternal micronutrient supplementation

In the adjusted linear regression analysis, Hb was significantly inversely associated with plasma sTfR; when the association between Hb and sTfR was stratified by sex the association remained significant only in boys (Table 4). Hb was positively associated with hepcidin in all children and in girls only. Hb was also positively associated with vitamin A in all children (Table 4). A

Table 2. Plasma biomarkers in school-age children in rural Matlab.

	All children (n = 540)	Fe30F (n = 188)	Fe60F (n = 185)	MM (n = 167)
sTfR (nmol/l)	44.83±10.96	45.34±11.38	44.67±11.61	44.45±9.69
Ferritin (µg/l)	59.82±31.35	62.04±30.10	59.57±28.55	57.59±35.42
Folate (nmol/l)	23.13±5.74	23.78±5.96	22.53±5.64	23.05±5.57
Vitamin B ₁₂ (pmol/l)	469.96±174.67	474.88±167.78	460.38±179.18	474.95±177.78
Hepcidin (µg/l)	9.57±5.92	9.85±6.65	9.60±5.69	9.23±5.29
Zinc (µmol/l)	13.33±2.54	13.34±2.55	13.36±2.64	13.29±2.43
Vitamin A (µmol/l)	1.03±0.22	1.03±0.22	1.04±0.23	1.01±0.21

Abbreviation: Fe30F, 30mg iron and 400µg folic acid; Fe60F, 60mg iron and 400µg folic acid; MM, Fe30F including 15 micronutrients; sTfR, soluble transferrin receptor.

Data is presented as mean ± standard deviation.

doi:10.1371/journal.pone.0161294.t002

positive tendency was obtained between Hb and Zn in girls, and a positive association was noted between Hb and B₁₂ in boys only.

To evaluate the influence of maternal supplementation, the associations between Hb and plasma biomarkers in 9 years old children were stratified by different supplementation groups (Table 4, and S4 Table). In the Fe60F group but not others (Fe30F and MM), Hb was strongly

Table 3. Analysis of covariance of child nutritional status and plasma biomarkers in different supplementation groups.

	Fe30F (n = 188)	Fe60F (n = 185)	P-value ¹	MM (n = 167)	P-value ²
Nutritional status					
HAZ at 4.5 years	-1.53±0.86	-1.55±0.88	0.59	-1.59±0.87	0.54
HAZ at 9 years	-1.27±0.86	-1.27±0.93	0.98	-1.40±0.88	0.19
WAZ at 4.5 years	-1.76±0.84	-1.78±0.83	0.58	-1.76±0.76	0.98
WAZ at 9 years	-1.68±1.04	-1.67±1.11	0.80	-1.74±0.96	0.38
BAZ at 4.5 years	-1.17±0.82	-1.18±0.76	0.97	-1.11±0.78	0.33
BAZ at 9 years	-1.47±1.09	-1.25±1.08	0.05	-1.23±1.06	0.04
Biomarkers					
Hb at 4.5 years	127.76±8.90	129.79±9.49	0.15	128.08±8.77	0.76
Hb at 9 years	123.54±12.50	124.26±13.45	0.52	123.56±14.03	0.98
Folate (nmol/l)	23.66±5.96	22.60±5.64	0.06	20.03±5.57	0.03
ESR (mm/1st hr)	16.15±10.95	15.76±10.43	0.75	14.25 ^b ±10.28	0.05
CRP (mg/L)	1.36±2.97	0.91±1.60	0.04	0.60±0.63	0.002
RDW-CV (%)	13.67±1.22	13.43±0.96	0.05	13.54±1.05	0.29
MCH (pg)	26.35±2.25	27.50±6.64	0.01	26.64±2.03	0.57

Abbreviation: sTfR, soluble transferrin receptor; Fe30F, 30mg iron and 400µg folic acid; Fe60F, 60mg iron and 400µg folic acid; MM, Fe30F including 15 micronutrients; Hb, hemoglobin, ESR, erythrocyte sedimentation rate, RDW, Red cell distribution width, MCH, mean corpuscular hemoglobin and CRP, plasma concentration of C-reactive protein.

¹Significant difference between Fe30F and Fe60F

²Significant difference between Fe30F and MM

Models were adjusted for socio economic status, body mass index, child sex, mother's occupation, mother's education levels and plasma concentration of C-reactive protein

Data are presented as mean ± standard deviation.

doi:10.1371/journal.pone.0161294.t003

Table 4. Linear regression analyses between hemoglobin concentrations and plasma biomarkers in all children and boys and girls separately.

Variables	All supplementation groups					
	All children (n = 540)		Boys (n = 263)		Girls (n = 277)	
	*Adj. β (95% CI)	P	#Adj. β(95% CI)	P	#Adj. β(95% CI)	P
sTfR (nmol/l)	-0.75(-1.48, 0.01)	0.04	-1.46(-2.38, -0.55)	0.002	0.46(-0.78, 1.70)	0.46
Ferritin (μg/l)	0.13(-0.12, 0.38)	0.31	0.20(-0.15, 0.56)	0.26	0.06(-0.30, 0.43)	0.73
Folate (nmol/l)	-0.11(-1.47, 1.26)	0.87	-1.01(-2.98, 0.96)	0.31	1.03(-0.92, 3.00)	0.30
Vitamin B ₁₂ (pmol/l)	0.02(-0.02, 0.06)	0.39	0.07(-0.001, 0.14)	0.05	-0.01(-0.07, 0.05)	0.64
Hepcidin (μg/l)	1.274(0.26, 3.31)	0.02	1.12(-1.44, 3.69)	0.38	1.92(0.09, 3.75)	0.04
Zinc (μmol/l)	1.65(-1.44, 4.73)	0.29	-0.85(-5.20, 3.50)	0.70	3.98(-0.48, 8.44)	0.08
Vitamin A (μmol/l)	2.16(0.88, 3.45)	0.001	3.02(1.04, 5.00)	0.003	1.70(0.01, 3.38)	0.04
	Fe60F					
	All children (n = 185)		Boys (n = 87)		Girls (n = 98)	
sTfR (nmol/l)	-0.79(-2.06, 0.48)	0.22	-1.06(-2.74, 0.62)	0.21	-0.5(-3.03, 1.53)	0.51
Ferritin (μg/l)	0.47(-0.02, 0.97)	0.06	0.72(-0.10, 1.53)	0.08	0.37(-0.27, 1.01)	0.25
Folate (nmol/l)	0.32(-2.22, 2.86)	0.80	-1.06(-4.84, 2.72)	0.57	3.12(-0.72, 6.96)	0.11
Vitamin B ₁₂ (pmol/l)	0.04(-0.04, 0.12)	0.27	0.19 (0.04, 0.34)	0.01	-0.02(-0.11, 0.07)	0.66
Hepcidin (μg/l)	2.88(0.35, 5.41)	0.02	0.80(-4.31, 5.91)	0.75	4.11(1.20, 7.02)	0.006
Zinc (μmol/l)	-1.05(-6.57, 4.47)	0.70	-5.25(-13.57, 3.07)	0.21	4.29(-4.00, 12.57)	0.30
Vitamin A (μmol/l)	3.13(0.89, 5.38)	0.006	6.04(2.44, 9.64)	0.001	1.11(-1.84, 4.06)	0.45

Abbreviation: Fe60F, 60mg iron and 400μg folic acid; sTfR, soluble transferrin receptor.

Data were given as regression coefficient (β) and 95% confidence intervals

*Adjusted for socio economic status, body mass index, child sex, mother’s occupation, mother’s education levels and plasma concentration of C-reactive protein

#Adjusted for socio economic status, body mass index, mother’s occupation, mother’s education levels and plasma concentration of C-reactive protein.

doi:10.1371/journal.pone.0161294.t004

positively associated with plasma hepcidin and vitamin A in all children. When these associations were stratified by sex, Hb was associated with vitamin A in boys and with hepcidin in girls (Table 4). Again in Fe60F group, the significant positive association was noted between Hb and Vitamin B₁₂ only in boys (Table 4).

Long term impact of maternal anemia in school-age children

Among mothers during pregnancy, about 28% and 35% of the women at GW14 and GW30, respectively, were found to be anemic, and 23.6% women were anemic at both time points. Percentage of anemia was higher in children at 9 years of age (15%) compared to 4.5 years of age (5%). Notably, thirteen children who were anemic at 4.5 years of age (45%) remained anemic at 9 years of age. Sixty nine children who were non-anemic at 4.5 years became anemic at 9 years. In adjusted model, the odds (OR) of being anemic in 9 years of age was 1.81 (95% CI 1.07, 3.05; P = 0.027) and 2.34 (1.37, 4.00; P = 0.002) fold higher if their mothers were anemic at either GW14 or GW30, respectively, compared to non-anemic mothers (Fig 2). The odds of having anemia at 9 years was even higher if the mothers were anemic at both time points (OR = 3.05, 95% CI 1.42, 6.14; P = 0.002). The odds increased, about 6 times (OR = 5.92, 95% CI 2.64, 13.25; P<0.001) when the children were also anemic at 4.5 years. However, no impact of maternal anemia was observed at 4.5 years of age (at GW14, OR = 0.45, 95% CI 0.15, 1.34; P = 0.15; at GW30, OR = 1.28, 95% CI 0.54, 3.04; P = 0.56 and at both time points, OR = 0.66, 95% CI 0.18, 2.43; P = 0.54).




Fig 2. The odds of having anemia at 9 years of age if the mother were anemic at GW-14 or GW-30 or if the children were anemic at 4.5 years of age. Abbreviation: Fe30F, 30mg iron and 400µg folic acid; Fe60F, 60mg iron and 400µg folic acid; MM, Fe30F including 15 micronutrients, GW, Gestational week.

doi:10.1371/journal.pone.0161294.g002

When the analyses was stratified by supplementation groups, the odds of having anemia at 9 years of age was 2.34 (95% CI 0.49, 11.04; P = 0.28), 3.95 (95% CI 1.09, 14.38; P = 0.03) and 4.15 (95% CI 1.25, 13.76; P = 0.02) times higher in the Fe30F, Fe60F and MM groups, respectively, if their mothers were anemic both at GW14 and GW30 (Fig 2). Again, the odds of having anemia at 9 years of age was 11.16 (95% CI 2.73, 45.59; P = 0.001), 6.35 (95% CI 1.50, 26.81; P = 0.01) and 3.50 (95% CI 0.76, 16.09; P = 0.10) times higher in the Fe30F, Fe60F and MM groups, respectively, if they were anemic at early childhood (4.5 years).

Discussion

In the present study, we found that maternal supplementation during pregnancy with MM had beneficial effects on child nutritional status (BAZ scores) that also decreased markers of inflammation (CRP, ESR) in 9 years old children.

In a larger cohort of MINIMat children (n = 1634), Khan et al showed that maternal MM supplementation (n = 897) increased the proportion of stunting in boys only at 4.5 years of age [34]. However, we did not find any impact of MM supplementation on the proportion of stunting or underweight children either at 4.5 or 9 years of age, not even in boys. Although, a long-term favorable effect of MM supplementation was seen in terms of fewer numbers of thin children at 9 years, but this was not evident at 4.5 years of age. Data on child growth beyond 5 years of age in relation to maternal supplementation are scarce. A meta-analysis of randomized controlled trials reported that maternal multi-micronutrient supplementation had a significant positive effect on head circumference of under-5 children without any effects on weight, height, WAZ, HAZ and WHZ [35]. In our study, supplementation with MM also markedly reduced markers of inflammation and infection in 9 year old children. The MM supplement of UNICEF/WHO/UNU contained important antioxidants such as vitamin E, vitamin C and selenium which are known to reduce inflammatory responses. It is possible that prenatal MM

supplementation improved the immune reserve of the growing fetus that persists and aid in combating infections in later childhood. Very few studies have reported effects of MM supplementation during pregnancy on the health outcomes of older children. Multi-vitamin supplementation of HIV-infected women during pregnancy and lactation was shown to be associated with reduced rate of all types of diarrhea among under-5 HIV-negative children [36]. A study in Nepal demonstrated that maternal vitamin A supplementation enhanced natural antibody concentrations in children at preadolescent age (9 to 13 years); the authors postulated that prenatal supplementation lead to a higher reservoir and sustained natural immunity in these children [37]. An increasing body of evidence suggests that maternal nutritional status including micronutrients, life-style, exposure to pollutants etc. from preconception through lactation causes fetal or neonatal epigenetic changes that might account for altered mechanisms of growth, metabolism and diseases observed later in life [38].

In the current study, majority of 4.5 (95%) and 9 years (85%) old children, Hb concentration was well within the normal range. The previous report on a larger cohort (n = 1354) of MINIMat children at 4.5 years of age showed similar finding, where 92% of the children had Hb levels within the normal range [39]. The positive association between Hb and plasma concentration of ferritin, B₁₂, hepcidin and vitamin A mainly in Fe60F group indicated a beneficial impact of the higher dose of antenatal iron on Hb concentration in 9 years old children. Vitamin A has an important role in iron mobilization into Hb of developing RBC [40,41]. Iron status impacts plasma and liver levels of vitamin A [42]. Supplementation with 60 mg iron and folate during pregnancy increased hepcidin concentration in iron deficient Tanzanian women (serum ferritin $\leq 12\mu\text{g/L}$) [43]. In rural Bangladesh, pregnant women with low iron deficiency, plasma concentrations of Zn, vitamin B₁₂, and α -tocopherol were positively associated with Hb levels [44]. We did not find similar associations between Hb and other plasma biomarkers in the Fe30F and MM groups. Hb concentrations did not differ by supplementation groups in children, this finding was in line with the original MINIMat trial where supplementation did not affect Hb concentration in pregnant women at GW30 [14].

The prevalence of anemia in 9 year old children was 15% which was slightly lower than the prevalence rate of 19% obtained in the recent Bangladesh National Micronutrient Survey conducted in school-age (6–11 years) children [4]. Notably, none of the children had folate deficiency; iron (8% or 0.2%) and vitamin B₁₂ deficiencies (0.9%) were also minimal suggesting that a major cause of anemia in children was not related to iron, folate or B₁₂ deficiency. These results are in line with the findings at 4.5 years of age as reported earlier on MINIMat children [39]. Presence of normocytic normochromic anemia in more than half (about 61%) of the anemic children could indicate presence of recent infections, deficiency of vitamin B₂ (riboflavin) and genetic disorders such as sickle cell anemia, red blood cell membrane disorders etc. that may cause hemoglobinopathies in children of this age group [45]. Microcytic anemia can be attributed to IDA, vitamin B₆ (pyridoxine) deficiency and thalassemia [45]. In the current study, the exact nature of anemia (whether genetic or B₂ or B₆ deficiency) could not be determined, however, parents were notified, and suspected children were referred to specialists.

Maternal anemia during pregnancy was a strong risk factor for children being anemic at 9 years of age. Several studies have demonstrated that maternal anemia was a risk factor for childhood anemia in under-5 children [41,46]. To the best of our knowledge, this is the first report that shows that this risk is also apparent at 9 years of age. In contrary to other studies, we did not find any association between maternal and childhood anemia at 4.5 years of age. The influence of maternal anemia during pregnancy seems to emerge in pre-adolescent age and not earlier in this cohort. An interesting finding of the present study was that Hb level at 4.5 years of age was a stronger risk factor than maternal Hb (~2 folds higher) for being anemic in 9 years of age. The risk of being anemic at 9 years was higher among the Fe30F group

(OR = 11), likely due to low Fe or absence of MM supplementation during pregnancy. The findings reflect that early childhood nutritional status is also very important besides prenatal nutrition for optimum health outcome at later age and highlighting the public health importance of these findings.

One of the limitations of the study was that we could not measure Hb by the cyanmethemoglobin method (the gold standard) although the hemocue technique is well-recognized and suitable for epidemiological surveys. Another drawback was that concentration of multiple micronutrients was not measured in 4.5 year old children or their mothers during pregnancy to compare with the current micronutrient status in school-age children. The strength of our study lies in the fact that we have determined important nutritional markers and indicators of inflammation (CRP) and infection (ESR). The statistical models were adjusted with SES, mother's education, mother's occupation, child BMI, child sex, and plasma CRP to rule out possible confounding factors. However, there may be some unmeasured social, environmental, dietary and other factors that may impact the outcomes. Another shortcoming of the current study was that we did not have reliable information on recent morbidity (past 1–2 months). However, we adjusted the statistical models with plasma CRP concentrations as a marker of recent infection.

In conclusion, our results indicate that maternal micronutrient supplementation confers beneficial health effects on pre-adolescent school-age children although the effects were not apparent at an earlier age of 4.5 years. Anemia during pregnancy and in early childhood appeared to be important risk factors for anemia in pre-adolescent age. This information might aid the public health services to take initiatives aimed at reducing anemia and micronutrient deficiencies in early childhood in addition to the programs being carried out for women during pregnancy. Further studies are needed to better understand the mechanisms by which maternal and early childhood nutritional status influence health in later childhood.

Supporting Information

S1 Table. External quality assurance program (VITAL EQA) of Centers for Disease Control and Prevention.

(DOCX)

S2 Table. Indicators used in this study to define anemia and micronutrient deficiencies in school-age children.

(DOCX)

S3 Table. Descriptive statistics of group B children who were included and those not included in the current study.

(DOCX)

S4 Table. Linear regression analyses between hemoglobin concentrations and plasma biomarkers in children in Fe30F and MMS supplementation groups.

(DOCX)

Acknowledgments

We gratefully acknowledge the participation of the women and their children.

Author Contributions

Conceptualization: RR YW.

Data curation: MAH SA EA AKR.

Formal analysis: MAH TM SA.

Funding acquisition: YW MK RR.

Investigation: AR EA AKR.

Methodology: RR YW.

Project administration: RR.

Resources: RR YW MK.

Supervision: RR YW.

Validation: EA AKR TM SA.

Visualization: MAH SA.

Writing - original draft: RR TM SA.

Writing - review & editing: RR TM MK E-CE SA.

References

1. Subramanian SV, Ackerson LK, Davey Smith G, John NA (2009) Association of maternal height with child mortality, anthropometric failure, and anemia in India. *JAMA* 301: 1691–1701. doi: [10.1001/jama.2009.548](https://doi.org/10.1001/jama.2009.548) PMID: [19383960](https://pubmed.ncbi.nlm.nih.gov/19383960/)
2. Ozaltin E, Hill K, Subramanian SV (2010) Association of maternal stature with offspring mortality, underweight, and stunting in low- to middle-income countries. *JAMA* 303: 1507–1516. doi: [10.1001/jama.2010.450](https://doi.org/10.1001/jama.2010.450) PMID: [20407060](https://pubmed.ncbi.nlm.nih.gov/20407060/)
3. Faruque AS, Ahmed AM, Ahmed T, Islam MM, Hossain MI, et al. (2008) Nutrition: basis for healthy children and mothers in Bangladesh. *J Health Popul Nutr* 26: 325–339. PMID: [18831228](https://pubmed.ncbi.nlm.nih.gov/18831228/)
4. NMS (2013) National Micronutrients Status Survey 2011–12. ICDDR,B, UNICEF, Bangladesh, GAIN and Institute of Public Health and Nutrition.
5. WHO (2008) Worldwide prevalence of anemia 1993–2005: WHO global database on anemia. Geneva.
6. Luo R, Zhang L, Liu C, Zhao Q, Shi Y, et al. (2011) Anaemia among students of rural China's elementary schools: prevalence and correlates in Ningxia and Qinghai's poor counties. *J Health Popul Nutr* 29: 471–485. PMID: [22106753](https://pubmed.ncbi.nlm.nih.gov/22106753/)
7. Institute of Public Health Nutrition (IPHN) MoHaFW, Government of the People's Republic of Bangladesh (2007) National Strategy for Anaemia Prevention and Control in Bangladesh
8. Sanou D, Ngnie-Teta I (2012) Risk Factors for Anemia in Preschool Children in Sub-Saharan Africa *InTech*. 171–183 p.
9. Pasricha SR, Black J, Muthayya S, Shet A, Bhat V, et al. (2010) Determinants of anemia among young children in rural India. *Pediatrics* 126: e140–149. doi: [10.1542/peds.2009-3108](https://doi.org/10.1542/peds.2009-3108) PMID: [20547647](https://pubmed.ncbi.nlm.nih.gov/20547647/)
10. Khan JR, Awan N, Misu F (2016) Determinants of anemia among 6–59 months aged children in Bangladesh: evidence from nationally representative data. *BMC Pediatr* 16: 3. doi: [10.1186/s12887-015-0536-z](https://doi.org/10.1186/s12887-015-0536-z) PMID: [26754288](https://pubmed.ncbi.nlm.nih.gov/26754288/)
11. Zerfu TA, Ayele HT (2013) Micronutrients and pregnancy; effect of supplementation on pregnancy and pregnancy outcomes: a systematic review. *Nutr J* 12: 20. doi: [10.1186/1475-2891-12-20](https://doi.org/10.1186/1475-2891-12-20) PMID: [23368953](https://pubmed.ncbi.nlm.nih.gov/23368953/)
12. Haider BA, Bhutta ZA (2015) Multiple-micronutrient supplementation for women during pregnancy. *Cochrane Database Syst Rev* 11: CD004905. doi: [10.1002/14651858.CD004905.pub4](https://doi.org/10.1002/14651858.CD004905.pub4) PMID: [26522344](https://pubmed.ncbi.nlm.nih.gov/26522344/)
13. Ahmed S, Moore SE, Kippler M, Gardner R, Hawlader MD, et al. (2014) Arsenic exposure and cell-mediated immunity in pre-school children in rural bangladesh. *Toxicol Sci* 141: 166–175. doi: [10.1093/toxsci/kfu113](https://doi.org/10.1093/toxsci/kfu113) PMID: [24924402](https://pubmed.ncbi.nlm.nih.gov/24924402/)
14. Persson LA, Arifeen S, Ekstrom EC, Rasmussen KM, Frongillo EA, et al. (2012) Effects of prenatal micronutrient and early food supplementation on maternal hemoglobin, birth weight, and infant mortality

- among children in Bangladesh: the MINIMat randomized trial. *JAMA* 307: 2050–2059. doi: [10.1001/jama.2012.4061](https://doi.org/10.1001/jama.2012.4061) PMID: [22665104](https://pubmed.ncbi.nlm.nih.gov/22665104/)
15. Gwatkin DR, Rustein S, Johnson K, Pande RP, Wagstaff A (2000) Socioeconomic differences in health, nutrition, and population in Bangladesh, HNP/Poverty Thematic Group Working Paper. The World Bank, Washington, DC.
 16. Hawlader MD, Ma E, Noguchi E, Itoh M, Arifeen SE, et al. (2014) *Ascaris lumbricoides* Infection as a Risk Factor for Asthma and Atopy in Rural Bangladeshi Children. *Trop Med Health* 42: 77–85. doi: [10.2149/tmh.2013-19](https://doi.org/10.2149/tmh.2013-19) PMID: [25237284](https://pubmed.ncbi.nlm.nih.gov/25237284/)
 17. Hawlader MD, Noguchi E, El Arifeen S, Persson LA, Moore SE, et al. (2014) Nutritional status and childhood wheezing in rural Bangladesh. *Public Health Nutr* 17: 1570–1577. doi: [10.1017/S1368980013001262](https://doi.org/10.1017/S1368980013001262) PMID: [23680045](https://pubmed.ncbi.nlm.nih.gov/23680045/)
 18. Ahmed S, Rekha RS, Ahsan KB, Doi M, Grandner M, et al. (2013) Arsenic exposure affects plasma insulin-like growth factor 1 (IGF-1) in children in rural Bangladesh. *PLoS One* 8: e81530. doi: [10.1371/journal.pone.0081530](https://doi.org/10.1371/journal.pone.0081530) PMID: [24303053](https://pubmed.ncbi.nlm.nih.gov/24303053/)
 19. Raqib R, Ahmed S, Sultana R, Wagatsuma Y, Mondal D, et al. (2009) Effects of in utero arsenic exposure on child immunity and morbidity in rural Bangladesh. *Toxicol Lett* 185: 197–202. doi: [10.1016/j.toxlet.2009.01.001](https://doi.org/10.1016/j.toxlet.2009.01.001) PMID: [19167470](https://pubmed.ncbi.nlm.nih.gov/19167470/)
 20. Doi M, Rekha RS, Ahmed S, Okada M, Roy AK, et al. (2011) Association between calcium in cord blood and newborn size in Bangladesh. *Br J Nutr* 106: 1398–1407. doi: [10.1017/S0007114511001747](https://doi.org/10.1017/S0007114511001747) PMID: [21736799](https://pubmed.ncbi.nlm.nih.gov/21736799/)
 21. Ahmed S, Mahabbat-e Khoda S, Rekha RS, Gardner RM, Ameer SS, et al. (2011) Arsenic-associated oxidative stress, inflammation, and immune disruption in human placenta and cord blood. *Environ Health Perspect* 119: 258–264. doi: [10.1289/ehp.1002086](https://doi.org/10.1289/ehp.1002086) PMID: [20940111](https://pubmed.ncbi.nlm.nih.gov/20940111/)
 22. Ahmed S, Ahsan KB, Kippler M, Mily A, Wagatsuma Y, et al. (2012) In utero arsenic exposure is associated with impaired thymic function in newborns possibly via oxidative stress and apoptosis. *Toxicol Sci* 129: 305–314. doi: [10.1093/toxsci/kfs202](https://doi.org/10.1093/toxsci/kfs202) PMID: [22713597](https://pubmed.ncbi.nlm.nih.gov/22713597/)
 23. WHO (2006) WHO Child Growth Standards: Methods and development: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age.
 24. De Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, et al. (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 85: 660–667. PMID: [18026621](https://pubmed.ncbi.nlm.nih.gov/18026621/)
 25. NCCLS. Reference and Selected Procedure for the Erythrocyte Sedimentation Rate (ESR) Test; Approved Standard.
 26. WHO (2001) Iron deficiency anaemia: assessment, prevention, and control. A guide for programme managers. Geneva.
 27. WHO (2011) Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System. Geneva.
 28. Kolbe-Busch S, Lotz J, Hafner G, Blanckaert NJ, Claeys G, et al. (2002) Multicenter evaluation of a fully mechanized soluble transferrin receptor assay on the Hitachi and cobas integra analyzers. the determination of reference ranges. *Clin Chem Lab Med* 40: 529–536. PMID: [12113300](https://pubmed.ncbi.nlm.nih.gov/12113300/)
 29. Akcam M, Ozdem S, Yilmaz A, Gultekin M, Artan R (2007) Serum ferritin, vitamin B(12), folate, and zinc levels in children infected with *Helicobacter pylori*. *Dig Dis Sci* 52: 405–410. PMID: [17211708](https://pubmed.ncbi.nlm.nih.gov/17211708/)
 30. IZINCG (2007) International Zinc Nutrition Consultative Group. Assessing population zinc status with serum zinc concentration.
 31. WHO (2011) Serum retinol concentrations for determining the prevalence of vitamin A deficiency in populations. Vitamin and Mineral Nutrition Information System. Geneva.
 32. Sdogou T, Tsentidis C, Gourgiotis D, Marmarinos A, Gkourogianni A, et al. (2015) Immunoassay-based serum hepcidin reference range measurements in healthy children: differences among age groups. *J Clin Lab Anal* 29: 10–14. doi: [10.1002/jcla.21719](https://doi.org/10.1002/jcla.21719) PMID: [24659514](https://pubmed.ncbi.nlm.nih.gov/24659514/)
 33. EIA-5258 E DRG Hepcidin-25 (bioactive)
 34. Khan AI, Kabir I, Ekstrom EC, Asling-Monemi K, Alam DS, et al. (2011) Effects of prenatal food and micronutrient supplementation on child growth from birth to 54 months of age: a randomized trial in Bangladesh. *Nutr J* 10: 134. doi: [10.1186/1475-2891-10-134](https://doi.org/10.1186/1475-2891-10-134) PMID: [22152147](https://pubmed.ncbi.nlm.nih.gov/22152147/)
 35. Lu WP, Lu MS, Li ZH, Zhang CX (2014) Effects of multimicronutrient supplementation during pregnancy on postnatal growth of children under 5 years of age: a meta-analysis of randomized controlled trials. *PLoS One* 9: e88496. doi: [10.1371/journal.pone.0088496](https://doi.org/10.1371/journal.pone.0088496) PMID: [24586335](https://pubmed.ncbi.nlm.nih.gov/24586335/)

36. Khavari N, Jiang H, Manji K, Msamanga G, Spiegelman D, et al. (2014) Maternal multivitamin supplementation reduces the risk of diarrhoea among HIV-exposed children through age 5 years. *Int Health* 6: 298–305. doi: [10.1093/inthealth/ihu061](https://doi.org/10.1093/inthealth/ihu061) PMID: [25173342](https://pubmed.ncbi.nlm.nih.gov/25173342/)
37. Palmer AC, Schulze KJ, Khatry SK, De Luca LM, West KP Jr. (2015) Maternal vitamin A supplementation increases natural antibody concentrations of preadolescent offspring in rural Nepal. *Nutrition* 31: 813–819. doi: [10.1016/j.nut.2014.11.016](https://doi.org/10.1016/j.nut.2014.11.016) PMID: [25933488](https://pubmed.ncbi.nlm.nih.gov/25933488/)
38. Capra L, Tezza G, Mazzei F, Boner AL (2013) The origins of health and disease: the influence of maternal diseases and lifestyle during gestation. *Ital J Pediatr* 39: 7. doi: [10.1186/1824-7288-39-7](https://doi.org/10.1186/1824-7288-39-7) PMID: [23343462](https://pubmed.ncbi.nlm.nih.gov/23343462/)
39. Henriksson H (2015) Prevalence of anemia and its association with socio-demographic factors and micronutrient deficiencies in 4.5-year old children in Matlab, Bangladesh: a cross-sectional follow-up study. Uppsala University: Uppsala University. 2015., 2046 p p.
40. Semba RD, Bloem MW (2002) The anemia of vitamin A deficiency: epidemiology and pathogenesis. *Eur J Clin Nutr* 56: 271–281. PMID: [11965502](https://pubmed.ncbi.nlm.nih.gov/11965502/)
41. Demirchyan A, Petrosyan V, Sargsyan V, Hekimian K (2016) Prevalence and determinants of anaemia among children aged 0–59 months in a rural region of Armenia: a case-control study. *Public Health Nutr* 19: 1260–1269. doi: [10.1017/S1368980015002451](https://doi.org/10.1017/S1368980015002451) PMID: [26329316](https://pubmed.ncbi.nlm.nih.gov/26329316/)
42. Rosales FJ, Jang JT, Pinero DJ, Erikson KM, Beard JL, et al. (1999) Iron deficiency in young rats alters the distribution of vitamin A between plasma and liver and between hepatic retinol and retinyl esters. *J Nutr* 129: 1223–1228. PMID: [10356091](https://pubmed.ncbi.nlm.nih.gov/10356091/)
43. Abioye AI, Aboud S, Premji Z, Etheredge AJ, Gunaratna NS, et al. (2016) Iron Supplementation Affects Hematologic Biomarker Concentrations and Pregnancy Outcomes among Iron-Deficient Tanzanian Women. *J Nutr* 146: 1162–1171. doi: [10.3945/jn.115.225482](https://doi.org/10.3945/jn.115.225482) PMID: [27121530](https://pubmed.ncbi.nlm.nih.gov/27121530/)
44. Shamim AA, Kabir A, Merrill RD, Ali H, Rashid M, et al. (2013) Plasma zinc, vitamin B(12) and alpha-tocopherol are positively and plasma gamma-tocopherol is negatively associated with Hb concentration in early pregnancy in north-west Bangladesh. *Public Health Nutr* 16: 1354–1361. doi: [10.1017/S1368980013000475](https://doi.org/10.1017/S1368980013000475) PMID: [23469947](https://pubmed.ncbi.nlm.nih.gov/23469947/)
45. Janus J, Moerschel SK (2010) Evaluation of anemia in children. *Am Fam Physician* 81: 1462–1471. PMID: [20540485](https://pubmed.ncbi.nlm.nih.gov/20540485/)
46. Zhao A, Zhang Y, Peng Y, Li J, Yang T, et al. (2012) Prevalence of anemia and its risk factors among children 6–36 months old in Burma. *Am J Trop Med Hyg* 87: 306–311. doi: [10.4269/ajtmh.2012.11-0660](https://doi.org/10.4269/ajtmh.2012.11-0660) PMID: [22855763](https://pubmed.ncbi.nlm.nih.gov/22855763/)