



DATA NOTE

The genome sequence of the channel bull blenny, *Cottoperca gobio* (Günther, 1861) [version 1; peer review: 2 approved]

Iliana Bista ^{1,2}, Shane A. McCarthy ^{1,2}, Jonathan Wood ¹, Zemin Ning¹, H. William Detrich III ³, Thomas Desvignes ⁴, John Postlethwait⁴, William Chow¹, Kerstin Howe ¹, James Torrance ¹, Michelle Smith¹, Karen Oliver¹, Vertebrate Genomes Project Consortium, Eric A. Miska^{2,5}, Richard Durbin^{1,2}

¹Wellcome Sanger Institute, Cambridge, CB10 1SA, UK

²Department of Genetics, University of Cambridge, Cambridge, CB2 3EH, UK

³Department of Marine and Environmental Sciences, Northeastern University Marine Science Center, Massachusetts, MA 01908, USA

⁴Institute of Neuroscience, University of Oregon, Eugene, OR 97403-1254, USA

⁵Gurdon Institute, University of Cambridge, Cambridge, CB2 1QN, UK

V1 First published: 24 Jun 2020, 5:148
<https://doi.org/10.12688/wellcomeopenres.16012.1>

Latest published: 24 Jun 2020, 5:148
<https://doi.org/10.12688/wellcomeopenres.16012.1>

Abstract

We present a genome assembly for *Cottoperca gobio* (channel bull blenny, (Günther, 1861)); Chordata; Actinopterygii (ray-finned fishes), a temperate water outgroup for Antarctic Notothenioids. The size of the genome assembly is 609 megabases, with the majority of the assembly scaffolded into 24 chromosomal pseudomolecules. Gene annotation on Ensembl of this assembly has identified 21,662 coding genes.

Keywords

Cottoperca gobio, channel bull blenny, genome assembly chromosomal, Notothenioidei

Open Peer Review

Reviewer Status

	Invited Reviewers	
	1	2
version 1		
24 Jun 2020	report	report

1. **Kevin Bilyk**, Montclair State University, Montclair, USA

2. **Chenhong Li** , Shanghai Ocean University, Shanghai, China

Liang Lu, Shanghai Ocean University, Shanghai, China

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Iliana Bista (ib8@sanger.ac.uk)

Author roles: **Bista I:** Conceptualization, Formal Analysis, Investigation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **McCarthy SA:** Conceptualization, Formal Analysis, Methodology, Writing – Review & Editing; **Wood J:** Data Curation, Methodology, Writing – Review & Editing; **Ning Z:** Formal Analysis, Methodology, Writing – Review & Editing; **Detrich III HW:** Conceptualization, Resources, Writing – Review & Editing; **Desvignes T:** Resources, Writing – Review & Editing; **Postlethwait J:** Conceptualization, Resources, Writing – Review & Editing; **Chow W:** Data Curation, Methodology, Writing – Review & Editing; **Howe K:** Data Curation, Writing – Review & Editing; **Torrance J:** Data Curation, Methodology, Writing – Review & Editing; **Smith M:** Investigation, Writing – Review & Editing; **Oliver K:** Investigation, Writing – Review & Editing; **Miska EA:** Conceptualization, Funding Acquisition, Supervision, Writing – Review & Editing; **Durbin R:** Conceptualization, Formal Analysis, Funding Acquisition, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Wellcome Trust through core funding to the Wellcome Sanger Institute (206194). IB, SAM and RD were supported by Wellcome grant 207492. HWD was supported by National Science Foundation grants OPP-0132032 (ICEFISH 2004 Cruise) and PLR-1444167. This work was also supported by grants from Wellcome (104640, 092096) to EAM. Publication number 29 from the ICEFISH Cruise of 2004 (HWD, Chief Scientist, RVIB Nathaniel B. Palmer). This is contribution 406 from the Marine Science Center at Northeastern University. JHP, HWD, and TD were supported by the National Science Foundation grant OPP-1543383. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2020 Bista I *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Bista I, McCarthy SA, Wood J *et al.* **The genome sequence of the channel bull blenny, *Cottoperca gobio* (Günther, 1861) [version 1; peer review: 2 approved]** Wellcome Open Research 2020, 5:148 <https://doi.org/10.12688/wellcomeopenres.16012.1>

First published: 24 Jun 2020, 5:148 <https://doi.org/10.12688/wellcomeopenres.16012.1>

Species taxonomy

Eukaryota; Metazoa; Chordata; Vertebrata; Gnathostomata; Actinopterygii; Teleostei; Clupeocephala; Percomorphaeae; Perciformes; Notothenioidei; Bovichtidae; Cottoperca; *Cottoperca gobio* (Günther, 1861) - synonym: *Cottoperca trigloides* (Balushkin, 2000), NCBI taxid: 56716.

Background

Cottoperca gobio (channel bull blenny) is a member of the Bovichtidae family of the Notothenioidei, a fish group endemic to the Southern Ocean. The Bovichtidae (thornfishes), are considered to be the most basally diverging family of notothenioids and are less adapted to life in the extreme cold in comparison to Antarctic members of the clade (Near *et al.*, 2015). *C. gobio* occupies the Patagonian regions of Chile and Argentina, and the area around the Falkland Islands. In contrast to Antarctic notothenioids (cryonotothenioids), the Bovichtidae do not produce antifreeze glycoproteins (AFGPs), a key adaptation to extreme Antarctic cold (Chen *et al.*, 1997; Cheng *et al.*, 2003) and their hemoglobins possess slightly higher oxygen affinity than most high-Antarctic species (Giordano *et al.*, 2006; Giordano *et al.*, 2009). Cytogenetic investigation of *C. gobio* showed that the karyotype of this species consists of $2n=48$ chromosomes (Pisano *et al.*, 1995). This condition, shared by other Bovichtidae, is considered to be the ancestral karyotype condition for all notothenioids (Mazzei *et al.*, 2006).

Here, we present a chromosomally complete genome sequence of *Cottoperca gobio* generated using specimens collected south of the Falkland Islands/Islands Malvinas. We trust that this genome sequence will be used to aid analysis of population structure and phylogeography of non-Antarctic and Antarctic notothenioid fish species, which are increasingly under threat due to climate change and human activities (Dornburg *et al.*, 2017).

Genome sequence report

The *C. gobio* genome was sequenced from a specimen collected under permits to fish in territorial waters of the Falkland Islands/Islands Malvinas issued by the United Kingdom, by the Falkland Islands Government, and by Argentina. The genome assembly for *C. gobio* (fCotGob3.1) is based on a combination of data from four technologies, including 75x coverage Pacific Biosciences (PacBio) single-molecule long reads (N50 14 kb), 54x coverage of Illumina data generated from a 10X Genomics Chromium library (estimated molecule length N50 43 kb), and BioNano Saphyr two-enzyme data (BspQI and BssSI). Additionally, 145x coverage of Illumina HiSeqX data were obtained from a Hi-C library prepared by Arima Genomics using tissue from a second individual (fCotGob2, spleen tissue).

The final assembly has a total length of 609 Mb, in 322 sequence scaffolds with a scaffold N50 of 25 Mb (Figure 1; Table 1). The majority (94.36%) of the assembly sequence was assigned to 24 chromosomal-level scaffolds using the Hi-C data

(Figure 2; Table 2). The assembly has a BUSCO (Simão *et al.*, 2015) gene completeness score of 93.4% using the actinopterygii reference set (with -sp zebrafish parameter). The chromosomes clearly show a one-to-one relationship with those in the Japanese medaka (*Oryzias latipes*) HdrR assembly GCA_002234675.1 (Figure 3 and Figure 4), with 3671 of the 3780 complete and single copy BUSCO genes present in both genomes found on homologous chromosomes (97.1%), and were thus named correspondingly. Analysis of conserved synteny detected no major interchromosomal rearrangements in the approximately 195 million years since the divergence of medaka and *C. gobio* lineages (Steinke *et al.*, 2006), but many intrachromosomal rearrangements (Figure 4). While not fully phased, the assembly deposited represents one haplotype. Contigs corresponding to the second haplotype have also been deposited.

Gene annotation

An Ensembl annotation was generated for the fCotGob3.1 assembly using RNA-seq data generated from 4 tissues (brain, muscle, ovary, and spleen). The annotation for assembly fCotGob3.1 was released in Ensembl under database version 99.31 (Hunt *et al.*, 2018) (for fish clade annotation information see 2019-09: fish clade gene annotation). The resulting Ensembl annotation includes 60,811 transcripts assigned to 21,662 coding and 2,823 non-coding genes (Channel bull blenny - Ensembl). RefSeq annotation is also available as NCBI *Cottoperca gobio* Annotation Release 100 (Table 2).

Methods

Specimen acquisition and nucleic acid extractions

Both specimens used to generate the genome assembly were collected south of the Falkland Islands/Islands Malvinas in 2004 (Lat Long: -52° 40', -59° 12') during the ICEFISH 2004 Cruise (International Collaborative Expedition to collect and study Fish Indigenous to Sub-Antarctic Habitats; led by H. W. Detrich (Detrich *et al.*, 2012)) of the *RVIB Nathaniel B. Palmer*. Following euthanasia, fresh blood was collected from specimen fCotGob3, and spleen tissue (used for Hi-C) was collected from specimen fCotGob2 and was flash frozen in liquid nitrogen. Blood was processed immediately, whereas flash frozen spleen was preserved in the -80 freezer until processing. For RNA sequencing, tissue samples from two specimens were used (fCotGob2 - spleen, and fCotGob1 - brain, skeletal muscle, ovary). The additional tissues (fCotGob1) were preserved in RNALater and kept frozen until extraction. The tissues were sampled by T. Desvignes, H. W. Detrich, and J. H. Postlethwait from a specimen captured northwest of the Falkland Islands in 2018 by the Falkland Islands Fisheries Department (Grass *et al.*, 2018).

High molecular weight (HMW) DNA from fresh blood cells was prepared using an agarose plug extraction protocol (Smith *et al.*, 2010). Blood DNA was initially stabilised in agarose plugs and then shipped to Sanger Institute where the final steps of the extraction were performed using a BioNano Tissue extraction protocol. Quality control (QC) of HMW DNA was




Figure 1. Genome assembly of *Cottoperca gobio*, fCotGob3.1. - BlobToolKit Snailplot, showing N50 metrics and BUSCO gene completeness. BlobToolKit plots are available at: [fCotGob3.1 - BlobToolKit](#).

performed using the Femto Pulse instrument (Agilent). Total RNA was extracted from approximately 20–40 mg of tissue, from brain, skeletal muscle, ovary and spleen tissues using the RNeasy Qiagen extraction kit (Qiagen). QC was performed using Qubit HS RNA kit, and Agilent Bioanalyzer Nano chips. Only extracts with RIN value >8 were used for sequencing.

Sequencing

PacBio continuous long read (CLR) and 10X Genomics linked read sequencing libraries were constructed according to manufacturers' instructions. Sequencing was performed by the Scientific Operations core at the Wellcome Sanger Institute on PacBio SEQUEL I and Illumina HiSeq X

instruments. Hi-C data were generated using the Arima Hi-C kit v1 by Arima Genomics. BioNano data were generated on Saphyr (dual enzyme) at Bionano Genomics. RNA-seq was performed on HiSeq 4000 with 150bp insert paired end (PE) libraries.

Genome assembly

An initial PacBio assembly was made using Falcon-unzip (Chin *et al.*, 2016) without repeat-masking during overlap detection with Dazzler. The contigs from this assembly were first scaffolded by comparing them to a second wtdbg (Ruan & Li, 2019) assembly using [cross_genome](#), then they were scaffolded further using the 10X data with [scaff10X](#), and then with BioNano two-enzyme hybrid scaffolding using

Table 1. Data information for *Cottoperca gobio*, fCotGob3.1 genome assembly.

Project accession information	
Assembly identifier	fCotGob3.1
Species	<i>Cottoperca gobio</i> (<i>Cottoperca trigloides</i>)
Specimens	fCotGob3 (PacBio, 10XG and BioNano), fCotGob2 (Hi-C and RNA-seq) fCotGob1 (RNA-seq)
NCBI taxonomy ID	56716
BioProject	PRJEB30272
Study accession	PRJEB19273
BioSample IDs	SAMEA104132835 (fCotGob1) SAMEA5365137 (fCotGob1.brain1) SAMEA5365124 (fCotGob1.gonad1) SAMEA5365123 (fCotGob1.muscle1) SAMEA104242971 (fCotGob2) SAMEA4872137 (fCotGob2.spleen1) SAMEA104242975 (fCotGob3)
Raw data accessions	
Pacific Biosciences SEQUEL I	ERR2219167 - ERR2219176
10X Genomics Illumina	ERR2639757 - ERR2639760
Hi-C Illumina	ERR4179340 - ERR4179344
BioNano	ERZ1392783 - ERZ1392785
RNA-seq	ERR3132340 (fCotGob1.brain1) ERR3132342 (fCotGob1.gonad1) ERR3132341 (fCotGob1.muscle1) ERR2639616 (fCotGob2.spleen1)
Genome assembly	
Assembly accession	GCA_900634415.1
Accession of alternate haplotype	GCA_900634435.1
Span (Mb)	609
Number of contigs	766
Contig N50 length (Mb)	5,939,854
Number of scaffolds	322
Scaffold N50 length (Mb)	25,156,145
Longest scaffold (Mb)	30.48
BUSCO genome score	C:93.4%, [S:90.5%, D:2.9%], F:1.3%, M:5.3%, n:4584

Solve v3.2.1. The original PacBio data were then used to fill gaps with PBJelly (English *et al.*, 2012) and polish with Arrow. The resulting assembly was then polished again using the 10X Illumina data, by mapping with bwa mem (Li, 2013), calling variants with freebayes (Garrison & Marth, 2012), and correcting

homozygous non-reference variants with **bcftools** consensus. Contiguity was increased further by filling gaps with the contigs from a second wtdgb assembly, which was made using PacBio reads corrected with Canu (Koren *et al.*, 2017). This assembly was re-polished with Arrow and freebayes, and retained




Figure 2. Hi-C contact map for the genome assembly of *Cottoperca gobio*, fCotGob3.1. Visualized in Juicebox (Durand *et al.*, 2016).

Table 2. Chromosomal pseudomolecules in the genome assembly fCotGob3.1, of species *Cottoperca gobio* - GCA_900634415.1.

Name	INSDC	RefSeq	Size (Mb)	GC%	Protein	Gene
1	LR131916.1	NC_041355.1	27.06	40.8	1,808	1,175
2	LR131927.1	NC_041356.1	12.92	41.9	792	681
3	LR131933.1	NC_041357.1	30.03	40.3	1,487	919
4	LR131934.1	NC_041358.1	28.95	40.7	1,629	1,007
5	LR131935.1	NC_041359.1	30.48	40.9	2,033	1,302
6	LR131936.1	NC_041360.1	27.68	40.9	1,823	1,143
7	LR131937.1	NC_041361.1	23.07	41	1,619	1,088
8	LR131938.1	NC_041362.1	23.43	41.2	1,836	1,194
9	LR131939.1	NC_041363.1	30.07	41	1,888	1,158
10	LR131917.1	NC_041364.1	27.44	40.8	1,407	992
11	LR131918.1	NC_041365.1	22.19	40.8	1,440	909
12	LR131919.1	NC_041366.1	22.9	40.6	1,424	850
13	LR131920.1	NC_041367.1	27.74	41	1,542	1,029
14	LR131921.1	NC_041368.1	25.7	40.6	1,627	1,134
15	LR131922.1	NC_041369.1	24.96	41	1,365	967
16	LR131923.1	NC_041370.1	26.58	41	1,811	1,094
17	LR131924.1	NC_041371.1	25.16	40.8	1,663	1,228
18	LR131925.1	NC_041372.1	14.93	41.8	1,018	690
19	LR131926.1	NC_041373.1	21.06	41.2	1,563	969
20	LR131928.1	NC_041374.1	17.6	41.4	964	649
21	LR131929.1	NC_041375.1	24.1	40.6	1,400	937
22	LR131930.1	NC_041376.1	22.61	41.3	1,415	1,026
23	LR131931.1	NC_041377.1	15.93	41.9	973	594
24	LR131932.1	NC_041378.1	22.44	41.1	1,229	1,184
Unplaced	-	.	34.34	41.6	2,093	1,676




Figure 3. Syntenic relationships of fCotGob3.1 assembly with Japanese medaka HdrR chromosomes, based on single copy orthologs. Visualised in Circos (Krzywinski *et al.*, 2009).

haplotigs were identified with Purge Haplotigs (Roach *et al.*, 2018). Finally, the assembly was scaffolded to chromosomes using Arima Hi-C data with Salsa (Ghurye *et al.*, 2017). The scaffolded assembly was checked for contamination and manually improved using gEVAL (Chow *et al.*, 2016). The manual curation included steps such as correcting mis-joins, improving concordance with all available data types, and Hi-C 2D map visualized in Juicebox to produce complete chromosomal units (Durand *et al.*, 2016). Curation resulted in 9 manual breaks, 114

manual joins and the removal of 102 regions representing false duplications, decreasing the scaffold count by 39% to 322 and increasing the scaffold N50 by 68% to 25.2 Mb. The chromosomal-level scaffolds were named based on conserved synteny to the medaka assembly (*Oryzias latipes*, Assembly accession GCA_002234675.1). The genome was further analysed within the BlobToolKit environment (Challis *et al.*, 2020). Software tools and versions used for assembly are listed in Table 3.




Figure 4. Examples of conserved synteny between Japanese medaka HdrR (purple) and fCotGob3.1 (pink) from chromosomes 1, 3, 6, and 16 (source: Ensembl).

Data availability

Underlying data

European Nucleotide Archive: *Cottoperca gobio* (channel bull blenny) genome assembly, fCotGob3.1. BioProject accession number PRJEB30272; <https://identifiers.org/ena.embl:PRJEB30272>.

The *C. gobio* genome sequencing is part of the Wellcome Sanger Institute's Vertebrate Sequencing project, and of the Vertebrate Genomes Project (VGP) ordinal references programme (Rhie *et al.*, 2020). All raw data and the assembly have been deposited in the ENA. Raw data and assembly accession identifiers are reported in Table 1.

Reporting guidelines

Not applicable.

Table 3. Software tools used for genome assembly.

Software tool	Version	Source
Falcon-unzip	falcon-2018.03.12-04.00	(Chin <i>et al.</i> , 2016)
wtdbg	1.1	(Ruan & Li, 2019)
cross_genome	2014-08-22	https://sourceforge.net/projects/phusion2/files/cross_genome/
PBJelly	PBSuite_15.8.24	(English <i>et al.</i> , 2012)
Canu	1.6	(Koren <i>et al.</i> , 2017)
Purge Haplotigs	v1	(Roach <i>et al.</i> 2018)
Juicebox		(Durand <i>et al.</i> , 2016; Robinson <i>et al.</i> , 2018)
scaff10x	1.0	https://github.com/wtsi-hpag/Scaff10X
Solve	Solve3.2.2_08222018	https://bionanogenomics.com/downloads/bionano-solve/
arrow	GenomicConsensus 2.2.2	https://github.com/PacificBiosciences/GenomicConsensus
Bwa-mem	0.7.17-r1188	(Li, 2013)
freebayes	v1.1.0-3-g961e5f3	(Garrison & Marth, 2012)
bcftools consensus	1.7	http://samtools.github.io/bcftools/bcftools.html

Consent

Not applicable.

Author contributions

RD, JHP, HWD, SAM, IB: designed the experiment. IB, MS, KO: generated data. HWD, TD, JHP: provided samples. SAM, IB, JW, ZN, RD: performed data analysis. JW, WC, KH, JT: performed data curation. VGP Consortium: provided guidance for methodology development. EAM, RD: supervised the work and provided funding. IB: wrote the manuscript. All authors reviewed and edited the final version of the manuscript.

Acknowledgements

We thank Arima Genomics for generating the Hi-C library, Bionano Genomics for generating Saphyr optical mapping data, Prof. Erich Jarvis for comments on the manuscript, and Richard Challis for help with BlobToolKit.

Author information

The Vertebrate Genomes Project Consortium includes Farooq O. Al-Ajli, Iliana Bista, Dave Burt, William Chow, Karen Clark, Hiram Clawson, Joanna Collins, Andrew J. Crawford, Joana Damas, Federica Di Palma, Mark Diekhans, Richard Durbin,

Olivier Fedrigo, Paul Flicek, Giulio Formenti, Arkarachai Functammasan, Erik Garrison, Jay Ghurye, M. Thomas P. Gilbert, Jennifer Marshall Graves, Dengfeng Guan, Bettina Haase, Leanne Haggerty, Brett T. Hannigan, Robert S. Harris, Alex Hastie, David Haussler, Jinna Hoffman, Kevin Howe, Kerstin Howe, Erich D. Jarvis, Warren E. Johnson, Juwan Kim, Heebal Kim, Sarah B. Kingan, Byung June Ko, Klaus-Peter Koepfli, Sergey Koren, Jonas Korch, Zev Kronenberg, Woori Kwak, Tanya M. Lama, Chul Lee, Joyce Lee, Harris Lewin, Kateryna D. Makova, Tomas Margues-Bonet, Fergal Martin, Patrick Masterson, Shane A. McCarthy, Paul Medvedev, Claudio V. Mello, Axel Meyer,

Mark Mooney, Jacquelyn Mountcastle, Robert W. Murphy, Eugene W. Myers, Luis Nassar, Gavin J.P. Naylor, Zemin Ning, Stephen J. O'Brien, Sadye Paez, Benedict Paten, Sarah Pelan, Trevor Pesout, Adam M. Phillippy, Martin Pippel, Damon-Lee Pointon, Arang Rhie, Oliver A. Ryder, Simona Secomandi, Siddarth Selvaraj, Beth Shapiro, Maria Simbirsky, Ying Sims, Michelle Smith, Ivan Sović, Emma C. Teeling, Constantina Theofanopoulou, Francoise Thibaud-Nissen, James Torrance, Alan Tracey, Marcela Uliano-Silva, Byrappa Venkatesh, Sonja C. Vernes, Brian P. Walenz, Tandy Warnow, Wesley C. Warren, Sylke Winkler, Jonathan Wood and Guojie Zhang.

References

- Balushkin AV: **Morphology, Classification, and Evolution of Notothenioid Fishes of the Southern Ocean (Notothenioidei, Perciformes).** *J Ichthyol.* 2000; **40**: S74–S109.
[Reference Source](#)
- Challis R, Richards E, Rajan J, *et al.*: **BlobToolKit - Interactive Quality Assessment of Genome Assemblies.** *G3.* 2020; **10**(4): 1361–1374.
[Publisher Full Text](#)
- Chen L, DeVries AL, Cheng CH: **Evolution of antifreeze glycoprotein gene from a trypsinogen gene in Antarctic notothenioid fish.** *Proc Natl Acad Sci U S A.* 1997; **94**(8): 3811–3816.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng CHC, Chen L, Near TJ, *et al.*: **Functional antifreeze glycoprotein genes in temperate-water New Zealand notothenioid fish infer an Antarctic evolutionary origin.** *Mol Biol Evol.* 2003; **20**(11): 1897–1908.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chin CS, Peluso P, Sedlazeck FJ, *et al.*: **Phased diploid genome assembly with single-molecule real-time sequencing.** *Nat Methods.* 2016; **13**(12): 1050–1054.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chow W, Brugger K, Caccamo M, *et al.*: **gEVAL - a web-based browser for evaluating genome assemblies.** *Bioinformatics.* 2016; **32**(16): 2508–2510.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Detrich W, Buckley B, Doolittle D, *et al.*: **Sub-Antarctic and High Antarctic Notothenioid Fishes: Ecology and Adaptational Biology Revealed by the ICEFISH 2004 Cruise of RVIB Nathaniel B. Palmer.** *Oceanography.* 2012; **25**(3): 184–187.
[Publisher Full Text](#)
- Dornburg A, Federman S, Lamb AD, *et al.*: **Cradles and museums of Antarctic teleost biodiversity.** *Nat Ecol Evol.* 2017; **1**(9): 1379–1384.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Durand NC, Robinson JT, Shamim MS, *et al.*: **Juicebox Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom.** *Cell Syst.* 2016; **3**(1): 99–101.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- English AC, Richards S, Han Y, *et al.*: **Mind the gap: upgrading genomes with Pacific Biosciences RS long-read sequencing technology.** *PLoS One.* 2012; **7**(11): e47768.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Garrison E, Marth G: **Haplotype-based variant detection from short-read sequencing.** In *arXiv [q-bio.GN]. arXiv.* 2012.
[Reference Source](#)
- Ghurye J, Pop M, Koren S, *et al.*: **Scaffolding of long read assemblies using long range contact information.** *BMC Genomics.* 2017; **18**(1): 527.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Giordano D, Grassi L, Parisi E, *et al.*: **Embryonic β -globin in the non-Antarctic notothenioid fish *Cottoperca gobio* (Bovichtidae).** *Polar Biol.* 2006; **30**(1): 75–82.
[Publisher Full Text](#)
- Giordano D, Boechi L, Vergara A, *et al.*: **The hemoglobins of the sub-Antarctic fish *Cottoperca gobio*, a phylogenetically basal species--oxygen-binding equilibria, kinetics and molecular dynamics.** *FEBS J.* 2009; **276**(8): 2266–2277.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Grass M, Busbridge T, Blake A, *et al.*: **Cruise report ZDLM3-02-2018, Ground fish survey, Falkland Islands Government, Directorate of Natural Resources, Fisheries, Stanley, Falkland Islands.** 2018.
- Hunt SE, McLaren W, Gil L, *et al.*: **Ensembl variation resources.** *Database (Oxford).* 2018; **2018**: bay119.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Koren S, Walenz BP, Berlin K, *et al.*: **Canu: scalable and accurate long-read assembly via adaptive k -mer weighting and repeat separation.** *Genome Res.* 2017; **27**(5): 722–736.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Krzywinski M, Schein J, Birol I, *et al.*: **Circos: an information aesthetic for comparative genomics.** *Genome Res.* 2009; **19**(9): 1639–1645.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li H: **Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.** In *arXiv [q-bio.GN]. arXiv.* 2013.
[Reference Source](#)
- Mazzei F, Ghigliotti L, Lecointre G, *et al.*: **Karyotypes of basal lineages in notothenioid fishes: the genus *Bovichtus*.** *Polar Biol.* 2006; **29**(12): 1071.
[Publisher Full Text](#)
- Near TJ, Dornburg A, Harrington RC, *et al.*: **Identification of the notothenioid sister lineage illuminates the biogeographic history of an Antarctic adaptive radiation.** *BMC Evol Biol.* 2015; **15**: 109.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pisano E, Ozouf-Costaz C, Hureau JC, *et al.*: **Chromosome differentiation in the subantarctic Bovichtidae species *Cottoperca gobio*.** (Günther, 1861) and *Pseudaphritis urvillii* (Valenciennes, 1832) (Pisces, Perciformes). *Antarctic Science / Blackwell Scientific Publications.* 1995; **7**(4): 381–386.
[Publisher Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, *et al.*: **Towards complete and error-free genome assemblies of all vertebrate species.** In *bioRxiv.* 2020; (p. 2020.05.22.110833).
[Publisher Full Text](#)
- Roach MJ, Schmidt SA, Borneman AR, *et al.*: **Purge Haplotigs: allelic contig reassignment for third-gen diploid genome assemblies.** *BMC Bioinformatics.* 2018; **19**(1): 460.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Robinson JT, Turner D, Durand NC, *et al.*: **Juicebox.js Provides a Cloud-Based Visualization System for Hi-C Data.** *Cell Syst.* 2018; **6**(2): 256–258.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ruan J, Li H: **Fast and accurate long-read assembly with wtdbg2.** In *bioRxiv.* (p. 530972). 2019.
[Publisher Full Text](#)
- Simão FA, Waterhouse RM, Ioannidis P, *et al.*: **BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.** *Bioinformatics.* 2015; **31**(19): 3210–3212.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Smith JJ, Stuart AB, Sauka-Spengler T, *et al.*: **Development and analysis of a germline BAC resource for the sea lamprey, a vertebrate that undergoes substantial chromatin diminution.** *Chromosoma.* 2010; **119**(4): 381–389.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Steinke D, Salzburger W, Meyer A: **Novel relationships among ten fish model species revealed based on a phylogenomic analysis using ESTs.** *J Mol Evol.* 2006; **62**(6): 772–784.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 06 November 2020

<https://doi.org/10.21956/wellcomeopenres.17562.r41211>

© 2020 Li C et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Chenhong Li 

Shanghai Universities Key Laboratory of Marine Animal Taxonomy and Evolution, Shanghai Ocean University, Shanghai, China

Liang Lu

Shanghai Ocean University, Shanghai, China

The authors reported a well-assembled genome of an interesting species. The methods used are sound and sufficient, and the datasets are easily accessible. The only thing I found is that they maybe want to elaborate the importance of sequencing this species and to use it as an outgroup for comparative genomics in the Antarctic notothenioids, in addition to conservation of this species per se. Or, if possible make a comparison with the published Antarctic fish, which would increase the interest much more and maybe number of citations as well.

Comments from my Doctoral student (not that I agree with them all):

1. Have the barcodes been trimmed when using 10X Illumina data for polishing?
2. The genes were annotated, but how their function was predicted was not clear.
3. The sample numbers, BioProject, BioSamples and the length of each chromosome can be lodged in the "data available" part, not in the main text.
4. The tools used were detailed in the text, which is redundant to include them in Table 3.
5. The biological background was touched in the introduction, but no any analyses were used to address any biological questions. (I understand that this is a data note though).

Is the rationale for creating the dataset(s) clearly described?

Partly

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Molecular phylogenetics, genomics, bioinformatics.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 05 August 2020

<https://doi.org/10.21956/wellcomeopenres.17562.r39467>

© 2020 Bilyk K. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Kevin Bilyk

Department of Biology, Montclair State University, Montclair, NJ, USA

The manuscript is reporting the newly available genome assembly for *Cottoperca gobio*, a basal notothenioid that diverged before the evolution of the cold-adaptations and cold-specialization that define the cryonotothenioids. One of the major challenges in studying the Antarctic notothenioids has been limited availability of genetic and genomic information from phylogenetically close temperate species, making comparative investigations difficult. The *C. gobio* genome plays an incredibly important role filling this gap, providing a temperate companion to the recent *Eleginops maclovinus* genome. These provide a far more appropriate baseline to evaluate the changes that have come with evolution in a polar environment than the traditional model fish species all of which are far more phylogenetically distant. The genome sequencing, assembly, and annotation are all well described and appear handled appropriately. Personally, I can only say that I am happy that this genome is now available and look forward to utilizing it in my own work over the coming months.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Comparative physiology, transcriptomics, genomics and bioinformatics.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
