

Short report

Open Access

Regulation of thymus-dependent and thymus-independent production of immunoglobulin G subclasses by $G\alpha_{12}$ and $G\alpha_{13}$

Song Jin Lee^{1,2}, Woo Hyung Lee¹, Chang Ho Lee³ and Sang Geon Kim*¹

Address: ¹College of Pharmacy and Research Institute of Pharmaceutical Sciences, Seoul National University, Seoul 151-742, Korea, ²Yuhan Corp., Daebang-dong, Dongjak-gu, Seoul 156-754, Korea and ³Department of Pharmacology and Institute of Biomedical Science, College of Medicine, Hanyang University, Seoul 133-791, Korea

Email: Song Jin Lee - lsj30664@snu.ac.kr; Woo Hyung Lee - lk404@snu.ac.kr; Chang Ho Lee - jennysue@hanyang.ac.kr; Sang Geon Kim* - sgk@snu.ac.kr

* Corresponding author

Published: 12 July 2008

Received: 16 May 2008

Journal of Molecular Signaling 2008, 3:12 doi:10.1186/1750-2187-3-12

Accepted: 12 July 2008

This article is available from: <http://www.jmolecularsignaling.com/content/3/1/12>

© 2008 Lee et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: A previous study from this laboratory showed that $G\alpha_{12}$ members participate in the production of inflammatory cytokines. In spite of the identification of B cell homeostasis responses regulated by $G\alpha_{13}$, the functional roles of $G\alpha_{12}$ members in the production of immunoglobulin (Ig) isotypes remained unknown. This study investigated whether $G\alpha_{12}$ members are involved in the Ig isotype antibody production with the purpose of establishing their functions in thymus-dependent and thymus-independent humoral responses.

Results: Mice lacking $G\alpha_{12}$ and/or $G\alpha_{13}$ showed an impaired antigen-specific antibody production promoted by challenge(s) of ovalbumin or trinitrophenyl-lipopolysaccharide (TNP-LPS), used for thymus-dependent and thymus-independent stimuli, respectively. Homozygous knockout (KO) of $G\alpha_{12}$ or double heterozygous KO of $G\alpha_{12}/G\alpha_{13}$ significantly reduced the antigen-specific total IgG level after multiple ovalbumin immunizations with decreases in the production of IgG1, IgG2a and IgG2b subclasses, as compared to wild type control. In contrast, IgM production was not decreased. Moreover, mice deficient in $G\alpha_{12}$ or partially deficient in $G\alpha_{13}$ or $G\alpha_{12}/G\alpha_{13}$ showed significantly low production of IgG2b in response to TNP-LPS. In TNP-LPS-injected mice, IgG1 and IgG2a productions were unaffected by the G protein KOs.

Conclusion: Our results demonstrate that both $G\alpha_{12}$ and $G\alpha_{13}$ are essentially involved in thymus-dependent and independent production of IgG subclasses, implying that the G-proteins contribute to the process of antigen-specific IgG antibody production.

Background

G-protein-coupled receptors (GPCR) regulate signal transduction in cells, participating in a variety of biological processes [1]. An activated GPCR makes a complex with a G-protein defined by α subunit. $G\alpha$ subunits consist with

$G\alpha_s$, $G\alpha_i$ or $G\alpha_q$ and $G\alpha_{12}$ [1]. The members of $G\alpha_{12}$ family consisting of $G\alpha_{12}$ and $G\alpha_{13}$ are critical mediators in regulating effectors or cellular responses. Their interactions with specific Rho guanine nucleotide exchanging factor result in small GTPase RhoA activation, leading to diverse

biological functions such as migration and maturation of B cells, morphology and motility of cells, and smooth muscle contraction [2-4].

Activated B cells proliferate and differentiate into immunoglobulin (Ig)-producing plasma cells or long-lived memory cells [5]. It is well recognized that Ig antibody (Ab) production by B cells is the critical process in the defense against infection. Immune cells express a variety of GPCRs, whose activations result in coupling of G-proteins for signal amplifications [1]. In particular, it has been shown that $G\alpha_{12}$ and $G\alpha_{13}$ regulate the homeostasis of marginal zone B cells, in which the G-proteins activated LSC, a p115RhoGEF, for humoral responses [2]. Our studies showed that activation of $G\alpha_{12}$ and/or $G\alpha_{13}$ leads to the induction of iNOS and COX-2 through NF- κ B, an essential transcription factor required for immune responses [6,7]. Moreover, sphingosine 1-phosphate (S1P), a representative lysophospholipid GPCR ligand in blood or lymph nodes, disseminates the signal to B cells via S1P₃ GPCR to position marginal zone B cells [8].

Despite the identified role of $G\alpha_{12}/G\alpha_{13}$ in the regulation of marginal zone B cell biology [2], the functions of $G\alpha_{12}/G\alpha_{13}$ in Ig production have not been completely identified. In view of complex network of the immune system and involvements of diverse cell types, we investigated whether the $G\alpha_{12}$ family members regulate the production of Ig classes and IgG subclasses with particular emphasis on their roles in thymus-dependent or thymus-independent immunity. In this study, Ab production was measured after challenges of two different types of antigens. To elicit thymus-dependent immune response, wild type (WT) or $G\alpha_{12}$ and/or $G\alpha_{13}$ heterozygous or homozygous knockout (KO) mice were immunized with ovalbumin (OVA). In another set, the animals were injected with trinitrophenyl-lipopolysaccharide (TNP-LPS) to promote thymus-independent Ig production. Through these *in vivo* analyses, it was found that $G\alpha_{12}$ and

$G\alpha_{13}$ have regulatory functions in producing IgG subclasses. This information may bring insight in understanding the role of $G\alpha_{12}$ members in humoral immune responses.

Results

To determine the possible role(s) of $G\alpha_{12}$ and/or $G\alpha_{13}$ in the regulation of thymus-dependent Ab production, the Ig titers were measured in WT mice or mice deficient in $G\alpha_{12}/G\alpha_{13}$ that had been immunized with OVA and booster-injected with the same antigen 2 weeks after, which was followed by OVA nebulizations (Fig. 1A). After the immunizations and nebulizations, WT mice showed normal OVA-specific IgG and IgM production (Fig. 1A, Table 1). In contrast, homozygous absence of the $G\alpha_{12}$ gene significantly impaired OVA-specific IgG production. When OVA-specific Ig subclass contents were assessed, $G\alpha_{12}$ deficiency decreased the production of IgG1, IgG2a and IgG2b subclasses compared to those in WT. Also, double heterozygous deletions of $G\alpha_{12}$ and $G\alpha_{13}$ significantly reduced OVA-specific IgG content with decreases in the levels of IgG1, IgG2a or IgG2b subclasses. The increased production of IgM was unaffected by the absence of $G\alpha_{12}$ and/or $G\alpha_{13}$. In this assay, we could not use the animals with homozygous $G\alpha_{13}$ KO because these mice die before birth [9]. Our data demonstrated that $G\alpha_{12}$ and $G\alpha_{13}$ are both required for thymus-dependent IgG1, IgG2a and IgG2b production.

LPS is a classic mitogenic stimulus that activates the innate B cell receptors [5]. It is known that LPS stimulation activates B cells without T cell help. In another set of experiments, we determined the roles of $G\alpha_{12}/G\alpha_{13}$ for the regulation of B cell immune response stimulated by TNP-LPS, a well-known thymus-independent antigen. In WT mice, total Ig production by TNP-LPS was greatly promoted 2 weeks after (Fig. 1B). The heterozygous or homozygous KO of $G\alpha_{12}$ inhibited TNP-LPS-specific IgG production in a gene dose-dependent manner. In the mice

Table 1: Production of Ig classes and IgG subclasses in WT mice or mice deficient in $G\alpha_{12}$ and/or $G\alpha_{13}$

	Thymus-dependent Ig production					Thymus-independent Ig production				
	WT	$G\alpha_{12} +/-$	$G\alpha_{12} -/-$	$G\alpha_{13} +/-$	$G\alpha_{12}/G\alpha_{13} +/-$	WT	$G\alpha_{12} +/-$	$G\alpha_{12} -/-$	$G\alpha_{13} +/-$	$G\alpha_{12}/G\alpha_{13} +/-$
IgG	++	++	+	+	+	++++	+++	+	++	+
IgG1	+++	+++	+	++	+	+	+	+	+	+
IgG2a	++	+	+	+	+	+	+	+	+	+
IgG2b	++	++	+	+	+	++++	+++	+	++	+
IgG3	t	t	t	t	t	+	+	+	+	++
IgM	+	+	+	++	+	+	+	+	+	+
IgA	t	t	t	t	t	t	t	t	t	t
IgE	t	t	t	t	t					Not detected

The relative Ig production was defined as follows: -, OD₄₅₀ < 0.15; +, OD₄₅₀ 0.15–0.4; ++, OD₄₅₀ 0.4–0.75; +++, OD₄₅₀ 0.75–1.0; and +++++, OD₄₅₀ > 1.0. The absorbance at 450 nm (OD₄₅₀) was measured using ELISA assays. Number of animals in each treatment group = 10

LPS-specific IgG1, IgG2a, and IgM contents were not changed by the gene KO's, suggesting that the change in TNP-LPS-specific IgG production might be due to that in IgG2b. Thus, Ig production in response to TNP-LPS was found to be regulated by $G\alpha_{12}$ and $G\alpha_{13}$. All of these results provide evidence that $G\alpha_{12}$ and $G\alpha_{13}$ are both required for regulating thymus-dependent and thymus-independent production of IgG subclasses.

Discussion

There are two distinctive pathways that induce B cell responses (that is, thymus-dependent and thymus-independent B cell activation) [10,11]. In the present study, we demonstrated for the first time that $G\alpha_{12}$ and $G\alpha_{13}$ are both required for thymus-dependent production of IgG subclasses. Significant changes in the humoral response, particularly in regulating IgG, by the lack(s) of $G\alpha_{12}$ and $G\alpha_{13}$ highlight the need to consider the G-protein pathway as one of important regulatory controls for T-cell dependent immune network.

Affinity maturation of B cells, which need Th1 cytokines, consists with the processes of clonal proliferation, somatic hypermutation, and selection [10]. Th1 cytokines such as interferon- γ (IFN- γ) stimulate IgG2a, IgG2b, and IgG3 production [11]. Therefore, our results showing that $G\alpha_{12}$ and $G\alpha_{13}$ deficiency notably decreased IgG2a and IgG2b subclasses in an OVA-immunized model suggest that the affinity maturation processes of B cells might be affected by $G\alpha_{12}$ and $G\alpha_{13}$. Th2 cytokine (e.g., interleukin-4) particularly induces Ig switching to IgG1 and IgE [12]. Our data indicated that the titer of anti-OVA IgG1, but not anti-OVA IgE, was lowered by $G\alpha_{12}/G\alpha_{13}$ deficiency, which suggested that maturation of B cell to IgG1-producing plasma cell might need the G-protein-mediated signaling processes presumably upon stimulation of Th2 cytokine(s). Collectively, the essential regulatory function of $G\alpha_{12}/G\alpha_{13}$ in producing IgG subclasses lends support to the hypothesis that helper T cell-dependent B cell activation by antigen requires the G-protein-mediated signaling pathway.

Another type of B cell activation is thymus-independent, which is triggered by viral particles, common bacterial antigens, and TLR ligands without T cell help [5]. B cell activation by thymus-independent antigens (e.g., LPS) causes the production of polyclonal antibody [13]. In our animal model, TNP-LPS-inducible production of IgG2b, but not IgG3, was decreased by the absence of $G\alpha_{12}$ and $G\alpha_{13}$, showing that the G-proteins might regulate thymus-independent IgG production by B cells. Because transforming growth factor- β stimulates IgG2b and IgA production [14], the decrease in IgG2b in TNP-LPS-injected mice deficient in $G\alpha_{12}/G\alpha_{13}$ might be associated with repression of transforming growth factor- β . This possibil-

ity is strengthened by our finding that $G\alpha_{12}/G\alpha_{13}$ regulate transforming growth factor- β production in the liver of mice challenged with dimethylnitrosamine (Lee et al, unpublished data).

$G\alpha_{12}$ regulates NF- κ B-mediated COX-2 induction by S1P in a process mediated by the JNK-dependent ubiquitination and degradation of I κ B α [7]. Because antigen-induced cell signaling requires NF- κ B and AP-1, decreased activation of the transcription factors may account for altered IgG production in the KO mice. Our results illustrating the role of $G\alpha_{12}$ and $G\alpha_{13}$ in IgG production may be explained by other possibilities: that is, (1) the role of $G\alpha_{12}/G\alpha_{13}$ in B cell reentry into the secondary lymphoid organ and (2) the regulatory role of $G\alpha_{12}/G\alpha_{13}$ in the specific step of Ab production.

Methods

OVA or TNP-LPS Immunizations

All experiments were conducted under the guidelines of the Institutional Animal Use and Care Committee at Seoul National University, Korea. WT and $G\alpha_{12}/G\alpha_{13}$ KO mice at the age of 8–10 weeks (25~30 g) ($G\alpha_{12}^{+/-}$, $G\alpha_{12}^{-/-}$, $G\alpha_{13}^{+/-}$, and $G\alpha_{12}/G\alpha_{13}^{+/-}$) were used for *in vivo* experiments. To induce OVA-specific Ab production, the mixture of 100 μ g endotoxin-free OVA (MP Biomedicals, Aurora, OH) dissolved in PBS and alum (Pierce, IL) was i.p. injected to the mice on day 1. On day 14, the mice were i.p. injected again with 100 μ g OVA. The mice were subsequently exposed to aerosol of 1% OVA solution for 30 min once a day on days 26, 27 and 28. In another set of experiments, a single dose of TNP-LPS (Biosearch Technologies, Novato, CA) was i.p. injected to WT and $G\alpha_{12}/G\alpha_{13}$ KO mice to assess the production of TNP-LPS-specific Ab. The animals were bled 2 weeks after.

Enzyme-Linked Immunosorbent Assay (ELISA)

ELISA assays were performed to determine antigen-specific total IgG, IgG1, IgG2a, IgG2b, IgG3, IgA, IgE and IgM in aliquots of diluted serum (1:2000) in a 96-well plate (Maxisorp, Nunc Co., Rochester, NY). Alkaline phosphatase-conjugated goat anti-mouse Ig isotype and IgG subclass (Southern Biotechnology Associates Inc., Birmingham, Ala), and *p*-nitrophenyl phosphate (phosphatase substrate) were used to assay titers.

Statistical Analysis

One way analysis of variance (ANOVA) procedures were used to assess significant differences among treatment groups. The Newman-Keuls test was used for comparisons of multiple group means.

Abbreviations

Ab: antibody; GPCRs: G-Protein-coupled receptors; G-proteins: GTP-binding proteins; OVA: ovalbumin; LPS:

lipopolysaccharide; TNP: trinitrophenyl; Ig: immunoglobulin; S1P: sphingosine 1-phosphate; WT: wild type; KO: knockout.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SJL, CHL, SGK designed research; SJL and WHL performed research; SJL and WHL analyzed data; and CHL and SGK wrote the paper. All authors read and approved of the final manuscript.

Acknowledgements

This work was supported by the Korea Science and Engineering Foundation (KOSEF) grant funded by the Korea government (MEST)(No.R11-2007-107-01001-0), South Korea.

References

1. Riobo NA, Manning DR: **Receptors coupled to heterotrimeric G proteins of the G12 family.** *Trends Pharmacol Sci* 2005, **26**:146-154.
2. Girkontaite I, Missy K, Sakk V, Harenberg A, Tedford K, Pötzel T, Pfeffer K, Fischer KD: **Lsc is required for marginal zone B cells, regulation of lymphocyte motility and immune responses.** *Nat Immunol* 2001, **2**:855-862.
3. Johnson EN, Drucey KM: **Heterotrimeric G protein signaling: role in asthma and allergic inflammation.** *J Allergy Clin Immunol* 2002, **109**:592-602.
4. Rieken S, Sassmann A, Herroeder S, Wallenwein B, Moers A, Offermanns S, Wettschureck N: **G12/G13 family G proteins regulate marginal zone B cell maturation, migration, and polarization.** *J Immunol* 2006, **177**:2985-2993.
5. Gray D, Gray M, Barr T: **Innate responses of B cells.** *Eur J Immunol* 2007, **37**:3304-3310.
6. Kang KW, Choi SY, Cho MK, Lee CH, Kim SG: **Thrombin induces nitric-oxide synthase via Galpha12/13-coupled protein kinase C-dependent I-kappaBalpha phosphorylation and JNK-mediated I-kappaBalpha degradation.** *J Biol Chem* 2003, **278**:17368-17378.
7. Ki SH, Choi MJ, Lee CH, Kim SG: **Galpha12 specifically regulates COX-2 induction by sphingosine 1-phosphate. Role for JNK-dependent ubiquitination and degradation of I-kappaBalpha.** *J Biol Chem* 2007, **282**:1938-1947.
8. Cinamon G, Zachariah MA, Lam OM, Foss FW Jr, Cyster JG: **Follicular shuttling of marginal zone B cells facilitates antigen transport.** *Nat Immunol* 2008, **9**:54-62.
9. Offermanns S, Mancino V, Revel JP, Simon MI: **Vascular system defects and impaired cell chemokinesis as a result of Galpha13 deficiency.** *Science* 1997, **275**:533-536.
10. Okada T, Cyster JG: **B cell migration and interactions in the early phase of antibody responses.** *Curr Opin Immunol* 2006, **18**:278-285.
11. Mosmann TR, Coffman RL: **TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties.** *Annu Rev Immunol* 1989, **7**:145-173.
12. Snapper CM, Mond JJ: **Towards a comprehensive view of immunoglobulin class switching.** *Immunol Today* 1993, **14**:15-17.
13. Coutinho A, Poltorack A: **Innate immunity: from lymphocyte mitogens to Toll-like receptors and back.** *Curr Opin Immunol* 2003, **15**:599-602.
14. Stavnezer J: **Regulation of antibody production and class switching by TGF-beta.** *J Immunol* 1995, **155**:1647-1651.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

