

Supplementary Material

Estimating the risk of incident risk of SARS-CoV-2 infection among healthcare workers residing in Egyptian quarantine hospitals

Sofía Jijón^{1,2†} (PhD), Ahmad Al Shafie^{3†} (MSc), Essam Hassan⁴, EMAE-MESuRS working group on nosocomial SARS-CoV-2 modeling, Laura Temime^{1,2} (PhD), Kévin Jean^{1,2,4} (PhD), Mohamed El-Kassas^{3*} (MD)*

¹ Laboratoire Modélisation, épidémiologie et surveillance des risques sanitaires (MESuRS), Conservatoire national des Arts et Métiers, Paris, France

² Unité PACRI, Institut Pasteur, Conservatoire national des Arts et Métiers, Paris, France

³ Endemic Medicine Department, Faculty of Medicine, Helwan University, Cairo, Egypt

⁴ Tropical Medicine Department, Faculty of Medicine, Fayoum University, Fayoum, Egypt

⁵ MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, Imperial College London, United Kingdom

Supplementary figures




Figure S1 Markov Chain Monte-Carlo diagnostics

Trace plot of the posterior distribution for the force of infection (λ), for Hosp1–3. In 10,000 iterations, the sample space is explored many times and no high serial correlation is apparent in the chain.