

Supplementary Figures

Bioprospecting Antarctic Microorganisms: Investigating *Hymenobacter psoromatis* PAMC26554 for Biochemical Potential

Hirotake Yamaguchi^{1†}, Ryoichi Yamada^{1†}, Jun Hyuck Lee², and Tae-Jin Oh^{1,3,4,5*}

¹Department of Life Science and Biochemical Engineering, Graduate School, SunMoon University, Asan 31460, Republic of Korea

²Research Unit of Cryogenic Novel Materials, Korea Polar Research Institute, Incheon 21990, Republic of Korea

³Bio Big Data-based Chungnam Smart Clean Research Leader Training Program, SunMoon University, Asan 31460, Republic of Korea

⁴Genome-based BioIT Convergence Institute, Asan 31460, Republic of Korea

⁵Department of Pharmaceutical Engineering and Biotechnology, SunMoon University, Asan 31460, Republic of Korea

[†]These authors contributed equally to this work

***Correspondence**

Prof. Tae-Jin Oh, E-mail: tjoh3782@sunmoon.ac.kr




Fig. S1. The NRPS-T1PKS-NRPS-like biosynthetic gene cluster in *Hymenobacter psoromatis* PAMC26554 is illustrated. The graphical output generated by antiSMASH for region 3 is displayed, including the following components: KR (ketoreductase), C (condensation domain), A (adenylation domain), TE (thioesterase), AT (acyltransferase), and AmT (Aminotran_3). The carrier protein domains—which comprise the acyl carrier protein domain for PKS and the peptidyl carrier protein domain for NRPS—are represented by light blue ovals.




Fig. S2. Comparison of chromatograms of EA extracts of YPM medium only (A) and EA extracts of PAMC26554 (B).




Fig. S3. ^{13}C NMR spectra of PAA.




Fig. S4. COSY spectra of PAA.