Check for updates

RESEARCH ARTICLE

REVISED Effects of paracetamol (acetaminophen) on gene expression and permeability properties of the rat placenta and fetal brain [version 2; peer review: 2 approved]

Liam M. Koehn^{*}, Yifan Huang[®], Mark D Habgood, Kai Kysenius, Peter J. Crouch, Katarzyna M Dziegielewska, Norman R Saunders[®]

Pharmacology & Therapeutics, University of Melbourne, Parkville, Victoria, 3010, Australia

* Equal contributors

V2 First published: 08 Jun 2020, 9:573 https://doi.org/10.12688/f1000research.24119.1 Latest published: 19 Aug 2020, 9:573 https://doi.org/10.12688/f1000research.24119.2

Abstract

Background: Paracetamol (acetaminophen) is widely used in pregnancy and generally regarded as "safe" by regulatory authorities. Methods: Clinically relevant doses of paracetamol were administered intraperitoneally to pregnant rats twice daily from embryonic day E15 to 19 (chronic) or as a single dose at E19 (acute). Control samples were from un-treated age-matched animals. At E19, rats were anaesthetised, administered a final paracetamol dose, uteruses were opened and fetuses exposed for sample collection. For RNA sequencing, placentas and fetal brains were removed and flash frozen. Fetal and maternal plasma and cerebrospinal fluid were assayed for α -fetoprotein and interleukin 1 β (IL1 β). Brains were fixed and examined (immunohistochemistry) for plasma protein distribution. Placental permeability to a small molecule (¹⁴C-sucrose) was tested by injection into either mother or individual fetuses; fetal and maternal blood was sampled at regular intervals to 90 minutes. **Results:** RNA sequencing revealed a large number of genes up- or down-regulated in placentas from acutely or chronically treated animals compared to controls. Most notable was down-regulation of three acute phase plasma proteins (α-fetoprotein, transferrin, transthyretin) in acute and especially chronic experiments and marked up-regulation of immune-related genes, particularly cytokines, again especially in chronically treated dams. IL1ß increased in plasma of most fetuses from treated dams but to variable levels and no IL1B was detectable in plasma of control fetuses or any of the dams. Increased placental permeability appeared to be only from fetus to mother for both ¹⁴C-sucrose and α -fetoprotein, but not in the reverse direction. In the fetal brain, gene regulatory changes were less prominent than in the placenta of treated fetuses and did not involve inflammatoryrelated genes; there was no evidence of increased blood-brain barrier permeability.



- 1. **Roland Bainton**, University of California, San Francisco (UCSF), San Francisco, USA
- 2. Patrizia Ferretti D, University College London, London, UK

Any reports and responses or comments on the article can be found at the end of the article.

Conclusion: Results suggest that paracetamol may induce an immune-inflammatory-like response in placenta and more caution should be exercised in use of paracetamol in pregnancy.

Keywords

placenta, transfer, inflammation, permeability, interleukin-1 β , IL1 β , α -fetoprotein, AFP, immune response

Corresponding author: Norman R Saunders (n.saunders@unimelb.edu.au)

Author roles: Koehn LM: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; Huang Y: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; Habgood MD: Conceptualization, Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; Kysenius K: Methodology, Writing – Review & Editing; Crouch PJ: Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; Dziegielewska KM: Conceptualization, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; Saunders NR: Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Part of this study was supported by the CASS Foundation, Victoria, Australia [7981]. KK was in receipt of an Early Career Researcher grant from The University of Melbourne [1859114].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Koehn LM *et al*. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Koehn LM, Huang Y, Habgood MD *et al.* Effects of paracetamol (acetaminophen) on gene expression and permeability properties of the rat placenta and fetal brain [version 2; peer review: 2 approved] F1000Research 2020, 9:573 https://doi.org/10.12688/f1000research.24119.2

First published: 08 Jun 2020, 9:573 https://doi.org/10.12688/f1000research.24119.1

REVISED Amendments from Version 1

Statistics: Added "We also tested our data using ANOVA followed by Tukey's posthoc test. The significance levels are the same as those we obtained with a t-test."

Table 1: Colour coding added, explanation in legend.

Table 2: Legend: Added "(p<0.05, see Methods)." Revised as suggested by Reviewer #2 re inflammatory immune-related genes. "Symbol - indicates no significant difference in fold changes; it does not indicate that there was no fold change. This table includes only genes with inflammatory and immunerelated functions and thus does not include some of the highly expressed genes in Table 1."

Table 3: Colour coding added, explanation in legend.

The effect of paracetamol exposure on placental gene expression (transcriptomic analysis): Number of upregulated genes corrected to 64. P-value added.

IL1ß concentration (ELISA): Clarification/correction of IL1ß results: "None of the dams in any of the treatment groups had a detectable level of IL1ß in their plasma (limit <5pg/ml) nor was IL1ß detected nor was IL1ß detected in the control untreated fetuses. In contrast, IL1ß in the plasma of many of the E19 fetuses whose mothers had been treated with paracetamol was detected. The levels were generally higher in fetuses of mothers treated chronically (acute 2/4, chronic low 7/16 and chronic high 10/19)."

Figure 5: Redrawn.

The effect of paracetamol exposure on E19 fetal brain gene expression (transcriptomic analysis): Added "Following both treatments 26/50 genes were up-regulated and 40/50 were down-regulated."

Figure 6: Legend rewritten to clarify n numbers. "Each data point is a single fetus."

Figure 7: Legend rewritten to clarify n numbers. "Each data point is a single fetus."

Figure 8: Text and legend modified to correct error in Figure 8A. Figure 8A revised to correct error in number of controls; also corrected in legend and text: "Treatment groups were: control (1/1b; n=2), acute (2/2b; n=1),.....".

Any further responses from the reviewers can be found at the end of the article

Abbreviations

AFP, α -fetoprotein; CSF, cerebrospinal fluid; DPM, disintegrations per minute; E, embryonic (note that by longstanding convention all gestational ages in rodents are referred to as embryonic, but in this study E19 is a fetal stage); IL1 β , Interleukin 1 β cytokine; i.p., intraperitoneal; i.v., intravenous; P, postnatal; RNA-Seq, RNA sequencing; SD, standard deviation; μ Ci, micro Curie.

Introduction

Paracetamol (acetaminophen) is commonly taken, either by prescription or self-medication, for the relief of pain and fever. In pregnancy it is the most widely consumed drug, with estimates of expectant mothers talking this medication ranging from 56% in Australia and the Americas (Wyszynski & Shields, 2016) to 76% in Europe (Dreyer *et al.*, 2015). The Australian Medicines Handbook (2019) states without qualification that paracetamol is safe for use in pregnancy and breast-feeding. However, epidemiological reports of behavioural effects in the offspring of mothers who took paracetamol during pregnancy are beginning to be published, suggesting a more cautious approach would be appropriate (see Bauer *et al.*, 2018 and Discussion).

In a recent study we have found that paracetamol, when administered to a pregnant dam at doses within the clinical range used in patients, transfers across the placenta to reach the fetus at about 40% of the levels of the drug in the maternal circulation (Koehn et al., 2019b). Thus, the placenta provides a degree of protection for the developing fetus but the mechanisms involved are not yet understood, nor are the effects that paracetamol may have on placental functions. We have therefore carried out an RNA sequencing (RNA-Seq) study of E19 placentas and brains from control (un-treated) rats and from rats treated with a single (acute) or multiple (chronic) doses of paracetamol. This RNA-Seq study yielded the unexpected outcome of widespread up-regulation of inflammatory and immunerelated genes in the placenta of the dam exposed to paracetamol over a prolonged period, with a much less pronounced effect on inflammatory-related genes following a single dose; however, many other genes showed a regulatory response following a single dose of paracetamol. Inflammatory responses during pregnancy have been linked to a range of clinical complications including pre-term birth, fetal cardiac conditions and neurological deficiencies (Challis et al., 2009; Fleiss et al., 2020; Huleihel et al., 2004; Romero et al., 2007; Salafia et al., 1989). High cytokine levels in blood have been linked to increased blood-brain barrier permeability (Anthony et al., 1997; Stolp et al., 2005a) and possibly leading to a range of health complications (Brochu et al., 2011; Nelson et al., 1998; Thornton et al., 2017). Inflammation in the placenta has also been linked to increased placental permeability, as shown in studies that identified a size-dependent increase in maternal-fetal nanoparticle transfer in mice (Tian et al., 2013).

In the present study, the inflammatory response in the placenta and the fetal brain following maternal paracetamol exposure was examined to see if it was associated with alterations in placental and blood-brain barrier permeability. Placental permeability was assessed using a low-molecular weight, hydrophobic molecule sucrose to determine the transfer in both directions: from the dam's circulation to the fetal circulation and from the fetal circulation back to the dam. Transfer of a large molecule, the endogenous fetal-derived plasma protein α -fetoprotein (AFP), across the placenta into maternal circulation was also investigated. Results from both of these markers indicate that placental transfer was potentially affected by paracetamol treatment, and demonstrated increased levels of AFP detected in blood plasma of dams treated with paracetamol. The inflammatory cytokine IL-1bß was measured in fetal and maternal plasma; it showed higher levels only in fetal plasma following maternal paracetamol treatment. Permeability of the fetal blood-brain barrier to both small (sucrose) and large (plasma protein) molecules was not affected in spite of increased IL-1ß levels in fetal circulation. The results presented here highlight responses to paracetamol use during pregnancy that appear to be tissue-specific and dependent on duration of treatment. The results are discussed in the context of the appropriate use of paracetamol during pregnancy.

Methods

Ethical statement

The animals used in this study were the Sprague Dawley strain of *Rattus norvegicus*. All animal experimentation was approved by the University of Melbourne Animal Ethics Committee (Ethics Permission AEC: 1714344.1) and conducted in compliance with Australian National Health and Medical Research Guidelines. All animals were assessed as healthy prior to commencement of experiments. Animals were monitored prior to and following every injection ensuring there was no abnormalities in weight (>10%), appearance (fur) or behaviour (vocalisation, respiration, movements). All efforts were made to ameliorate any suffering of animals. They were handled by experienced researchers in such a way as to minimise stress prior to being anaesthetised.

Animals

These were supplied by the University of Melbourne Biological Research Facility and subjected to a 12 hour light/dark cycle with ad libitum access to food (dry pellets of a fixed formulation diet for laboratory rats and mice fortified with vitamins and minerals to meet the requirements of breeding animals after the diet is autoclaved or irradiated, supplied by Speciality Feeds, Western Australia) and water. Animals were housed in groups of 2-4 (adult) per cage (25cm × 35cm × 25cm on Breeders Choice paper bedding, made from 99% recycled paper; it is biodegradable with no added chemicals). Age groups investigated (at treatment completion) were embryonic day 19 (E19) pups of both sexes and dams, which were all primigravida 350-400g body weight) and non-pregnant female adults (175-230g body weight). E19 was chosen because this is a stage of development when adequate volumes of blood and cerebrospinal fluid (CSF) can be obtained for analysis from fetal rats without pooling (Dziegielewska et al., 1981) and individual pups can be injected intraperitoneally while still inside the uterine horn and kept viable for periods of time. Animal numbers were based on previous experience of such experiments and were the minimum number required to detect a significant difference between groups at p <0.05. Animals were selected for treatment groups to ensure weights were statistically similar between direct comparisons. Where possible, equal numbers of male and female fetuses were used. Animals on gestational day E19 were allocated to experiments by animal house staff, who had no knowledge of the particular experiments to be performed. The experimenters had no role in the selection of the animals, thus avoiding selection bias. The numbers (n) of animals used for each experiment are indicated in the relevant Methods or Results section and where appropriate in legends. Two litters in the sucrose permeability studies were excluded from the study. One mother died under anaesthesia. In the other case the fetuses were observed to be in poor physiological state, which would have affected the results.

Drugs and markers

Paracetamol (acetaminophen $\geq 99.0\%$, Sigma-Aldrich) was applied either at a high dose of 15 mg/kg (higher limit in the

range used clinically, Australian Medicines Handbook, 2019 and Koehn *et al.*, 2019b) or a dose in the lower clinical range of 3.75mg/kg. Paracetamol was dissolved in sterile 0.9% sodium chloride solution for injection. For passive permeability experiments [U- ¹⁴C]-labelled sucrose (Amersham International, CFB146) was injected in sterile 0.9% sodium chloride solution. Details are described in our previous study (Koehn *et al.*, 2019b). Estimates of protein (AFP) permeability were obtained from western blot analysis of fetal and maternal plasma, as described below.

Transcriptomic analysis: RNA-Seq

All experiments took place between 09.00 and 15.00h. Placentas and fetal brains from dams subjected to three treatment regimes were analysed in this study (n=4 for each tissue from each dam).

- (i) an E15 pregnant dam was given an intraperitoneal
 (i.p.) injection twice daily with 15mg/kg of paracetamol
 (dissolved in sterile 0.9% sodium chloride solution)
 over four days. On the 5th day (E19) the dam was given a final injection of the drug. This experiment is referred to as "chronic";
- (ii) an E19 dam was given a single i.p injection of 15mg/ml paracetamol and is referred to as "acute"; and
- (iii) an E19 untreated dam (referred as control).

In experiments (i) and (ii), 30 minutes after the last injection of the drug the tissue samples (placentas, fetal brains) were collected (n=4 for each dam).

For RNA-Seq analysis, placental tissue was sampled as a cross section of the chorio-allantoic placental disc, following removal of the externally attached umbilical and maternal circulatory vessels. Brain samples of the cortex were dissected out as described before (Koehn *et al.*, 2019a). Samples were collected under RNase free conditions and immediately frozen in liquid nitrogen and transferred to -80 °C for storage. RNA extraction was completed using the RNeasy Plus Mini Kits and QIAshredder (Qiagen, catalogue number 74134) for placenta and using the RNeasy Plus Micro Kits (Qiagen, catalogue number 74004) for fetal cortex, following manufacturers specifications. RNA quantity and purity were determined using a NanoDrop ND-1000 UV-VIS spectrophotometer (Thermo Scientific).

RNA samples were transported on dry ice to the Australian Genome Research Facility (AGRF) in Melbourne for Illumina, next-generation sequencing. Runs were 100bp single reads, providing raw FASTAq data. Data were processed using the Galaxy platform and their online software packages (Afgan *et al.*, 2018). Default parameters were used unless directly specified. Alignment was conducted using HISAT2 (Galaxy version 2.1.0) using the reference genome for rat (rn6; accession number GCA_000001895.4) and the reverse strand setting. For transcript quantification and differential expression analysis, three different methods were employed. In the first, pathway transcripts were assembled with cufflinks (Galaxy version 2.2.1.2) using the reference annotation for rat RefGene (genome) obtained from

UCSC Main. Relevant data were passed through Cuffmerge (Galaxy version 2.2.1.1) and analysed for differential expression between groups of interests using Cuffdiff (Galaxy version 2.2.1.5). For the second and third pathway, counts were aligned using HTSeq-counts (Galaxy version 0.9.1) using the reverse strand setting. Generate Count Matrix (Galaxy Australia version 1.0) produced a matrix form of the data, which were then fed through either DEseq2 (Galaxy version 2.11.40.6) or EdgeR (likelihood ratio; Galaxy version 3.24.1) to receive differential expression analysis between treatment groups. Statistically different expression levels between relevant treatment groups were selected if present in two of the three datasets above the statistical threshold of p < 0.05 for the adjusted P value of Cuffdiff (Padj), DEseq2 (q value) or EdgeR (FDR). This method of statistical selection minimizes the known false positives and false negatives that can be obtained due to the analysis pathway selected, ensuring all results can be found between multiple pipelines (see Seyednasrollah et al., 2016; Soneson & Delorenzi, 2013). Gene synonym names were produced via bioDBnet (Mudunuri et al., 2009). Pathway analysis was conducted using DAVID Bioinformatics Resources (version 6.8), with benjamini false discovery rate correction (Huang et al., 2009a; Huang et al., 2009b).

Interleukin 1 β (IL1 β) enzyme-linked immunosorbent assay (ELISA)

IL1B cytokine concentrations in rat plasma were determined using ELISA specific for rat IL1B (R&D systems, Quantikine kit, catalogue number RDSRLB00, monoclonal mouse anti-rat IL1B) following the manufacturer's protocol. Plasma samples were diluted 1:2 and 50µL of each sample was added to the same volume of assay diluent. Standard dilutions were assayed in duplicate. The plate was incubated at room temperature for two hours, then washed extensively. 100μ of rat IL1 β conjugate was added and incubated for a further two hours. After additional thorough washes, the plate was incubated for 30minutes in 100µL of substrate solution then developed with 100µl of stop solution. Plates were read using a FlexStation 3 Multimode Microplate Reader (wavelength 450nm, using 570nm to correct for any optical imperfections in the plate) within 30 minutes of the addition of the stop solution. Cytokine concentrations were determined by comparison with the standard curve produced from each run.

Permeability across the placenta

All permeability experiments were conducted on E19 dams and fetuses. Two chronic paracetamol treatment regimes were used. Time-mated E15 pregnant dams were injected i.p. twice daily with either a 15mg/kg (referred to as "high") or 3.75mg/kg (referred to as "low") dose of paracetamol (dissolved in sterile 0.9% sodium chloride solution) over four days ("chronic" experiments). On the 5th day, at E19, these were compared to age-matched animals that were not treated (controls). Numbers (n) of individual experiments are indicated below and are included in the legends of corresponding figures in the Results section. ¹⁴C-sucrose permeability. Animals were treated either with a "low" dose (3.75mg/kg) or "high" dose (15mg/kg) of paracetamol over four days starting at E15 following the same protocol as above. On the 5 th day the pregnant dams (E19) were anaesthetised i.p. with 25% w/v urethane, (Sigma, 1ml per 100g body weight) and placed supine on a 35°C heating plate and an endotracheal cannula inserted prior to sampling. Left femoral artery and vein were cannulated. All injections were by slow infusion to the femoral vein; the cannula was flushed with 2ml of heparinized (Hospira Inc, 5000 units per ml) saline. Maternal blood samples were taken from the femoral artery; blood volume was maintained by intraarterial injection of equivalent volumes of 1ml heparinized sodium chloride solution. Blood (right cardiac ventricle), CSF (cisterna magna) and brains (cortex) were sampled from each fetus. Sampling was concluded when the state of the placental circulation (normal condition: umbilical veins pink with oxygenated blood) was deemed insufficient, usually around 90 minutes (see Koehn et al., 2019b for details). CSF samples were examined microscopically for traces of red blood cells and discarded if contaminated (Habgood et al., 1992). Maternal blood was also collected at the end of the experiment. Blood samples were centrifuged (5000rpm, five minutes). Plasma supernatant was removed and stored at -20°C until used.

Two sets of permeability experiments were conducted:

(i) Fetal to maternal placental barrier permeability: pregnant animals treated with paracetamol as described above and control, untreated dams were terminally anaesthetized and an arterial cannula inserted into maternal circulation. Once the uterine horns were exposed, individual fetuses still within their amniotic sacs were injected serially with ¹⁴C-sucrose as described in Koehn *et al.* (2019b). Each fetus was taken at 30 minutes post injection. Maternal blood samples were collected at the same time as fetuses were consecutively removed for blood sampling. Maternal to fetal plasma levels ratios of ¹⁴C-sucrose were used as a measure of fetal to maternal placental transfer and calculated as follows:

Fotal to maternal placental transfer —	Maternal plasma at time $y(DPM / \mu l)$
reiai iomaiernai piaceniai iransjer =	Average fetal plasma up to time $y(DPM / \mu l)$
	y = maternal plasma sampling time

One control litter (n=6); one litter from a chronically treated dam with a low dose 3.75 mg/kg (n=5) and two litters from two chronically treated dams with a high dose (15 mg/kg, n=5 for each litter) were used.

(ii) Maternal to fetal placental barrier permeability: pregnant animals treated with paracetamol as described above and control untreated dams were terminally anaesthetized and ¹⁴C-sucrose was infused into the maternal circulation as detailed for paracetamol permeability studies above. Fetal samples were taken serially between 30 minutes and 90 minutes post injection. Blood

Maternal to fatal placental transfer -	fetal plasma at time $x(DPM / \mu l)$
maternario jetat placenta transfer =	Average maternal plasma up to time $x(DPM / \mu l)$
	x = fetal plasma sampling time

One control litter (n=8) and one litter from a chronically treated dam with high dose (15mg/kg, n=6) were used.

Permeability of a fetal specific protein, AFP- western blotting. Levels of AFP in both the maternal blood samples and in fetal samples obtained from experiments of paracetamol treated dams as described above, were estimated using western blotting and antibodies to human AFP (DAKO).

All plasma samples were diluted 10-fold in isotonic saline (0.9%) prior to sample preparation. Samples were run using a total of 9µL of dam and 2µL of diluted fetal sera, denatured in 4x sample buffer (62.2 mM Tris, 5% (v/v) glycerol, 2% (w/v) SDS, and 0.0025% (w/v) bromophenol blue), heated to 95°C for five minutes and centrifuged briefly to remove potential particular matter. 12µL of each sample was loaded onto a 4-12% NuPAGE Novex Bis-Tris Midi gel (Life Technologies) and proteins were resolved at 200V for 40 minutes immersed in MES SDS running buffer (Life Technologies). Gel-resolved proteins were transferred onto PVDF membranes using iBlot gel transfer stacks (iBlot 2; Life Technologies) as per manufacturer's instructions. Membranes were incubated for one hour at room temperature in PBS-T blocking buffer (PBS supplemented with 0.05% (v/v) Tween-20 [Chemsupply]) and 5% (w/v) skim milk powder. Membrane was incubated with AFP primary antibody (AFP, rabbit polyclonal, 1:1000, DAKO, catalogue number A0008, RRID AB_2650473) diluted in the blocking buffer and incubated overnight at 4°C. After three PBS-T washes, the membrane was incubated for two hours at room temperature in horseradish peroxidase-conjugated goat anti-rabbit IgG (Cell Signaling; 1:5000, catalogue number 7074) secondary antibody. Immunoreactive protein bands were visualised by adding 1mL of Enhanced Chemiluminescence mixture (ECL Advance, GE Healthcare) onto membranes and detecting luminescence using a FujiFilm LAS-3000 imager at three and 75 second exposures. Densitometric quantitation of immunoreactivity was performed using ImageJ 2-bit, v1.46 run on OSx 10.14 Mojave on 8-bit TIFF file images. All samples that were directly compared were run on the same gel. Serum from an age-matched non-pregnant female was used as a negative control, while an E19 pregnant dam that was not injected with paracetamol was used as a positive control. Both samples were included on every gel.

Permeability of the fetal blood-brain barrier

Blood-brain barrier permeability in the fetus was estimated using (i) radioactive sucrose as an example of a small molecular marker permeability and (ii) plasma protein (immunohistochemistry), as an example of a large molecular marker permeability (Habgood *et al.*, 1993; Johansson *et al.*, 2008; Stolp *et al.*, 2005b). Fetal blood, CSF and brain samples were obtained from the same placental permeability experiments described above. ¹⁴*C*-sucrose permeability. For estimation of transfer from mother to fetus, pregnant animals treated with paracetamol (as above) were anaesthetized i.p. with urethane. Starting at 30 minutes after the last maternal injection, embryos were individually extracted. For estimation of transfer from fetal blood to fetal brain and CSF, the fetuses were exposed and injected i.p. with ¹⁴C-sucrose.

In both types of experiment fetal blood and CSF were sampled as described previously (Koehn *et al.*, 2019b). Fetal brain samples were taken by opening the cerebral hemispheres to expose the lateral ventricles and a sample of the parietal cortex was removed, taking care to avoid the choroid plexuses. Brain or CSF to plasma ratios of ¹⁴C-sucrose radioactive counts were used as an estimate of the transfer of sucrose across the blood brain barriers. These were calculated using the equation:

 $Brain or CSF transfer = \frac{Brain or CSF DPM / \mu l}{Plasma DPM / \mu l} \times 100\%$

Treatment groups investigated were control, no paracetamol (n=13), chronic low dose (3.75mg/kg, n=11) and chronic high dose (15mg/kg, n=11) in fetuses that were injected directly. In experiments in which the ¹⁴C-sucrose was injected into the treated mothers, numbers of pups used were control (n=8), acute (n=10) and chronic high dose (n=6).

Immunohistochemistry. Individual fetal brains were fixed in Bouin's fixative for 24-48h then dehydrated through graded alcohols, cleared in chloroform and embedded into paraffin wax blocks. These blocks were cut serially into coronal 5µm sections (Leica microtome). Selected sections were heated for 30 minutes (60°C) then washed twice with Histolene (Fronine) for 10 minutes, then five minutes. The sections were rehydrated through graded alcohols for five minutes each (100%, 100%, 95%, 70%) and washed in phosphate buffered saline with 0.2% Tween20 for five minutes. Peroxidase and protein blockers (DAKO) were added to sections and incubated at room temperature for two hours each to block non-specific binding. The primary antibody, plasma protein (anti-rat whole serum, SIGMA, catalogue number R5129, rabbit polyclonal) diluted 1:3000 in a blocker (0.5% fish gelatine and PBS + Tween20), was applied to the slides and incubated overnight at 4°C. After three washes of PBS + Tween20 for five minutes each, the secondary (swine anti-rabbit, DAKO, catalogue number Z0196, polyclonal) and tertiary antibodies (rabbit PAP, SIGMA, catalogue number P1291) both diluted 1:200 were each added and incubated for two hours at room temperature with washes between incubations. Sections were developed with DAB (Diaminobenzidine) using DAKO DAB+ kit (catalogue number K3468) according to manufacturer's directions and washed in running water for five minutes. Sections were dehydrated through a series of graded alcohols (70%, 95% for five minutes, then 100% for 10 minutes), then 3x five minutes in histolene washes. Slides were then mounted using DPX mounting medium (Fronine). Stained sections were examined under a compound microscope (Olympus, BX50) fitted with a digital camera (Olympus DP70). One control slide was included with every round of immunostaining and had the primary antibody omitted but was otherwise treated in the same way. These were always

blank. A total of 11 brains with at least two brains per treatment group were prepared and serially sectioned and mounted on glass slides. Each slide contained 6–8 sections, every 10th slide was stained with haematoxylin and eosin for general morphology. One or two adjacent slides per brain were immunostained for plasma protein from comparable brain regions.

Liquid scintillation counting

Plasma (10 μ L), CSF and every injectate (1 μ L of 1:10 dilution) were weighed and transferred into scintillation vials. In all experiments the radioactivity in the injectate was measured to confirm the uniformity of the injected material. Soluene350 (0.5ml, PerkinElmer) was added to the brain samples and incubated overnight at 36°C. Prior to measurement, two drops of glacial acetic acid (Sigma) were added to brain vials to neutralize the strongly alkaline Soluene350. All samples were mixed with 5ml of scintillation fluid (Emulsifier-safe, PerkinElmer) and measured on the liquid scintillation counter (Tri-Carb 4910 TR, PerkinElmer). Counting was conducted in disintegrations per minute (DPM) for five minutes each with luminescence correction on. Vials containing control, non-radioactive tissues processed identically were also counted simultaneously to establish background counts (which were subtracted from all radioactive samples). Counts were normalized to the sample weight and expressed as DPM per μ L or μ g of sample. Results are described as concentration ratios, defined as a % of the counts (per μ L or μ g) in the compartment of interest (brain, CSF, maternal or fetal plasma) divided by the counts (per μ L) in the plasmacompartment of comparison (see also Koehn *et al.*, 2019b).

Statistics

RNA-Seq data analysis is detailed above, with significance set at p <0.05. For all other experimentation, statistical differences between treatment groups were determined by unpaired Student t-tests using Prism 6.2 (Graphpad Software Inc) with significance set at p <0.05. We also tested our data using ANOVA followed by Tukey's posthoc test; this approach yielded the same outcomes.

Results

E19 placentas and brains from three treatment groups were compared for transcriptomic analysis using RNAseq datasets: (i) untreated controls (n=4), (ii) acutely paracetamol treated (n=4) and (iii) chronically paracetamol treated (n=4) dams (see Methods), providing a three-way comparison for each tissue (Figure 1 and Table 1–Table 5).



Figure 1. Number of up-regulated and down-regulated genes in the E19 placenta and brain following chronic maternal treatment with paracetamol. Transcript numbers for Chronic/control, Acute/control and Chronic/acute comparisons. Controls were from untreated animals. For details of chronic and acute dosage schedules see Methods. Data derived from RNA-Seq analysis. Overlapping segments represent shared genes.

:h paracetamol.	
following treatment wit	
es in the E19 placenta	
ıd down-regulated gen	
lop 50 up-regulated ar	
Table 1.	

	ontrol)	FC	-2332	-1698	-1415	006-	-639	-474	-452	-356	-278	-249	-236	-166	-162	-77	-57	-43	-28
	ed (chronic/c	Gene	Afp	Apoc2	Rbp4	dodA	Fgb	Apoa2	Apoa1	Ttr	Fgg	Spp2	Serpina1	Clca4	Apom	F2	τţ	Mttp	Cubn
	Down-regulat	QI	NM_012493	NM_001085352	NM_013162	NM_019287	NM_020071	NM_013112	NM_012738	NM_012681	NM_012559	NM_053577	NM_022519	NM_201419	NM_019373	NM_022924	NM_001013110	NM_001107727	NM_053332
	trol)	FC	42	23	13	7.75	5.62	3.89	3.20	3.12	2.99	2.87	2.86	2.77	2.77	2.77	2.71	2.51	2.50
	chronic/cont	Gene	Cc/3	Cxcl2	<i>q1</i> 11	Il1rn	Grem 1	Lcn2	Fcer1g	LOC684107	Plek	Fgr	Itgam	Ifit3	Cd53	Thbs4	Cd14	Tnfrsf1b	Kcnc4
enta	Up-regulated (Ð	NM_013025	NM_053647	NM_031512	NM_022194	NM_019282	NM_130741	NM_001131001	NM_001317798	NM_001025750	NM_024145	NM_012711	NM_001007694	NM_012523	NM_017133	NM_021744	NM_130426	NM_001122776
E19 Plac	'ol)	FC	-54	-10		- 100	-16	-16	-16	-15	-14	-12	-12	-10	-8.80	-8.78	-7.72	-7.72	-7.69
-	d (acute/conti	Gene	Clca4	Afp	Fgg	Apom	Apoc2	Spp2	Apoa1	F2	τf	Mttp	Serpina1	Maob	Myh14	Agt	Apoa4	Gucy2c	Cubn
	Down-regulate	Ð	NM_201419	NM_012493	NM_012559	NM_019373	NM_001085352	NM_053577	NM_012738	NM_022924	NM_001013110	NM_001107727	NM_022519	NM_013198	NM_001100690	NM_134432	NM_012737	NM_013170	NM_053332
		FC	4.68	3.98	3.84	3.45	3.37	2.95	2.84	2.66	2.57	2.44	2.40	2.36	2.35	2.31	2.28	2.27	2.27
																0			
	d (acute/control)	Gene	Aoc3	Chodl	Fcr16	100290595	LOC684107	Ccdc77	Dox/1	Cbx2	Ctsm	Mboat1	qddN	Pr18a2	Adipoq	LOC10255731	Prl8a4	LOC683313	Psg16
	Up-regulated (acute/control)	ID Gene	NM_031582 Aoc3	NM_001105894 Chodl	NM_001164726 Fcrl6	NM_001106063 LOC290595	NM_001317798 LOC684107	NM_001191752 Ccdc77	NM_199233 Dox/1	NM_001107071 Cbx2	NM_181378 Ctsm	NM_001109120 Mboat1	NM_031545 Nppb	NM_022846 Pr/8a2	NM_14744 Adipoq	NM_001329892 LOC102557319	NM_021580 Pr/8a4	NM_001101007 LOC683313	NM_001025679 Psg16
	Up-regulated (acute/control)	ID Gene	1 NM_031582 Aoc3	2 NM_001105894 Chodl	3 NM_001164726 Fcr/6	4 NM_001106063 LOC290595	5 NM_001317798 LOC684107	6 NM_001191752 Ccdc77	7 NM_199233 Dox/1	8 NM_001107071 Cbx2	9 NM_181378 Ctsm	10 NM_001109120 <i>Mboat1</i>	11 NM_031545 Nppb	12 NM_022846 Pr/8a2	13 NM_144744 Adipog	14 NM_001329892 LOC10255731	15 NM_021580 Pr/8a4	16 NM_001101007 LOC683313	17 NM_001025679 Psg16

	ontrol)	FC	-17	-17	-16	-15	-14	-6.89	-6.36	-6.06	-5.39	-4.30	-4.18	-3.77	-3.50	-3.40	-3.35	-3.29	-3.12
	ted (chronic/co	Gene	Apoa4	Gucy2c	Maob	Myh14	Agt	Amn	Orai2	Ctsc	Palm	Lfng	Creb313	Cldn2	Uap111	Aqp8	RGD1566401	Phgdh	Gipc2
	Down-regulat	ID	NM_012737	NM_013170	NM_013198	NM_001100690	NM_134432	NM_001108061	NM_001170403	NM_017097	NM_130829	NM_133393	NM_001012115	NM_001106846	NM_001134516	NM_019158	NR_131064	NM_031620	NM_001037210
	trol)	FC	2.36	2.33	2.33	2.27	2.26	2.25	2.21	2.17	2.17	2.17	2.17	2.16	2.14	2.09	2.07	2.07	2.03
	chronic/cont	Gene	Nppb	Nfkbia	Lamc3	Cenpf	S100a8	Ramp3	Cxcl1	Sv2a	Ccdc77	Serpinb1a	Scg5	S100a9	Clcn4	Tnfsf13	Rac2	Tyrobp	Cybrd1
enta	Up-regulated (Ð	NM_031545	NM_001105720	NM_001107830	NM_001100827	NM_053822	NM_020100	NM_030845	NM_057210	NM_001191752	NM_001031642	NM_013175	NM_053587	NM_022198	NM_001009623	NM_001008384	NM_212525	NM_001011954
E19 Plac	rol)	Ę	-6.18	-5.24	-4.59	-3.52	-3.52	-3.34	-3.29	-3.26	-3.11	-2.99	-2.89	-2.82	-2.82	-2.71	-2.71	-2.59	-2.56
-	ed (acute/cont	Gene	Ctsc	Aqp8	Amn	Cldn2	Uap 111	Vil1	Atp7b	Lfng	Lfng	Prr7	Tgfbi	LOC310926	PIIP	PIs1	Marcks11	Scand1	Scd
	Down-regulate	Ð	NM_017097	NM_019158	NM_001108061	NM_001106846	NM_001134516	NM_001108224	NM_012511	NM_133393	NM_031620	NM_001109116	NM_053802	NM_001025002	NM_022533	NM_001108178	NM_030862	NM_001108599	NM_139192
	-	ñ	2.25	2.22	2.21	2.17	2.09	2.07	2.03	2.03	1.98	1.97	1.96	1.95	1.95	1.94	1.94	1.91	1.91
	d (acute/control	Gene	Prl6a1	Frem2	Slco2a1	Clcn4	Psg29	Tnfsf13	Pri4a1	Kcnc4	Ncam1	Ppp1r3c	Adcy3	Zfp9	FInc	Hsd11b1	111 <i>Tf</i>	Ednrb	Cenpf
	Up-regulate	ID	NM_022176	NM_001245978	NM_022667	NM_022198	NM_001025641	NM_001009623	NM_017036	NM_001122776	NM_031521	NM_001012072	NM_130779	NM_001127635	NM_001191862	NM_017080	NM_001015011	NM_017333	NM_001100827
			18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34

						E19 Pla	icenta					
	Up-regulated	d (acute/contro	()	Down-regulate	ed (acute/cont	rol)	Up-regulated (chronic/cont	trol)	Down-regulat	ed (chronic/co	ontrol)
	DI	Gene	FC	ID	Gene	FC	Ð	Gene	FC	DI	Gene	FC
	NM_001191918	C1 qtnf2	1.89	NM_024160	Cyba	-2.49	NM_017080	Hsd11b1	1.97	NM_001014193	Rsrp1	-3.07
	NM_031022	Cspg4	1.88	NR_131064	RGD1566401	-2.48	NM_022954	Fat2	1.97	NM_001025002	LOC310926	-3.03
	NM_017198	Pak1	1.87	NM_012862	Mgp	-2.47	NM_017051	Sod2	1.96	NM_139192	Scd	-3.03
	NM_172073	Τρbpa	1.85	NM_001109627	Epop	-2.47	NM_001009681	Oas/	1.95	NM_022533	PIIp	-2.74
	NM_001191915	Gpr50	1.85	NM_001014193	Rsrp1	-2.46	NM_130779	Adcy3	1.92	NM_139087	Cgref1	-2.64
	NM_001034010	Tril	1.79	NM_17927	Serpinf1	-2.45	NM_001033691	Irf7	1.92	NM_001108178	PIs1	-2.63
	NM_134385	Pr18a9	1.79	NM_053565	Socs3	-2.45	NM_001107887	Cd163	1.92	NM_001047891	RGD1310507	-2.63
	NM_053360	Sh3kbp1	1.78	NM_001108971	Myorg	-2.38	NM_133624	Gbp2	1.91	NM_177927	Serpinf1	-2.61
	NM_030994	Itga 1	1.78	NM_001170584	Pex5	-2.31	NM_012924	Cd44	1.89	NM_153736	Prl3a1	-2.56
	NM_001109141	Kctd15	1.76	NM_139087	Cgref1	-2.31	NM_001007691	Prss23	1.87	NM_001008890	Hbe1	-2.51
	NM_001135877	Taf71	1.76	NM_001037210	Gipc2	-2.22	NM_031764	Ddr2	1.85	NM_030827	Lrp2	-2.47
	NM_001106515	Fermt1	1.75	NM_001191974	Chst13	-2.20	NM_001134858	Synm	1.85	NM_172030	Entpd2	-2.29
	NM_001100984	Ncf2	1.75	NM_001037659	Mpp 1	-2.17	NM_001106420	Adamts12	1.83	NM_001109627	Epop	-2.27
	NM_001034932	C1 qtnf6	1.72	NM_001108552	Trim2	-2.16	NM_031022	Cspg4	1.83	NM_001014790	Rarres1	-2.26
	NM_001271283	Golm2	1.72	NM_001012470	Irf2bpl	-2.14	NM_207606	Kirrel1	1.83	NM_019231	Mapk13	-2.24
	NM_001106402	Lhfp12	1.71	NM_001107052	Arl4d	-2.11	NM_019341	Rgs5	1.82	NM_001014088	Eepd1	-2.22
もそう	hange of transcript nu change compared to c c treated animals. Noti	mbers in placentas control (p<0.05, see e that only 10/50 g	treated Method enes wer	with paracetamol (chrc ls). Colours indicate ger re upregulated followin	onic, acute or cont nes that were upr ot both treatment	trol, n=4 i. egulated s but 34/5	n each group). For det (green) in both acute . 50 were downregulate	tails of dosage s and chronic tre- ed following bot	schedule ated anii ch treatm	s see <i>Methods</i> . Data fr mals and downregulationts.	om RNA-Seq anal ed (red) in both ac	/sis. FC cute and

Page 10 of 37

		E19 Pla	centa: i	mmune/	inflammation-related	d genes			
Up-reg	julated (chro	nic/cont	rol)		Up-re	egulated (acute	control)	
ID	Gene	Ch / Co	Ac / Co	Ch / Ac	ID	Gene	Ch / Co	Ac / Co	Ch / Ac
NM_013025	Ccl3	42	-	34	NM_013095	Smad3	-	1.61	-
NM_053647	Cxcl2	23	-	17	NM_001025721	Colec12	-	1.59	-
NM_031512	Il1b	13	-	17	NM_001008328	Parp3	-	1.50	-
NM_022194	Il1rn	7.75	-	6.19	NM_017113	Grn	-	1.48	-
NM_130741	Lcn2	3.89	-	3.91	NM_133380	Il4r	-	1.46	0.78
NM_001131001	Fcer1g	3.20	-	3.60	NM_031514	Jak2	-	1.35	-
NM_024145	Fgr	2.87	-	2.17	NM_173328	Lgr4	-	1.32	-
NM_012711	Itgam	2.86	-	2.92	NM_001107754	Traf6	-	1.31	-
NM_021744	Cd14	2.71	-	-	NM_001107063	Cdc42ep4	-	1.28	-
NM_130426	Tnfrsf1b	2.51	-	1.89	NM_001191552	Nsd2	-	1.24	-
NM_053822	S100a8	2.26	-	4.72	NM_001017385	Kdelr1	-	1.22	-
NM_030845	Cxcl1	2.21	-	-	NM_012925	Cd59	-	1.21	-
NM_001031642	Serpinb1a	2.17	-	1.68	NM_001106715	Pum2	-	1.19	-
NM_001008384	Rac2	2.07	-	2.83					
NM_001009681	Oasl	1.95	-	-					
NM_001009689	Cdc42ep2	1.79	-	-					
NM_001134555	C1r	1.71	-	-					
NM_012673	Thy1	1.66	-	-					
NM_001136124	Ifitm3	1.56	-	-					
NM_001013062	Thbs1	1.55	-	-					
NM_001106314	Ifitm1	1.52	-	-					
NM_053535	Enpp1	1.47	-	-					
NM_138881	Rsad2	1.45	-	-					
NM_138844	Unc13d	1.44	-	-					
NM_013016	Sirpa	1.43	-	-					
NM_001166403	RT1-T24-3	1.43	-	-					
NM_001191760	Dock11	1.37	-	1.24					
NM_001271227	Tfe3	1.36	-	-					
NM_001100565	Peli1	1.31	-	-					
NM_001108101	Irak3	1.30	-	1.33					
NM_023092	Myo1c	1.30	-	-					
NM_031140	Vim	1.30	-	-					

Table 2. Changes in gene regulation in E19 placenta following maternal treatment with paracetamol.

	E1	9 Placen	ta: imr	nune/in	flammation-related g	enes			
Up-regu	ulated (chronic	c/contro	I)		Up-reg	gulated (acute	/control)	
ID	Gene	Ch / Co	Ac / Co	Ch / Ac	ID	Gene	Ch / Co	Ac / Co	Ch / Ac
NM_017256	Tgfbr3	1.26	-	-					
NM_133624	Gbp2	1.91	-	1.86					
NM_021655	Chga	1.70	-	1.71					
NM_013069	Cd74	1.61	-	1.79					

Down-reg	gulated (chron	ic/contr	ol)		Down-r	egulated (acut	e/contr	ol)	
ID	Gene	Ch / Co	Ac / Co	Ch / Ac	ID	Gene	Ch / Co	Ac / Co	Ch / Ac
NM_020071	Fgb	-639	-	-	NM_001109535	Rab20	-	-1.93	1.89
NM_013112	Apoa2	-474	-	-	NM_001108207	Tnfrsf21	-	-1.75	-
NM_001007729	Cxcl4	-1.55	-	-	NM_053587	S100a9	2.16	-1.71	3.70
NM_001106095	Lig4	-1.53	-	-	NM_173838	Fzd5	-	-1.53	-
NM_001008322	Shmt2	-1.46	-	-	NM_001025707	Tfeb	-	-1.40	-
NM_030826	Gpx1	-1.39	-	-	NM_012939	Ctsh	-	-1.40	-
NM_001108741	Appl2	-1.26	-	-	NM_001025672	Pspc1	-	-1.34	-
NM_001127390	Washc1	-1.25	-	-	NM_013157	Ass1	-	-1.34	-
NM_001108069	Stk11	-1.23	-	-	NM_001031653	Polr3d	-	-1.23	-
NM_001134974	Trim27	-1.20	-	-	NM_001024967	Tmem106a	-	-1.23	-
					NM_019357	Ezr	-	-1.22	-
					NM_053743	Cdc37	-	-1.18	-

Up-regulated and down-regulated inflammatory and immune-related gene changes in E19 placenta following no treatment (co, controls), acute (ac,) or chronic (ch) maternal paracetamol treatment; see *Methods* for details of dosage schedule. Data from RNA-Seq analysis. Numbers are fold changes for comparisons indicated (Ch/Co, Ac/Co, Ch/Ac). The chronic/acute comparison indicates significant differences in regulation between the two dosage regimes (P<0.05, see Methods). In all cases expression was greater with chronic treatment. This table includes only genes with inflammatory and immune-related functions and thus includes some of the highly expressed genes in Table 1.

The effect of paracetamol exposure on placental gene expression (transcriptomic analysis)

As illustrated in Figure 1, following maternal exposure to paracetamol (either acute or chronic), there was a large number of genes that significantly altered their expression in the E19 placentas in two-way comparisons to control tissue, with much fewer that changed between the two treatment groups (chronic/ acute). Most genes were uniquely regulated, either up or down, depending on treatment duration, with relatively few that were common to both treatment regimes (64 up-regulated and 57 down-regulated). In contrast, in a three-way comparison, only one gene, *Nfkbia* (NF-kappa-B inhibitor alpha), was shared in all three comparisons (Figure 1). NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines.

The expression of 121 transcripts (the sum of up-regulated and down-regulated genes in the chronic/acute comparison) was significantly different between acute and chronic treatment groups, suggesting an effect of treatment duration. Of these genes, 34 were significantly up-regulated in chronically treated animals when compared to either the acute treatment group or the control group and eight were down-regulated (Figure 1).

Comparing datasets of placentas from chronically treated dams with untreated control dams, the expression of 737 genes was significantly different (either up or down p<0.05, see Methods) (Figure 2). The top 50 up-regulated and down-regulated genes in E19 placentas are displayed for both acute and chronic treatment groups compared to controls in Table 1. In the E19 placentas, many of the top genes up-regulated following chronic treatment were related to immune-response and inflammation (Table 1, Table 2 and Table 5). It is difficult to determine the extent to which a statistically difference in gene expression is also functionally significant. It is perhaps worth noting that fewer genes were up-regulated two-fold or more with either acute or chronic paracetamol treatment (25 and 34 genes, respectively) compared to the number that were down-regulated two-fold or more (58 and 61, respectively). In addition, the degree of down-regulation was appreciably greater for many of these genes compared with those that were up-regulated. This was particularly evident for the chronically treated group

Table 3. Top 50 up-regulated and down-regulated genes in the E19 fetal brain following paracetamol treatment.

a) hotelinen musik	() Province in the second of t			-) 1004		E19 B	Brain	uter of others bound	-	Donna kined	stands and stands and stands	1
-d N	regulated	l (chronic/contr	o)	Down-regula	ated (chronic/contr	(lo	Up-regulat	ed (acute/contr	(lo	Down-regul	ited (acute/contr	ol)
ID		Gene	FC	ID	Gene	FC	ID	Gene	FC	DI	Gene	FC
NM_00125	8011	Snhg11	3.41	NM_053304	Col1a1	-10	NM_001258011	Snhg11	4.13	NM_053304	Col1a1	-16
NM_01295	6	Gfra1	3.38	NM_001003978	Gspt1	-9.21	NM_001108951	AK5	3.57	NM_032085	Col3a1	-6.99
NM_00117	0531	Rasgrf1	2.76	NM_001365151	Crmp1	-7.56	NM_012959	Gfra1	3.43	NM_001365151	Crmp1	-4.70
NM_01292	50	Camk2a	2.57	NM_032085	Col3a1	-6.98	NM_012920	Camk2a	3.28	NM_001008890	Hbe1	-3.37
NR_03761	4	Vof16	2.47	NM_001013853	Hba-a3	-5.10	NM_001109003	Cpne4	3.09	NM_033234	Чbb	-3.26
NM_0011	05878	Polq	2.30	NM_057212	Tmem158	-4.86	NM_012671	Tgfa	3.03	NM_001127523	Mis18a	-3.04
NM_0011	05880	Zbtb20	2.28	NM_130829	Palm	-4.70	NM_012519	Camk2d	2.94	NM_001109627	Epop	-3.02
NR_1310(54	RGD1566401	2.27	NM_001025041	Ier5l	-3.95	NM_001170531	Rasgrf1	2.80	NM_001025041	Ier5	-3.01
NM_0011	91970	Thsd7a	2.25	NM_001191905	Scrt2	-3.83	NM_001127492	Sphkap	2.73	NM_001108464	Prr18	-2.90
NM_0011	91752	Ccdc77	2.20	NM_001108599	Scand1	-3.72	NM_001191970	Thsd7a	2.70	NM_001111269	Hbb-bs	-2.81
NM_0011	07310	Spock3	2.15	NM_001108464	Prr18	-3.71	NR_131064	RGD1566401	2.60	NM_001191905	Scrt2	-2.74
NM_0011	72305	Prkcb	2.12	NM_001109627	Epop	-3.24	NM_030993	Ddn	2.59	NM_057212	Tmem158	-2.73
NR_1265	81	Lnc215	2.10	NM_001109116	Prr7	-3.24	NM_017318	Ptk2b	2.40	NM_001108599	Scand1	-2.57
NM_0218	53	Kcnt1	2.06	NM_001106014	RGD1560394	-3.10	NM_001107671	Plcxd3	2.40	NM_001113223	LOC100134871	-2.57
NM_0012	71191	Sgo2	2.00	NM_033234	Чрр	-3.06	NM_001108335	Rasal1	2.36	NM_001106299	Ahsp	-2.57
NM_0535	05	Slc24a3	1.98	NM_001111269	Hbb-bs	-2.89	NM_001105880	Zbtb20	2.34	NM_013197	Alas	-2.57
NM_2068	347	Pfkp	1.98	NM_030862	Marcksl1	-2.62	NM_001271404	Hs6st3	2.34	NR_027324	H19	-2.55

						E19	Brain					
Up-regulated (chronic	(chronic	c/contr	(lo	Down-regula	ated (chronic/cont	rol)	Up-regulat	ed (acute/contro	ol)	Down-regulat	ted (acute/contr	(lo
ID Gei	Gel	ne	Ę	DI	Gene	FC	IJ	Gene	FC	QI	Gene	FC
NM_017007 >G	Š A	ad1	1.98	NM_001107307	Cilp2	-2.47	NM_013192	Kcnj6	2.32	NM_030862	Marcksl1	-2.51
NM_013192 Kc	Kc	nj6	1.97	NM_001109270	Fam110d	-2.46	NM_001001508	Plppr4	2.28	NM_012651	Slc4a1	-2.38
NM_053336 Aq	Ă	ger	1.97	NM_001008890	Hbe1	-2.46	NM_021853	Kcnt1	2.26	NM_001109116	Prr7	-2.12
NR_111959 N	2	liat	1.96	NR_027324	H19	-2.37	NM_134363	Slc12a5	2.23	NM_001115013	Selenom	-2.02
NM_031003	A	bat	1.94	NM_001106066	Pgls	-2.36	NM_001289778	Map7d2	2.20	NM_181380	Rtn4rl2	-2.02
NM_001136241 N	z	gef	1.91	NM_001107745	C1qtnf4	-2.33	NM_012713	Prkcb	2.19	NM_001168650	Sox12	-1.99
NM_012713	ā	rkcb	1.89	NM_001113223	LOC100134871	-2.29	NM_001191752	Ccdc77	2.17	NM_001106014	RGD1560394	-1.99
NR_130129 T	F	incr	1.85	NM_001168650	Sox12	-2.25	NM_001134837	Trps1	2.14	NM_019291	Car2	-1.97
NM_053817	z	rxn3	1.83	NM_013197	Alas2	-2.23	NM_030871	Pde1a	2.13	NM_001037659	Mpp1	-1.95
NM_001202552 Eif-	Eif	4ebp3	1.82	NM_001037659	Mpp1	-2.21	NM_053505	Slc24a3	2.10	NM_022300	Basp1	-1.93
NM_001106895	Τŧ	ap2d	1.82	NM_001164043	Mapkapk5	-2.20	NM_001136241	Ngef	2.09	NM_001107951	Dmrta2	-1.9(
NM_001014207	F	af1d	1.80	NM_022300	Basp1	-2.17	NM_017007	Gad1	2.08	NM_001106066	Pgls	-1.9(
NM_024371 S	S	lc6a1	1.79	NM_001109621	Tpgs1	-2.14	NM_206847	Pfkp	2.07	NM_031114	S100a10	-1.86
NM_031783		Nefl	1.78	NM_001107603	Fbxl15	-2.11	NM_001135779	Gabra2	2.04	NM_001107307	Cilp2	-1.86
NM_153630	2	Valcn	1.75	NM_181380	Rtn4rl2	-2.07	NM_031003	Abat	2.04	NM_001100741	Col6a2	-1.8
NM_001271366	2	Aki67	1.74	NM_019386	Tgm2	-2.07	NM_001172305	Prkcb	2.03	NM_017271	Nudc	-
NM_017275		nck	1.74	NM_017271	Nudc	-2.03	NM_001108542	Arhgef28	2.03	NM_001109621	Tpgs1	-1.77

						E19 Bra	Ē					
	Up-regulated	(chronic/cont	'ol)	Down-regula	ated (chronic/contr	(lo	Up-regulated	l (acute/cont	rol)	Down-regulate	d (acute/cont	rol)
	ID	Gene	FC	Ð	Gene	FC	QI	Gene	FC	Ð	Gene	ñ
35	NM_001107659	Sema5a	1.72	NM_001071776	Bola1	-2.01	NM_001025664	Wsb1	1.98	NM_001169138	Thbs2	-1.76
36	NM_022631	Wnt5a	1.72	NM_001130570	Scrt1	-2.00	NM_001106895	Tfap2d	1.98	NM_001164043	Mapkapk5	-1.76
37	NM_001107638	Scml4	1.72	NM_001013163	Bmyc	-2.00	NM_173121	Brinp3	1.98	NM_001109270	Fam110d	-1.76
38	NM_031046	Itpr2	1.71	NM_001107951	Dmrta2	-2.00	NM_001107310	Spock3	1.98	NM_001107745	C1qtnf4	-1.75
39	NM_001029911	Cit	1.71	NM_031677	FhI2	-1.95	NM_057130	Hrk	1.98	NM_001195482	Ndufaf8	-1.73
40	NM_031730	Kcnd2	1.70	NM_021678	Camk2n2	-1.95	NM_001107473	Zim1	1.97	NM_001130570	Scrt1	-1.72
41	NM_001012235	Impact	1.70	NM_019291	Car2	-1.93	NM_017275	Pnck	1.96	NM_019354	Ucp2	-1.72
42	NM_057130	Hrk	1.69	NM_001134570	Nupr2	-1.93	NM_001107881	Plxnd1	1.95	NM_031677	FhI2	-1.71
43	NM_001134553	Ubn2	1.67	NM_001195482	Ndufaf8	-1.93	NM_17481	Slco3a1	1.95	NM_001025137	Ier5	-1.70
44	NM_133302	Adarb2	1.66	NM_019326	Neurod2	-1.87	NM_153630	Nalcn	1.93	NM_001108955	Fjx1	-1.70
45	NM_001107881	Plxnd1	1.66	NM_001106299	Ahsp	-1.87	NM_001271191	Sgo2	1.93	NM_019326	Neurod2	-1.70
46	NM_052807	Igf1r	1.65	NM_001107574	Znhit2	-1.87	NM_001107659	Sema5a	1.92	NM_001107603	Fbxl15	-1.69
47	NM_001037139	Pcdhga2	1.64	NM_012651	Slc4a1	-1.86	NM_001106498	Pak6	1.91	NM_001107708	Olfm13	-1.67
48	NM_001107201	Prox1	1.64	NM_001127523	Mis18a	-1.85	NM_001114656	7 mjd7	1.88	NM_001134570	Nupr2	-1.67
49	NM_001107281	KIf12	1.64	NM_012932	Crmp1	-1.84	NM_001108242	Slc9a7	1.87	NM_031838	Rps2	-1.66
50	NM_001107425	Nfat5	1.63	NM_001142652	Neurl1b	-1.82	NM_053817	Nrxn3	1.87	NM_019905	Anxa2	-1.66
^c old ch	nange of transcript nun	ubers followina cł	nronic, ac	ute or control maternal	treatment with paracet	camol, n=4	in each aroup. For det	ails of dosage so	hedules se	e <i>Methods</i> . Data from R	NA-Sea analvsis.	=C = fold

concrements or conscript numbers romoving encount netering treatment with paracecanior, in-4 in each group. For actain or boade extrements see *methods*. For a method in port acute and chronic treated animals and downregulated (red) in both acute and chronic treated animals. Note that 26/50 genes were upregulated following both treatments and 40/50 were downregulated following both treatments.

E19 Brain: immune/inflammation-related genes											
Up-re	Up-regulated (acute/control)										
ID	Gene	Ch / Co	Ac / Co	Ch / Ac	ID	Gene	Ch / Co	Ac / Co	Ch / Ac		
NM_053336	Ager	1.97	-	-	NM_017318	Ptk2b	-	2.40	-		
NM_001202552	Eif4ebp3	1.82	-	-	NM_001004444	Zbtb1	-	1.52	-		
NM_022631	Wnt5a	1.72	-	-	NM_001108614	Lime1	-	1.51	-		
NM_001105734	Dusp10	1.54	-	-	NM_001107770	Sppl2a	-	1.44	-		
NM_017187	Hmgb2	1.47		-	NM_001105733	Cacnb4	-	1.31	-		
					NM_001107678	Dhx36	-	1.16	-		
Down-regulated (chronic/control)					Down-regulated (acute/control)						
Down-r	regulated (c	hronic/co	ntrol)		Down-	regulated (acute/coi	ntrol)			
Down-ı ID	regulated (c	hronic/con	ntrol) Ac / Co	Ch / Ac	Down-	regulated (Gene	acute/cou	ntrol) Ac / Co	Ch / Ac		
Down-1 ID NM 001033968	r <mark>egulated (c</mark> Gene Baa6	hronic/con Ch / Co -1.52	ntrol) Ac / Co	Ch / Ac	Down-1 ID NM 030858	regulated (Gene Smad7	acute/cou Ch / Co	Ac / Co	Ch / Ac		
Down-1 ID NM_001033968 NM_001100565	regulated (c Gene Bag6 Peli1	hronic/con Ch / Co -1.52 -1.27	ntrol) Ac / Co -	Ch / Ac	Down-1 ID NM_030858 NM_022701	regulated (Gene Smad7 Flot1	acute/con Ch / Co -	Ac / Co -1.45 -1.20	Ch / Ac		
Down-H ID NM_001033968 NM_001100565 NM_001008322	egulated (cl Gene Bag6 Peli1 Shmt2	hronic/con Ch / Co -1.52 -1.27 -1.21	ntrol) Ac / Co - -	Ch / Ac - -	Down- ID NM_030858 NM_022701 NM_017278	regulated (Gene Smad7 Flot1 Psma1	acute/coi Ch / Co - -	Ac / Co -1.45 -1.20 -1.17	Ch / Ac - -		
Down-r ID NM_001033968 NM_001100565 NM_001008322 NM_133388	egulated (cl Gene Bag6 Peli1 Shmt2 Rbm14	Ch / Co -1.52 -1.27 -1.21 -1.21 -1.19	Ac / Co - - - -	Ch / Ac - - - -	Down- ID NM_030858 NM_022701 NM_017278 NM_001128083	Gene Smad7 Flot1 Psma1 Trim8	Ch / Co Ch / Co - - -	Ac / Co -1.45 -1.20 -1.17 -1.16	Ch / Ac - - - -		
Down-r ID NM_001033968 NM_001100565 NM_001008322 NM_133388 NM_138710	egulated (cl Gene Bag6 Peli1 Shmt2 Rbm14 Dab2ip	Ch / Co -1.52 -1.27 -1.21 -1.19 -1.18	Ac / Co - - - - -	Ch / Ac - - - - -	Down-I ID NM_030858 NM_022701 NM_017278 NM_001128083 NM_001106332	Gene Smad7 Flot1 Psma1 Trim8 Otub1	acute/coi Ch / Co - - - - -	Ac / Co -1.45 -1.20 -1.17 -1.16 -1.16	Ch / Ac - - - - - -		
Down-H ID NM_001033968 NM_001100565 NM_001008322 NM_133388 NM_138710 NM_001039914	egulated (cl Gene Bag6 Peli1 Shmt2 Rbm14 Dab2ip Akirin2	Ch / Co -1.52 -1.27 -1.21 -1.21 -1.19 -1.18 -1.17	ntrol) Ac / Co - - - - - - - -	Ch / Ac - - - - - - - - - - -	Down- ID NM_030858 NM_022701 NM_017278 NM_001128083 NM_001106332 NM_021655	Gene Smad7 Flot1 Psma1 Trim8 Otub1 Chga	Ch / Co Ch / Co - - - - - - -	Ac / Co -1.45 -1.20 -1.17 -1.16 -1.16 -1.14	Ch / Ac 		

Table 4. Changes in gene regulation in E19 fetal brain following maternal treatment with paracetamol.

Up-regulated and down-regulated inflammatory and immune-related gene changes in the E19 fetal brain following no treatment (Co, controls), acute (Ac,) or chronic (Ch) maternal paracetamol treatment; see *Methods* for details of dosage schedule. Data from RNA-Seq analysis. Numbers are fold changes for comparisons indicated (Ch/Co, Ac/Co; Ch/Ac). Compared to the placenta, in the fetal brain many fewer inflammatory and immune-related genes showed regulatory changes and there were no significant differences between acute and chronic treatments. - indicates no significant difference in fold changes, not that there was no fold change.

compared to the control group, with five genes down-regulated greater than 500-fold (*Afp, apoc2, rbp4, apob and fgb,* Table 1). In addition, 10/50 genes were up-regulated following both treatments but 34/50 were down-regulated following both treatments. Thus overall the down-regulatory effects of paracetamol were much more pronounced than the up-regulatory effects.

Genes that showed a regulatory response in placentas of animals following both acute and chronic treatment with paracetamol are listed in Table 5. Seven of these down-regulated genes showed a fold change of more than two, which was greater in the chronically treated placentas. Other changes were so small that they are unlikely to be of much functional significance.

The inflammatory response

In the placenta of chronically treated rats there was a notable up-regulation of immune response related genes compared to the acutely treated group (Table 2). Figure 3 illustrates an analysis from biological Gene Ontology (GO) categories of immune response genes (A) subdivided into the innate (B) and adaptive (C) immune systems in the chronically treated animals. In the placenta these included GO biological processes such as neutrophil chemotaxis (p=4.7E-05) and innate immune response (p=0.045). Figure 3 illustrates that the number of significantly up-regulated genes was substantially more than the number of down-regulated genes and that most of these were in the innate immune system category, with a small number in the adaptive immune system. A list of inflammatory and immunerelated genes that were up-regulated in the placenta following chronic treatment is shown in Table 2. Overall, some 36 genes showed a statistically significant up-regulation. These included 15 genes that were up-regulated two-fold or more. As can be seen from Table 2, the third most up-regulated gene in the placenta following chronic paracetamol exposure was Il1ß. Figure 4 illustrates the number of $II1\beta$ gene transcripts in the three treatment groups in the fetal brain and placenta. There was a prominent increase in $II1\beta$ transcripts in the placentas from the dams treated with chronic paracetamol and no difference between the datasets of placentas from the control and

E19 Place	E19 Brain							
Up-regulated (chronic/con	Up-regulated (chronic/control & acute/control)							
ID Gene Ch	/ Co Ac /	Co Ch / Ac	ID	Gene	Ch / Co	Ac / Co	Ch / Ac	
NM_001009623 Tnfsf13 2	.09 2.0	7 -	NM_001172305	Prkcb	2.12	2.03	-	
NM_001034010 Tril 1	.63 1.79	9 -	NM_012713	Prkcb	1.89	2.19	-	
NM_001033691 Irf7 1	.92 1.52	2 -	NM_052807	Igf1r	1.65	1.49	-	
NM_017269 Ptprj 1	.70 1.5	2 -	NM_053374	Il18bp	1.60	1.52	-	
NM_001106123 Mrc1 1	.64 1.4	5 -	NM_001106757	Cfp	1.57	1.53	-	
NM_019211 Rasgrp1 1	.45 1.4	1 -	NM_001276715	Prkd1	1.39	1.41	-	
NM_001008886 RT1-S3 1	.31 1.4	0 -	NM_001079894	Plekha1	1.37	1.48	-	
NM_019140 Ptprs 1	.31 1.4	0 -	NM_001106095	Lig4	1.33	1.33	-	
NM_012512 B2m 1	.37 1.3	7 -	NM_013187	Plcg1	1.26	1.24	-	
NM_019195 Cd47 1	.24 1.34	4 -	NM_012747	Stat3	1.24	1.25	-	
NM_133395 Serinc5 1	.26 1.29	9 -						
NM_052807 Igf1r 1	.26 1.20	6 -						
Dermonande te distancia (es		h = (= = = t = = 1)	Deres an and a				4	
Down-regulated (chronic/co	Down-regulat	ea (chronic/		acute/con	ch (Ac			
ID Gene Cn		CO Ch / AC		Gene		AC / CO	Ch / Ac	
NM_012738 Apod 1	-16	-	NM_032085	Col3a l	-6.98	-6.99	-	
NM_022924 F2 -	-15) -	NM_001109116	Prr7	-3.24	-2.12	-	
NM_012737 Apod4 -	-/./	2 -	NM_030826	GpxT	-1.67	-1.52	-	
NM_017097 Ctsc -6	-6.1	8 -	NM_030859	Mdk	-1.33	-1.39	-	
NM_133393 Ljrig -4	-3.2	0 -	NM_053761	Zyx	-1.33	-1.39	-	
NM_001109116 P/17 -2		9 -	NM_212509	NJKDII I	-1.43	-1.38	-	
NM_022751 Cyba -2	2.03 -2.4	9 - 0	NM_022257	Masp I	-1.44	-1.38	-	
NM_033351 FCgrt -1	.85 -1.0	8 - 4	NM_001277283	Traine 27	-1.32	-1.30	-	
NM_001035922 Itil12 -1	.41 -1.5	4 -	NM_001134974	Irim27	-1.40	-1.35	-	
NM_001006969 IIJ3 -1	.50 -1.5	-	NM_022546	Dapk3	-1.58	-1.35	-	
NM_133293 Gata3 -1	.53 -1.4	2 -	NM_001106164	Cmtm3	-1.40	-1.35	-	
NM_001025126 //avira1 1	.55 -1.5	Z -	NM_001108153	Semara	-1.38	-1.33	-	
NM_001025136 Hexim1 -1	.34 -1.2	4 -	NM_172045	Ppp1r14b	-1.36	-1.32	-	
			NM_130411	Corola	-1.41	-1.31	-	
				SHZUZ	-1.52	-1.51	-	
			NM_001004080	USII	-1.55	-1.50	-	
			NM_053727	NJII3	-1.21	-1.29	-	
			NM_012931	BCal 1	-1.25	-1.24	-	
			NM_001107063	Cac42ep4	-1.20	-1.23	-	
			NIM 010250	rSIIID4	-1.20	-1.20	-	
			NNA 001021052	Dolrad	-1.20	-1.20	-	
				ruiisu	-1.1/	-1.19	-	
			NM 0010/7000	Vthdf2	-1.25	-1.19	_	

 Table 5. Inflammatory and immune-related gene regulation in both acute and chronic treatment with paracetamol.

Only genes that showed a response in placentas from E19 animals (left panels) and fetal brains (right panels) following both acute and chronic maternal treatment with paracetamol are shown; see *Methods* for details of dosage schedule. Data from RNA-Seq analysis. Numbers are fold changes for comparisons indicated (Ch/Co, Ac/Co; Ch/Ac). There were no significant differences for these genes between acute and chronic treatments, although there were small fold changes (data not shown).



Figure 2. Number of up-regulated and down-regulated genes in the E19 placenta and brain following chronic maternal treatment of paracetamol. Transcript numbers in placenta and brain from chronically (15mg/kg) treated compared to control, untreated animals. For details of chronic dosage schedule see Methods. Data derived from RNA-Seq analysis. Overlapping segments represent shared genes.



Figure 3. Pathway analysis from the Biological Gene Ontology categories (BP:GO). (A) "immune response", **(B)** "innate immune system" and **(C)** "adaptive immune system". The number of genes significantly up-regulated (green) and significantly down-regulated (red) are shown for adult brain, E19 brain and E19 placenta, as determined by RNA-Seq. Results are displayed for chronic and acute paracetamol treated rats (n=4). For details of chronic and acute dosage schedules see Methods.



Figure 4. IL1ß transcript counts. Transcripts per million in E19 fetal placenta and brain from control (n=4), acute (n=4) or chronically treated dams (n=4) as determined by RNA-Seq. (HTSeq-counts, EdgeR). Means \pm SD. * p <0.05.

acutely treated mothers, both showing very low numbers. $IL1\beta$ is a prototypical marker for inflammation and immune response, with up-regulation in the chronically treated placenta of 13.3 fold change; it could thus be a potential indicator of fetal harm. The response in the placenta following a single acute dose of paracetamol was much more muted (Table 2).

Amongst the down-regulated genes in the placenta (Table 1) were several transcripts for plasma proteins (AFP, transthyretin and transferrin, see Discussion) that have been shown to down-regulate under inflammatory conditions (negative acute phase response, Heinrich *et al.*, 1990; Hu *et al.*, 2019; Mackiewicz *et al.*, 1990). Two of these were markedly down-regulated in the acute experiments and further down-regulated in the chronic experiments (Table 1). This suggest that the response of these plasma protein genes was rapid in onset and continuing over several days in the presence of chronic treatment. In contrast, the up-regulatory response of cytokine genes only became prominent in the placentas of animals chronically exposed to paracetamol (Table 1; Figure 4).

IL1\beta concentration (ELISA). In order to see if the increase in transcript numbers for *IL1\beta* in placentas from dams treated chronically with paracetamol (Figure 4) translated into an increase in its protein concentration, the levels of this cytokine in plasma of both the dams and pups were measured using a commercially available ELISA kit (see Methods). Results are illustrated in Figure 5. None of the dams in any of the treatment groups had a detectable level of IL1ß in their plasma (limit <5pg/ml) nor was IL1ß detected in the control untreated fetuses. In contrast, IL1ß in the plasma of many of the E19 fetuses whose mothers had been treated with paracetamol was detected. The levels were generally higher in fetuses of mothers treated chronically (acute 2/4, chronic low 7/16 and chronic high 10/19).

The effect of paracetamol exposure on E19 fetal brain gene expression (transcriptomic analysis)

Transcriptomic analysis of the E19 fetal brain was carried out in material collected from the same animals as was prepared for placental analysis, thus allowing a direct comparison between the response of the two tissues to paracetamol treatment of the mother.

As illustrated in Figure 1, following maternal exposure to paracetamol, there was a large number of genes that significantly altered their expression in the fetal brain.

As shown in Figure 2, comparing the dataset for fetal brains from chronically treated dams with untreated control dams, there was a total 1128 genes with significantly different transcript numbers in the E19 brain. The top 50 up-regulated and down-regulated genes in the E19 brain are shown for both acute and chronic treatment groups compared to controls in Table 3. Following both treatments 26/50 genes were up-regulated and 40/50 were down-regulated. Additionally, the level of



IL1β concentration in plasma

Figure 5. Quantification of IL1β concentration in dam and fetal rat plasma. Samples from control (4 fetuses from 2 dams), acute (4 fetuses from 4 dams) or chronic paracetamol treated dams at low dose (3.75mg/kg, 16 fetuses from 5 dams) or high dose (15mg/kg, 19 fetuses from 7 dams). Measured by ELISA (R&D Quantikine). Means ± SD.

down-regulation was greater for most transcripts than up-regulation following both acute and chronic paracetamol treatment, for example *Colla1* (collagen type 1 alpha 1 chain) and *Col3a1* (collagen type 3 alpha 1 chain), see Table 3. There will be a further analysis of the brain data in a later publication (Koehn *et al., unpublished reports*) that will deal with expression of ABC efflux transporters and related enzymes as these may play a role in the extent to which paracetamol enters the brain at different stages of development (Koehn *et al.*, 2019b).

Comparison of the inflammatory response in E19 placenta and brain following maternal paracetamol treatment

In addition to effects of the length of exposure to the drug on gene expression in individual tissues, the regulation in brain and placenta was very different following the same treatment.

Only 98 genes were significantly regulated in both tissues, equating to 5.5% of the transcripts that changed their expression (Figure 2).

In the E19 placenta many of the top genes up-regulated following chronic treatment were related to immune-response and inflammation, including *Il1β*, which was 3 rd highest (Table 1). In contrast, in the brain, very few transcripts for *Il1β* (Figure 4) or other cytokines (Table 4) could be detected and there was no difference in transcripts for *Il1β* between the treatment groups (Figure 4). Table 4 lists the inflammatory and immunerelated genes that were up- or down-regulated significantly in the E19 brain. The changes were very small compared to the placenta in both the innate immune and the adaptive immune category (Figure 3). Table 5 shows immune/inflammatory related genes that showed a regulatory change in both the placenta and the brains from both acutely and chronically treated fetuses. No changes in plasma protein transcript numbers were detected in the fetal brain (see Discussion). This, together with lack of up-regulation of the inflammatory cytokine $Il1\beta$, as seen in the placentas, indicates that an inflammatory response was elicited by paracetamol in the placenta but little or none in the fetal brain. We do not have information if other organs not investigated in this study, such as the liver, could also have been affected.

Placental permeability

In order to investigate if a prolonged exposure to paracetamol and resulting inflammatory response could affect the permeability of the placenta, two sets of permeability experiments were conducted using a small molecular size marker, ¹⁴C-sucrose (see Methods). These were designed to examine the transfer from the mother to the fetus but also from the fetus back to the mother. Results are illustrated in Figure 6 and Figure 7.

Fetal to maternal transfer of ¹⁴C-sucrose. To investigate the placental transfer of sucrose from fetus back to the dam following maternal paracetamol exposure, sucrose was injected directly into the pups still within their amniotic sacs (see Methods). Two litters were injected in mothers that had been treated with chronic high doses of paracetamol and one litter from a mother treated with chronic low dose paracetamol. These were compared with one litter from an untreated control mother. Plasma samples from both the fetuses and dam were collected and ratio of ¹⁴C- sucrose estimated (see Methods). The results are shown in Figure 6. All three of the litters from mothers treated with chronic paracetamol (either high or low dose) showed slightly higher permeability from the fetus back to the mother than in the control dam. However, the ratios are extremely low, making accurate comparison difficult.



Figure 6. Fetal (E19) to maternal transfer of ¹⁴C-sucrose. Following maternal paracetamol treatment fetuses, still within their amniotic sacs, were directly injected (i.p.) with ¹⁴C-sucrose and plasma from both fetus and dam were collected to calculate maternal/average fetal plasma ratio (%) over time. Treatment groups: control (n=6), chronic low dose (3.75mg/kg, n=5) and chronic high dose (15mg/kg, n=5); n refers to number of fetuses. Each data point is a single fetus.



Figure 7. Maternal to fetal (E19) transfer of ¹⁴C-sucrose. Mothers were treated with paracetamol. ¹⁴C-sucrose was injected (i.p) into the mothers. Blood samples from individual fetuses were collected at the same time as maternal samples. Treatment groups were: untreated (control, n=8) and paracetamol injected (chronic dose 15mg/kg, n=6) dams. n refers to number of fetuses. Each data point is a single fetus. Transfer calculated as fetal/maternal plasma ratio (%).

Maternal to fetal transfer of ¹⁴C-sucrose. In order to investigate if the rate of transfer of a small molecular marker from dam to fetus across the placental barrier was affected following paracetamol exposure, dams either untreated (control) or treated with chronic high (15mg/kg) doses of paracetamol were given a final intravenous (i.v.) injection of ¹⁴C-sucrose 30 minutes

before removing their fetuses (Figure 7). Blood samples from dams were time matched to the removal and blood collection from each fetus (see Koehn *et al.*, 2019b). The transfer from the mother to the fetus in the paracetamol treated dams was slightly less than that in the control animal. The much higher ratios obtained in the maternal to fetal transfer experiments (Figure 7)

compared to the fetal to maternal transfer (Figure 6) are due to the differences in volume of distribution, hence dilution of the marker, when sucrose is injected into the mother or into the fetuses.

Detection of AFP in fetal and maternal plasma. In order to investigate if exposure to paracetamol can also influence the transfer of a protein from the fetal circulation into the maternal blood across the placenta, western blot analysis was made of fetal and maternal plasma samples using cross-reacting antibodies specific for AFP (see Methods). Figure 8A shows the blot that contained both the fetal and maternal samples together with one negative control (non-pregnant female rat). Densitometry measurements are illustrated in Figure 8B together with maternal/fetal ratios. There was no detectable band in the non-pregnant control sample and all maternal samples showed a much lower level of the protein than fetal samples. The levels of the protein in fetal samples did not appear to change between the control and any of the treatment groups (Figure 8B), but in maternal samples, AFP levels were higher in all chronically treated dams compared to un-treated controls. This was reflected in the ratios of AFP in maternal to fetal plasma (Figure 8B, right panel) in which all of the chronically treated animals had ratios that were above those in untreated controls and in one acutely treated animal. Prolonged exposure to the drug increased AFP transfer from fetus to dam by about three times compared to the control animals.

Permeability of the fetal blood brain barrier

Two different molecular size markers (¹⁴C-sucrose and plasma proteins) were used to assess any changes in blood-brain barrier permeability following chronic paracetamol treatment of the dam. The samples were obtained from the same experiments as the placental permeability studies.

Transfer of ¹⁴C-sucrose into the brain and CSF following different paracetamol treatment regimes. To investigate the transfer of ¹⁴C-sucrose into the fetal brain after maternal paracetamol exposure, fetal blood, brain and CSF samples from dams untreated (control) or treated acutely or chronically with either low (3.75mg/kg) or high (15mg/kg) doses of paracetamol were measured. As shown in Figure 9, there was no significant difference in the transfer into the brain and CSF between any treatment groups (Figure 9).

Entry into the brain and CSF when the ¹⁴C-sucrose was injected directly into the fetus was also investigated and results are illustrated in Figure 9. Here too there were no significant



Figure 8. Estimations of α-fetoprotein (AFP) concentrations in fetuses (E19) and dams. A) Western blots of AFP in plasma from dams and fetuses in different treatment groups. Numbers for dam blots are samples from individual animals; numbers in fetal plasma blots indicate individual fetuses from corresponding dams. Treatment groups were: control (1/1b; n=2), acute (2/2b; n=1), chronic low dose (3/3b; n=2), chronic high dose (4/4b; n=3) and non-pregnant control (5; n=1). **B)** Estimations of AFP in dam and fetal plasma (densitometry units from blots in (**A**) and fetal to maternal transfer of AFP expressed as dam/plasma AFP ratio (%). Note: each point represents an individual animal. Note that all chronic treated dams had higher plasma levels of AFP than the un-treated control pregnant dams; AU are ordinate arbitrary densitometry units.



Figure 9. Transfer of ¹⁴C-sucrose into the E19 brain (A) and cerebrospinal fluid (CSF; B) following paracetamol treatment. Fetuses were exposed to ¹⁴C-sucrose either directly (fetal i.p. injection) or indirectly (maternal i.v. injection). Treatment groups investigated were control, no paracetamol (n=13), chronic low dose (3.75mg/kg, n=11) and chronic high dose (15mg/kg, n=11) in fetuses that were injected directly. In experiments in which the ¹⁴C-sucrose was injected into the mothers; n numbers were control (n=8), acute (n=10) and chronic high dose (n=6) in the mothers. Means ± SD.

differences in the entry of sucrose into brain and CSF between the three treatments.

paracetamol exposure of the dam, the inflammatory response in the placenta nor the increased levels on IL1ß in fetal blood.

However, the fetuses that were directly exposed to sucrose (i.p injection) showed a lower level of transfer into the brain and CSF compared to those that were exposed indirectly (i.v. injection to dam), around 10% compared to 40%, respectively. This reflects differences in distribution volume following the different routes of injection as well as the time involved in samples collection.

Blood brain barrier integrity for endogenous plasma protein. Transfer of large molecule plasma proteins into the fetal brain following maternal paracetamol exposure was studied using immunohistochemistry and antibodies to rat serum proteins (see Methods). Brains were matched with plasma samples containing detectable IL1ß levels as estimated by ELISA (Figure 5). The distribution of the proteins in E19 brains from control, acute and chronic high dose (15mg/kg) paracetamol treated dams is illustrated in micrographs in Figure 10. There was no evidence of a "leak" of protein in any of the vessels in the fetal brains examined. In all sections stained from all brains investigated, immunostaining was exclusively localised in the blood vessels, choroid plexus stroma and precipitated CSF and there was no visible difference in the brain morphology between treatment groups.

Thus, the results clearly show that the blood-brain barrier, at least to plasma proteins and to sucrose, was not affected by

Discussion

In order for a drug taken by a pregnant mother to reach the fetal brain it has to cross both the placental and the blood-brain barriers. Any changes to normal functioning of these interfaces could have detrimental effects on fetal health and pregnancy outcomes. We have therefore analysed the transcriptomic changes in rat E19 placentas and brains following paracetamol treatment of the dams. Paracetamol is one of the most commonly used medications in pregnancy (Dreyer et al., 2015; Wyszynski & Shields, 2016). Pregnant rats were treated with paracetamol acutely and chronically and compared to controls (no treatment). The doses used were within the clinically recommended range (0.5g to 4g in 24 hours in adults). In the case of the chronic treatment, this corresponded to a relatively prolonged period of pregnancy in the rat (E15-19, about 25% of gestation). This was followed by investigating placental transfer of small and large molecules from the dam to the fetus and from the fetus back to the maternal circulation, to see whether paracetamol exposure altered barrier function. Finally, the permeability of the blood-brain barrier was analysed in the fetuses of paracetamol treated and untreated dams.

From the results it was apparent that some form of acute phase response was elicited as transcripts for several plasma proteins were down-regulated in placentas of both acute and chronic treated animals (Table 1). These proteins were AFP, transthyretin



Figure 10. Histology of E19 fetal brains. A) Hematoxylin and eosin coronal section of E19 neocortex of fetus from mother treated with chronic high dose paracetamol. **B**) Adjacent section from same brain as **A** immunostained for plasma proteins. **C**) High power image from **B** (box). **D**) High power immunostained image of E19 neocortex of fetus from mother treated with acute high dose paracetamol. Note that all cerebral vessels appear intact with protein immunostained deposits all within blood vessel lumen, indicating that paracetamol treatment has not affected their barrier permeability to plasma proteins. Bars in **A** & **B** are 1mm; bars in **C** & **D** are 100µm.

and transferrin (Vranckx et al., 1989), fibrinogen beta chain (Birch & Schreiber, 1986) and apolipoproteins Apoal-4, several of which are known to respond to inflammation as negative acute phase proteins (Tu et al., 1987). Since a marked response was already apparent after a single dose of paracetamol, it seems that this was a rapid response to paracetamol, which was sustained and increased when the treatment was chronic. A summary of transcript numbers for AFP, transferrin and transthyretin, together with numbers for $II1\beta$ for comparison, is presented in Table 6 for both the brain and the placenta. These clearly show that some form of acute phase response was taking place in the placenta following paracetamol treatment; however, other typical acute phase response-related cytokines were not up-regulated (such as TNF α or IL6). Transcript numbers in the brain did not change, demonstrating that the acute phase response was tissue specific and restricted to the placenta.

Several immune and inflammatory-related genes were up-regulated in placentas of animals treated with the chronic doseregime, but much less so in the placentas of acutely treated animals (Table 1, Table 2 and Table 5). The key inflammatory cytokine, IL18, was shown to be present in the blood of a high proportion of fetuses of mothers exposed to both acute and chronic treatment with paracetamol. The levels were variable in different fetuses but generally higher in the chronically treated animals. No IL1 β could be detected in either the maternal blood of paracetamol treated animals or in fetuses of control untreated animals. This confirms that paracetamol was indeed eliciting an inflammatory response but only on the fetal side of the placental circulation. Thiele *et al.* (2015) reported that pregnant mice treated with either 50 or 250mg/kg paracetamol showed some immune responses in the uterus and some morphological changes in the placenta. However, they did not investigate possible immune responses in the placenta and the doses of paracetamol were much larger than the ones we used and were well above the clinical range.

In order to determine if prolonged paracetamol exposure of the dam could affect some aspect of placental function, we have estimated placental permeability to a small molecular marker, sucrose and to large plasma protein AFP in both directions i.e. from the dam to the fetuses and from the fetuses back to the dam. The results showed that there was a small and variable increase in permeability to ¹⁴C-sucrose and of AFP permeability in the direction from fetus to mother (Figure 6 and Figure 8). There may also have been a small decrease in sucrose permeability

E19 placenta												
	Control				Acute				Chronic			
Sample	Afp	Tf	Ttr	IL1β	Afp	Tf	Ttr	IL1β	Afp	Tf	Ttr	IL1β
1	10372	2906	783	2.7	3.5	31	0.7	1.3	2.7	35	1.4	11
2	16873	3789	1100	2.5	3.0	35	2.6	2.2	3.5	28	1.6	30
3	1263	366	65	2.3	1571	413	95	3.3	3.5	17	1.7	46
4	72	56	4.6	3.8	8.4	29	1.6	2.2	2.9	45	1.1	58
E19 brain												
	Control				Acute				Chronic			
Sample	Afp	Tf	Ttr	IL1β	Afp	Tf	Ttr	IL1β	Afp	Tf	Ttr	IL1β
1	0.4	2.8	0.2	0.1	0.2	1.2	2.3	0.1	0.2	2.9	0.8	0.2
2	1.7	2.1	0.5	0.1	0.1	1.2	0.4	0.1	0.1	3.2	1.1	0.1
3	0.2	5.0	0.4	0.2	0.1	1.9	0.2	0.1	0.1	3.5	0.9	0.1
4	0.1	33	0.2	0.1	0.2	1 /	03	0.1	0.2	28	10	0.1

Table 6. Transcript numbers in E19 placenta and fetal brain for negative acute phase plasma proteins and IL1 $\beta.$

In placental samples transcript counts per million for the three negative acute phase proteins were smaller in chronically treated animals compared to controls and all $IL1\beta$ numbers were greater than in controls. There was some variation in values between individual placentas which was more obvious in samples from acutely treated animal indicating that the response was potentially time-dependent. In all brain samples the transcript numbers were very low, with no evidence of an inflammatory response. This indicates that the response in the placenta was tissue specific.

from mother to fetus (Figure 7) but due to small numbers this is inconclusive.

Placental inflammation induced by lipopolysaccharide (LPS) injection in pregnant rats has been reported to induce maternal serum and placental cytokines and increased maternal serum AFP (Hu *et al.*, 2019). In those experiments LPS did not increase the expression of AFP in fetal liver, maternal liver or placenta, but did reduce the fetal serum AFP levels, a pattern implying a possibility of increased transfer of AFP from the fetus to the mother, thus depleting it from fetal circulation. We did not find any difference in fetal AFP levels but this discrepancy could be due to either the duration and severity of the response or sensitivity of the methods used.

Permeability of the fetal blood-brain barrier to both sucrose and plasma protein was also investigated. In contrast to the placenta, there was no evidence of a change in brain barrier permeability to either marker in fetuses of dams treated with paracetamol. This is relevant to earlier studies in which inflammation induced by LPS was shown to result in a breakdown of the blood-brain barrier that was age-dependant (Stolp *et al.*, 2005a; Stolp *et al.*, 2005b). However, it is likely that E19 is at a developmental stage when the response to LPS is not yet developed, as shown in a similar study in a marsupial species, *Monodelphis domestica* (Stolp *et al.*, 2005a).

Limitations of the study

The study has been carried out in pregnant rats at a single gestational age (E19). This stage of brain development in rats at E19 is approximately equivalent to 22-24 weeks gestation in humans (Clancy et al., 2001), corresponding to the earliest age of viability (Fischer et al., 2009; Stoll et al., 2010). The rat and human placentas are both classed as hemochorial (Blood et al., 2007; Dawe et al., 2007) but there are differences in morphology, in particular that the rat placenta has more morphological layers between the fetal and maternal circulations. However, that might mean that the relatively small changes in placental permeability from fetus to mother shown here might be more prominent in the human. The responses of these two species to an inflammatory event are similar with respect to the three plasma proteins AFP, transferrin and transthyretin (prealbumin); as in this study, these proteins have been reported to be acute phase negative proteins under inflammatory conditions (Heinrich et al., 1990; Hu et al., 2019; Mackiewicz et al., 1990). This supports the suggestion that these findings should be taken account of when advising pregnant women about the use of paracetamol. Given the unexpected findings of up-regulation of inflammatory cytokines and down-regulation of some acute phase plasma proteins, we are in the process of carrying out RNA-Seq replication studies and extending the range of cytokines estimated in fetal and maternal blood. Unfortunately, these experiments have been delayed by the COVID-19 emergency, which has

closed our laboratories for an indefinite period. In view of the potential significance of our findings for the use of paracetamol in pregnancy, particularly the high frequency of its use, we feel it is justified to present these findings for peer review, in their present form.

Clinical relevance

Paracetamol (acetaminophen) is generally considered "safe" to use in pregnancy and lactation (Australian Medicines Handbook, 2019; Briggs et al., 2017) although it is one of the most commonly overdosed drugs, including in pregnancy (Rayburn et al., 1984). However, some authors urge caution in its use because of evidence of adverse effects (Brune et al., 2015). It has been reported that as many as nearly 80% of pregnant women in some populations ingest paracetamol (Dreyer et al., 2015). The findings of the present study, although based solely on experiments in rats, should be taken account of when advising pregnant patients on the use of paracetamol in pregnancy. The clinical situation is not straightforward because in addition to taking paracetamol to relieve pain, it may also be taken to reduce an increase in body temperature accompanying an infection (often respiratory), but there is evidence of an association between infection/fever and adverse outcomes for pregnancies; this seems to be a particular problem when infection/fever occurs at the beginning of the 3rd trimester (Hagberg et al., 2015). Thus, continued but limited use of paracetamol to control severe pain and to reduce body temperature at critical stages of pregnancy would seems to be appropriate but not the widespread use for lesser indications that is implied by the reports that most pregnant women take paracetamol.

Increased transfer of sucrose and AFP from fetal circulation into maternal circulation, as demonstrated in the present study, suggests that other molecules/metabolites could potentially also reach the maternal circulation. There are several clinical implications, including that increased AFP levels detected in pregnant women are used to detect potential neural tube closure defects, although this test is done earlier in pregnancy and we have as yet no evidence of paracetamol affecting placental permeability this early in pregnancy. Further investigation is required to see if there are similar effects in the placentas of patients who have taken paracetamol. If the effect is indeed confined to the fetal side of the placenta it will be clinically difficult to determine such an effect in pregnant patients, particularly if it turns out to be variable as in our rat experiments, although transfer of AFP from fetal to maternal circulation might be a useful indicator.

Data availability Underlying data

RNA-Seq data on NCBI, Accession number PRJNA633629: https://identifiers.org/ncbi/bioproject:PRJNA633629

Figshare: Effects of paracetamol on rat placenta and fatal brain. https://doi.org/10.26188/5ebff4c2781a0 (Koehn *et al.*, 2020)

This project contains the following underlying data:

- 200514 ELISA raw data.xlsx (raw data for the IL-1 β ELISA)
- 200514 sucrose permeability data.xlsx (brain, CSF and plasma levels of sucrose in pregnant rats and fetuses)
- RA708 chronic high dose paracetamol.zip (plasma protein and H&E stained sections in Figure 10, A: RA708-50-04 HE x4.jpg, B: RA708-46-05 PP x4.jpg, C: RA708-46-05 PP x40.jpg
- RA677 actute high dose paracetamol.zip (plasma protein stained section in Figure 10, D: RA677-41-03 x40 B.jpg)
- 20191204 AFP loExp 1.tif (original unedited western blot image for Figure 8)

Data are available under the terms of the Creative Commons Attribution 4.0 International license (CC-BY 4.0).

Acknowledgements

RNA-Sequencing was undertaken by The Australian Genome Research Facility, Peter MacCallum Cancer Centre, Melbourne, Victoria.

References

Afgan E, Dannon Baker D, Batut B, et al.: The Galaxy Platform for Accessible, Reproducible and Collaborative Biomedical Analyses: 2018 Update. Nucleic Acids Res. 2018; 46(W1): W537-W544.

PubMed Abstract | Publisher Full Text | Free Full Text

Anthony DC, Bolton SJ, Fearn S, *et al.*: Age-related effects of interleukin-1 beta on polymorphonuclear neutrophil-dependent increases in bloodbrain barrier permeability in rats. *Brain*. 1997; **120**(Pt 3): 435–444. PubMed Abstract | Publisher Full Text

Australian Medicines Handbook Pty Ltd: Adelaide SA, Australia. 2019. Reference Source

Bauer AZ, Kriebel D, Herbert MR, et al.: Prenatal paracetamol exposure and child neurodevelopment: A review. Horm Behav. 2018; 101: 125–147. PubMed Abstract | Publisher Full Text

Birch HE, Schreiber G: Transcriptional regulation of plasma protein

synthesis during inflammation. J Biol Chem. 1986; 261(18): 8077–8080. PubMed Abstract

Blood DC, Studdert VP, Gay CC: Saunders Comprehensive Veterinary Dictionary, 3 ed. © Elsevier, Inc. 2007. Reference Source

Reference source

Briggs GG, Freeman RK, Towers CV, *et al.*: **Drugs in Pregnancy and Lactation: A Reference Guide to Fetal and Neonatal Risk**. 11th Ed. Lippincott Williams & Wilkins (LWW). 2017.

Reference Source

Brochu ME, Girard S, Lavoie K, *et al.*: **Developmental regulation of the neuroinflammatory responses to LPS and/or hypoxia-ischemia between preterm and term neonates: An experimental study.** *J Neuroinflammation.* 2011; 8: 55.

PubMed Abstract | Publisher Full Text | Free Full Text

Brune K, Renner B, Tiegs G: Acetaminophen/paracetamol: A history of errors, failures and false decisions. *Eur J Pain*. 2015; **19**(7): 953–65. PubMed Abstract | Publisher Full Text

Challis JR, Lockwood CJ, Myatt L, et al.: Inflammation and pregnancy. Reprod Sci. 2009; 16(2): 206-215.

PubMed Abstract | Publisher Full Text

Clancy B, Darlington RB, Finlay BL: Translating developmental time across mammalian species. Neuroscience. 2001; **105**(1): 7–17. PubMed Abstract | Publisher Full Text

Dawe GS, Tan XW, Xiao ZC: Cell migration from baby to mother. Cell Adh Migr. 2007; 1(1): 19-29.

PubMed Abstract | Free Full Text

Dreyer NA, Blackburn SC, Mt-Isa S, et al.: Direct-to-Patient Research: Piloting a New Approach to Understanding Drug Safety During Pregnancy. JMIR Public Health Surveill. 2015; 1(2): e22.

PubMed Abstract | Publisher Full Text | Free Full Text

Dziegielewska KM, Evans CA, Lai PC, et al.: Proteins in Cerebrospinal Fluid and Plasma of Fetal Rats During Development. Dev Biol. 1981; 83(1): 193-200 PubMed Abstract | Publisher Full Text

Fischer N, Steurer MA, Adams M, et al.: Survival rates of extremely preterm infants (gestational age < 26 weeks) in Switzerland: impact of the Swiss guidelines for the care of infants born at the limit of viability. Arch Dis Child Fetal Neonatal Ed. 2009; 94(6): F407–F413.

PubMed Abstract | Publisher Full Text

Fleiss B, Gressens P, Stolp HB: Cortical gray matter injury in encephalopathy of prematurity: link to neurodevelopmental disorders. Front Neurol. 2020; 11: 567.

PubMed Abstract | Publisher Full Text | Free Full Text

Habgood MD, Sedgwick JE, Dziegielewska KM, et al.: A Developmentally Regulated Blood-Cerebrospinal Fluid Transfer Mechanism for Albumin in Immature Rats. J Physiol. 1992; 456: 181–192. PubMed Abstract | Publisher Full Text | Free Full Text

Habgood MD, Knott GW, Dziegielewska KM, *et al.*: **The Nature of the Decrease** in **Blood-Cerebrospinal Fluid Barrier Exchange During Postnatal Brain Development in the Rat.** *J Physiol.* 1993; **468**: 73–83. **PubMed Abstract | Publisher Full Text | Free Full Text**

Hagberg H, Mallard C, Ferriero DM, et al.: The role of inflammation in perinatal brain injury. Nat Rev Neurol. 2015; 11(4): 192-208. PubMed Abstract | Publisher Full Text | Free Full Text

Heinrich PC, Castell JV, Andus T: Interleukin-6 and the acute phase response. Biochem J. 1990; 265(3): 621-636.

PubMed Abstract | Publisher Full Text | Free Full Text

Hu J, Zhang J, Chan Y, et al.: A rat model of placental inflammation explains the unexplained elevated maternal serum alpha-fetoprotein associated with adverse pregnancy outcomes. J Obstet Gynaecol Res. 2019; 45(10): 1980-1988.

PubMed Abstract | Publisher Full Text

Huang DW, Sherman BT, Lempicki RA: **Systematic and integrative analysis of large gene lists using DAVID Bioinformatics resources.** *Nature Protoc.* 2009a; **4**(1): 44–57.

PubMed Abstract | Publisher Full Text

Huang DW, Sherman BT, Lempicki RA: Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res. 2009b; 37(1): 1-13.

PubMed Abstract | Publisher Full Text | Free Full Text

Huleihel M, Golan H, Hallak M: Intrauterine infection/inflammation during pregnancy and offspring brain damages: possible mechanisms involved. Reprod Biol Endocrinol. 2004: 2: 17.

PubMed Abstract | Publisher Full Text | Free Full Text

Johansson PA, Dziegielewska KM, Liddelow SA, et al.: The blood-CSF Barrier Explained: When Development Is Not Immaturity. Bioessays. 2008; 30(3): 237-48.

PubMed Abstract | Publisher Full Text

Koehn L, Dziegielewska KM, Mollgard K, et al.: Developmental differences in the expression of ABC transporters at rat brain barrier interfaces following chronic exposure to diallyl Sulfide. Sci Rep. 2019a; 9(1): 5998. PubMed Abstract | Publisher Full Text | Free Full Text

Koehn L, Habgood M, Huang Y, et al.: Determinants of drug entry into the developing brain [version 1; peer review: 3 approved]. F1000Res. 2019b; 8: 1372. PubMed Abstract | Publisher Full Text | Free Full Text

Koehn L, Huang Y, Habgood M, *et al*.: Effects of paracetamol on rat placenta and fatal brain. University of Melbourne. 2020. http://www.doi.org/10.26188/5ebff4c2781a0

Koehn L, Huang Y, Habgood MD, et al.: Efflux transporter transcriptome in rat placenta and developing brain: response to paracetamol. Unpublished Reports

Mackiewicz A, Ganapathi MK, Schultz D, et al.: Transforming growth factor beta 1 regulates production of acute-phase proteins. Proc Natl Acad Sci U S A. 1990; **87**(4): 1491-1495.

PubMed Abstract | Publisher Full Text | Free Full Text

Mudunuri U, Che A, Yi M, et al.: bioDBnet: the biological database network. Bioinformatics. 2009; 25(4): 555-556

PubMed Abstract | Publisher Full Text | Free Full Text

Nelson KB, Dambrosia JM, Grether JK, et al.: Neonatal cytokines and coagulation factors in children with cerebral palsy. Ann Neurol. 1998; 44(4): 665-675

PubMed Abstract | Publisher Full Text

Rayburn W, Aronow R, DeLancey B, et al.: Drug overdose during pregnancy: an overview from a metropolitan poison control center. Obstet Gynecol. 1984; 64(5): 611-614. **PubMed Abstract**

Romero R, Gotsch F, Pineles B, et al.: Inflammation in pregnancy: its roles in reproductive physiology, obstetrical complications, and fetal injury. Nutr Rev. 2007; 65(12 Pt 2): S194-202.

PubMed Abstract | Publisher Full Text

Salafia CM, Mangam HE, Weigl CA, et al.: Abnormal fetal heart rate patterns and placental inflammation. *Am J Obstet Gynecol.* 1989; **160**(1): 140–147. PubMed Abstract | Publisher Full Text

Seyednasrollah F, Rantanen K, Jaakkola P, et al.: ROTS: reproducible RNA-seq biomarker detector—prognostic markers for clear cell renal cell cancer. Nucleic Acids Res. 2016; 44(1): e1. PubMed Abstract | Publisher Full Text | Free Full Text

Soneson C, Delorenzi M: A comparison of methods for differential expression analysis of RNA-seq data. *BMC Bioinformatics*. 2013; **14**: 91. PubMed Abstract | Publisher Full Text | Free Full Text

Stoll BJ, Hansen NI, Bell EF, et al.: Neonatal outcomes of extremely preterm infants from the NICHD Neonatal Research Network. Pediatrics. 2010; 126(3): 443-456.

PubMed Abstract | Publisher Full Text | Free Full Text

Stolp HB, Dziegielewska KM, Ek CJ, et al.: Breakdown of the blood-brain barrier to proteins in white matter of the developing brain following systemic inflammation. Cell Tissue Res. 2005a; 320(3): 369-378. PubMed Abstract | Publisher Full Text

Stolp HB, Dziegielewska KM, Ek CJ, et al.: Long-term changes in blood-brain barrier permeability and white matter following prolonged systemic inflammation in early development in the rat. *Eur J Neurosci.* 2005b; 22(11): 2805-2816

PubMed Abstract | Publisher Full Text

Thiele K, Solano ME, Huber S, et al.: Prenatal Acetaminophen Affects Maternal Immune and Endocrine Adaptation to Pregnancy, Induces Placental Damage, and Impairs Fetal Development in Mice. Am J Pathol. 2015: 185(10): 2805-2818.

PubMed Abstract | Publisher Full Text

Thornton C, Leaw B, Mallard C, et al.: Cell Death in the Developing Brain after Hypoxia-Ischemia. Front Cell Neurosci. 2017; 11: 248. PubMed Abstract | Publisher Full Text | Free Full Text

Tian X, Zhu M, Du L, et al.: Intrauterine inflammation increases maternofetal transfer of gold nanoparticles in a size-dependent manner in murine pregnancy. Small. 2013; 9(14): 2432-2439.

PubMed Abstract | Publisher Full Text

Tu GF, De Jong F, Apostolopoulos J, et al.: Effect of acute inflammation on rat apolipoprotein mRNA levels. Inflammation. 1987; 11(2): 241-251 PubMed Abstract | Publisher Full Text

Vranckx R, Savu L, Cohen A, *et al*.: Inflammatory competence of fetal rat: acute-phase plasma protein response of the fetus treated by turpentine *in* utero. Inflammation. 1989; 13(1): 79-90. PubMed Abstract | Publisher Full Text

Wyszynski DF, Shields KE: Frequency and type of medications and vaccines used during pregnancy. Obstet Med. 2016; 9(1): 21-27. PubMed Abstract | Publisher Full Text | Free Full Text

Open Peer Review

Current Peer Review Status: 💉

Version 2

Reviewer Report 07 September 2020

https://doi.org/10.5256/f1000research.28770.r69768

© **2020 Ferretti P.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Patrizia Ferretti 匝

Stem Cells and Regenerative Medicine Section, UCL Institute of Child Health, University College London, London, UK

I thank the authors for carefully considering my comments. I am happy with their replies, hence I endorse indexing of their interesting manuscript.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: stem cells, tissue repair, developmental biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 20 July 2020

https://doi.org/10.5256/f1000research.26605.r64436

© **2020 Ferretti P.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

?

Patrizia Ferretti 匝

Stem Cells and Regenerative Medicine Section, UCL Institute of Child Health, University College London, London, UK

This paper addresses an important and neglected question of potential negative effects of

paracetamol during pregnancy. It examines gene expression changes in placenta and foetal brain and also presents out a number of functional studies to establish whether placenta permeability or brain barriers in the fetus are affected by the treatment. The study reports significant changes in the expression of a number of genes, including genes associated with the immune response, and validates changes in one of these genes, Il1ß, at the protein level in the placenta. While in some respects this study is still preliminary, the information presented here is valuable for underpinning future studies.

The authors clearly explain their choice of the end point (E19) selected, but not the reason for starting the chronic treatment at E15, rather than earlier, when important developmental events occur and teratogenic effects might be more likely and significant.

Specific Comments:

P6: It is not clear why only the t-test was used when comparing multiple groups, as ANOVA followed by a post-hoc test should have been used.

On p7 the authors say "...65 up-regulated and 57 down-regulated....", but Fig. 1 indicates 64 upregulated genes, consistent wih the total of 121 up and down-regulated transcripts indicated in the right column.

The authors indicate that expression of 737 genes is significantly affected by chronic treatment, but do not show the level of significance. Does this mean that p is <0.05 (but never <0.01 or smaller) for all transcripts?

Table 1 and 3. It would be helpful to colour code genes that change in both acute and chronic treatment groups and use thicker vertical lines between groups for ease of visualization.

Table 2 includes genes that are not in the top 50 shown in Table 1, and this should be clearly stated (at a first glance the Table seemed a bit redundant). As for Table 1, the level of significance should be indicated. The Table could be made it easier to read if the "up-regulated (acute/control)" genes were shown below the "up-regulated (chronic/control)", rather than in adjacent columns, or were clearly separated using a thicker vertical line. In addition, it is confusing to have a column "chronic/control" under the "up-regulated (acute/control)" list. This seems to have been done to accommodate S100a rather than inserting it under each comparison. Please check carefully that the difference indicated in different Tables are the same (e.g. S100a8 has a FC 2.25 in Tab1 and FC 2.26 in Table 2). "Il1b" should be changed to "Il1ß". A pie chart of the inflammatory genes to complement Table 2 and Fig. 3 would be useful.

P17, left column, top and Fig. 4. There is clearly variability, but to give numbers of fetuses where II1ß levels could be detected over total numbers assayed for all groups would be more accurate and informative (e.g. acute 2/4, chronic low 7/16 and chronic high 10/19) than including these number only for the chronic high group, which appears to be wrongly given as 19/39, while the number of fetuses indicated for this group in Fig. 5 legend is 19.

Figs 6 and 7 do not include error bars and no statistical analysis of these data seems to have been performed. It should be clearly indicated whether there was no statistical difference among groups at any time point studied.

P20, left column, top. The statement: "AFP levels were higher in all treated dams compared to an un-treated control." should be revised, as Fig. 8 shows an AFP increase only in chronically treated dams. It is a pity that the number of dams is too small to assess the significance of this observation and that no housekeeping protein was used to normalize AFP expression. If B is a densitometry of the gel in A, where according to the western blot labelling and the legend there is only 1 control for both dam and fetus, why are there 2 samples indicated in the controls in the charts? Given the low sample numbers and variability, particularly in fetal AFP levels, expressing the data as ratio is not appropriate.

Is the work clearly and accurately presented and does it cite the current literature? Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others? $\ensuremath{\mathsf{Yes}}$

If applicable, is the statistical analysis and its interpretation appropriate? Partly

Are all the source data underlying the results available to ensure full reproducibility? $\ensuremath{\mathsf{Yes}}$

Are the conclusions drawn adequately supported by the results? Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: stem cells, tissue repair, developmental biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response () 02 Aug 2020

Norman Saunders, University of Melbourne, Parkville, Australia

We should like to thank Professor Ferretti for her detailed review of this paper and in particular for her helpful suggestions for clarification and improvement of presentation of the results of this study. We provide responses below to all of the matters raised by Professor Ferretti.

This paper addresses an important and neglected question of potential negative effects of paracetamol during pregnancy. It examines gene expression changes in placenta and foetal brain and also presents out a number of functional studies to establish whether placenta

permeability or brain barriers in the fetus are affected by the treatment. The study reports significant changes in the expression of a number of genes, including genes associated with the immune response, and validates changes in one of these genes, Illß, at the protein level in the placenta. While in some respects this study is still preliminary, the information presented here is valuable for underpinning future studies.

The authors clearly explain their choice of the end point (E19) selected, but not the reason for starting the chronic treatment at E15, rather than earlier, when important developmental events occur and teratogenic effects might be more likely and significant.

The chronic treatment was limited to the last week of pregnancy in the rats, starting at E15, because it involved twice daily intraperitoneal (IP) injections of the drug and we were concerned that a longer treatment was an unreasonable imposition on the animals. In addition, it is well known that pregnant rats, if unduly stressed, are prone to aborting their fetuses. Also as mentioned in the 1st paragraph of the Discussion, the period of treatment covers about 25% of the gestational period of the rat, which is the main period that we are currently interested in. Oral medication would have been an alternative but it gives much less control over the amount of drug administered than when administered by IP injection. To have any control of oral administration requires monitoring of blood levels of a drug (itself an invasive process). This was not available at the time of this study but has now been developed for future studies. In relation to possible teratogenic effects, this was not an aim of the study and would, as the Reviewer indicates, require administration earlier in pregnancy. There is only limited information on possible teratogenic effects of paracetamol, probably because it came into clinical use long before specific regulatory requirements (e.g. FDA) for teratogenic testing in animals came into force.

Studies involving limited epidemiological data have concluded that there is no evidence of an association between paracetamol ingestion and congenital malformations (Briggs et al. 2019, pp8-11).

Specific Comments:

P6: It is not clear why only the t-test was used when comparing multiple groups, as ANOVA followed by a post-hoc test should have been used.

For the analysis of RNA-Seq data, the t-test is part of the packages we used and includes built in posthoc corrections for multiple comparisons. For the data on Il1ß the advice we have from our departmental statistical expert is that t-tests are appropriate for this type of research: (Lew M.J. (2019) A Reckless Guide to P-values. In: Bespalov A., Michel M., Steckler T. (eds) Good Research Practice in Non-Clinical Pharmacology and Biomedicine. Handbook of Experimental Pharmacology, vol 257. Springer, Cham.).

Nevertheless in view of the Reviewer's comment we have run the data through ANOVA followed by Tukey's posthoc test. The significance levels are the same as those we obtained with a *t*-test. We have added this information to the Methods subsection "Statistics".

On p7 the authors say "...65 up-regulated and 57 down-regulated....", but Fig. 1 indicates 64 up-regulated genes, consistent with the total of 121 up and down-regulated transcripts indicated in the right column.

We thank the Reviewer for drawing our attention to this error which has been corrected.

The authors indicate that expression of 737 genes is significantly affected by chronic treatment, but do not show the level of significance. Does this mean that p is <0.05 (but never <0.01 or smaller) for all transcripts?

As indicated in the Methods section on "Statistical Analysis" we used P<0.05 for two of the three analyses used. We focussed on genes with large fold changes as these are more likely to be of functional significance than would be indicated by a higher level of statistical significance.

Table 1 and 3. It would be helpful to colour code genes that change in both acute and chronic treatment groups and use thicker vertical lines between groups for ease of visualization.

We thank the Reviewer for this suggestion. The Tables have been modified accordingly. The treatment groups are now separated by a gap. The colour coding highlights some interesting differences in the number of genes that responded in the different treatment groups. A note of this has been added to the Table legends and in the text.

Table 2 includes genes that are not in the top 50 shown in Table 1, and this should be clearly stated (at a first glance the Table seemed a bit redundant).

This Table shows only inflammatory and immune-related genes and therefore some genes in the top 50 in Table 1 do not appear here. This is now indicated in the legend.

As for Table 1, the level of significance should be indicated.

P<0.05 added to legend.

The Table could be made it easier to read if the "up-regulated (acute/control)" genes were shown below the "up-regulated (chronic/control)", rather than in adjacent columns, or were clearly separated using a thicker vertical line.

This change would make a very long 2 column table. We prefer the helpful suggestion that the columns should be separated which we have done with a narrow blank column

In addition, it is confusing to have a column "chronic/control" under the "up-regulated (acute/control)" list. This seems to have been done to accommodate S100a rather than inserting it under each comparison.

Unfortunately in the editorial process of preparing the pdf from the submitted Table spreadsheets some of the down-regulated genes have been sliced off and put incorrectly under the up-regulated categories. We are puzzled by this as the proof we received to check was correct. I have discussed this with the Editorial Office who have indicated that they will make sure this does not occur in the next version. Please check carefully that the difference indicated in different Tables are the same (e.g. S100a8 has a FC 2.25 in Table1 and FC 2.26 in Table 2).

This was due to a difference in rounding, which has now been corrected.

"Il1b" should be changed to "Il1ß".

Il1b is the notation used in the gene data base ncbi.nim.nih.gov, we would prefer to retain this notation in tables.

A pie chart of the inflammatory genes to complement Table 2 and Fig. 3 would be useful.

We generally find that pie charts are not helpful and would prefer not to make this addition.

P17, left column, top and Fig. 4. There is clearly variability, but to give numbers of fetuses where Il1ß levels could be detected over total numbers assayed for all groups would be more accurate and informative (e.g. acute 2/4, chronic low 7/16 and chronic high 10/19) than including these number only for the chronic high group, which appears to be wrongly given as 19/39, while the number of fetuses indicated for this group in Fig. 5 legend is 19.

We think that the Reviewer is probably referring to Fig 5. We agree that the way of representing these data that the Reviewer has suggested is clearer. This has been incorporated into the text (bottom P30). Figure 5 has been modified to make it clearer that values were obtained from 4 control fetuses. The legend has been re-written to make it clearer how many dams and fetuses were involved in this part of the study.

Figs 6 and 7 do not include error bars and no statistical analysis of these data seems to have been performed. It should be clearly indicated whether there was no statistical difference among groups at any time point studied.

Each point is a single fetus. The n values represent the number of fetuses in each treatment group. The legend has been rewritten to explain this more clearly.

P20, left column, top. The statement: "AFP levels were higher in all treated dams compared to an un-treated control." should be revised, as Fig. 8 shows an AFP increase only in chronically treated dams.

This has been revised to state that there was an increase in dams' AFP only in the chronically treated animals.

It is a pity that the number of dams is too small to assess the significance of this observation and that no housekeeping protein was used to normalize AFP expression. If B is a densitometry of the gel in A, where according to the western blot labelling and the legend there is only 1 control for both dam and fetus, why are there 2 samples indicated in the controls in the charts? Given the low sample numbers and variability, particularly in fetal AFP levels, expressing the data as ratio is not appropriate. We agree that it is a pity that the numbers were very small, but we were constrained by the effects of being shut out of our laboratories for several months because of the coronavirus emergency. In general the only way to obtain accurate AFP values is to measure the actual concentrations of the protein. We are very aware that Western blots are only semi-quantitative at best. We attempted to make the gels from which we took measurements as comparable as possible within each age group by using similar volumes of plasma (or diluted sample). The concentrations of plasma proteins vary between different animals and are not related to each other therefore using albumin as a reference protein would not provide more clarity. We thank the reviewer for drawing our attention to the discrepancy in control adult numbers in the western blots (A) and in the densitometry readings (B). This has been corrected.

Competing Interests: The authors have no competing interests

Reviewer Report 07 July 2020

https://doi.org/10.5256/f1000research.26605.r64439

© **2020 Bainton R.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Roland Bainton

Department of Anesthesia and Perioperative Care, University of California, San Francisco (UCSF), San Francisco, CA, USA

Overview.

The authors present a well-controlled immaculately-conceived and artfully interpreted paper on proteomic and genomic responses of the mammalian placenta to the most commonly administered drug in the world, paracetemol.

The results of this paper are quite profound and long overdue. There are few are very few studies that attempt as complete an evaluation of drug response and in an organ specific manner. This authorship team has exploited the most pertinent drug interfaces for chemoprotection of the fetus to glimpse the system of toxicologicologic protection of the developing fetus and brain. They exploit two well known pharmacologically highly regulated barrier interfaces: the placenta and BBB. As experts on barrier development they have the right expertise to measure the developmental role and robustness of these understudied barrier interfaces to the drug paracetemol. Tylenol (as known by the US brand name) is a ubiquitous pharmacologic agent used world-wide for the abrogation of pain and systemic suppression of inflammation. While deemed one of the safest medications ever invented because of its common utilization by every age group and gender, and its long standing well-described clinical toxicities suggests that it has been vetted for safety over and over. But with the right question and under the correct experimental circumstance profound novel sensitivities in the physiology of mammals can be discovered. Such is the insight of this manuscript.

Of note.

The paper is very complete. They demonstrate both acute and chronic changes to the placenta transcriptome with strong statistical relevance. Interestingly the chronic and acute genetic changes have few if any overlapping genes suggesting that long term toxicologic homeostasis may have very different effects to fetal development than single dosing. Thus, as noted by the authors, the use for the control of acute inflammatory responses may be warranted, but chronic ingestion any substance should be view with caution when the developing fetus is concerned.

Conclusion.

This paper is well conceived, clearly written and expertly interpreted. Safety profiles of drugs are in flux and whether vertebrate homeostatic metabolic responses to drug exposure, acute or chronic, is truly benign is an open question. These authors clearly demonstrate, by the discovery of soluble protein changes in dosing of paracetamol, that there is more to learn about drug toxicology, in particular at the chemoprotective interfaces of the body, in this case the placenta and BBB. While the consequences of these proteomic changes are unclear they are corroborated by profound compensations in the transcriptional profiles of the placenta. Interestingly, the placental barrier does the lion share of compensation as the BBB of pups is nearly unchanged. This is a reassuring finding for the developing brain, but leaves many unanswered questions about how the fetus may affect maternal physiology (as noted by the authors).

The implications of this study are profound and not only for the use of paracetamol. In this paper they describe a road map for the study of all drugs that could have maternal fetal interactions and provide the physiologic and genomic insights to back up their assertions. Indeed their proposed experiments in pregnant women to follow up on their findings would be very important to the management of pregnancy and to the field of maternal/fetal physiology as a whole.

I love this paper. BRAVO!

Major Issues None.

Minor Issues. None.

Typos. None.

Is the work clearly and accurately presented and does it cite the current literature? $\ensuremath{\mathsf{Yes}}$

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others? $\ensuremath{\mathsf{Yes}}$

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility? $\ensuremath{\mathsf{Yes}}$

Are the conclusions drawn adequately supported by the results? Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Blood brain barrier, genomic toxicology, biomarkers of stress and tissue injury, blood diagnostics, metabolic compensations of the CNS

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response () 03 Aug 2020

Norman Saunders, University of Melbourne, Parkville, Australia

We thank Dr Bainton for his detailed and perceptive commentary on our study of paracetamol/acetaminophen in pregnancy and in the newborn period. We particularly appreciate his view that our findings raise serious concerns about the use of this drug in pregnancy. At this stage we have only animal data that raises concerns, but we plan to follow up with human studies insofar as this is possible. We hope that our findings will give pause for thought by the regulatory authorities and doctors who regard paracetamol/acetaminophen as "safe" to be used in pregnancy and breast feeding, especially as the concept of "being safe" for any drug is a dubious one, and particularly for one that is used so frequently. We also appreciate the Reviewer's comment that our approach provides a "roadmap" for studies of the many drugs that are prescribed in pregnancy about which there is little or no evidence on entry across the placenta and into the fetal brain. We are currently undertaking studies of psychotropic and anti-epileptic drugs as well new drugs introduced for the treatment of cystic fibrosis. Of course, the best outcome would be to find that little or no drug crosses the placenta and enters into the fetal brain.

Dr Bainton's comments are very important in helping to maintain this type of *in vivo* study.

Competing Interests: The authors have no competing interests to declare

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000 Research