

# Controllable drug uptake and nongenomic response through estrogen-anchored cyclodextrin drug complex

Juan-Juan Yin,<sup>1,2</sup> Stepan P Shumyak,<sup>2</sup> Christopher Burgess,<sup>2</sup> Zhi-Wei Zhou,<sup>2</sup> Zhi-Xu He,<sup>3</sup> Xue-Ji Zhang,<sup>4</sup> Shu-Ting Pan,<sup>2,5</sup> Tian-Xin Yang,<sup>6</sup> Wei Duan,<sup>7</sup> Jia-Xuan Qiu,<sup>5</sup> Shu-Feng Zhou<sup>2</sup>

<sup>1</sup>Xiaolan People's Hospital, Southern Medical University, Zhongshan, Guangdong, People's Republic of China; <sup>2</sup>Department of Pharmaceutical Sciences, College of Pharmacy, University of South Florida, Tampa, FL, USA; <sup>3</sup>Guizhou Provincial Key Laboratory for Regenerative Medicine, Stem Cell and Tissue Engineering Research Center and Sino-US Joint Laboratory for Medical Sciences, Guizhou Medical University, Guiyang, Guizhou; <sup>4</sup>Research Center for Bioengineering and Sensing Technology, University of Science and Technology Beijing, Beijing; <sup>5</sup>Department of Oral and Maxillofacial Surgery, the First Affiliated Hospital of Nanchang University, Nanchang, Jiangxi, People's Republic of China; <sup>6</sup>Department of Internal Medicine, University of Utah and Salt Lake Veterans Affairs Medical Center, Salt Lake City, UT, USA; <sup>7</sup>School of Medicine, Deakin University, Waurn Ponds, VIC, Australia

Correspondence: Shu-Feng Zhou  
Department of Pharmaceutical Sciences,  
College of Pharmacy, University of South  
Florida, Tampa, FL 33612, USA  
Tel +1 813 974 6276  
Fax +1 813 905 9885  
Email szhou@health.usf.edu

Jia-Xuan Qiu  
Department of Oral and Maxillofacial Surgery,  
The First Affiliated Hospital of Nanchang  
University, Yongwai Main St, Nanchang 330006,  
Jiangxi, People's Republic of China  
Tel +86 791 8869 5069  
Fax +86 791 869 2745  
Email qiujiaxuan@163.com

**Abstract:** Breast cancer is a leading killer of women worldwide. Cyclodextrin-based estrogen receptor-targeting drug-delivery systems represent a promising direction in cancer therapy but have rarely been investigated. To seek new targeting therapies for membrane estrogen receptor-positive breast cancer, an estrogen-anchored cyclodextrin encapsulating a doxorubicin derivative Ada-DOX (CDE<sub>1</sub>-Ada-DOX) has been synthesized and evaluated in human breast cancer MCF-7 cells. First, we synthesized estrone-conjugated cyclodextrin (CDE<sub>1</sub>), which formed the complex CDE<sub>1</sub>-Ada-DOX via molecular recognition with the derivative adamantane-doxorubicin (Ada-DOX) ( $K_d=1,617\text{ M}^{-1}$ ). The structure of the targeting vector CDE<sub>1</sub> was fully characterized using <sup>1</sup>H- and <sup>13</sup>C-nuclear magnetic resonance, mass spectrometry, and electron microscopy. CDE<sub>1</sub>-Ada-DOX showed two-phase drug-release kinetics with much slower release than Ada-DOX. The fluorescence polarization analysis reveals that CDE<sub>1</sub>-Ada-DOX binds to recombinant human estrogen receptor  $\alpha$  fragments with a  $K_d$  of 0.027  $\mu\text{M}$ . Competition assay of the drug complex with estrogen ligands demonstrated that estrone and tamoxifen competed with CDE<sub>1</sub>-Ada-DOX for membrane estrogen receptor binding in MCF-7 cells. Intermolecular self-assembly of CDE<sub>1</sub> molecules were observed, showing tail-in-bucket and wire-like structures confirmed by transmission electronic microscopy. CDE<sub>1</sub>-Ada-DOX had an unexpected lower drug uptake (when the host-guest ratio was  $>1$ ) than non-targeting drugs in MCF-7 cells due to ensconced ligands in cyclodextrins cavities resulting from the intermolecular self-assembly. The uptake of CDE<sub>1</sub>-Ada-DOX was significantly increased when the host-guest ratio was adjusted to be less than half at the concentration of CDE<sub>1</sub> over 5  $\mu\text{M}$  due to the release of the estrone residues. CDE<sub>1</sub> elicited rapid activation of mitogen-activated protein kinases (p44/42 MAPK, Erk1/2) in minutes through phosphorylation of Thr202/Tyr204 in MCF-7 cells. These results demonstrate a targeted therapeutics delivery of CDE<sub>1</sub>-Ada-DOX to breast cancer cells in a controlled manner and that the drug vector CDE<sub>1</sub> can potentially be employed as a molecular tool to differentiate nongenomic from genomic mechanism.

**Keywords:** breast cancer, drug vector, functionalized, membrane estrogen receptor, polysaccharide, targeted drug delivery

## Introduction


The targeted drug-delivery systems formed through various types of intermolecular forces have attracted great interest because of their therapeutic potential in drug development and cancer treatment.<sup>1,2</sup> Many elegant drug-delivery systems based on such non-bonding host-guest interactions have been developed.<sup>3-6</sup> Cyclodextrins (CDs) are among the best nonimmunogenic vector candidates for self-assembly of targeted drug-delivery systems<sup>7,8</sup> due to their unique hydrophobic hollow-cavity-containing structure, excellent biocompatibility, chemical modifiability of hydroxyl groups in the primary face, and strong ability to entrap drug molecules through molecular

recognition.<sup>7-9</sup> Ligand-attached CDs facilitate drug delivery to tumor cells that often express abundant target receptors on the tumor cell membrane. For example, multivalent CD-based glycoclusters specifically target clinically relevant sugar receptors;<sup>10</sup> a recently synthesized series of drug complexes based on CDs functionalized with folic acid, arginylglycylaspartic acid (RGD tripeptide), and hyaluronic acid, in which hydrophobic therapeutics were entrapped, have been found to possess higher targeting efficiency and better drug affinity to target receptor-positive cancer cells (data from arginylglycylaspartic acid- and hyaluronic acid-conjugated CDs are unpublished).<sup>11</sup> Ligand-attached CDs represent promising multimodal drug-delivery vectors.

However, CD-based estrogen-anchored therapeutic delivery systems have never been investigated, even though estrogens are crucial in the initiation, development, and progression of human breast cancer.<sup>12-14</sup> Overexpression of estrogen receptors (ERs) is observed in approximately 70% of human breast cancers.<sup>12,15</sup> In addition to residing in

the nucleus (“nuclear ERs” [nERs], ie, ER $\alpha$  and ER $\beta$ ), ERs can also reside on the membrane and endoplasmic reticulum (so-called “membrane ERs” [mERs]).<sup>16-19</sup> Targeting the overexpressed mERs for the delivery of therapeutics represents a promising and effectual strategy for breast cancer therapy.<sup>20-22</sup> In addition, the estrogen-conjugated CD vector is expected to preserve and elicit the estrogen response and act as a new antiestrogen for additional adjuvant endocrine treatments of ER-dependent breast cancer via targeting the ER-dependent pathways besides the targeting drug-delivery modality.<sup>21</sup>

Herein, we report an mER targeting estrone (E<sub>1</sub>)-conjugated CD vector CDE<sub>1</sub> and prepared drug-delivery complex CDE<sub>1</sub>-Ada-DOX (Figure 1A) through molecular recognition of a doxorubicin derivative adamantane-doxorubicin (Ada-DOX) and CDE<sub>1</sub>. In this system, E<sub>1</sub> was first conjugated to CDs (CDE<sub>1</sub>) to vectorize the encapsulated drug Ada-DOX which comprises an adamantane (Ada) molecule covalently linked to doxorubicin.<sup>11</sup> The molecular moiety of adamantine



**Figure 1** Schematic of estrone targeting vector based on cyclodextrin.

**Notes:** (A) The chemical structure of the CDE<sub>1</sub>-Ada-DOX inclusion complex. (B) Reaction scheme for the synthesis of ligand-attached cyclodextrin vectors (CDE<sub>1</sub> and CDPg), intermediates (CDOTs, CDNH<sub>2</sub><sup>28</sup>), and mER-targeting supramolecule CDE<sub>1</sub>-Ada-DOX. (a) Dissolve  $\beta$ -CD in H<sub>2</sub>O at the presence of NaOH (3eq) followed by adding acetonitrile solution of p-TsCl (1 eq), react for 6 hours at rt. (b) (1) Neutralized by adding 2 eq of HCl, followed by recrystallization with cold acetone; (2) React in NH<sub>3</sub>·H<sub>2</sub>O for 3–7 days at 70°C. (c) (1) React in pyridine (or DMF) for 48 hours with ligand (2) excess NaBH<sub>4</sub> in methanol solution. (d) React in anhydrous CH<sub>2</sub>Cl<sub>2</sub> at presence of Et<sub>3</sub>N and N<sub>2</sub> overnight at rt. (e) Molecular recognition with CDE<sub>1</sub> or CDPg. L\* interpreted as: ligand as estrone (E<sub>1</sub>); progesterone (Pg). The CDE<sub>1</sub>-Ada-DOX inclusion complex is a supermolecule generated through hydrophobic interaction and molecular recognition.

**Abbreviations:** CDE<sub>1</sub>, estrone-conjugated cyclodextrin; CDNH<sub>2</sub>, mono-6-deoxy-6-aminob-cyclodextrin; CDOTs, mono-6-deoxy-6-(p-tolylsulfonyl)-b-cyclodextrin; CDPg, progesterone-conjugated cyclodextrin; mER, membrane estrogen receptor; Ada-Dox, adamantane-doxorubicin; rt, room temperature.

forms stable drug inclusion with CDs with high affinity and low cytotoxicity.<sup>23,24</sup> The drug uptake of CDE<sub>1</sub>-Ada-DOX and non-targeting CD-Ada-DOX have been examined in time course altered with host–guest ratios; notably, the drug release and uptake of Ada-DOX from CDE<sub>1</sub>-Ada-DOX in MCF-7 cells was in a controllable manner due to the unique intermolecular self-assembly of and the resulting stealth or release of the E<sub>1</sub> residues in CDE<sub>1</sub>. Finally, we investigated whether the synthesized “estrogen-like” CDE<sub>1</sub> could activate nonclassical rapid response through activated ER-mediated pathway in MCF-7 cells, and the potential application in differentiation of genomic and nongenomic pathways.<sup>25–27</sup>

## Methods

For chemical synthesis, characterization, drug release, TEM microscopy, drug uptake, and Western blot, see [Supplementary materials](#).

### Binding affinity of CDE<sub>1</sub> with Ada-DOX in aqueous solution determined using the fluorescence titration method

We next determined how strong the host molecule CDE<sub>1</sub> interacted with the guest molecule Ada-DOX using the fluorescence titration method as described previously.<sup>29</sup> The effect of varying CDE<sub>1</sub> concentrations on the fluorescence intensity of Ada-DOX was examined in order to determine the association constant ( $K_d$ ) between CDE<sub>1</sub> and Ada-DOX. The concentration of Ada-DOX was set at 50  $\mu$ M in the presence of CDE<sub>1</sub> at escalating concentrations of 0, 0.17, 0.26, 0.35, 0.44, 0.53, 0.62, 0.71, 0.79, and 0.88 mM, and the fluorescence intensity was monitored using a Synergy™ H4 Hybrid Microplate Reader (BioTek Inc., Winooski, VT, USA). The solvent used was dimethylformamide:H<sub>2</sub>O (50/50, v/v). The fluorescence of Ada-DOX was measured with  $\lambda_{ex}$  at 490 nm and  $\lambda_{em}$  ranging from 500 to 700 nm with an escalating step of 2 nm. The  $K_d$  value was calculated using the above-mentioned approach by nonlinear fitting for various models.

### Effects of the phosphorylation of p44/42 mitogen-activated protein kinase (Erk1/2) at Thr202/Tyr204 in MCF-7 cells

To examine whether CDE<sub>1</sub> elicited nongenomic events in MCF-7 cells, the cells were treated with CDE<sub>1</sub> or other drugs at 1  $\mu$ M for 5, 10, 15, 30, and 60 minutes to detect p44/42 mitogen-activated protein kinase (MAPK) (Erk1/2) phosphorylation at Thr202/Tyr204 using Western blotting analysis in MCF-7 cells. The experiments were repeated in triplicate.

### Binding determination of the CDE<sub>1</sub>-Ada-DOX inclusion complex to recombinant human ER $\alpha$ fragments using the fluorescence polarization method

Fluorescence polarization (also called fluorescence anisotropy) is a versatile solution-based technique that has been widely used to investigate molecular interactions (eg, ligand–receptor binding), enzymatic activity, and nucleic acid hybridization.<sup>48</sup> Quantitatively, fluorescence polarization (mP) is defined as the difference of the emission light intensity parallel and perpendicular to the excitation light plane normalized by the total fluorescence emission intensity. The binding of the CDE<sub>1</sub>-Ada-DOX inclusion complex to recombinant human ER $\alpha$  fragments consisting of amino acid residues 1–116 at the C-terminus (His tag C-terminus, Molecular Mass=12,200 Da) (catalogue number: ab153776; Abcam Plc, Cambridge, UK) was investigated using the fluorescence polarization method as described previously.<sup>31</sup> Briefly, human ER $\alpha$  fragments were reconstituted in phosphate-buffered saline to the final concentration of 0.8  $\mu$ M, and CDE<sub>1</sub>-Ada-DOX complex samples at concentrations from 0.04  $\mu$ M to 1.26  $\mu$ M were added to the protein solution. The samples were mixed well at room temperature and subject to analysis immediately. mP was measured using the Synergy H4 Hybrid Multi-Mode Microplate reader at  $\lambda_{ex}$ =485/20 nm and  $\lambda_{em}$ =620/10 nm with xenon flash as the light source. The acquisition parameters were set as follows: 200 flashes and positioning delay 100 msec. The experiment was repeated in triplicate. The mP values were plotted against an increasing concentration of CDE<sub>1</sub>-Ada-DOX and the equilibrium binding constant ( $K_d$ ) was calculated using nonlinear least squares that fit the curve data by Prism 6.03 program (GraphPad Software, Inc., La Jolla, CA, USA).

## Results and discussion

In this study, E<sub>1</sub> was successfully conjugated with  $\beta$ -CD to generate CDE<sub>1</sub> as a new drug vector to target mERs of breast cancer cells. At the same time, progesterone, having a similar structure to E<sub>1</sub>, was also conjugated to CDs as the substrate extension of the conjugation reaction (Figure 1B). CDE<sub>1</sub> accommodates the hydrophobic Ada-DOX through host–guest interactions to form the CDE<sub>1</sub>-Ada-DOX supramolecule for drug delivery since the geometry and hydrophobicity of the adamantyl group allows an excellent fit into the CD’s torus inner cavity. The structures of the synthetic compounds have been confirmed using multiple spectral methods which include <sup>1</sup>H nuclear magnetic resonance, <sup>13</sup>C nuclear magnetic resonance, high-resolution

matrix-assisted laser desorption/ionization-time of flight mass spectroscopy, fluorescence spectroscopy, and circular dichroism spectroscopy (Figure 2). The 1:1 stoichiometric inclusion complex CDE<sub>1</sub>-Ada-DOX was readily prepared by the coprecipitation method through molecular recognition.


The host-guest binding affinity of CDE<sub>1</sub> with Ada-DOX ( $K_d$ ) was measured using the fluorescence titration method as described previously.<sup>29</sup> With an increasing amount of CDE<sub>1</sub> added to the dimethylformamide and H<sub>2</sub>O (1:1) solution containing Ada-DOX, the fluorescence intensity was measured at varying emission wavelengths from 500 to 700 nm with 600 nm as the maximum peak, which decreased in a concentration-dependent manner (Figure 2C). The  $K_d$  values were calculated using the fluorescence intensity versus CDE<sub>1</sub> concentration curve. This data showed that the one-phase and two-phase exponential decay models were the best fit. The calculated  $K_d$  was 0.93, 1.09, 0.77, and 0.73 mM when the emission wavelengths were 550, 574, 600, and 634 nm, respectively. The fluorescence quenching effect during the titration of CDE<sub>1</sub> to Ada-DOX indicated that host-guest binding occurs.

Moreover, we examined the releasing kinetics of Ada-DOX from the CDE<sub>1</sub>-Ada-DOX inclusion complex in phosphate-buffered saline against 50% fetal bovine serum solution using a validated fluorescence method.<sup>11</sup> The release kinetics of Ada-DOX and CDE<sub>1</sub>-Ada-DOX from the formulations clearly demonstrated that the drug released from CDE<sub>1</sub>-Ada-DOX was significantly slower (>50%) than Ada-DOX within the incubation time period (Figure 3). Detachment of Ada-DOX from the CDE<sub>1</sub>-Ada-DOX inclusion complex showed sustained releasing kinetics. By fitting the released drug data versus time, the kinetic parameters were calculated. For Ada-DOX, the  $K_{da}$  and  $K_{db}$  were 1.04 and 0.015 hours<sup>-1</sup>, with elimination half-life ( $t_{1/2\alpha}$ ) and  $t_{1/2\beta}$  of 0.67 and 45.02 hours, respectively. For CDE<sub>1</sub>-Ada-DOX, the  $K_{da}$  and  $K_{db}$  were 0.02 and 0.02 hours<sup>-1</sup>, with  $t_{1/2\alpha}$  and  $t_{1/2\beta}$  of 35.40 and 35.30 hours, respectively. The accumulative and absolute released drug for Ada-DOX and CDE<sub>1</sub>-Ada-DOX were also compared using two-way analysis of variance. At 2 hours, the cumulated drug release was 6.5% and 20.5% for CDE<sub>1</sub>-Ada-DOX and Ada-DOX, respectively ( $P>0.05$ ). At 4 hours, the values were almost unchanged (6.6% versus 23.5%,  $P<0.05$ ). At 6 hours, the values were increased to 11.1% and 28.4%, respectively ( $P<0.05$ ). At 21 hours, drug-release values were increased to 20.9% and 49.1%, respectively ( $P<0.001$ ). The measured drug released was 46.1% for CDE<sub>1</sub>-Ada-DOX and 99.0% for Ada-DOX over


75 hours of dialysis ( $P<0.001$ ). These data clearly show that the release of Ada-DOX from CDE<sub>1</sub>-Ada-DOX is sustained compared with Ada-DOX over 75 hours. The sustained release of Ada-DOX from the inclusion complex would facilitate continuous drug uptake and long-term proliferation inhibition of the cancer cells.

In addition to the sustained drug release of CDE<sub>1</sub>-Ada-DOX, it has been clearly demonstrated that anchored ligands in the drug-delivery systems providing specific drug-cell surface interactions are crucial in determining the ultimate drug internalization by cancer cells.<sup>8</sup> The cellular uptake of the mER-targeting drug complex CDE<sub>1</sub>-Ada-DOX and non-targeting compound CD-Ada-DOX was investigated to check for the targeting efficiency; flow cytometric analysis was performed in MCF-7 cells by taking advantage of intrinsic fluorescence emission from Ada-DOX. MCF-7 cells were treated with CDE<sub>1</sub>-Ada-DOX and non-targeting drug CD-Ada-DOX at different host-guest molecular molar ratios and drug concentrations in time course (Figure 4). The control (MCF-7 cells with no drug treated) and CDE<sub>1</sub> itself did not elicit obvious fluorescence and showed very low levels of autofluorescence. The addition of CDE<sub>1</sub> at 1, 3, or 5  $\mu$ M to the MCF-7 cells slightly increased the fluorescence. Interestingly, excess CDE<sub>1</sub> quenched the fluorescence intensity of Ada-DOX at 1  $\mu$ M when formulated in ratios of 1:1, 1:3, and 1:5 (Ada-DOX:CDE<sub>1</sub>) with 12.5% and 27.2% decrease for 1:3 and 1:5, respectively, in comparison to 1:1 CDE<sub>1</sub>-Ada-DOX complex (Figure 4A). CDE<sub>1</sub>-Ada-DOX had an unexpected lower cellular uptake than non-targeting CD-Ada-DOX complex in MCF-7 cells at 1  $\mu$ M with a 1:1 host-guest molar ratio (Figure 4B), while the uptake of CDE<sub>1</sub>-Ada-DOX was enhanced when the host-guest molecule molar ratio and total concentration were altered. The cells were exposed to different CD or CDE<sub>1</sub> inclusion complexes for 2, 4, and 6 hours, respectively, with host and guest molar ratio set at 1:1. The drug uptake in MCF-7 cells treated with CD-Ada-DOX with host-guest ratio of 1:1 (CDE<sub>1</sub> concentration = 1  $\mu$ M) for 2, 4, and 6 hours was 3.34-, 3.55-, and 3.61-fold of that in cells treated with CDE<sub>1</sub>-Ada-DOX with host-guest ratio of 1:1 with CDE<sub>1</sub> concentration at 1  $\mu$ M, respectively ( $P<0.001$ ). The internalization efficacy of the targeting complex CDE<sub>1</sub>-Ada-DOX rebounded greatly compared with the drug without ligand attached (CD-Ada-DOX) as the host-guest molecular molar ratio increased to 1:2 with CDE<sub>1</sub> concentration of 1  $\mu$ M (Figure 4C). The drug uptake in MCF-7 cells treated with CD-Ada-DOX with host-guest ratio of 1:2 at 1  $\mu$ M of CDE<sub>1</sub> for 2, 4, and 6 hours was 1.62-, 1.36-, and 1.57-fold of that in cells treated with CDE<sub>1</sub>-Ada-DOX with host-guest ratio of





**Figure 2** Characterization of CDE<sub>1</sub>. **Notes:** (A) The <sup>1</sup>H-NMR spectrum of CDE<sub>1</sub> (800 MHz, d<sub>6</sub>-DMSO, temperature = 298 K). (B) The HR-MALDI-TOF spectrum of CDE<sub>1</sub>-Ada-DOX (Con<sub>Ada-DOX</sub> = 50 μM) in DMSO at different CDE<sub>1</sub> concentrations of 0.17, 0.26, 0.35, 0.44, 0.53, 0.62, 0.71, 0.79, and 0.88 mM, with increasing step of 0.08 mM. (C) Fluorescence emission spectra of CDE<sub>1</sub>-Ada-DOX. Samples in fluorescence measurements are excited at λ<sub>ex</sub> = 485 nm. \*10<sup>3</sup> indicates the y-axis value needs to be multiplied by 1,000. **Abbreviations:** CDE<sub>1</sub>, estrone-conjugated cyclodextrin; DMSO, dimethyl sulfoxide; HR-MALDI-TOF, high-resolution matrix-assisted laser desorption/ionization-time of flight; NMR, nuclear magnetic resonance; Ada-DOX, adamantane-doxorubicin.



**Figure 3** Drug release profile of CDE<sub>1</sub>-Ada-DOX.


**Notes:** (A) The cumulative drug-releasing profile of Ada-DOX and CDE<sub>1</sub>-Ada-DOX over 75 hours. The best fit for the released drug level over time curve is a two-phase association model and the  $K_{da}$ ,  $K_{dp}$ ,  $t_{1/2a}$ , and  $t_{1/2p}$  values were obtained. (B) The absolute drug releasing profile of Ada-DOX and CDE<sub>1</sub>-Ada-DOX over 75 hours. (C) The cumulative drug-releasing profile of Ada-DOX and CDE<sub>1</sub>-Ada-DOX over 75 hours. (D) The absolute drug-releasing profile of Ada-DOX and CDE<sub>1</sub>-Ada-DOX over 75 hours. The release of Ada-DOX was determined by a dialysis method against 50% fetal bovine serum and 50% phosphate buffered saline solution, and the released Ada-DOX was quantified using a microplate reader at  $\lambda_{ex}=490$  nm and  $\lambda_{em}=600$  nm. A calibration curve was prepared using different concentrations of free Ada-DOX. \* $P<0.05$ , \*\*\* $P<0.001$ . Data are presented as the mean  $\pm$  SD from three independent experiments.

**Abbreviations:** CDE<sub>1</sub>, estrone-conjugated cyclodextrin;  $t_{1/2}$ , elimination half-life; Ada-DOX, adamantane-doxorubicin.


1:1 (CDE<sub>1</sub> concentration = 1  $\mu$ M), respectively ( $P<0.001$ ). The drug uptake of CDE<sub>1</sub>-Ada-DOX exceeded that of CD-Ada-DOX when the concentration of the guest drug was continually raised. When the host-guest molecular ratio was altered to 1:2 with a CDE<sub>1</sub> concentration of 5  $\mu$ M, the drug uptake of MCF-7 cells treated with CDE<sub>1</sub>-Ada-DOX was higher (69.9%) than cells treated with CD-Ada-DOX ( $P<0.01$ ) (Figure 4D and E). These results suggest that the targeted CDE<sub>1</sub>-Ada-DOX inclusion complex improved the uptake of Ada-DOX in comparison to the non-targeted CD-Ada-DOX inclusion complex when CDE<sub>1</sub> and Ada-DOX were formulated in appropriate ratios and drug concentrations to release ample E<sub>1</sub> ligands from CDE<sub>1</sub>.

With higher guest drug concentrations or altered host-guest molecular molar ratios, the release of the targeting moieties from the complex CDE<sub>1</sub>-Ada-DOX have been enhanced through binding to the mERs and have consequently facilitated the drug internalization process in a controlled manner. These interesting findings led us to propose that estrogen

residues covalently bonded with the CD are stealthy under certain circumstances. In order to further substantiate the presence of stealthy ligands in CDE<sub>1</sub> and CDE<sub>1</sub>-Ada-DOX, transmission electronic microscopy (TEM) and scanning electronic microscopy (SEM) examinations were conducted. Representative TEM and SEM images of CDE<sub>1</sub> and CDE<sub>1</sub>-Ada-DOX are shown in Figure 5. The intermolecular assembly of CDE<sub>1</sub> exhibited a tail-in-bucket structure and wire-like morphology for CDE<sub>1</sub>, which shows the conjugated estrogen residing inside the CD cavity of the adjacent CDE<sub>1</sub> molecule. Figure 5A–C illustrate the long, tangled, uniform CDE<sub>1</sub> wires. The estrogen residues in CDE<sub>1</sub> act as linkers of CDE<sub>1</sub> molecules due to intermolecular recognition. Figure 5D–F show the TEM images of CDE<sub>1</sub>-Ada-DOX under the same experimental conditions as CDE<sub>1</sub>. CDE<sub>1</sub>-Ada-DOX particles showed an unorganized structure under the same preparation conditions compared with the CDE<sub>1</sub> containing no drug payload. The observation was consistent and reproducible in the TEM/SEM examinations. These findings indicate that



**Figure 4** Flow cytometric analysis of the drug uptake by MCF-7 cells after incubation with various drug formulations for 2–6 hours. **Notes:** (A) The fluorescence intensity of various samples of CDE<sub>1</sub> and CDE<sub>1</sub>-Ada-DOX with host–guest molar concentration ratio of 1:1, 3:1, and 5:1 at 1 μM of Ada-DOX in MCF-7 cells. The control is obtained by the autofluorescence from the MCF-7 cells without any drug exposure. (B–E) The drug uptake indicated as the fluorescence intensity of various samples of CDE<sub>1</sub>-Ada-DOX and the non-targeting drug complex CD-Ada-DOX with 1:1, 1:2, or 1:5 of host–guest molar concentration ratio in MCF-7 cells under CDE<sub>1</sub> concentration of 1 or 5 μM. Data are presented as the mean ± standard deviation from three independent experiments. \*\**P*<0.01; \*\*\**P*<0.001. The fluorescence is tested with E<sub>x</sub> of 490 nm and E<sub>m</sub> of 600 nm. **Abbreviations:** CD, cyclodextrin; CDE<sub>1</sub>, estrone-conjugated cyclodextrin; Ctrl, control; h, hours; Ada-DOX, adamantane-doxorubicin.



**Figure 5** The TEM/SEM characterization of CDE<sub>1</sub> and CDE<sub>1</sub>-Ada-DOX.

**Notes:** A solution of CDE<sub>1</sub> or CDE<sub>1</sub>-Ada-DOX at 1.8 mM in water:dimethylformamide (1:1, v/v) was dried in vacuum oven at 35°C overnight and then subject to TEM/SEM analysis. (A and B) The TEM image of CDE<sub>1</sub> (magnification: ×10,000 for A; ×40,000 for B) shows the wire-like morphology for CDE<sub>1</sub> due to self-assembly. The estrogen residue in CDE<sub>1</sub> acts as the linker in intermolecular recognition. (C) The SEM image of CDE<sub>1</sub> (magnification: ×40,000) shows the long, tangled, and uniform CDE<sub>1</sub> wires resulting from self-assembly. (D–F) The TEM image of CDE<sub>1</sub>-Ada-DOX (magnification: ×4,000 for D; ×10,000 for E; ×30,000 for F). (G) The structure of CDE<sub>1</sub>-Ada-DOX. (H and I) Presumable structures of CDE<sub>1</sub> which do not exist. (J) The tail-in-bucket and wire-like structure of CDE<sub>1</sub> resulting from self-assembly, which were confirmed by TEM and SEM examinations.

**Abbreviations:** CDE<sub>1</sub>, estrone-conjugated cyclodextrin; SEM, scanning electron microscope; TEM, transmission electron microscope; Ada-Dox, adamantane-doxorubicin.

the conjugated E<sub>1</sub> residues of CDE<sub>1</sub> were entrapped in the CD cavities of CDE<sub>1</sub> in the absence of a guest molecule. Furthermore, when the guest molecule such as Ada-DOX was added into CDE<sub>1</sub>, the competition between exogenous and the intermolecular recognition with the CD cavity resulted in the consequent release of a certain amount of E<sub>1</sub>

residues, in turn disrupting intramolecular self-assembly, disassembling the wire-like and tail-in-bucket structure of CDE<sub>1</sub>. Moreover, circular dichroism analysis indicated that Ada-DOX interacted with CDE<sub>1</sub> (Figure 2D) and caused conformational changes at the CD cavity binding site followed by chiral microenvironment changes for the whole



drug complex supramolecule. The circular dichroism spectra provide supportive evidence for the morphology differences of CDE<sub>1</sub> and CDE<sub>1</sub>-Ada-DOX complexes.

Herein, we consider that targeting mERs expressed on MCF-7 cells is an effective means of affecting the uptake of the drug complex into these cells since the drug complex is taken up by ER-mediated endocytosis. Our results showed that CDE<sub>1</sub> preserved the binding ability after E<sub>1</sub> conjugated to CD. More importantly, drug uptake can be augmented significantly if the estrogen molecules in the complex are released to ensure the estrogen residue tailed outside the CD cavity. The intermolecular recognition between the covalent attached estrogen residue of one CDE<sub>1</sub> and the CD cavity of another CDE<sub>1</sub> molecule results in the host-guest molar ratio-dependent difference in drug uptake since the targeting moiety has been entrapped, and in turn reduces or loses its affinity to mERs. When an appropriate guest molecule approaches the CDE<sub>1</sub>, the estrogen residues are pushed out and released. The competition between complexation of the drug and intermolecular inclusion of the estrogen in the CD cavity leads to altered drug uptake. In other words, the equilibrium of the host-guest molecules and the complex is the critical factor for controlling and optimizing the release kinetics of the targeting and drug moieties, mER binding, drug uptake efficiency, dissimilarity, and biological responses.

Furthermore, the cholesterol levels have been monitored in both cancerous cells (MCF-7 and A549 cells) and normal cells (MCF-10A and T80 cells) to investigate cholesterol depletion of lipid rafts on the cell membrane after drug exposure of CDE<sub>1</sub> and the drug complexes, since CD derivatives have been reported to be able to extract cholesterol from bilayer membrane by the CDs cavity, and modulate the activity of multiple signaling pathways.<sup>30</sup> Cholesterol and estrogens have structural similarity. It was shown that the cholesterol level was not significantly affected after CDE<sub>1</sub> treatment due to the preoccupation of the CD cavity by E<sub>1</sub> residues from the intramolecular self-assembly (Figure 6).

Therefore, the magnitude of drug binding and internalization in cancer cells could be modulated by disrupting the intramolecular self-assembly of CDE<sub>1</sub>, and changing the drug exposure levels and composition (eg, host-guest molecular molar ratio) in a controllable manner based on our findings.

To confirm the binding ability of CDE<sub>1</sub>-Ada-DOX to mERs in breast cancer cells, we performed the competition assay with CDE<sub>1</sub>-Ada-DOX using E<sub>1</sub> and a selective ER modulator, tamoxifen, as the inhibitors in MCF-7 cells. The


results are shown in Figure 7A. The flow cytometric analysis showed that E<sub>1</sub> at 5–50 μM inhibited CDE<sub>1</sub>-Ada-DOX uptake in a concentration-dependent manner in MCF-7 cells. E<sub>1</sub> at 5, 10, and 50 μM diminished the uptake of CDE<sub>1</sub>-Ada-DOX by 5.1%, 4.3%, and 8.3%, respectively, in MCF-7 cells ( $P < 0.001$ ). Furthermore, tamoxifen also concentration-dependently reduced the uptake of CDE<sub>1</sub>-Ada-DOX in MCF-7 cells. Tamoxifen at 1, 5, and 10 μM inhibited the uptake of CDE<sub>1</sub>-Ada-DOX by 7.0%, 20.5%, and 19.2%, respectively, in MCF-7 cells ( $P < 0.001$ ). The results demonstrate that both E<sub>1</sub> and tamoxifen competed with CDE<sub>1</sub>-Ada-DOX for mER binding in MCF-7 cells in a concentration-dependent manner with higher competing potency for tamoxifen compared to E<sub>1</sub>.

Binding to mERs on tumor cells by tamoxifen or E<sub>1</sub> prevented the entry of other ER ligands such as CDE<sub>1</sub>-Ada-DOX targeting complex to the binding pocket. These findings provide further evidence that estrogen residues from CDE<sub>1</sub>-Ada-DOX complex recognize and bind to mER/mERs on MCF-7 cells.

Additionally, we used a fragment of human ERα containing the ligand binding domain (recombinant human ERα fragments consisting of amino acid residues 1–116 at the C-terminus, His tag C-terminus, Mr = 12,200 Da; Abcam Plc) to examine whether CDE<sub>1</sub>-Ada-DOX could bind to it using the fluorescence polarization approach.<sup>31</sup> The polarization (mP) data over the concentration of the CDE<sub>1</sub>-Ada-DOX inclusion complex in the absence or presence of human ERα fragments at a fixed concentration of 0.08 μM are shown in Figure 7B–D. The mP values were increased when the concentration of the CDE<sub>1</sub>-Ada-DOX inclusion complex was increased without adding the ERα fragments, with a  $K_d$  of 0.018 μM.

When the CDE<sub>1</sub>-Ada-DOX inclusion complex at escalating concentrations from 0.04 to 1.26 μM was mixed with human ERα fragments, the mP values were increased with a  $K_d$  of 0.027 μM. These results demonstrate the interaction of CDE<sub>1</sub>-Ada-DOX with human ERα fragments containing the ligand-binding domain.

In addition to the targeting drug-delivery modality of the novel “estrogen-like” molecule CDE<sub>1</sub>, the cellular response triggered by CDE<sub>1</sub> in a manner different from the classical nER-mediated pathway was investigated by Western blotting assay (Figure 8). Estrogens bind to nuclear ERα and ERβ, triggering the classical pathway of estrogen-dependent action and finally eliciting remarkable genomic responses. The action of nuclear ERs includes binding lipophilic hormone molecule in cytoplasm, translocation of the ligand-ER complex to the nucleus, dimerization,




**Figure 6** Cholesterol depletion from the cell membrane of human breast cancer MCF-7 cells, human normal lung T-80 cells and human normal breast MCF-10A cells, human lung cancer A549 cells.

**Notes:** Cholesterol depletion from the cell membrane ([A] human breast cancer MCF-7 cells, [B] human normal breast MCF-10A cells, [C] human normal lung T-80 cells, [D] human lung cancer A549 cells) after the treatment with saturated CDE<sub>1</sub> solution; natural ligand E<sub>1</sub> (40 µM); positive control Methyl-CD and beta-CD; DOX at 0.5, 2.5 µM; AD at 10 and 100 µM; and the drug complex CDE<sub>1</sub>-Ada-DOX in different host-guest ratio of 1:1, 1:3, and 1:5, respectively, at the host concentration of 10 µM. Cell cultures were treated for 2 hours at 37°C to deplete cholesterol. Cells were then washed twice in serum-free media and lysed. Cholesterol levels were detected using EnzyChrom™ cholesterol assay kit (BioAssay Systems, Hayward, CA, USA). CDE 1/10 means the CDE<sub>1</sub> concentration is 1/10 of the saturated CDE<sub>1</sub> solution. \**P*<0.05, \*\**P*<0.01, \*\*\**P*<0.001. Data are presented as the mean ± SD from three independent experiments.

**Abbreviations:** AD, adamantane; beta-CD, β-cyclodextrin; CD, cyclodextrin; CDE<sub>1</sub>, estrone-conjugated cyclodextrin; Ctrl, control; DOX, doxorubicin; E<sub>1</sub>, estrone; Methyl-CD, methyl-β-cyclodextrin; NS, nonsignificant; Ada-Dox, adamantane-doxorubicin.

interaction with estrogen-specific response elements in the promoter areas of target genes, and finally initiating gene transcription.<sup>18,19,32</sup> The effects of steroid hormone action dependent on ER occur within hours or even days. On the other hand, some ligands can elicit rapid nongenomic signaling cascades in a much shorter time (from seconds to minutes) upon estrogen binding.<sup>18,19,33,34</sup> These rapid nongenomic effects of estrogens result in calcium mobilization, cyclic adenosine monophosphate stimulation, phospholipase C activation, inositol phosphate generation, and activation of membrane-associated signaling pathways, including protein kinase A, phosphatidylinositol-3 kinase, and MAPK (p44/42 MAPK, also called extracellular-signal-regulated kinase

[Erk1/2]) signaling pathways.<sup>18,19</sup> Importantly, crosstalk via second messengers between mER- and nER-initiated signaling responses can regulate transcriptional activation of multiple target genes in a coordinated manner.<sup>18,19</sup> The estrogens E<sub>1</sub> and E<sub>2</sub>, the selective ER modulator tamoxifen, and the synthetic estrogen 17α-ethinyl estradiol were used as positive control to trigger the rapid response of Erk1/2; adamantane and 1-adamantanol were used as structural fraction controls to the drug complex CDE<sub>1</sub>-Ada-DOX. It was demonstrated that CDE<sub>1</sub> resulted in rapid phosphorylation response of p44/42 MAPK (p-Erk1/2) within 30 minutes in ER-positive cancerous cells such as MCF-7 cells, A549 cells, OVCAR3 cells, and even in the normal lung T80 cells,




**Figure 7** Competition of drug uptake and binding experiment about CDE<sub>1</sub>-Ada-DOX.

**Notes:** (A) The competition for the drug uptake of CDE<sub>1</sub>-Ada-DOX in the presence of E<sub>1</sub> at 0.1–50 μM or tamoxifen at 0.1–10 μM in MCF-7 cells. Cells were incubated at 37°C with CDE<sub>1</sub>-Ada-DOX at 2 μM for 6 hours in the presence of E<sub>1</sub> or tamoxifen at different concentrations. Data are presented as the mean ± SD from three independent experiments. \*\*\*P<0.001. (B) Binding of CDE<sub>1</sub>-Ada-DOX to recombinant human ERα fragments using the fluorescence polarization method. Recombinant human ERα fragments consisting of amino acid residues 1–116 at the C-terminus of ERα were reconstituted in phosphate buffered saline to the final concentration of 0.8 μM, and the CDE<sub>1</sub>-Ada-DOX inclusion complex samples at concentrations from 0.04 to 1.26 μM were added to the protein solution. The mP values were determined using a microplate reader at λ<sub>ex</sub>=485/20 nm and λ<sub>em</sub>=620/10 nm. (C) The mP values were plotted against an increasing concentration of CDE<sub>1</sub>-Ada-DOX without adding human ERα fragments, and the K<sub>d</sub> was calculated using nonlinear least squares fitting the curve data. The mP values were increased when the concentration of the CDE<sub>1</sub>-Ada-DOX inclusion complex was increased without adding the ERα fragments, with a K<sub>d</sub> of 0.018 μM. (D) When the CDE<sub>1</sub>-Ada-DOX inclusion complex at escalating concentrations from 0.04 to 1.26 μM was mixed with human ERα fragments, the mP values were increased with a K<sub>d</sub> of 0.027 μM. Data are the mean ± SD from three independent experiments.

**Abbreviations:** CDE<sub>1</sub>, estrone-conjugated cyclodextrin; Ctrl, control; E<sub>1</sub>, estrone; ER, estrogen receptor; SD, standard deviation; Ada-Dox, adamantane-doxorubicin.

while no significant response was observed in normal breast MCF-10A cells. Cells treated with CDE<sub>1</sub> at 1 μM showed a maximum p44/p42 MAPK phosphorylation at Thr202/Tyr204 at 15 minutes drug exposure and increases by 22.2%, 25.5%, and 59.2% at 30 minutes for MCF-7, A549 and OVCAR3 cells respectively, compared with controls (P<0.05). These results show that treatment of MCF-7 cells with CDE<sub>1</sub> activated the mER-mediated signaling pathway as indicated by the significantly increased phosphorylation of p44/42 MAPK (Erk1/2) rapidly within 5–30 minutes at Thr202/Tyr204 in MCF-7 cancer cells. Estrogens such as E<sub>2</sub> induce a number of rapid signaling events in cells that express mERs such as GPR30,<sup>19</sup> except in classical pathways.<sup>18,19,32</sup>

Different stimuli including mitogens, growth factors, cytokines, virus infection, ligands for heterotrimeric G protein-coupled receptors, transforming agents, and carcinogens can activate the p44/42 MAPK (Erk1/2) pathway.<sup>35–37</sup> There is evidence that E<sub>2</sub>-induced p44/42 MAPK (Erk1/2) activation requires GPR30, and occurs via transactivation of the epidermal growth factor receptor.<sup>38,39</sup> In the Raf–MEK–MAPK/Erk pathway, receptor tyrosine kinases and G protein-coupled receptors activate Ras, which in turn activates c-Raf.<sup>35,37</sup> Activation of c-Raf involves phosphorylation at multiple residues including Ser338, Tyr341, Thr491, Ser494, Ser497, and Ser499. p21-activated protein kinase can phosphorylate c-Raf at Ser338 and the Src family phosphorylates Tyr341



**Figure 8** Western blot analysis of CDE<sub>1</sub>-Ada-DOX and its analogs on A549, MCF-7, OVCAR-3, T-80 and MCF-10A cells.

**Notes:** Western blot analysis shows the time course (0, 5, 10, 15, 30, and 60 minutes) of p44/42 MAPK (Erk1/2) phosphorylation at Thr202/Tyr204 by tamoxifen, Ada-DOX, 17 $\alpha$ -ethinyl estradiol, CDE<sub>1</sub>-Ada-DOX, adamantane, E<sub>2</sub>,  $\beta$ -cyclodextrin, E<sub>1</sub>, and CDE<sub>1</sub> at 1  $\mu$ M in MCF-7, A549, and OVCAR cancerous cells (**A**); and in T-80 and MCF-10A human normal cells (**B**). In the experiments, Western immunoblotting was performed using rabbit antibodies against p-p44/42 MAPK (Erk1/2) at Thr202/Tyr204 and  $\beta$ -actin was used as the internal control. Data are from three independent experiments. Data are presented as the mean  $\pm$  SD from three independent experiments.

**Abbreviations:** CDE<sub>1</sub>, estrone-conjugated cyclodextrin; E<sub>1</sub>, estrone; E<sub>2</sub>, estradiol; Ada-Dox, adamantane-doxorubicin.

of c-Raf.<sup>35,37</sup> Activated c-Raf activates MAPK kinase (called MKK, MEK, or MAP2K) at Ser217/221 located in the activation loop of subdomain VIII, and MEK1/2 activate p44 and p42 through phosphorylation of activation loop residues Thr202/Tyr204 and Thr185/Tyr187, respectively.<sup>37</sup> p44/42 MAPK (Erk1/2) are negatively regulated by a family of dual-specificity (Thr/Tyr) MAPK phosphatases, along with MEK inhibitors such as U0126 and PD98059.<sup>40</sup>

In this study, treatment of MCF-7 cells with the new synthetic CDE<sub>1</sub> resulted in rapid phosphorylation of p44/42 MAPK (Erk1/2) in minutes. These findings suggest that CDE<sub>1</sub> can interact with and activate mERs and might be used as a probe for studying mER-mediated nongenomic events in comparison with nER-mediated genomic responses that involve DNA binding and gene expression initiation; it also needs a much longer time. It was demonstrated that adamantane and 1-adamantanol molecules do not affect p44/42 MAPK phosphorylation in tumor cells significantly, while upregulation of p-Erk1/2 from the drug complex CDE<sub>1</sub>-Ada-DOX results from CDE<sub>1</sub> and partially from the Ada-DOX since Ada-DOX treatment also cause slight pErk1/2 over-expression.

It is well known that estrogens activate the rapid, non-classical signaling cascades via mERs, and there is crosstalk between mER-mediated nonclassical and nER-mediated classical pathways.<sup>39,41,42</sup> Deregulation of both pathways plays important roles in the pathogenesis of cancer and other diseases.<sup>43</sup> It is recognized that selective ER ligands are needed to delineate the role of extranuclear and nuclear ERs in disease development and therapeutics where ERs represent important therapeutic targets. Several estrogen conjugates, including CD-E<sub>2</sub>, E<sub>2</sub>-BSA, E<sub>2</sub>-peroxidase, and E<sub>2</sub>-dendrimers, have been reported and used to probe the nonclassical and classical cascades.<sup>44-47</sup> CD derivatives are generally considered cell-membrane impermeable. Therefore, CDE<sub>1</sub> can be employed as a molecular tool to differentiate nongenomic response from genomic response based on the findings.

## Conclusion

We synthesized the novel estrogen-anchored conjugate CDE<sub>1</sub> and the corresponding drug inclusion with the doxorubicin derivative Ada-DOX (CDE<sub>1</sub>-Ada-DOX). The structures



of these new compounds were confirmed with rigorous spectral methods. A tail-in-bucket and wire-like structure of CDE<sub>1</sub> via intermolecular self-assembly was observed by TEM and SEM examination; in contrast, CDE<sub>1</sub>-Ada-DOX exhibited unorganized structure due to disruption of self-assembly in the presence of guest molecules. The binding of K<sub>d</sub> between CDE<sub>1</sub> and Ada-DOX through hydrophobic interactions was determined to be 0.77 mM by fluorescence titration. CDE<sub>1</sub>-Ada-DOX showed sustained and two-phase exponential drug-release kinetics over 75 hours. Notably, for the mER-targeted CDE<sub>1</sub>-Ada-DOX inclusion complex, the critical factor for drug uptake efficiency in MCF-7 cells relied on the equilibrium between the host-guest and drug complex. By altering the ratios between the host and the guest molecules and the breakdown of the self-assembly nanostructure, CDE<sub>1</sub>-Ada-DOX delivered the anticancer drug into MCF-7 cells in a controlled manner. E<sub>2</sub> and tamoxifen suppressed the drug uptake in MCF-7 cells treated with CDE<sub>1</sub>-Ada-DOX through competition for mER binding. Moreover, CDE<sub>1</sub>-Ada-DOX binds to recombinant human ER $\alpha$  fragments with a K<sub>d</sub> of 0.027  $\mu$ M determined by fluorescence polarization. The treatment of MCF-7 cells with CDE<sub>1</sub>-Ada-DOX elicited rapid activation of MAPKs (p44/42 MAPK, Erk1/2) in minutes through phosphorylation of Thr202/Tyr204. These results demonstrate a targeted delivery of the DOX derivative Ada-DOX to mER-positive breast cancer cells using CDE<sub>1</sub> as the drug carrier vectors in a controlled manner. The estrogen conjugates elicit nongenomic (but not genomic) events in MCF-7 cells. CDE<sub>1</sub> can be used as a powerful probe to explore the classical and nonclassical steroid-mediated pathways that are critical in the initiation, development, and progression of certain type of cancer (breast and ovarian cancer).

## Acknowledgments

The authors are grateful for the support from the National Natural Science Foundation of China (Grant No 81372383), and the startup fund from College of Pharmacy, University of South Florida. This work has been supported in part by the Florida Center of Excellence for Drug Discovery and Innovation at the University of South Florida. The authors thank Dr S Karoly and Dr A Garces at the Lisa Muma Weitz Advanced Microscopy Core Laboratory for their support in data acquisition of flow cytometry and electronic microscopy. We also thank Dr R Sprunt at Moffitt Cancer Center for the MALDI-TOF mass spectroscopy.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Schneider HJ. Binding mechanisms in supramolecular complexes. *Angew Chem Int Ed Engl*. 2009;48(22):3924–3977.
- Kubik S. Molecular cages and capsules with functionalized inner surfaces. *Top Curr Chem*. 2012;319:1–34.
- Del Valle EMM. Cyclodextrins and their uses: a review. *Process Biochem*. 2004;39(9):1033–1046.
- Zhang X, Wang C. Supramolecular amphiphiles. *Chem Soc Rev*. 2011;40(1):94–101.
- Benito JM, Gómez-García M, Ortiz Mellet C, Baussanne I, Defaye J, García Fernández JM. Optimizing saccharide-directed molecular delivery to biological receptors: design, synthesis, and biological evaluation of glycodendrimer-cyclodextrin conjugates. *J Am Chem Soc*. 2004;126(33):10355–10363.
- Paliwal SR, Paliwal R, Pal HC, et al. Estrogen-anchored pH-sensitive liposomes as nanomodule designed for site-specific delivery of doxorubicin in breast cancer therapy. *Mol Pharm*. 2012;9(1):176–186.
- Zhang J, Ma PX. Cyclodextrin-based supramolecular systems for drug delivery: recent progress and future perspective. *Adv Drug Deliv Rev*. 2013;65(9):1215–1233.
- Davis ME, Brewster ME. Cyclodextrin-based pharmaceuticals: past, present and future. *Nat Rev Drug Discov*. 2004;3(12):1023–1035.
- Yin JJ, Zhou ZW, Zhou SF. Cyclodextrin-based targeting strategies for tumor treatment. *Drug Deliv Transl Res*. 2013;3(4):364–374.
- André S, Kaltner H, Furuie T, Nishimura SI, Gabius HJ. Persubstituted cyclodextrin-based glycoclusters as inhibitors of protein-carbohydrate recognition using purified plant and mammalian lectins and wild-type and lectin-gene-transfected tumor cells as targets. *Bioconjug Chem*. 2004;15(1):87–98.
- Yin JJ, Sharma S, Shumyak SP, et al. Synthesis and biological evaluation of novel folic acid receptor-targeted,  $\beta$ -cyclodextrin-based drug complexes for cancer treatment. *PLoS One*. 2013;8(5):e62289.
- Deblois G, Giguère V. Oestrogen-related receptors in breast cancer: control of cellular metabolism and beyond. *Nat Rev Cancer*. 2013;13(1):27–36.
- Manavathi B, Dey O, Gajulapalli VN, Bhatia RS, Bugide S, Kumar R. Derailed estrogen signaling and breast cancer: an authentic couple. *Endocr Rev*. 2013;34(1):1–32.
- Yue W, Yager JD, Wang JP, Jupe ER, Santen RJ. Estrogen receptor-dependent and independent mechanisms of breast cancer carcinogenesis. *Steroids*. 2013;78(2):161–170.
- Eroles P, Bosch A, Pérez-Fidalgo JA, Lluch A. Molecular biology in breast cancer: intrinsic subtypes and signaling pathways. *Cancer Treat Rev*. 2012;38(6):698–707.
- Liang J, Shang Y. Estrogen and cancer. *Annu Rev Physiol*. 2013;75:225–240.
- Dahlman-Wright K, Cavailles V, Fuqua SA, et al. International Union of Pharmacology. LXIV. Estrogen receptors. *Pharmacol Rev*. 2006;58(4):773–781.
- Kampa M, Pelekanou V, Notas G, Stathopoulos EN, Castanas E. The estrogen receptor: two or more molecules, multiple variants, diverse localizations, signaling and functions. Are we undergoing a paradigm-shift as regards their significance in breast cancer? *Hormones (Athens)*. 2013;12(1):69–85.
- Soltysik K, Czekaj P. Membrane estrogen receptors – is it an alternative way of estrogen action? *J Physiol Pharmacol*. 2013;64(2):129–142.
- Renoir JM, Marsaud V, Lazennec G. Estrogen receptor signaling as a target for novel breast cancer therapeutics. *Biochem Pharmacol*. 2013;85(4):449–465.
- Ali S, Buluwela L, Coombes RC. Antiestrogens and their therapeutic applications in breast cancer and other diseases. *Annu Rev Med*. 2011;62:217–232.
- Nilsson S, Koehler KF, Gustafsson JA. Development of subtype-selective oestrogen receptor-based therapeutics. *Nat Rev Drug Discov*. 2011;10(10):778–792.
- Grillaud M, Russier J, Bianco A. Polycationic adamantane-based dendrons of different generations display high cellular uptake without triggering cytotoxicity. *J Am Chem Soc*. 2014;136(2):810–819.

24. Bohm I, Isenbügel K, Ritter H, Branscheid R, Kolb U. Cyclodextrin and adamantane host-guest interactions of modified hyperbranched poly(ethylene imine) as mimetics for biological membranes. *Angew Chem Int Ed Engl*. 2011;50(34):7896–7899.
25. Silva E, Kabil A, Kortenkamp A. Cross-talk between non-genomic and genomic signalling pathways – distinct effect profiles of environmental estrogens. *Toxicol Appl Pharmacol*. 2010;245(2):160–170.
26. Marino M, Galluzzo P, Ascenzi P. Estrogen signaling multiple pathways to impact gene transcription. *Curr Genomics*. 2006;7(8):497–508.
27. Parveen S, Sahoo SK. Long circulating chitosan/PEG blended PLGA nanoparticle for tumor drug delivery. *Eur J Pharmacol*. 2011;670(2–3):372–383.
28. Tang W, Ng SC. Facile synthesis of mono-6-amino-6-deoxy-alpha-, beta-, gamma-cyclodextrin hydrochlorides for molecular recognition, chiral separation and drug delivery. *Nat Protoc*. 2008;3(4):691–697.
29. Anand R, Malanga M, Manet I, et al. Citric acid- $\gamma$ -cyclodextrin cross-linked oligomers as carriers for doxorubicin delivery. *Photochem Photobiol Sci*. 2013;12(10):1841–1854.
30. Kabouridis PS, Janzen J, Magee AL, Ley SC. Cholesterol depletion disrupts lipid rafts and modulates the activity of multiple signaling pathways in T lymphocytes. *Eur J Immunol*. 2000;30(3):954–963.
31. Parker GJ, Law TL, Lenoach FJ, Bolger RE. Development of high throughput screening assays using fluorescence polarization: nuclear receptor-ligand-binding and kinase/phosphatase assays. *J Biomol Screen*. 2000;5(2):77–88.
32. Nilsson S, Mäkelä S, Treuter E, et al. Mechanisms of estrogen action. *Physiol Rev*. 2001;81(4):1535–1565.
33. Chaudhri RA, Olivares-Navarrete R, Cuenca N, Hadadi A, Boyan BD, Schwartz Z. Membrane estrogen signaling enhances tumorigenesis and metastatic potential of breast cancer cells via estrogen receptor- $\alpha$ 36 (ER $\alpha$ 36). *J Biol Chem*. 2012;287(10):7169–7181.
34. Albanito L, Madeo A, Lappano R, et al. G protein-coupled receptor 30 (GPR30) mediates gene expression changes and growth response to 17 $\beta$ -estradiol and selective GPR30 ligand G-1 in ovarian cancer cells. *Cancer Res*. 2007;67(4):1859–1866.
35. Meloche S, Pouyssegur J. The ERK1/2 mitogen-activated protein kinase pathway as a master regulator of the G1- to S-phase transition. *Oncogene*. 2007;26(22):3227–3239.
36. Johnson GL, Lapadat R. Mitogen-activated protein kinase pathways mediated by ERK, JNK, and p38 protein kinases. *Science*. 2002;298(5600):1911–1912.
37. Dent P. Crosstalk between ERK, AKT, and cell survival. *Cancer Biol Ther*. 2014;15(3):245–246.
38. Filardo EJ, Quinn JA, Bland KI, Frackelton AR Jr. Estrogen-induced activation of Erk-1 and Erk-2 requires the G protein-coupled receptor homolog, GPR30, and occurs via *trans*-activation of the epidermal growth factor receptor through release of HB-EGF. *Mol Endocrinol*. 2000;14(10):1649–1660.
39. Banerjee S, Chambliss KL, Mineo C, Shaul PW. Recent insights into non-nuclear actions of estrogen receptor-alpha. *Steroids*. 2014;81:64–69.
40. Böttner M, Thelen P, Jarry H. Estrogen receptor-beta: tissue distribution and the still largely enigmatic physiological function. *J Steroid Biochem Mol Biol*. 2014;139:245–251.
41. Wehling M, Lösel R. Non-genomic steroid hormone effects: membrane or intracellular receptors? *J Steroid Biochem Mol Biol*. 2006;102(1–5):180–183.
42. Watson CS, Alyea RA, Jeng YJ, Kochukov MY. Nongenomic actions of low concentration estrogens and xenoestrogens on multiple tissues. *Mol Cell Endocrinol*. 2007;274(1–2):1–7.
43. Deroo BJ, Korach KS. Estrogen receptors and human disease. *J Clin Invest*. 2006;116(3):561–570.
44. Kim HY, Sohn J, Wijewickrama GT, et al. Click synthesis of estradiol-cyclodextrin conjugates as cell compartment selective estrogens. *Bioorg Med Chem*. 2010;18(2):809–821.
45. Clark S, Rainville J, Zhao X, Katzenellenbogen BS, Pfaff D, Vasudevan N. Estrogen receptor-mediated transcription involves the activation of multiple kinase pathways in neuroblastoma cells. *J Steroid Biochem Mol Biol*. 2014;139:45–53.
46. Bulayeva NN, Gametchu B, Watson CS. Quantitative measurement of estrogen-induced ERK 1 and 2 activation via multiple membrane-initiated signaling pathways. *Steroids*. 2004;69(3):181–192.
47. Adlanmerini M, Solinhac R, Abot A, et al. Mutation of the palmitoylation site of estrogen receptor- $\alpha$  in vivo reveals tissue-specific roles for membrane versus nuclear actions. *Proc Natl Acad Sci U S A*. 2014;111(2):E283–E290.
48. Lea WA, Simeonov A. Fluorescence polarization assays in small molecule screening. *Expert Opin Drug Discov*. 2011;6(1):17–32.

## International Journal of Nanomedicine

### Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-nanomedicine-journal>

Dovepress

Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.