



Cochrane
Library

Cochrane Database of Systematic Reviews

Larviciding to prevent malaria transmission (Review)

Choi L, Majambere S, Wilson AL

Choi L, Majambere S, Wilson AL.

Larviciding to prevent malaria transmission.

Cochrane Database of Systematic Reviews 2019, Issue 8. Art. No.: CD012736.

DOI: [10.1002/14651858.CD012736.pub2](https://doi.org/10.1002/14651858.CD012736.pub2).

www.cochranelibrary.com

Larviciding to prevent malaria transmission (Review)

Copyright © 2019 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	3
BACKGROUND	6
Figure 1.	7
OBJECTIVES	8
METHODS	8
RESULTS	11
Figure 2.	12
Figure 3.	14
DISCUSSION	16
AUTHORS' CONCLUSIONS	17
ACKNOWLEDGEMENTS	17
REFERENCES	19
CHARACTERISTICS OF STUDIES	26
DATA AND ANALYSES	42
Analysis 1.1. Comparison 1 Larviciding versus no larviciding, Outcome 1 Malaria incidence (cluster-randomized controlled trial (cRCT)).	44
Analysis 1.2. Comparison 1 Larviciding versus no larviciding, Outcome 2 Malaria incidence with subgrouping by extent of aquatic habitat (non-randomized study (NRS)).	44
Analysis 1.3. Comparison 1 Larviciding versus no larviciding, Outcome 3 Parasite prevalence (cRCTs).	45
Analysis 1.4. Comparison 1 Larviciding versus no larviciding, Outcome 4 Parasite prevalence (NRS).	45
Analysis 1.5. Comparison 1 Larviciding versus no larviciding, Outcome 5 Parasite prevalence with subgrouping by extent of aquatic habitat.	46
Analysis 1.6. Comparison 1 Larviciding versus no larviciding, Outcome 6 Mean haemoglobin level.	47
ADDITIONAL TABLES	47
APPENDICES	49
WHAT'S NEW	50
CONTRIBUTIONS OF AUTHORS	50
DECLARATIONS OF INTEREST	50
SOURCES OF SUPPORT	51
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	51
INDEX TERMS	51

[Intervention Review]

Larviciding to prevent malaria transmission

Leslie Choi^{1a}, Silas Majambere², Anne L Wilson^{3,4b}

¹Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK. ²Mosquito Consulting, Moss, Norway.

³Department of Biosciences, Durham University, Durham, UK. ⁴Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, UK

^aThese authors contributed equally to this work. ^bThese authors contributed equally to this work

Contact address: Leslie Choi, Department of Clinical Sciences, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, L3 5QA, UK. leslie.choi@lstmed.ac.uk.

Editorial group: Cochrane Infectious Diseases Group

Publication status and date: Unchanged, published in Issue 9, 2019.

Citation: Choi L, Majambere S, Wilson AL. Larviciding to prevent malaria transmission. *Cochrane Database of Systematic Reviews* 2019, Issue 8. Art. No.: CD012736. DOI: [10.1002/14651858.CD012736.pub2](https://doi.org/10.1002/14651858.CD012736.pub2).

Copyright © 2019 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration. This is an open access article under the terms of the [Creative Commons Attribution-Non-Commercial](https://creativecommons.org/licenses/by-nc/4.0/) Licence, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

ABSTRACT

Background

Larviciding refers to the regular application of chemical or microbial insecticides to water bodies or water containers to kill the aquatic immature forms of the mosquito (the larvae and pupae).

Objectives

To summarize research evidence evaluating whether larviciding with chemical or microbial insecticides prevents malaria transmission.

Search methods

We searched the Cochrane Infectious Diseases Group Specialized Register; the Cochrane Central Register of Controlled Trials (CENTRAL), published in the Cochrane Library; MEDLINE; Embase; CAB Abstracts; LILACS; the World Health Organization International Clinical Trials Registry Platform (WHO ICTRP); ClinicalTrials.gov; and the ISRCTN registry up to 6 June 2019.

Selection criteria

We included cluster-randomized controlled trials (cRCTs), interrupted time series (ITS), randomized cross-over studies, non-randomized cross-over studies, and controlled before-and-after studies (CBAs) that compared larviciding with no larviciding.

Data collection and analysis

We independently assessed trials for eligibility and risk of bias, and extracted data. We assessed the certainty of evidence using the GRADE approach.

Main results

Four studies (one cRCT, two CBAs, and one non-randomized cross-over design) met the inclusion criteria. All used ground application of larvicides (people hand-delivering larvicides); one evaluated chemical and three evaluated microbial agents. Studies were carried out in The Gambia, Tanzania, Kenya, and Sri Lanka. Three studies were conducted in areas where mosquito aquatic habitats were less extensive (< 1 km²), and one where habitats were more extensive (> 1 km²; a cross-over study from The Gambia).

Larviciding to prevent malaria transmission (Review)

Copyright © 2019 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

For aquatic habitats of less than 1 km², one cRCT randomized eight villages in Sri Lanka to evaluate chemical larviciding using insect growth regulator; and two CBA studies undertaken in Kenya and Tanzania evaluated microbial larvicides. In the cRCT, larviciding across all villages was associated with lower malaria incidence (rate ratio 0.24, 4649 participants, low-certainty evidence) and parasite prevalence (risk ratio (RR) 0.26, 5897 participants, low-certainty evidence) compared to no larviciding. The two CBA studies reported lower malaria prevalence during the intervention period (parasite prevalence RR 0.79, 95% confidence interval (CI) 0.71 to 0.89; 70,902 participants; low-certainty evidence). The Kenyan study also reported a reduction in the incidence of new malaria cases (RR 0.62, 95% CI 0.38 to 1.01; 720 participants; very low-certainty evidence).

For aquatic habitats of more than 1 km², the non-randomized cross-over trial using microbial larvicides did not detect an effect for malaria incidence (RR 1.58, 95% CI 0.94 to 2.65; 4226 participants), or parasite prevalence (RR 1.15, 95% CI 0.41 to 3.20; 3547 participants); both were very low-certainty evidence. The Gambia trial also reported the mean haemoglobin level, and there was no difference across the four comparisons (mean difference -0.13, 95% CI -0.40 to 0.13; 3586 participants).

We were unable to summarize or pool entomological outcomes due to unreported and missing data.

Authors' conclusions

Most controlled studies on larviciding have been performed with microbial agents. Ground larviciding for non-extensive larval habitats may have an effect on malaria transmission, and we do not know if there is an effect in large-scale aquatic habitats. We found no studies using larviciding application techniques that could cover large aquatic habitats, such as aerial spraying using aircraft.

16 September 2019

Up to date

All studies incorporated from most recent search

All published trials found in the last search (6 Jun, 2019) were included, and we did not identify any ongoing trials.

PLAIN LANGUAGE SUMMARY

Larviciding to control malaria

What was the aim of this review?

Larviciding is the regular application of microbial or chemical insecticides to water bodies or water containers. The aim of larviciding is to reduce the adult population of mosquitoes by killing the aquatic immature forms, so that fewer will develop into adults. This should reduce the number of mosquitoes that bite and infect humans with malaria.

Key messages

All four studies included in this review distributed larvicides manually. Hand larviciding of small mosquito habitats may be effective in preventing malaria. Only one study was conducted in an area where larval habitats spanned a large area and this study found no effect of larviciding.

What was studied in the review?

We searched for trials that evaluated the impact of larviciding, using a microbial agent or chemical insecticide on malaria transmission. We considered effects on both human health outcomes and on mosquito populations.

What were the main results of the review?

Evidence from three studies shows that larviciding may decrease at least one malaria disease outcome in some studies, and this was in areas where the mosquito aquatic habitats were less than 1 km² (low-certainty evidence). We do not know if larviciding in large water bodies shows an impact on malaria based on results from one study in The Gambia (very low-certainty evidence).

How up to date is the review?

We searched for relevant trials up to 6 June 2019.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Summary of findings table 1

Larviciding versus no larviciding where mosquito aquatic habitats are < 1 km²

Patient or population: people at risk of malaria

Setting: areas where mosquito aquatic habitats are < 1 km² (one RCT carried out in Sri Lanka, and two CBA studies carried out in Kenya and Tanzania (Yapabandara 2001; Fillinger 2009; Maheu-Giroux 2013a)).

Intervention: larviciding

Comparison: no larviciding

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Rate or risk with no larviciding	Rate or risk with larviciding				
Malaria incidence	23 episodes per 100 person-years	5 episodes per 100 person-years	Rate ratio 0.24	4649 person-years (1 cluster-RCT)	⊕⊕⊕⊕ Low ^{a,b} due to imprecision	Larviciding may decrease malaria incidence
Parasite prevalence	4 per 100	1 per 100	RR 0.26	5868 (1 cluster-RCT)	⊕⊕⊕⊕ Low ^c due to imprecision	Larviciding may decrease parasite prevalence
	12 per 100	9 per 100 (9 to 11)	RR 0.79 (0.71 to 0.89)	70,902 (2 controlled before-and-after studies)	⊕⊕⊕⊕ Low ^d due to non-randomized design	

*The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

Abbreviations: **CI:** confidence interval; **RCT:** randomized controlled trial; **ICC:** intracluster correlation coefficient; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded two levels for imprecision: the rate ratio and CIs reported in the study were not adjusted for clustering. Sensitivity analysis with a mean cluster population of 675 showed the most conservative estimate of an ICC of 0.1 gave a rate ratio of 0.24 (95% CI 0.05 to 1.08) whereas the least conservative estimate of an ICC of 0.01 gave a rate ratio of 0.24 (95% CI 0.14 to 0.4). This created uncertainty around the point estimate.

^bAn additional study measured incidence of new infections. As this study was not a RCT, it was not combinable. However, the study showed a large effect consistent with the results of the RCT (RR 0.62, 95% CI 0.38 to 1.01) (Fillinger 2009). On GRADE assessment, the point estimate of 0.62 was very low-certainty evidence. This was due to the study being a non-randomized controlled trial, therefore baseline GRADE assessment started at 'low'. Further downgraded one level for imprecision due to wide CIs.

^cDowngraded two levels for imprecision: the odds ratio and CIs reported in the study were not adjusted for clustering. Sensitivity analysis with a mean cluster population of 675 showed the most conservative estimate of an intracluster coefficient of 0.1 gave a RR of 0.26 (95% CI 0.03 to 2.42) whereas the least conservative estimate of an ICC of 0.01 gave an odds ratio of 0.08 (95% CI 0.02 to 0.46). The wide range in CIs generated very serious uncertainty around the point estimate.

^dNon-RCTs, so baseline GRADE assessment started at 'low', therefore no further downgrading required for risk of bias.

Summary of findings 2. Summary of findings table 2

Larviciding versus no larviciding where mosquito aquatic habitats are > 1 km²

Patient or population: people at risk of malaria

Setting: areas where the extent of mosquito aquatic habitats are > 1 km² (one non-randomized cross-over study in The Gambia (Majambere 2010)).

Intervention: larviciding

Comparison: no larviciding

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Rate or risk with no larviciding	Rate or risk with larviciding				
Malaria incidence	23 episodes per 100 child-years	36 episodes per 100 child-years (22 to 61)	RR 1.58 (0.94 to 2.65)	1793 child-years (1 non-randomized cross-over trial)	⊕⊕⊕⊕ Very low ^{a,b} due to inconsistency and imprecision	We are uncertain of the effects on malaria incidence.
Parasite prevalence	14 per 100	16 per 100 (6 to 45)	RR 1.15 (0.41 to 3.20)	3574 (1 non-randomized cross-over trial)	⊕⊕⊕⊕ Very low ^{a,b} due to inconsistency and imprecision	We are uncertain of the effects on parasite prevalence.

***The risk in the intervention group** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI). The assumed risk of the comparison group is calculated from the total number of events/total number of participants in the control arms of the trials contributing to the meta-analysis.

Abbreviations: **CI:** confidence interval; **RR:** risk ratio.

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded one level for inconsistency: both comparisons indicated an effect favouring no larviciding, but there was considerable quantitative heterogeneity ($I^2 = 77\%$).

^bDowngraded two levels for imprecision: very wide CIs.

BACKGROUND

Description of the condition

Malaria is caused by the *Plasmodium* parasite, which is transmitted by female *Anopheles* mosquitoes. There are five *Plasmodium* species that cause disease in humans; however, the most important species in terms of disease burden are *Plasmodium falciparum*, which is prevalent in sub-Saharan Africa, and *Plasmodium vivax*, which is more common in Asia and South America. There were an estimated 219 million malaria cases and 435,000 deaths worldwide due to malaria in 2017 (WHO 2018). Sub-Saharan Africa carries a disproportionately high share of the malaria burden, with 92% of cases and 93% of malaria deaths in 2017 (WHO 2018). As well as direct effects on health, malaria is a major cause of poverty and underdevelopment in many countries, due to household and health system costs, absenteeism from school or work, reduced productivity, and premature death (Chima 2008). Malaria-endemic countries are, on average, poorer by more than five-fold and have lower rates of economic growth than non-malaria endemic countries, with a mean growth of per-capita gross domestic product (GDP) of 0.4% per year versus 2.3% between 1965 and 1990 (Sachs 2002).

Vector control tools, such as long-lasting insecticidal nets (LLINs) and indoor residual spraying (IRS) of insecticides, play a major role in malaria control, alongside diagnosis and effective treatment of malaria cases, and chemoprevention in some population groups. Scale-up of vector control, diagnosis, and treatment averted 663 million clinical cases of malaria between 2000 and 2015 (Bhatt 2015). However, progress against malaria is stalling and a high burden of morbidity and mortality still remains (WHO 2017; WHO 2018). The World Health Organization (WHO) set out ambitious targets in the Global Technical Strategy to eliminate malaria in at least 35 countries by 2030 (WHO 2015a).

Description of the intervention

Larviciding refers to the regular application of microbial or chemical insecticides to water bodies or water containers to kill the

aquatic immature forms of the mosquito (the larvae and pupae) (Tusting 2013).


Malaria vectors lay their eggs in standing water and the eggs develop through a series of life stages (larvae and pupae) into adults. The type of standing water selected by ovipositing females depends on the species in question and can be natural or man-made, temporary or permanent (Bruce-Chwatt 1985). For example, *Anopheles stephensi* prefers containers such as water tanks, some species prefer brackish habitats (*Anopheles aquasalis* in Latin America), while others prefer riceland habitats (*Anopheles arabiensis*).

There are several different types of larvicide, including chemical larvicides (using conventional insecticides, such as temephos, or insect growth regulators, such as pyriproxyfen, methoprene, and diflubenzuron), microbial larvicides (such as *Bacillus thuringiensis israeliensis* (Bti) and *Bacillus sphaericus* (Bs)) and oils. Larvivorous fish have also been used as a form of malaria control. Larvicides have varying modes of action. For example surface films, such as mineral oils and alcohol-based surface products, suffocate the mosquito larvae and pupae by covering the surface of a water body. This is different from synthetic organic chemicals, such as organophosphates, which inhibit cholinesterase and affect the central nervous system of the mosquito. Insect growth regulators interfere with insect metamorphosis and prevent adult emergence from the pupal stage. Microbial larvicides function by bacterial proteins binding to the larval gut, which cause the larvae to stop eating and die (WHO 2013).

How the intervention might work

Larviciding aims to reduce malaria transmission by targeting the immature stages (larvae and pupae) of the anopheline mosquito, to reduce the number of mosquitoes that reach adulthood. By reducing adult vector populations in this way, larviciding is expected to reduce the transmission of *Plasmodium* species by anopheline mosquitoes, and reduce morbidity and mortality from malaria (Figure 1).

Figure 1. Logic model of the proposed effect of larviciding on various entomological and epidemiological outcomes. EIR: entomological inoculation rate.



Many of the principles behind vector control come from the theory of vectorial capacity developed by George Macdonald in the 1950s (Macdonald 1957). Vectorial capacity describes the total number of potentially infectious bites that would eventually arise from all the mosquitoes biting a single perfectly infectious (i.e. all mosquito bites result in infection) human on a single day. Vectorial capacity can be linked to the basic reproduction ratio of a disease which is the estimated number of secondary infections potentially transmitted by a single infected individual in a totally susceptible population (Black 1968). The basic reproduction number represents the theoretical estimate of the intensity of transmission. The George-Macdonald model shows that vectorial capacity is most sensitive to changes in adult mosquito survival, which led to the prioritization of IRS and LLINs as vector control

tools in the 1950s. However, the vectorial capacity model does not adequately consider the aquatic stages of the vector and so the potential of larviciding is likely to have been underestimated (Brady 2016). Models show that larval source management (LSM) reduces mosquito population density linearly with coverage if adult mosquitoes avoid laying eggs in treated habitats, but quadratically if eggs are laid in treated habitats and the effort is therefore wasted (Smith 2013). This would mean that if the most productive habitats are targeted, larviciding could be highly effective even without extensive coverage. Larviciding may also operate against both indoor and outdoor (e.g. *An arabiensis*) biting and resting mosquitoes, unlike LLINs and IRS. This is beneficial, since in some settings anthropophilic vectors are able to sustain transmission even with high coverage of LLINs or IRS, or both (Bayoh 2010;

Russell 2010; Lwetoijera 2014), and several studies have also shown evidence of behavioural adaptation of vectors towards early evening biting which may reduce the effectiveness of indoor interventions (Gatton 2013). Thus larviciding may be effective against 'residual malaria transmission', which is generally defined as transmission that exists despite universal coverage of LLINs or IRS to which vector populations are fully susceptible (Durnez 2013; Killeen 2014).

Why it is important to do this review

There is a need for new tools in malaria vector control if the goals set by the WHO Global Technical Strategy are to be achieved (WHO 2018). Malaria vector control currently relies largely on LLINs and IRS. Although the WHO recommends the use of LSM (including larviciding) as a supplementary control measure (WHO 2013), larviciding is not widely used by malaria control programmes. This is despite historical and contemporary successes with the use of larviciding for vector control. Programmatic application of Paris Green, an arsenic-based compound toxic to larvae, contributed to the elimination of species belonging to the *Anopheles gambiae* complex in Egypt and Brazil (Soper 1943; Shousha 1948). Larviciding is routinely practiced by mosquito control programmes in the USA and Europe (Becker 1997; Floore 2006). Larviciding has also been hugely successful against other vector-borne diseases; for example, *Bti* and temephos were used to control species of the *Simulium damnosum* complex – vectors of onchocerciasis – in Brazil and the continent of Africa as a supplement to mass drug administration (MDA) (Sékétéli 2002; Gustavsen 2011).

Larviciding has the potential to overcome several challenges currently facing malaria vector control. First, larviciding is able to target outdoor resting and biting mosquitoes that are less affected by LLINs and IRS. Second, it could be used to tackle residual foci of malaria where high coverage of LLINs and IRS is not sufficient to eliminate malaria. Last, larviciding could be used together with other interventions as part of an insecticide resistance management strategy. Insecticide resistance has been reported in all major malaria vectors and involves all classes of insecticide (but particularly pyrethroids) and may threaten the effectiveness of insecticide-based vector control (WHO 2012a). The distribution and intensity of insecticide resistance has been increasing over time. Of 80 malaria-endemic countries reporting insecticide resistance monitoring data since 2010, 68 reported resistance to at least one insecticide class and 57 reported resistance to two or more insecticide classes (WHO 2018). The WHO Global Plan for Insecticide Resistance Management recommends the use of insecticide-based and non-insecticide-based interventions targeting both immature and adult mosquitoes as an insecticide resistance management strategy (WHO 2012a). This is also aligned with Integrated Vector Management (IVM), an adaptive, evidence-based, and multi-sectorial approach to vector control, which is recommended by the WHO for more effective, sustainable, and ecologically sound vector control (WHO 2008).

A Cochrane Review of LSM for controlling malaria was published in 2013 (Tusting 2013). This contributed to WHO deliberations that led to the recommendation of LSM as a supplementary malaria vector control intervention, and a WHO operational manual on LSM (WHO 2012b; WHO 2013). Although all LSM interventions have the aim of reducing mosquito larvae, the ways they are carried out are very different and effectiveness is likely to differ. For example, habitat

modification (a permanent alteration to the environment such as drainage of aquatic habitats) is different to regular application of chemical or microbial larvicides to a water body. Due to the diversity of forms of LSM, a new assessment of larviciding alone is justified, thus splitting the original Cochrane Review on LSM (Tusting 2013).

OBJECTIVES

To summarize and appraise experimental and quasi-experimental studies evaluating the effect of larviciding with chemical or microbial insecticides on malaria transmission.

METHODS

Criteria for considering studies for this review

Types of studies

- Cluster-randomized controlled trials (cRCTs) with: the unit of randomization being a cluster, and at least two clusters per arm. As larvicides are distributed at a community level, we did not expect to find trials with individual randomization.
- Randomized and non-randomized cross-over trials with: the unit of randomization being a cluster, at least two clusters per arm, and a suitable washout period during which malaria or entomological indices have returned to baseline levels. As larvicides are distributed at a community level, we did not expect to find trials with individual randomization.
- Controlled before-and-after studies (CBAs) with: a contemporaneous control group, and at least two sites per arm.
- Interrupted time series (ITS) studies with: a clearly defined point in time when the intervention occurred, and at least three data points before and three during or after cessation of larviciding.

We excluded studies if:

- the intervention was applied for less than one year in sites with perennial (year-round) transmission (as reported by the study authors); or less than one transmission season (defined as the period from the onset of rains until one month afterwards) in sites with seasonal transmission (as reported by the study authors);
- the follow-up periods for the intervention and control periods were not identical.

Types of participants

All people living in a rural or urban malarious area that is at any level of endemicity, including both stable and unstable transmission.

We planned to include and analyse studies specific to special groups, such as refugees and soldiers, separately from other studies but none were identified.

Types of interventions

Intervention

Larviciding using chemicals (insecticides and insect growth regulators), microbial agents, or oils. We excluded plant products, because formulations have not been standardized and studies are thus not comparable. We also excluded biological larviciding using larvivorous fish, covered in a separate Cochrane Review (Walshe 2017).

Larviciding to prevent malaria transmission (Review)

Control

Not receiving larviciding interventions as described above. Any co-interventions such as LLINs, IRS, topical repellents, spatial repellents, environmental manipulation, environmental modification, MDA, and case management must have been received in both control and intervention arms.

Types of outcome measures

Studies must have reported at least one primary outcome for inclusion.

Primary outcomes

- Clinical malaria incidence: we used site-specific definitions, provided they include: demonstration of malaria parasites by blood smear or a rapid diagnostic test (RDT), or both; and clinical symptoms including fever or history of fever, detected passively or actively.
- Malaria parasitaemia incidence: measured as a count per person unit time of infections or new infections, both defined as parasitaemia confirmed by blood smear microscopy or RDT. New infections were defined as either infection in participants who were negative for parasites at an earlier survey or infection in participants who were cleared of parasites using drug treatment at an earlier survey.
- Malaria parasite prevalence: proportion of surveyed people with confirmed parasitaemia.

Secondary outcomes

Entomological

- Adult mosquito density measured by a technique previously shown to be appropriate for the vector (measured using human baits, light traps, knock-down catches, baited huts, or other methods). Adult mosquito density would most likely have been reported as bites/person/night for human landing catches and mosquitoes/traps/night for trap catches or pyrethrum spray catches.
- Sporozoite rate measured as the number of caught adult mosquitoes positive for malaria sporozoites. Sporozoites can be detected through molecular or immunological methods.
- Entomological inoculation rate (EIR): the estimated number of bites by infectious mosquitoes per person per unit time. This was measured using the human biting rate (the number of mosquitoes biting a person over a stated time period measured directly using human baits or indirectly using light traps, knock-down catches, baited huts, or other methods of biting rate determination) multiplied by the sporozoite rate.

Epidemiological

- Incidence of severe malaria: we used site-specific definitions, provided they include (a) and either (b) or (c):
 - * (a) demonstration of parasitaemia by blood smear;
 - * (b) symptoms of cerebral malaria including coma or prostration or multiple seizures, or both;
 - * (c) severe life-threatening anaemia ([WHO 2015b](#)).
- Malaria-related deaths.
- Mean haemoglobin levels (g/dL).
- Anaemia prevalence defined using WHO cut-offs ([WHO 2011](#)).
- Hospital admissions for malaria.

Adverse events

Any indicators of adverse events of the intervention, including the following.

- Non-target effects such as the larvicide killing other animals in the water body.
- Reports of poisoning in humans due to exposure to larviciding chemicals.
- Environmental impacts such as changes to the biodiversity and ecosystem due to the use of larvicides.

Search methods for identification of studies

We attempted to identify all relevant trials regardless of language or publication status (published, unpublished, in press, and in progress).

Electronic searches

We searched the following databases up to 6 June 2019, using the search terms and strategy described in [Appendix 1](#):

- Cochrane Infectious Diseases Group Specialized Register;
- Cochrane Central Register of Controlled Trials (CENTRAL) published in the Cochrane Library (Issue 6, 2019)
- MEDLINE (Pubmed, from 1966);
- Embase (OVID, from 1974);
- CAB Abstracts, from 1973 (Web of Science);
- Latin American Caribbean Health Sciences Literature (LILACS) (BIREME, from 1982).

We also searched [ClinicalTrials.gov](#), the WHO International Clinical Trials Registry Platform (ICTRP; www.who.int/trialsearch), and the International Standard Randomized Controlled Trials Number (ISRCTN) registry (www.isrctn.com/) for trials in progress, using "malaria", "mosquito", and "larvicide*" as search terms.

Searching other resources

[Tusting 2013](#) handsearched the US Armed Forces Pest Management Board Defense Pest Management Literature Retrieval System and the Tropical Diseases Bulletin using the terms: malaria or mosquito and larvicides up to the end of 2010 and incorporated the results into the Cochrane Review 'Mosquito larval source management for controlling malaria'. We had planned to update this search but decided it was unlikely any new studies that would fit the review's inclusion criteria would be found.

We contacted researchers in the field to identify unpublished data, and checked the reference lists of studies identified using electronic searches.

Data collection and analysis

Selection of studies

Two review authors (LC and AW) independently assessed the titles and abstracts of trials identified by the literature searches. We obtained the full-text articles of any potentially relevant articles. The same two review authors assessed the full-text articles of potentially relevant studies for inclusion using an eligibility form based on predetermined inclusion criteria. We resolved any disagreements by discussion and consensus, with arbitration by a third review author (SM), when necessary. We ensured that multiple

Larviciding to prevent malaria transmission (Review)

publications of the same trial were included only once. We listed studies excluded after full-text assessment, together with their reasons for exclusion, in the [Characteristics of excluded studies](#) table. We illustrated the study selection process in a PRISMA flow chart ([Moher 2009](#)).

Data extraction and management

Two review authors (LC and AW) independently extracted information from the trials using pre-piloted electronic data extraction forms. SM was a primary investigator and author of one included study. He was not involved in the screening, data extraction or risk of bias assessment, and analysis for this particular study. When differences in extracted data arose, the two review authors discussed these differences to reach consensus and involved a third review author (SM), where necessary. For missing data, we contacted the original study author(s) for clarification.

We extracted the following data.

- Trial design: type of trial; method of participant selection; adjustment for clustering (for cRCTs); sample size; method of blinding of participants and personnel.
- Participants: trial settings and population characteristics; recruitment rates; withdrawal and loss to follow-up.
- Intervention: description of intervention (active ingredient, dose, formulation, method, frequency and timing of application, buffer zone between clusters); quality control of the larvicide (e.g. WHO Pesticide Evaluation Scheme (WHOPES) approved); quality assurance of implementation of larviciding; co-interventions; description of control; duration of follow-up; passive or active case detection; coverage of larvicide (as reported by the study authors) and co-interventions (e.g. vector control, vaccines, chemoprophylaxis, diagnosis, and treatment); duration of the activity of the larvicide; compliance (with application of larvicide and co-interventions).
- Outcomes: definition of outcome; diagnostic method or surveillance method; number of events; number of participants or unit time; time point at which outcome was assessed in relation to larviciding implementation, statistical power; unit of analysis; incomplete outcomes or missing data.
- Other:
 - * primary and secondary vector(s) species; vector(s) behaviour (nature, stability, and extent (number and size) of aquatic habitats, proximity of aquatic habitats to human habitation, adult habitat, peak biting times, exophilic/endophilic, exophagic/endophagic, anthropophilic/zoophilic); method of mosquito collection(s); phenotypic insecticide resistance (based on WHO definitions if WHO cylinder assays, Centers for Disease Control and Prevention (CDC) bottle bioassays, intensity assays or synergist assays were performed while the trial was running); genotypic insecticide resistance profile (either performed during the trial or if the trial references data from previous studies done on the same local vector population within the previous five years); insecticide and larvicide resistance detected in the larvae (as reported by study authors);
 - * malaria endemicity; eco-epidemiological setting; population proximity and density; *Plasmodium* species.

For dichotomous outcomes, we extracted the number of participants experiencing each outcome and the number of

participants in each treatment group. For count data outcomes, we extracted the number of outcomes in the treatment and control groups, the total person time at risk in each group or the rate ratio, and a measure of variance (e.g. standard error). For continuous outcomes, we extracted the mean and a measure of variance (e.g. standard deviation).

For cRCTs we recorded the number of clusters randomized; number of clusters analyzed; measure of effect (such as risk ratio (RR), odds ratio (OR), rate ratio, or mean difference (MD)) with confidence intervals (CI) or standard deviations; number of participants; and the intracluster correlation coefficient (ICC) value.

For non-randomized studies (NRS), we extracted adjusted measures of intervention effects that attempt to control for confounding.

Assessment of risk of bias in included studies

Two review authors (LC and AW) independently assessed the risk of bias for each cRCT using the Cochrane 'Risk of bias' tool and the five additional criteria listed in Section 16.3.2 of the *Cochrane Handbook for Systematic Reviews of Interventions* that relate specifically to cRCTs ([Higgins 2011a](#); [Higgins 2011b](#)). For assessing the risk of bias for randomized cross-over trials, we used the Cochrane 'Risk of bias' tool also and the additional criteria listed in Section 16.4.3 of the *Cochrane Handbook for Systematic Reviews of Interventions* that relate specifically to randomized and non-randomized cross-over trials ([Higgins 2011a](#)). We planned to assess non-randomized controlled studies and ITS for risk of bias using the Cochrane Effective Practice and Organisation of Care (EPOC) 'Risk of bias' tool. We resolved any discrepancies through discussion or by consulting a third review author (SM). We judged studies at low, high, or unclear risk of bias, and used summary graphs ('Risk of bias' summary and 'Risk of bias' graph) to display results.

Measures of treatment effect

We compared intervention and control data using RRs if the outcome was dichotomous. Where effect sizes from studies were presented as an OR, we converted these to RRs following the methodology stated in Section 12.5.4.4 of the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011a](#)). We presented rate data as rate ratios. We calculated the MD for continuous measures. We used adjusted measures of effect to summarize treatment effect from NRS. We presented all results with their associated 95% CIs.

We aimed to report any accounts of possible adverse effects. We appreciated that the specified inclusion criteria were not designed to detect effects on animals in the water, people exposed to the larvicides, and the ecosystem overall, and we intended to note this in the discussion, but there were no adverse events.

Unit of analysis issues

For cRCTs, we planned to extract adjusted measures of effect where possible. If the study authors did not perform any adjustment for clustering, we planned to adjust the raw data using an ICC value. If the study did not report an ICC value, we contacted the study authors, obtained this from similar studies, or estimated the ICC. When we estimated the ICC, we performed sensitivity analyses to investigate the robustness of our analyses ([Richardson 2016](#)).

For cross-over trials, we applied the principles stated in Sections 16.4.4 and 16.4.5 of the *Cochrane Handbook for Systematic Reviews of Interventions* that relate specifically to randomized and non-randomized cross-over trials (Higgins 2011a).

Dealing with missing data

In case of missing data, we intended to apply available-case analysis and to only include data on the known results. The denominator would have been the total number of participants who had data recorded for the specific outcome. For outcomes with no missing data, we planned to perform analyses on an intention-to-treat basis. We intended to include all participants randomized to each group in the analyses and analyse participants in the group to which they were randomized.

Assessment of heterogeneity

We inspected forest plots for overlapping CIs and assessed statistical heterogeneity in each meta-analysis using the I^2 statistic and Chi^2 test. We regarded heterogeneity as moderate if I^2 statistic values were between 30% and 60%; substantial if they were between 50% and 90%; and considerable if they were between 75% and 100%. We regarded a Chi^2 test statistic with a P value ≤ 0.10 as indicative of statistically significant heterogeneity. We explored clinical and methodological heterogeneity through consideration of the trial populations, methods, and interventions, and by visualization of trial results.

Assessment of reporting biases

If there were 10 or more trials included in each meta-analysis, we intended to investigate reporting biases (such as publication bias) using funnel plots. We planned to assess funnel plot asymmetry both visually and using formal tests (Harbord 2006), and explore possible reasons for asymmetry.

Data synthesis

We analyzed data using Review Manager 5 (Review Manager 2014). We used a fixed-effect meta-analysis to combine data if heterogeneity was absent. If there was considerable heterogeneity, we combined data using a random-effects meta-analysis and reported a mean treatment effect (RRs and ORs for dichotomous outcomes and rate ratio for count data). We decided whether to use a fixed- or random-effects model based on the consideration of clinical and methodological heterogeneity between trials.

We combined data across follow-up time points for each included study.

Certainty of the evidence

We assessed the certainty of evidence using the GRADE approach (Guyatt 2011). We rated each primary epidemiological outcome (malaria incidence and prevalence) as described by Balschem 2011.

- High: we are very confident that the true effect lies close to that of the estimate of the effect.
- Moderate: we are moderately confident in the effect estimate. The true effect is likely to be close to the estimate of the effect.
- Low: our confidence in the effect estimate is limited. The true effect may be substantially different from the estimate of the effect.

- Very low: we have very little confidence in the effect estimate. The true effect is likely to be substantially different from the estimate of effect.

RCTs started as high-certainty evidence but we downgraded the certainty of the evidence if there were valid reasons within the following five categories: risk of bias, imprecision, inconsistency, indirectness, and publication bias. We upgraded the certainty of the evidence for studies where there was a large effect, a dose-response effect, and if all plausible residual confounding would reduce a demonstrated effect or would suggest a spurious effect if the was no effect (Balschem 2011). We presented the GRADE assessments in a 'Summary of findings' tables.

Subgroup analysis and investigation of heterogeneity

We initially analyzed all types of larvicide (e.g. surface films, synthetic organic chemicals, insect growth regulators, and microbial larvicides) together. If there was a sufficient number of studies then we grouped these and analyzed them separately.

We explored reasons for substantial heterogeneity using subgroup analysis. We intended to perform the following subgroup analyses.

- Seasonality of malaria:
 - * perennial, defined as year-round transmission;
 - * seasonal as reported by study authors in the manuscript or defined as 75% or more of all malaria episodes occurring in six or fewer months of the year (Roca-Feltrer 2009);
 - * epidemic, defined as a sharp rise in malaria incidence, higher than typical levels.
- Extent of aquatic habitat:
 - * container habitat;
 - * habitats smaller than 1 km² (excluding containers);
 - * habitats larger than 1 km².
- Continent:
 - * Africa;
 - * non-Africa.

We only performed a subgroup analysis based on the extent of aquatic habitat as there were insufficient studies to perform the other subgroup analyses.

Sensitivity analysis

We performed sensitivity analysis on the primary outcome to determine the effect of exclusion of trials at high risk of bias (for allocation concealment and incomplete outcome data) on the overall results. If the ICC value was estimated, we undertook sensitivity analyses to investigate the impact of varying the ICC value on meta-analysis results.


RESULTS

Description of studies

Results of the search

We identified 2510 reports using electronic searches. We removed one duplicate and screened all remaining 2509 abstracts against the review's inclusion criteria. Abstract screening resulted in 98 unique reports for full-text screening (Figure 2).

Figure 2. Study flow diagram.



Included studies

Design

Five reports detailing four separate studies met the inclusion criteria and these are described in the [Characteristics of included studies](#) table. One study was a cRCT ([Yapabandara 2001](#)), one was a non-randomized cross-over trial ([Majambere 2010](#)), and two were CBAs ([Fillinger 2009](#); [Maheu-Giroux 2013a](#)).

Transmission

Three studies were each conducted in sub-Saharan Africa with one in Kenya where *P falciparum* was present and malaria transmission was moderate ([Fillinger 2009](#)), one in Gambia where *P falciparum* was present and malaria transmission was seasonal ([Majambere 2010](#)), and one in Tanzania where *P falciparum* accounted for more than 90% of cases and malaria transmission was perennial with peaks in incidence after the two rainy seasons a year ([Maheu-Giroux 2013a](#)). The remaining study was conducted in Sri Lanka where both *P falciparum* and *P vivax* were present ([Yapabandara 2001](#)). [Yapabandara 2001](#) did not report on the level of malaria transmission.

Habitat and vectors

The four studies targeted a range of aquatic habitats and vector species and all applied larvicides by hand. In the Elahera gem-mining area situated in Matale District, Sri Lanka, [Yapabandara 2001](#) targeted shallow pits dug by gem miners that harbour *An culicifacies* and *An subpictus*. In The Gambia, investigators larvicided large flooded areas of the floodplain of the lower reaches of the Gambia River which are ideal larval habitats for *An gambiae* s.s., *An melas*, and *An arabiensis* ([Majambere 2010](#)). [Fillinger 2009](#) targeted aquatic habitats in the valley bottoms in the Kenyan highlands. These habitats harbour predominantly *An gambiae* s.s. and *An funestus* and are becoming more important as papyrus swamps are deforested to create agricultural land. [Maheu-Giroux 2013a](#) evaluated the effect of the Urban Malaria Control Programme (UMCP) in Dar es Salaam, Tanzania. Here there were numerous aquatic habitats including natural habitats (swamps, river beds, springs), agricultural habitats (rice paddies, ridge and furrow agriculture) and artificial non-agricultural habitats (drains, construction pits, etc.). Habitats harboured predominantly *An gambiae* s.l. but *An funestus* and *An coustani* were also present, along with large numbers of culicines.

Interventions

A summary of the interventions implemented is given in [Table 1](#). There are two main microbial larvicides: *Bs* and *Bti*, and these exist in two formulations – water dispersible and corn cob granule. Two studies used both water-dispersible and corn cob granule formulations of *Bti* ([Majambere 2010](#); [Maheu-Giroux 2013a](#)). [Fillinger 2009](#) used a water dispersible formulation of *Bs* for six months but switched to corn cob *Bti* after six months due to a lack of residual effect. [Maheu-Giroux 2013a](#) also used a corn cob *Bs* formulation to treat closed aquatic habitats that predominantly harboured culicines. All studies that used *Bti* or *Bs* used commercial formulations manufactured by Valent BioSciences LLC. [Yapabandara 2001](#) used the insect growth regulator pyriproxyfen as a larvicide in Sri Lanka.

The frequency of application varied across the studies that used a microbial larvicide. Two studies applied the larvicide at weekly

intervals ([Fillinger 2009](#); [Majambere 2010](#)). In the Tanzanian study, open habitats were treated every week with *Bti* whereas closed habitats were treated every three months with *Bs* ([Maheu-Giroux 2013a](#)). [Yapabandara 2001](#) applied pyriproxyfen on three occasions during the study, one in December 1994, the second between June and July 1995 and the last application at the end of November 1995. Only one study reported on the duration of the activity of the larvicide ([Fillinger 2009](#)).

In Sri Lanka, local volunteers helped field staff to access villages and locate gem pits and assisted with administering the intervention ([Yapabandara 2001](#)). In The Gambia, field applicators were recruited from communities within each zone to make use of their local knowledge of the environment ([Majambere 2010](#)). They were supervised by one field supervisor in each zone and trained for one month before larviciding. In the Kenyan study, larviciding was implemented by project staff ([Fillinger 2009](#)). The UMCP in Dar es Salaam utilized community-owned resource people, each assigned to a particular neighbourhood, to deliver the larvicides ([Maheu-Giroux 2013a](#)).

In the Sri Lankan study, the control arm received no interventions ([Yapabandara 2001](#)). Both CBA studies had two arms: the control arm received standard practice vector control (insecticide-treated nets (ITNs) in [Fillinger 2009](#) and predominantly untreated bednets in [Maheu-Giroux 2013a](#)), while the intervention arm received larviciding plus standard practice vector control. The cross-over trial had two intervention arms; standard practice vector control (ITNs), and standard practice vector control (ITNs) plus larviciding ([Majambere 2010](#)). There were two units per arm, called zones. Zones 1 and 3 had the larviciding in the first year and served as control in the second year. Zones 2 and 4 received the interventions in the reverse order.

[Fillinger 2009](#) reported an increase in ITN use from 4.8% (95% CI 3.0% to 6.6%; range in control valleys 1% to 9% and in intervention valleys 2% to 6%) at baseline to 40.8% (95% CI 36.7% to 45.0%; range in control valleys 24% to 51% and in intervention valleys 25% to 51%) during the intervention year. [Majambere 2010](#) also reported an increase in net use during the study period, from between 6.1% to 38.3% in 2006 to between 37.2% to 81.4% in 2007.

Outcomes

Two studies measured clinical outcomes in children aged six months to 10 years only ([Fillinger 2009](#); [Majambere 2010](#)). The other two studies measured outcomes in participants of all ages. Two studies measured the incidence of clinical malaria ([Yapabandara 2001](#); [Majambere 2010](#)). Four studies measured the prevalence of *Plasmodium* infection ([Yapabandara 2001](#); [Fillinger 2009](#); [Majambere 2010](#); [Maheu-Giroux 2013a](#)). [Fillinger](#) and colleagues also reported the incidence of new *Plasmodium* infections ([Fillinger 2009](#)). They used children with no parasites at the first cross-sectional survey of the season who had become infected two months later to calculate the incidence rate of new parasite infections over the two-month follow-up. One study measured mean haemoglobin concentration ([Majambere 2010](#)).

Three studies reported EIR ([Fillinger 2009](#); [Majambere 2010](#); [Maheu-Giroux 2013a](#)). Three studies presented adult mosquito density as a biting rate ([Yapabandara 2001](#); [Fillinger 2009](#); [Maheu-Giroux 2013a](#)). One study presented this outcome as totals caught in

Larviciding to prevent malaria transmission (Review)

traps (Majambere 2010). One study measured sporozoite rate (Majambere 2010).

Excluded studies

We excluded 90 full-text articles for the following reasons (see Characteristics of excluded studies table).

- No relevant outcomes (40 articles).
- Study design did not match inclusion criteria (29 articles).
- Intervention did not match inclusion criteria (eight articles).
- Full text not available (four articles).
- Duplicate but under a different journal (three articles).
- Conference abstract (two articles).

- Protocol (three articles).
- Protocol of a study not performed (one article).

We found three studies awaiting classification (see Characteristics of studies awaiting classification table). We contacted the authors of Fuseini 2017, Javadian 1974, and Zhou 2013 for additional data to determine whether the studies would meet the review's inclusion criteria but we did not receive the necessary information.

Risk of bias in included studies

Judgement of the risk of bias in the included studies is summarized in Figure 3. We listed individual risk of bias assessments in the Characteristics of included studies table.

Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Baseline characteristics	Contamination	Incorrect analysis	Baseline outcome measurements	Other bias
Fillinger 2009	-	-	+	+	-	?	+	+	+	+	
Maheu-Giroux 2013a	-	-	+	+	+	?	?	?	+	+	
Majambere 2010	-	-	?	+	-	?	-	+	?	-	+
Yapabandara 2001	?	?	+	?	?	?			-		?

Allocation

Three studies were not randomized trials, and, therefore, we judged them at high risk of selection bias (Fillinger 2009; Majambere 2010; Maheu-Giroux 2013a). We judged the cRCT to have an unclear risk of selection bias as the trial authors did not explicitly state whether they conducted random sequence generation or allocation concealment (Yapabandara 2001).

Blinding

We judged studies to have a low (Yapabandara 2001; Fillinger 2009; Maheu-Giroux 2013a), or unclear (Majambere 2010) risk of bias for performance bias and low (Fillinger 2009; Majambere 2010; Maheu-Giroux 2013a) or unclear (Yapabandara 2001) detection bias.

Incomplete outcome data

Two studies were at high risk of attrition bias. Fillinger 2009 reported absences from cross-sectional surveys and the magnitude of these absences differed by study arm. There was also over 10% loss to follow-up in study groups in Majambere 2010. Maheu-Giroux 2013a had low risk of attrition bias and Yapabandara 2001 was at unclear risk.

Selective reporting

All studies had unclear risk of reporting bias.

Other potential sources of bias

We considered the risk of additional types of bias in the studies (baseline characteristics, contamination, incorrect analysis, and baseline outcome). We judged Majambere 2010 at high risk of bias due to significant differences in baseline characteristics between the zones. For example, the prevalence of *P falciparum* infections was much higher in zone 1 (38.4%) compared to the others (range 9.5% to 16.8%).

Although not a form of bias, Yapabandara 2001 used an inappropriate analysis technique which did not adjust for the clustered nature of the data in their analysis. Unadjusted estimates from cRCTs contribute disproportionately to the pooled result in meta-analysis since they receive too much weight.

Effects of interventions

See: [Summary of findings for the main comparison Summary of findings table 1](#); [Summary of findings 2 Summary of findings table 2](#)

Primary epidemiological outcomes

Clinical malaria incidence

Cluster-randomized controlled trial

Yapabandara 2001 found a reduction of 76% in the incidence of clinical malaria when communities in four villages received larviciding compared to those in four villages that did not receive larviciding. The study authors did not adjust for clustering. Using an ICC value of 0.1 (a conservative estimate), gave wide CIs ranging from a 95% reduction to an 8% increase (Analysis 1.1). Using an ICC value of 0.01 resulted in a smaller range (rate ratio 0.24, 95% CI 0.14 to 0.40).

Non-randomized studies

Two NRS investigated the impact of larviciding on malaria incidence. Majambere 2010 measured clinical malaria incidence and Fillinger 2009 measuring malaria parasitaemia incidence. They were subgrouped by the extent of aquatic habitats due to considerable heterogeneity (Analysis 1.2; I^2 statistic = 89%). Majambere 2010 reported two time points. The first time point in 2006 showed an increase in the risk of clinical malaria in the study group receiving larviciding by 101% compared to the control arm which did not receive larviciding (95% CI 51% to 168% increase). The second time point showed no effect of larviciding on the incidence of clinical malaria (RR 1.18, 95% CI 0.79 to 1.78).

Malaria parasitaemia incidence

Non-randomized studies

Fillinger 2009 found a reduction in the risk of new infections in the study group receiving larviciding of 38% compared to the control arm which did not receive larviciding (95% CI 62% decrease to 1% increase).

Parasite prevalence

Cluster-randomized controlled trial

Yapabandara 2001 reported two separate time periods postintervention for parasite prevalence. Similar to the malaria incidence outcome measured in this study, sensitivity analysis to take account of clustering showed some uncertainty around the precision. The most conservative ICC gave a very imprecise result, a RR of 0.25 (95% CI 0.03 to 2.42), while the least conservative gave a RR of 0.08 (95% CI 0.02 to 0.46), somewhat more precise but CIs were still wide.

Non-randomized studies

All three NRSs reported the effect of larviciding on parasite prevalence (Analysis 1.5). Fillinger 2009 and Maheu-Giroux 2013a reported an adjusted OR. When converted to a RR and pooled, the effect size showed a reduction of 21% in parasite prevalence in areas receiving larviciding compared to areas not receiving larviciding (95% CI 11% to 29% reduction).

Majambere 2010 reported an unadjusted OR. We took the two northern zones as one comparison and the two southern zones as another. We also took each follow-up year as separate comparisons. This led to four separate comparisons for Majambere 2010. The pooled analysis across the three NRS taking the least conservative estimate for the ICC in Majambere 2010 gave a RR of 0.88 with CIs crossing 1 (95% CI 0.66 to 1.16). However, when we took the most conservative estimate for the ICC for Majambere 2010, the pooled RR showed a reduction in parasite prevalence associated with larviciding (RR 0.80, 95% CI 0.71 to 0.89). This was because less weighting was given in the meta-analysis for the Majambere 2010 comparisons as the ICC increased.

Due to concerns of a high risk of bias for baseline imbalance, we also conducted a sensitivity analysis excluding two comparisons from Majambere 2010 (Analysis 1.4). Excluding the northern zones of Majambere 2010 and taking an estimated ICC of 0.01, the pooled result suggested a lower 21% in parasite prevalence in the area that received larviciding compared to the area that did not receive larviciding (95% CI 11% to 29% lower).

Larviciding to prevent malaria transmission (Review)

Subgroup analyses

There was moderate heterogeneity in [Analysis 1.4](#) when the comparisons from [Majambere 2010](#) with an estimated ICC of 0.01 were pooled with the other two NRS ([Fillinger 2009](#); [Maheu-Giroux 2013a](#)) ($I^2 = 59\%$; $P = 0.003$). This was explained when we subgrouped the studies by extent of the aquatic habitat in [Analysis 1.5](#). The subgroup analysis showed there was a reduction in parasite prevalence when ground application of larvicides was conducted in areas where the extent of aquatic habitats were smaller than 1 km² (RR 0.79, 95% CI 0.71 to 0.89). The analysis did not show an effect of larviciding on parasite prevalence when larvicides were administered by ground application in areas where the extent of aquatic habitats exceeded 1 km² (RR 1.15, 95% CI 0.41 to 3.20). We did not conduct the other subgroup analyses planned due to the small number of studies identified.

Secondary outcomes

Entomological

Several studies reported EIR in the intervention and control arms (summarized in [Table 2](#)). Entomological outcomes could not be pooled due to issues with the way the figures were reported in the manuscripts. All studies reported a mean number with 95% CIs except for [Majambere 2010](#). Both [Fillinger 2009](#) and [Maheu-Giroux 2013a](#) analyzed the data by using a model to adjust for confounders, whereas [Majambere 2010](#) reported the raw data with no analysis. [Fillinger 2009](#) and [Maheu-Giroux 2013a](#) reported a statistically significant lower EIR in areas receiving larviciding compared to control areas. For the first year in [Majambere 2010](#), there was no difference in EIR in the northern zones (each EIR 0) and an increase in EIR in the southern intervention zone (EIR 5.82) compared to the southern control zone (EIR 3.13). For the second year of the study, there was barely any difference in EIR between the northern intervention zone (EIR 2.32) and the northern control zone (EIR 2.24). The southern zones for 2007 showed a large difference between arms with an EIR of 17 in the southern control zone compared to 3.91 in the southern intervention zone.

Epidemiological

The mean haemoglobin level reported by study arm by [Majambere 2010](#) was converted into an MD. The pooled result from the four comparisons showed no statistically significant difference in the mean haemoglobin level of children living in areas that received larviciding compared to areas that received no larviciding (MD – 0.13, 95% CI –0.40 to 0.13). We adjusted using an ICC of 0.01 for this analysis as the CIs were already very wide.

DISCUSSION

Summary of main results

See [Summary of findings for the main comparison](#) and [Summary of findings 2](#). We included four studies: one cRCT in Sri Lanka, and three NRSs in sub-Saharan Africa.

Primary outcomes

Malaria incidence

The cRCT reported a protective efficacy against malaria of 76% and adjusting using a conservative ICC of 0.1 gave the same effect estimate but wide CIs (95% CI 0.05 to 1.08) ([Yapabandara 2001](#)).

The pooled estimate of the two comparisons from the cross-over trial showed a statistically non-significant 58% increase (95% CI 6% decrease to 165% increase) in the incidence of malaria in children in the group that received larviciding compared to those who did not receive larviciding ([Majambere 2010](#)). However, the trial authors found that the year of study was a potential effect modifier which generated uncertainty around this pooled analysis.

[Fillinger 2009](#) reported a protective efficacy of 38% against incidence of new parasite infections in the study group receiving larviciding compared to the control arm (95% CI 62% reduction to 1% increase).

Parasite prevalence

[Yapabandara 2001](#) reported on the effect of larviciding on parasite prevalence. Utilizing the most conservative ICC value gave a statistically non-significant protective efficacy of 74% (95% CI 97% reduction to 142% increase). Pooled estimates from two NRS showed a significant protective efficacy of larviciding against parasite prevalence of 21% (95% CI 11% to 29% reduction). The extent of aquatic habitat explained the moderate heterogeneity present when all NRS studies were pooled for this outcome.

Secondary outcomes

[Majambere 2010](#) was the only study to measure mean haemoglobin level. The pooled analysis showed there was no statistically significant difference between the two study arms (MD –0.13, 95% CI –0.40 to 0.13).

For entomological outcomes, three studies reported on the impact of larviciding on EIR. Due to missing data and differences in how EIR was measured in each study, it was not possible to pool across studies. Two studies reported a statistically significant reduction in EIR when an area received larviciding compared to an area that did not receive larviciding ([Fillinger 2009](#); [Maheu-Giroux 2013a](#)). The entomological results from [Majambere 2010](#) were mixed.

Certainty of the evidence

We appraised the certainty of evidence using the GRADE approach. The GRADE assessments are presented in [Summary of findings for the main comparison](#) and [Summary of findings 2](#).

Three studies (one cRCT, two NRS) evaluated the efficacy of ground application of larvicides where aquatic habitats were less than 1 km². One cRCT provided low-certainty evidence that ground application of larvicides could have had a large impact on malaria incidence. The analysis of this study was not adjusted for clustering and so the CIs may have been misleadingly narrow, and thus we downgraded by two levels due to imprecision. The same trial also reported a large beneficial effect of ground larviciding on parasite prevalence but again this was not adjusted for clustering. We downgraded by two levels to low-certainty evidence. We judged the certainty of the evidence for the pooled estimate for the prevalence of malaria from NRS to be low.

One study evaluated the efficacy of ground application of larvicides where the extent of aquatic habitats was more than 1 km². We judged the certainty of the evidence for the pooled OR for the incidence of malaria as very low, as we downgraded by one level due to unexplained heterogeneity and by two levels due to imprecision. The level of certainty for the pooled estimate for

parasite prevalence was also very low, as we downgraded by two levels for imprecision and by one level for inconsistency.

Overall completeness and applicability of evidence

Larviciding is a context-dependent intervention which requires knowledge of malaria transmission dynamics, vector ecology, behaviour, and the extent of water bodies and population density in proposed target areas. It is currently listed as a supplementary intervention for malaria control ([WHO 2012b](#)), compared to LLINs and IRS which are potentially more broadly applicable to different settings.

All studies included in this review looked at the efficacy of ground application of larvicides. Our review provided low-certainty evidence of the efficacy of larviciding where habitats could be feasibly treated by hand, such as drainage channels, irrigation channels, ponds, and pools. The certainty of evidence was downgraded due to imprecision; however, point estimates of effect from the three studies consistently showed a high reduction in malaria incidence and prevalence.

There was very low-certainty evidence on the efficacy against malaria of ground larvicide application where aquatic habitats were extensive, such as flood plains. The larval habitats treated in the Majambere study were huge as marshland areas stretched for several kilometres along the River Gambia and some larval habitats were up to 2 km wide ([Bogh 2003](#); [Majambere 2008](#)). Not all aquatic habitats were treated in this study due to deep water which prevented access by ground staff, especially during high tides. Another reason for the lack of effect in this study could have been that mosquitoes can fly long distances in this part of The Gambia ([Bogh 2007](#)), potentially leading to spillover of mosquitoes from non-intervention areas into study zones treated with larvicide. Because there was only one study with very low-certainty evidence conducted in such a setting using ground application of larvicides, it was difficult to draw any conclusions.

LSM, primarily larviciding, is the basis of contemporary mosquito control in large parts of the USA (Mosquito Abatement Districts) and Europe (e.g. Rhine valley, Germany) ([Becker 1997](#); [Floore 2006](#)). Programmes in the USA and Europe routinely use aerial application technologies such as planes and helicopters which enable higher coverage of extensive larval habitats compared to ground larviciding, and also target cryptic aquatic habitats. There is also a vast body of historical literature on the programmatic use of LSM including larviciding. For example, LSM was the primary intervention responsible for the eradication of *An gambiae* from Brazil ([Soper 1943](#)) and in Wadi Haifa, Egypt ([Najera 2001](#)). Unfortunately, this evidence could not be included in this review due to the study designs employed and extensive literature searches only identified four studies. Many large-scale field trials were excluded from this review as they only measured entomological outcomes, while epidemiological outcomes are typically required to demonstrate the public health benefit of an intervention ([Wilson 2015](#)). Larviciding, as with other wide area vector control interventions like environmental management, does not lend itself to cRCTs since the cost of studies with sufficient numbers of large clusters would be prohibitive.

Potential biases in the review process

We identified no potential biases in the review process. SM is a trial author of one of the included studies but was not involved in the screening or data extraction of this study.

Agreements and disagreements with other studies or reviews

There is an existing Cochrane Review that included all LSM interventions ([Tusting 2013](#)). Aside from that review, we are not aware of any other systematic reviews on larviciding for malaria control.

AUTHORS' CONCLUSIONS

Implications for practice

The WHO currently recommends larviciding and other larval source management (LSM) interventions as a supplementary malaria control intervention. Unlike insecticide-treated nets (ITNs) and indoor residual spraying (IRS) which target indoor vectors, LSM could potentially target outdoor as well as indoor transmission. As a result, many programmes in the elimination phase are now considering LSM including larviciding to tackle the remaining foci of malaria transmission. This review supports the use of ground larviciding for non-extensive larval habitats. We do not know if larviciding by hand in extensive habitats, largely inaccessible on foot or where water is tidal has any effect on malaria based on the results of one study of very low-certainty evidence. Operational research could strengthen the evidence base in these particular settings, with an aim of identifying effective methods for distributing larvicides over large areas

Most countries do not have the capacity or capability to conduct larviciding. If malaria control programmes are to implement larviciding, then support will be required to assess feasibility, and implement, and monitor and evaluate the intervention.

Implications for research

The findings of this review indicate low-certainty evidence of benefit from controlled studies; however, the reality is that few, if any, studies will be conducted in the coming years. Further evidence on the effects of larviciding should be generated through monitoring and evaluation of programmatic implementation using concurrent control areas, perhaps in pragmatic stepped wedge designs.

Although not evaluated or discussed in this review, evaluation of new technologies for identifying aquatic habitats (such as high-resolution imaging) and aerial application of larvicides in malaria-endemic areas may well be relevant to further refine larviciding strategies.

ACKNOWLEDGEMENTS

The Academic Editor is Dr Jimee Hwang.

We are grateful to Vittoria Lutje, Information Specialist with the Cochrane Infectious Diseases Group (CIDG), for help with the literature search strategy.

Leslie Choi is supported by the Research, Evidence and Development Initiative (READ-It) project. Anne Wilson was

supported for this work by a subcontract from READ-It. The CIDG editorial base and READ-It are funded by UK aid from the UK government for the benefit of low- and middle-income countries (project number 300342-104). The views expressed do not necessarily reflect the official policies of the UK government.

Leslie Choi is supported by PIIVeC, the Partnership for Increasing the Impact of Vector Control. PIIVeC is funded by the Medical

Research Council of the UK (grant number MR/P027873/1) through the Global Challenges Research Fund.

This work was partly supported through a grant from the Global Malaria Programme, WHO (WHO Global Malaria Programme Agreement for Performance of Work (APW) Grant 2017 (number 709319)). The views expressed in this review have not been influenced by, or necessarily reflect, WHO policy.

REFERENCES

References to studies included in this review

Fillinger 2009 {published data only}

Fillinger U, Ndenga B, Githeko A, Lindsay SW. Integrated malaria vector control with microbial larvicides and insecticide-treated nets in western Kenya: a controlled trial. *Bulletin of the World Health Organization* 2009;**87**(9):655-65.

Maheu-Giroux 2013a {published data only}

Geissbühler Y, Kannady K, Chaki PP, Emidi B, Govella NJ, Mayagaya V, et al. Microbial larvicide application by a large-scale, community-based program reduces malaria infection prevalence in urban Dar es Salaam, Tanzania. *PLOS One* 2009;**4**(3):e5107.

* Maheu-Giroux M, Castro MC. Impact of community-based larviciding on the prevalence of malaria infection in Dar es Salaam, Tanzania. *PLOS One* 2013;**8**(8):e71638.

Majambere 2010 {published data only}

Majambere S, Pinder M, Fillinger U, Ameh D, Conway DJ, Green C, et al. Is mosquito larval source management appropriate for reducing malaria in areas of extensive flooding in The Gambia? A cross-over intervention trial. *American Journal of Tropical Medicine and Hygiene* 2010;**82**(2):176-84.

Yapabandara 2001 {published data only}

Yapabandara AM, Curtis CF, Wickramasinghe MB, Fernando WP. Control of malaria vectors with the insect growth regulator pyriproxyfen in a gem-mining area in Sri Lanka. *Acta Tropica* 2001;**80**(3):265-76.

References to studies excluded from this review

Abdalmagid 2012 {published data only}

Abdalmagid MA, Elkhalfifa SM, Ismail AB, Ismail SM, Abdalrahman AH, Jamal AE. The mosquito dunk (*Bacillus thuringiensis israeliensis*) as a control agent against mosquito larvae in Khartoum State, Sudan. *Sudanese Journal of Public Health* 2012;**7**(2):51-5.

Afrane 2016 {published data only}

Afrane YA, Mweresa NG, Wanjala CL, Gilbreath TM III, Zhou G, Lee MC, et al. Evaluation of long-lasting microbial larvicide for malaria vector control in Kenya. *Malaria Journal* 2016;**15**(1):577.

Ansari 2005 {published data only}

Ansari MA, Razdan RK, Sreehari U. Laboratory and field evaluation of Hilmilin against mosquitoes. *Journal of the American Mosquito Control Association* 2005;**21**(4):432-6.

Balaraman 1983 {published data only}

Balaraman K, Balasubramanian M, Jambulingam P. Field trial of *Bacillus thuringiensis* H-14 (VCRC B-17) against *Culex* and *Anopheles* larvae. *Indian Journal of Medical Research* 1983;**77**:38-43.

Balaraman 1987 {published data only}

Balaraman K, Gunasekaran K, Pillai PK, Manonmani AM. Field trial with different formulations of *Bacillus sphaericus* for mosquito control. *Indian Journal of Medical Research* 1987;**85**:620-5.

Bertram 1950 {published data only}

Bertram DM. Larvicidal treatments with DDT and 'Gammexane' in Upper Assam, with particular reference to their effect on *Anopheles minimus*. *Annals of Tropical Medicine & Parasitology* 1950;**44**(3):255-9.

Bhalwar 1995 {published data only}

Bhalwar R, Deshpande VR, Sandhu HS, Gokarn AG. Randomised, controlled, blinded field trial on the efficacy of biocide formulation (*Bacillus* spp.) in the control of mosquito vectors. *Medical Journal Armed Forces India* 1995;**51**(1):4-8.

Biswas 1997 {published data only}

Biswas D, Ghosh SK, Dutta RN, Mukhopadhyay AK. Field trial of bacticide on larval populations of two species of vector mosquitoes in Calcutta. *Indian Journal of Malariology* 1997;**34**(1):37.

Bond 2004 {published data only}

Bond JG, Marina CF, Williams T. The naturally derived insecticide spinosad is highly toxic to *Aedes* and *Anopheles* mosquito larvae. *Medical and Veterinary Entomology* 2004;**18**(1):50-6.

Brescia 1947 {published data only}

Brescia F. Effect of concentration of DDT in oil aerosols on toxicity to mosquito larvae. *Journal of Economic Entomology* 1947;**40**(3):441.

Castro 2002 {published data only}

Castro SD, Colombi E, Flores LN, Canales D. Biolarvicide *Bacillus sphaericus*-2362 (GRISELESF) for the control of malaria in a health area of the Republic of Honduras [Aplicación del biolarvicide *Bacillus sphaericus*-2362 (GRISELESF) para el control de la Malaria en un área de salud de la República de Honduras]. *Revista Cubana de Medicina Tropical* 2002;**54**(2):134-41.

Castro 2009 {published data only}

Castro MC, Tsuruta A, Kanamori S, Kannady K, Mkude S. Community-based environmental management for malaria control: evidence from a small-scale intervention in Dar es Salaam, Tanzania. *Malaria Journal* 2009;**8**(1):57.

Chaki 2011 {published data only}

Chaki PP, Dongus S, Fillinger U, Kelly A, Killeen GF. Community-owned resource persons for malaria vector control: enabling factors and challenges in an operational programme in Dar es Salaam, United Republic of Tanzania. *Human Resources for Health* 2011;**9**(1):21.

Chen 1985 {published data only}

Chen SF, Xiao YC, Lü JF. Open solid plate medium cultured *Bacillus thuringiensis* and its mosquitocidal effect in field application. *Journal of Parasitology & Parasitic Diseases* 1985;**3**(3):195.

Chen 1988 {published data only}

Chen SF. Comparison of insecticidal effects against mosquito larvae between *Bacillus sphaericus* strain 1593 and *Bacillus thuringiensis* H-14 B. t. israelensis. *Chinese Journal of Parasitology & Parasitic Diseases* 1988;**6**(4):312.

Claborn 2002 {published data only}

Claborn DM, Masuoka PM, Klein TA, Hooper T, Lee A, Andre RG. A cost comparison of two malaria control methods in Kyunggi Province, Republic of Korea, using remote sensing and geographic information systems. *American Journal of Tropical Medicine and Hygiene* 2002;**66**(6):680-5.

Derua 2017 {published data only}

Derua YA, Kahindi SC, Mosha FW, Kweka JE, Atieli HE, Zhou G, et al. Microbial larvicides for mosquito control: impact of long lasting formulation of *Bacillus thuringiensis* var. *israelensis* and *Bacillus sphaericus* on non target organisms in western Kenya highlands. *American Journal of Tropical Medicine and Hygiene* 2017;**97**:58.

Djènantin 2014 {published data only}

Djènantin A, Pennetier C, Zogo B, Soukou KB, Ole-Sangba M, Akogbéto M, et al. Field efficacy of Vectobac GR as a mosquito larvicide for the control of Anopheline and Culicine mosquitoes in natural habitats in Benin, West Africa. *PLOS One* 2014;**9**(2):e87934.

El Safi 1986 {published data only}

El Safi SH, Haridi AM. Field trial of the insect growth regulator, Dimilin, for control of *Anopheles pharoensis* in Gezira, Sudan. *Journal of the American Mosquito Control Association* 1986;**2**(3):374.

Farashiani 2000 {published data only}

Farashiani E, Ladonni H. Laboratory and field evaluation of triflumuron and a larvicide against Anopheline larvae on rice field in South of Iran. *Journal of Entomological Society of Iran* 2000;**19**(1/2):45-55.

Fillinger 2003 {published data only}

Fillinger U, Knols BG, Becker N. Efficacy and efficiency of new *Bacillus thuringiensis* var. *israelensis* and *Bacillus sphaericus* formulations against Afrotropical anophelines in Western Kenya. *Tropical Medicine & International Health* 2003;**8**(1):37-47.

Galardo 2013 {published data only}

Galardo AK, Zimmerman R, Galardo CD. Larval control of *Anopheles (Nyssorhynchus) darlingi* using granular formulation of *Bacillus sphaericus* in abandoned gold-miners excavation pools in the Brazilian Amazon Rainforest. *Revista da Sociedade Brasileira de Medicina Tropical* 2013;**46**(2):172-7.

Giurcă 1978 {published data only}

Giurcă I, Tâcu V, Petrescu S. Action of some chemical products – insecticides and detergents on the larvae of mosquitoes of *Anopheles labranchiae atroparvus* and *Culex pipiens autogenicus* species. *Revista de Igiene, Bacteriologie, Virusologie, Parazitologie, Epidemiologie, Pneumoftiziologie. Pneumoftiziologia* 1978;**23**(4):237-46.

Haq 2004 {published data only}

Haq S, Bhatt RM, Vaishnav KG, Yadav RS. Field evaluation of biolarvicides in Surat city, India. *Journal of Vector Borne Diseases* 2004;**41**(3/4):61.

Houten 1980 {published data only}

Houten AT, Aminah NS, Suroso T, Seregeg IG. Effects of diflubenzuron (OMS 1804) against malaria vectors *Anopheles sundaicus* breeding in lagoons in Bali, Indonesia. *WHO/VBC/80* 1980;**795**:1-7.

Imbahale 2012 {published data only}

Imbahale SS, Githeko A, Mukabana WR, Takken W. Integrated mosquito larval source management reduces larval numbers in two highland villages in western Kenya. *BMC Public Health* 2012;**12**(1):362.

Johnson 1947 {published data only}

Johnson CG, Walton WH. The effect of area dosage, solution concentration and drop size of sprayed solutions and emulsions of DDT against mosquito larvae. *Bulletin of Entomological Research* 1947;**38**(3):405-30.

Julvez 1987a {published data only}

Julvez J, Galtier J, Halidi A, Henry M, Mouchet J. Epidemiology of malaria and the antimalarial campaign in Mayotte (Comoro archipelago, Indian Ocean). Development of the situation between 1976 and 1986. Outlook. *Bulletin de la Societe de Pathologie Exotique et de ses Filiales* 1987;**90**(3 Pt 2):505-19.

Julvez 1987b {published data only}

Julvez J, Galtier J, Halidi A, Henry M, Mouchet J. Epidemiology of malaria and the antimalarial campaign in Mayotte (Comoro archipelago, Indian Ocean). Development of the situation between 1976 and 1986. Outlook. *Bulletin de la Societe de Pathologie Exotique et de ses Filiales* 1987;**80**(3 bis):505-19.

Kanda 1995 {published data only}

Kanda TO, Bunnag DA, Deesin VA, Deesin TH, Leemingsawat SO, Komalamisra NA, et al. Integration of control measures for malaria vectors in endemic areas of Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* 1995;**26**(1):154-63.

Karanja 1994 {published data only}

Karanja DM, Githeko AK, Vulule JM. Small-scale field evaluation of the monomolecular surface film 'Arosurf MSF' against *Anopheles arabiensis* Patton. *Acta Tropica* 1994;**56**(4):365-9.

Konradsen 1999 {published data only}

Konradsen F, Steele P, Perera D, Van Der Hoek W, Amerasinghe PH, Amerasinghe FP. Cost of malaria control in Sri Lanka. *Bulletin of the World Health Organization* 1999;**77**(4):301.

Larviciding to prevent malaria transmission (Review)

Kramer 2014 {published data only}

Kramer RA, Mboera LE, Senkoro K, Lesser A, Shayo EH, Paul CJ, et al. A randomized longitudinal factorial design to assess malaria vector control and disease management interventions in rural Tanzania. *International Journal of Environmental Research and Public Health* 2014;**11**(5):5317-32.

Kumar 1994 {published data only}

Kumar A, Sharma VP, Sumodan PK, Thavaselvam D, Kamat RH. Malaria control utilizing *Bacillus sphaericus* against *Anopheles stephensi* in Panaji, Goa. *Journal of the American Mosquito Control Association* 1994;**10**(4):534-9.

Kumar 1998 {published data only}

Kumar A, Sharma VP, Sumodan PK, Thavaselvam D. Field trials of biolarvicide *Bacillus thuringiensis* var. israelensis strain 164 and the larvivorous fish *Aplocheilus blocki* against *Anopheles stephensi* for malaria control in Goa, India. *Journal of the American Mosquito Control Association* 1998;**14**(4):457-62.

Kumar 2013 {published data only}

Kumar AN, Murugan K, Shobana K, Abirami D. Isolation of *Bacillus sphaericus* screening larvicidal, fecundity and longevity effects on malaria vector *Anopheles stephensi*. *Scientific Research and Essays* 2013;**8**(11):425-31.

Kusumawathie 2008 {published data only}

Kusumawathie PH, Wickremasinghe AR, Karunaweera ND, Wijeyaratne MJ. Costs and effectiveness of application of *Poecilia reticulata* (guppy) and temephos in anopheline mosquito control in river basins below the major dams of Sri Lanka. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2008;**102**(7):705-11.

Ladoni 1986 {published data only}

Ladoni H, Zaim M, Motabar M. The effect of *Bacillus thuringiensis* on the anopheline mosquitoes in Southern Iran. *Journal of Entomological Society of Iran* 1986;**8**(1/2):45-52.

Lee 1990 {published data only}

Lee HL, Salleh H. Preliminary field trial of permethrin against *Anopheles* and *Culex* larvae in Malaysia. *Mosquito-Borne Diseases Bulletin* 1990;**7**(4):127-30.

Liu 2009 {published data only}

Liu Z, Sun C, Wang W, Shen P. Efficacy of mosquito larvicides against mosquito larvae in field. *Chinese Journal of Hygienic Insecticides & Equipments* 2009;**15**(6):439-41.

Lunin 1979 {published data only}

Lunin SV. Effect of the preparation altozar having juvenile hormone activity on mosquito larvae during treatment of temporary bodies of water in the northern Baikal region USSR. *Biologicheskije Nauki (Moscow)* (8): 50-53 1979;**0**:50.

Mahdi 1967 {published data only}

Mahdi A, Taha AM, Ezz ME. Preliminary small scale field trial for evaluation of aerial spraying of Malathion LV and Abate against *A. pharoensis* in the Nile delta 1966. *Journal of the Egyptian Public Health Association* 1967;**42**(1):8-13.

Maheu-Giroux 2013b {published data only}

Maheu-Giroux M, Castro M. Do malaria vector control measures impact disease-related behaviors and knowledge? Evidence from a large-scale larviciding intervention in Tanzania. *Tropical Medicine & International Health* 2013;**18**:71.

Maheu-Giroux 2013c {published data only}

Maheu-Giroux M, Castro MC. Do malaria vector control measures impact disease-related behaviour and knowledge? Evidence from a large-scale larviciding intervention in Tanzania. *Malaria Journal* 2013;**12**(1):422.

Maheu-Giroux 2014 {published data only}

Maheu-Giroux M, Castro MC. Cost-effectiveness of larviciding for urban malaria control in Tanzania. *Malaria Journal* 2014;**13**(1):477.

Marina 2014 {published data only}

Marina CF, Bond JG, Muñoz J, Valle J, Novelo-Gutiérrez R, Williams T. Efficacy and non-target impact of spinosad, Bti and temephos larvicides for control of *Anopheles spp.* in an endemic malaria region of southern Mexico. *Parasites & Vectors* 2014;**7**(1):55.

McCann 2017a {published data only}

McCann RS, van den Berg H, Diggle PJ, van Vugt M, Terlouw DJ, Phiri KS, et al. Assessment of the effect of larval source management and house improvement on malaria transmission when added to standard malaria control strategies in southern Malawi: study protocol for a cluster-randomised controlled trial. *BMC Infectious Diseases* 2017;**17**(1):639.

McCann 2017b {published data only}

McCann RS, van den Berg H, van Vugt M, Terlouw DJ, Phiri KS, Diggle PJ, et al. Community-led implementation of integrated malaria control in southern Malawi. *American Journal of Tropical Medicine and Hygiene* 2017;**49**:421-2.

Meng 1996 {published data only}

Meng XX. Observations on the mosquito-killing effects of a mixed fermentative preparation of *Bacillus sphaericus* and *Bacillus thuringiensis*. *Chinese Journal of Parasitic Disease Control* 1996;**9**(3):204.

Minakawa 2007 {published data only}

Minakawa N, Sonye G, Futami K, Kaneko S, Mushinzimana E, Fillinger U. A large-scale field trial to evaluate the efficacy of bacillus larvicides for controlling malaria in western Kenya: study design and methods. *Tropical Medicine and Health* 2007;**35**(2):41-5.

Mossadegh 1973 {published data only}

Mossadegh A, Motabar M, Javadian E. Evaluation and investigation of the effect of oil spraying on the interruption of malaria disease. *Bulletin de la Société de Pathologie Exotique* 1973;**66**(1):92-100.

Msellemu 2016 {published data only}

Msellemu D, Namango HI, Mwakalinga VM, Ntamatungiro AJ, Mlacha Y, Mtema ZJ, et al. The epidemiology of residual *Plasmodium falciparum* malaria transmission and infection

Larviciding to prevent malaria transmission (Review)

burden in an African city with high coverage of multiple vector control measures. *Malaria Journal* 2016;**15**(1):288.

Müller 1984 {published data only}

Müller P. Characteristics of *Bacillus thuringiensis* var. israelensis and its effect on mosquito larvae (Diptera: Culicidae). *Angewandte Parasitologie* 1984;**25**(3):157-63.

Obopile 2018 {published data only}

Obopile M, Segoea G, Waniwa K, Ntebela DS, Moakofhi K, Mottleng M, et al. Did microbial larviciding contribute to a reduction in malaria cases in eastern Botswana in 2012–2013?. *Public Health Action* 2018;**8**(1):S50-4.

Ouedraogo 2017 {published data only}

Ouedraogo A, Diarra A, Kargougou D, Barry A, Ouédraogo NI, Tiono A, et al. Impact of biolarvicides on the malaria morbidity in schoolchildren living in Ouagadougou, Burkina Faso. *American Journal of Tropical Medicine and Hygiene* 2017;**49**:307.

Parvez 2003 {published data only}

Parvez SD, Al Wahaibi SS. Comparison of three larviciding options for malaria vector control. *Eastern Mediterranean Health Journal* 2003;**9**(4):627-36.

Perich 1990 {published data only}

Perich MJ, Boobar LR, Stivers JC, Rivera LA. Field evaluation of four biorational larvicide formulations against *Anopheles albimanus* in Honduras. *Medical and Veterinary Entomology* 1990;**4**(4):393-6.

Prabhu 2011 {published data only}

Prabhu K, Murugan K, Nareshkumar A, Bragadeeswaran S. Larvicidal and pupicidal activity of spinosad against the malarial vector *Anopheles stephensi*. *Asian Pacific Journal of Tropical Medicine* 2011;**4**(8):610-3.

Pridantseva 1980 {published data only}

Pridantseva EA, Popova NA, Myrzin AS, Lukiyanichikov VP. Dimilin for control of larvae of the mosquito *Anopheles maculipennis messeae* Fall. and the rice midge in the Krasnodar region. *Meditsinskaya Parazitologiya i Parazitarnye Bolezni* 1980;**49**(3):53-7.

Ranjbar 2012 {published data only}

Ranjbar M, Gorgij K, Mohammadi M, Haghdoost AA, Ansari-Moghaddam A, Nikpour F, et al. Efficacy of applying self-assessment of larviciding operation, Chabahar, Iran. *Malaria Journal* 2012;**11**(1):329.

Rettich 1973 {published data only}

Rettich F. Effect of fenitrothion on mosquito larvae under laboratory and field conditions. Conference on disinfection and disinsection, with international participation 1973:59.

Rifaat 1974 {published data only}

Rifaat MA, Khalil HM, Gad AM, Sadek S. Effect of sublethal concentrations of the insecticides DDT, Abate and Sevin applied to 3rd stage larvae of *Anopheles pharoensis* on malaria cycle in the adult mosquito. *Journal of the Egyptian Public Health Association* 1974;**49**(6):329-40.

Sharma 1983 {published data only}

Sharma SK, Kalra NL, Bhargava YS. Small scale field trials with *Bacillus thuringiensis* variety israelensis H-14 strain against larvae of anopheline and culicine mosquitoes. *Journal of Communicable Diseases* 1983;**15**(4):223.

Sharma 1989 {published data only}

Sharma VP, Sharma RC. Community based bioenvironmental control of malaria in Kheda District, Gujarat, India. *Journal of the American Mosquito Control Association* 1989;**5**(4):514-21.

Sharma 2003 {published data only}

Sharma SN, Shukla RP, Mittal PK, Adak T, Kumar A. Efficacy of a new formulation of *Bacillus thuringiensis* var israelensis (Bti) in laboratory and field conditions of Kumaun foothills of Uttarakhand, India. *Journal of Communicable Diseases* 2003;**35**(4):290-9.

Shililu 2003 {published data only}

Shililu JI, Tewolde GM, Brantly E, Githure JI, Mbogo CM, Beier JC, et al. Efficacy of *Bacillus thuringiensis* israelensis, *Bacillus sphaericus* and temephos for managing *Anopheles* larvae in Eritrea. *Journal of the American Mosquito Control Association* 2003;**19**(3):251-8.

Shimada 2007 {published data only}

Shimada M, Ichinose I, Kaneko S, Minakawa N. Profile of the Nagasaki University Kenya Research Station and Activities. *Tropical Medicine and Health* 2007;**35**:29-31.

Skovmand 1997 {published data only}

Skovmand O, Bauduin S. Efficacy of a granular formulation of *Bacillus sphaericus* against *Culex quinquefasciatus* and *Anopheles gambiae* in West African countries. *Journal of Vector Ecology* 1997;**22**(1):43-51.

Skovmand 1999 {published data only}

Skovmand O, Sanogo E. Experimental formulations of *Bacillus sphaericus* and *B. thuringiensis* israelensis against *Culex quinquefasciatus* and *Anopheles gambiae* (Diptera: Culicidae) in Burkina Faso. *Journal of Medical Entomology* 1999;**36**(1):62-7.

Skovmand 2011 {published data only}

Skovmand O, Ouedraogo TD, Sanogo E, Samuelsen H, Toé LP, Bosselmann R, et al. Cost of integrated vector control with improved sanitation and road infrastructure coupled with the use of slow-release *Bacillus sphaericus* granules in a tropical urban setting. *Journal of Medical Entomology* 2011;**48**(4):813-21.

Some 1994 {published data only}

Some ES. Effects and control of highland malaria epidemic in Uasin Gishu District, Kenya. *East African Medical Journal* 1994;**71**(1):2-8.

Song 2013 {published data only}

Song GZ, Ma MH, Qu J, Liu DH, Cao GS, Yuan QD, et al. A study on the technical measures and effect evaluation for the mosquito sustainable management in the demonstration area of Dongying city of Shandong Province. *Chinese Journal of Hygienic Insecticides & Equipments* 2013;**19**(6):498-501.

Larviciding to prevent malaria transmission (Review)

Srivastava 1996 {published data only}

Srivastava R, Tilak VW, Mukherjee S, Yadav JD. Field trial of *Bacillus thuringiensis* var *israelensis* pellet formulation in the control of mosquitoes. *Medical Journal Armed Forces India* 1996;**52**(4):233-5.

Tăcu 1977 {published data only}

Tăcu V, Giurcă I, Petrescu S. The larvicide effect of some local and foreign chemical products (insecticides and detergents) on the *Anopheles labranchiae atroparvus* and *Culex pipiens autogenicus*. *Archives Roumaines de Pathologie Experimentales et de Microbiologie* 1977;**36**(2):181.

Tchicaya 2009 {published data only}

Tchicaya ES, Koudou BG, Keiser J, Adja AM, Cissé G, Tanner M, et al. Effect of repeated application of microbial larvicides on malaria transmission in central Côte d'Ivoire. *Journal of the American Mosquito Control Association* 2009;**25**(3):382-5.

Tchicaya 2010 {published data only}

Tchicaya E, Koudou BG, Utzinger J. Effect of repeated application of microbial larvicides on malaria transmission in central Côte d'Ivoire. *Malaria Journal* 2010;**9**(S2):68.

Teng 2005 {published data only}

Teng H, Lu L, Wu Y, Fang J. Evaluation of various control agents against mosquito larvae in rice paddies in Taiwan. *Journal of Vector Ecology* 2005;**30**(1):126.

Usenbaev 2006 {published data only}

Usenbaev NT, Ezhov MN, Zvantsov AB, Annarbaev A, Zhoroev AA, Almerikov K. An outbreak of *Plasmodium vivax* malaria in Kyrgyzstan. *Meditssinskaia Parazitologiya i Parazitarnye Bolezni* 2006;**1**:17-20.

Vasuki 1992 {published data only}

Vasuki V, Rajavel AR. Influence of short time exposure to an insect growth regulator, hexaflumuron, on mortality and adult emergence of vector mosquitoes. *Memorias do Instituto Oswaldo Cruz* 1992;**87**(2):275-83.

Wang 1983 {published data only}

Wang MX. Experimental observation of urea derivatives for their disinfection effect on larva of mosquitoes. *Zhonghua Liu Xing Bing Xue za Zhi* 1983;**4**(6):354-8.

Xu 1980 {published data only}

Xu QF. Laboratory and field evaluation of the mosquito larvicide effect of *Bacillus thuringiensis* var. *israelensis* (author's translation). *Zhonghua Yu Fang Yi Xue za Zhi [Chinese Journal of Preventive Medicine]* 1980;**14**(4):193-6.

Xu 1983 {published data only}

Xu QF. Studies on the prolongation of mosquito larvicide effect of *Bacillus thuringiensis* var *israelensis*. *Zhonghua Yu Fang Yi Xue za Zhi [Chinese Journal of Preventive Medicine]* 1983;**17**(6):333-5.

Xu 2004 {published data only}

Xu Y, Liang T, He S. Study on the effect of abate granules against the mosquito larvae in rivers in Urban. *Chinese Journal of Vector Biology and Control* 2004;**15**(4):299-300.

Yapabandara 2002 {published data only}

Yapabandara AM, Curtis CF. Laboratory and field comparisons of pyriproxyfen, polystyrene beads and other larvicidal methods against malaria vectors in Sri Lanka. *Acta Tropica* 2002;**81**(3):211-23.

Yapabandara 2004 {published data only}

Yapabandara AM, Curtis CF. Control of vectors and incidence of malaria in an irrigated settlement scheme in Sri Lanka when using the insect growth regulator pyriproxyfen. *Journal of the American Mosquito Control Association* 2004;**20**(4):395-400.

Yapabandara 2005 {published data only}

Yapabandara M. Review of the efficacy and persistence of pyriproxyfen 0.5% granules on *Anopheles sp* and *Aedes sp* breeding sites in Sri Lanka.. Proceedings of the Fifth International Conference on Urban Pests. Malaysia, 2005 Jul.

Zhou 2010 {published data only}

Zhou G, Liu Q, Jiang J, Liu X, Li H, Zhang Y, et al. Effects of malaria vector mosquito control approaches in Yongcheng city, Henan province, 2008. *Zhongguo Meijie Shengwuxue ji Kongzhi Zazhi [Chinese Journal of Vector Biology and Control]* 2010;**21**(4):346-9.

Zhou 2016 {published data only}

Zhou G, Wiseman V, Atieli HE, Lee MC, Githeko AK, Yan G. The impact of long-lasting microbial larvicides in reducing malaria transmission and clinical malaria incidence: study protocol for a cluster randomized controlled trial. *Trials* 2016;**17**(1):423.

Zohdy 1982 {published data only}

Zohdy NZ, Matter MM. Effect of *Bacillus thuringiensis* var. *israelensis* on some Egyptian mosquito larvae. *Journal of the Egyptian Society of Parasitology* 1982;**12**(2):349-57.

References to studies awaiting assessment
Fuseini 2017 {published data only}

Fuseini G, Phiri WP, Nlang JA, Efiri PB, Akum A, Smith J, et al. Effectiveness of community-based larviciding program on malaria vector abundance on Bioko Island, Equatorial Guinea. *American Journal of Tropical Medicine and Hygiene* 2017;**49**:537.

Javadian 1974 {published data only}

Javadian E, Acheson MA. An evaluation of the effect of larviciding operations in rural areas near Abadan, Iran. *Iranian Journal of Public Health* 1974;**3**(1):54-61.

Zhou 2013 {published data only}

Zhou G, Afrane YA, Dixit A, Atieli HE, Lee MC, Wanjala CL, et al. Modest additive effects of integrated vector control measures on malaria prevalence and transmission in western Kenya. *Malaria Journal* 2013;**12**(1):256.

Additional references

Balshem 2011

Balshem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *Journal of Clinical Epidemiology* 2011;**54**(4):401-6.

Bayoh 2010

Bayoh NM, Mathias DK, Odiere MR, Mutuku FM, Kamau L, Gimnig JE, et al. *Anopheles gambiae*: historical population decline associated with regional distribution of insecticide-treated bed nets in western Nyanza Province, Kenya. *Malaria Journal* 2010;**9**:62.

Becker 1997

Becker N. Microbial control of mosquitoes: management of the upper Rhine mosquito population as a model programme. *Parasitology Today* 1997;**13**:485-7.

Bhatt 2015

Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature* 2015;**526**(7572):207-11.

Black 1968

Black RH. Manual of epidemiology and epidemiological services in malaria programmes, 1968. apps.who.int/iris/bitstream/10665/40401/1/924154015X.pdf (accessed 11 July 2017).

Bogh 2003

Bogh C, Clarke S, Jawara M, Thomas CJ, Lindsay SW. Localized breeding of the *Anopheles gambiae* complex (Diptera: Culicidae) along the River Gambia, West Africa. *Bulletin of Entomological Research* 2003;**93**(4):279-87.

Brady 2016

Brady OJ, Godfray HC, Tatem AJ, Gething PW, Cohen JM, McKenzie FE, et al. Vectorial capacity and vector control: reconsidering sensitivity to parameters for malaria elimination. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2016;**110**(2):107-17.

Bruce-Chwatt 1985

Bruce-Chwatt LJ. Essential Malariology. 2nd Edition. London (UK): William Heineman Medical Books Ltd, 1985.

Bøgh 2007

Bøgh C, Lindsay SW, Clarke SE, Dean A, Jawara M, Pinder M, et al. High spatial resolution mapping of malaria transmission risk in the Gambia, west Africa, using LANDSAT TM satellite imagery. *American Journal of Tropical Medicine and Hygiene* 2007;**76**(5):875-81.

Chima 2008

Chima RI, Goodman CA, Mills A. The economic impact of malaria in Africa: a critical review of the evidence. *Health Policy* 2008;**63**(1):17-36.

Durnez 2013

Durnez L, Coosemans M. Residual transmission of malaria: an old issue for new approaches. In: Manguin S editor(s). *Anopheles Mosquitoes – New Insights into Malaria Vectors*. Rijeka (Croatia): InTech, 2013.

Floore 2006

Floore T. Mosquito Larval Control Practices: Past and Present. *Journal of the American Mosquito Control Association* 2006;**22**:527-33.

Gatton 2013

Gatton ML, Chitnis N, Churcher T, Donnelly MJ, Ghani AC, Godfray HC, et al. The importance of mosquito behavioural adaptations to malaria control in Africa. *Evolution* 2013;**67**(4):1218-30.

Gustavsen 2011

Gustavsen K, Hopkins A, Sauerbrey M. Onchocerciasis in the Americas: from arrival to (near) elimination. *Parasites & Vectors* 2011;**4**:205.

Guyatt 2011

Guyatt GH, Oxman AD, Schünemann HJ, Tugwell P, Knottnerus A. GRADE guidelines: a new series of articles in the Journal of Clinical Epidemiology. *Journal of Clinical Epidemiology* 2011;**64**(4):380-2.

Harbord 2006

Harbord RM, Egger M, Sterne JA. A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Statistics in Medicine* 2006;**25**(20):3443-57.

Higgins 2011a

Higgins JP, Deeks JJ, Altman DG. Chapter 16: Special topics in statistics. In: Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0* (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Higgins 2011b

Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;**343**:d5928.

Killeen 2014

Killeen GF. Characterizing, controlling and eliminating residual malaria transmission. *Malaria Journal* 2014;**13**:330.

Lwetoijera 2014

Lwetoijera DW, Harris C, Kiware SS, Dongus S, Devine GJ, McCall PJ, et al. Increasing role of *Anopheles funestus* and *Anopheles arabiensis* in malaria transmission in the Kilombero Valley, Tanzania. *Malaria Journal* 2014;**13**:331.

Macdonald 1957

Macdonald G. *The Epidemiology and Control of Malaria*. London (UK): Oxford University Press, 1957.

Majambere 2008

Majambere S, Fillinger U, Sayer DR, Green C, Lindsay SW. Spatial distribution of mosquito larvae and the potential for targeted larval control in The Gambia. *American Journal of Tropical Medicine and Hygiene* 2008;**79**(1):19-27.

Moher 2009

Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLOS Medicine* 2009;**6**:e1000097.

Najera 2001

Najera JA. Malaria control: achievements, problems and strategies. *Parassitologia* 2001;**43**(1-2):1-89.

Review Manager 2014 [Computer program]

Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Richardson 2016

Richardson M, Garner P, Donegan S. Cluster randomised trials in Cochrane reviews: evaluation of methodological and reporting practice. *PLOS One* 2016;**11**(3):e0151818.

Roca-Feltrer 2009

Roca-Feltrer A, Schellenberg JR, Smith L, Carneiro I. A simple method for defining malaria seasonality. *Malaria Journal* 2009;**8**:276.

Russell 2010

Russell TL, Lwetoijera DW, Maliti D, Chipwaza B, Kihonda J, Charlwood JD, et al. Impact of promoting longer-lasting insecticide treatment of bed nets upon malaria transmission in a rural Tanzanian setting with pre-existing high coverage of untreated nets. *Malaria Journal* 2010;**9**:187.

Sachs 2002

Sachs JD. Macroeconomics and health: investing in health for economic development. *Revista Panamericana de Salud Pública* 2002;**12**(2):143-4.

Shousha 1948

Shousha AT. The eradication of *Anopheles gambiae* from Upper Egypt 1942-1945. *Bulletin of the World Health Organization* 1948;**1**(2):309-42.

Smith 2013

Smith DL, Perkins TA, Tusting LS, Scott TW, Lindsay SW. Mosquito population regulation and larval source management in heterogeneous environments. *PLOS One* 2013;**8**(8):e71247.

Soper 1943

Soper FL, Wilson DB. *Anopheles gambiae* in Brazil: 1930 - 1940. New York (NY): Rockefeller Foundation, 1943:1-262.

Sékétéli 2002

Sékétéli A, Adeoye G, Eyamba A, Nnoruka E, Drameh P, Amazigo UV, et al. The achievements and challenges of the

African Programme for Onchocerciasis Control (APOC). *Annals of Tropical Medicine & Parasitology* 2002;**96**(Suppl 1):S15-28.

Tusting 2013

Tusting LS, Thwing J, Sinclair D, Fillinger U, Gimnig J, Bonner KE, et al. Mosquito larval source management for controlling malaria. *Cochrane Database of Systematic Reviews* 2013, Issue 8. [DOI: [10.1002/14651858.CD008923.pub2](https://doi.org/10.1002/14651858.CD008923.pub2)]

Walshe 2017

Walshe DP, Garner P, Abdel-Hameed Adeel AA, Pyke GH, Burkot T. Larvivorous fish for preventing malaria transmission. *Cochrane Database of Systematic Reviews* 2017, Issue 12. [DOI: [10.1002/14651858.CD008090.pub3](https://doi.org/10.1002/14651858.CD008090.pub3)]

WHO 2008

World Health Organization. WHO position statement on integrated vector management, 2008. apps.who.int/iris/bitstream/10665/69745/1/WHO_HTM_NTD_VEM_2008.2_eng.pdf (accessed 11 July 2017).

WHO 2011

World Health Organization Vitamin and Mineral Nutrition Information System. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity, 2011. apps.who.int/iris/bitstream/10665/85839/3/WHO_NMH_NHD_MNM_11.1_eng.pdf?ua=1. Geneva: World Health Organization, (accessed 11 July 2017).

WHO 2012a

World Health Organization. Global plan for insecticide resistance management in malaria vectors, 2012. apps.who.int/iris/bitstream/10665/44846/1/9789241564472_eng.pdf?ua=1. Geneva: World Health Organization, (accessed 11 July 2017).

WHO 2012b

World Health Organization. Interim Position Statement The role of larviciding for malaria control in sub-Saharan Africa, April 2012. www.who.int/malaria/publications/atoz/interim_position_statement_larviciding_sub_saharan_africa.pdf?ua=1 (accessed prior to 5 June 2019).

WHO 2013

World Health Organization. Larval source management. A supplementary measure for malaria vector control. An operational manual. apps.who.int/iris/bitstream/10665/85379/1/9789241505604_eng.pdf?ua=1. Geneva, Switzerland: WHO, (accessed 11 July 2017).

WHO 2015a

World Health Organization. Global Technical Strategy for Malaria 2016-2030. Geneva: World Health Organization, 2015.

WHO 2015b

World Health Organization. Guidelines for the Treatment of Malaria. 3rd Edition. Geneva: World Health Organization, 2015.

WHO 2017

World Health Organization. World Malaria Report 2017. Geneva: World Health Organization, 2017.

Larviciding to prevent malaria transmission (Review)

WHO 2018

World Health Organization. World Malaria Report 2018. Geneva: World Health Organization, 2018.

Wilson 2015

Wilson AL, Boelaert M, Kleinschmidt I, Pinder M, Scott TW, Tusting LS, et al. Evidence-based vector control? Improving the quality of vector control trials. *Trends in Parasitology* 2015;**31**:380-90.

References to other published versions of this review
Choi 2017

Choi L, Wilson A. Larviciding to control malaria. *Cochrane Database of Systematic Reviews* 2017, Issue 7. [DOI: [10.1002/14651858.CD012736](https://doi.org/10.1002/14651858.CD012736)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]

Fillinger 2009

Methods	<p>Study design: controlled before-and-after study</p> <p>Unit of allocation: clusters (valleys)</p> <p>Number of units: 3 valleys per arm</p> <p>Outcome assessment/surveillance type: 6 paired cross-sectional surveys in cohort of children conducted during the long rains (April–June) and short rains (November–January) each year. Blood smears were collected from all children in each cohort to be assessed by microscopy for parasite identification and density. On each occasion, each pair of surveys were carried out 2 months apart. Thus, parasite infection status was assessed during 6 consecutive rainy seasons (the first 6 surveys at baseline, the following 6 during the intervention) from April 2004 to January 2007.</p> <p>Length of follow-up: April 2004 to January 2007 (survey 1–3 at baseline, survey 4–6 during the intervention)</p> <p>Adjustment for clustering: yes</p>
Participants	<p>Number of participants: 120 children in each valley (360:360)</p> <p>Population characteristics: children aged 6 months to 10 years. Approximately equal numbers of boys and girls.</p> <p>Withdrawal and loss to follow-up: some absences from cross-sectional surveys. Children said to be due to travelling in all cases. Figure 1 in the study paper reported numbers of each survey. Numbers present appeared to be consistently higher by above 10% in the control group compared to the intervention group.</p>
Interventions	<p>Larvicide:</p> <p>Active ingredient and dosage: commercial strains of <i>Bs</i> or <i>Bti</i></p> <p>Formulation: water-dispersible (<i>Bs</i>) and granule formulations (<i>Bti</i>)</p> <p>Manufacturer: Valent BioSciences LLC, USA</p> <p>Quality control of the larvicide: not reported</p> <p>Duration of the activity of the larvicide: <i>Bs</i> was used for the first 6 months of the intervention but, due to a lack of a residual effect, it was replaced with <i>Bti</i>, which is cheaper and forestalls the development of resistance due to its more complex mode of action.</p> <p>Method of application: not reported</p> <p>Frequency of application: applied to 3/6 valleys beginning in July 2005. Larvicide was applied to all water bodies at weekly intervals.</p> <p>Coverage: not reported</p>

Larviciding to prevent malaria transmission (Review)

Fillinger 2009 (Continued)

Buffer size between clusters: valleys at least 1 km apart

Cointerventions: ITNs, mainly long-lasting insecticidal nets

Types of nets used: PermaNet, Vestergaard Frandsen, Lausanne, Switzerland; Olyset, Sumitomo Chemical, Tokyo, Japan

Delivery method: not delivered as part of study – from 2005 onwards, ITNs were intensively promoted and supplied by government health facilities and non-governmental organizations. In addition, in July and August 2006, joint measles–malaria campaigns combining the distribution of ITNs with measles vaccinations were conducted.

Coverage: not reported

Net use: ITN use in sentinel households increased from 4.8% (95% CI 3.0% to 6.6%; range in control valleys 1–9% and in intervention valleys 2–6%) at baseline to 40.8% (95% CI 36.7% to 45.0%; range in control valleys 24–51% and in intervention valleys 25–51%) during the intervention year. The probability that ITNs were used was the same in the control and intervention group (OR 1.06, 95% CI 0.56 to 1.99) in both years but increased in both groups during the intervention (OR 13.58, 95% CI 7.30 to 25.28).

Outcomes
Primary:

Incidence of new *Plasmodium* infections in children aged 0.5–10 years, measured as children who were negative at the first paired survey but positive at the second

Prevalence of *Plasmodium* infections in children aged 0.5–10 years

Secondary:

Annual EIR

Mean human biting rate

Anopheline late instar larval abundance

Anopheline adult abundance

Location profile

Study location: Kakamega and Vihiga highlands, western Kenya. Hilly areas characterized by steep-sided valleys with flat bottoms and plateaus, where most homes were built. Small streams ran along the valley bottoms and papyrus swamps were common. Altitude of the highland valley communities ranged from 1453 to 1632 m.

Malaria endemicity: moderately endemic

EIR: annual EIR of *An gambiae s.l.* and *An funestus s.l.* combined was 10–12 infectious bites per person in both groups at baseline

Population proximity/density: densely populated districts

Plasmodium species: *Plasmodium falciparum*

Vector profile

Primary (and secondary) vector species: *An gambiae s.s.* (*An funestus*, *An arabiensis*, *An rufipes*)

Vector behaviour (nature, stability, adult habitat, peak biting times, exophilic/endophilic, exophagic/endophagic, anthropophilic/zoophilic): not reported

Phenotypic resistance profile: not reported

Genotypic resistance profile: not reported

Method of mosquito collection:

Larval surveys done weekly in all valleys. The presence or absence of anopheline and culicine larvae was recorded in all aquatic habitats. Purposive dipping was used to sample larvae (10 dips per site). Larvae were categorized as early stage (first and second instars) and late stage (third and fourth in-

Fillinger 2009 (Continued)

stars). In 10 randomly selected sentinel sites per valley, weekly larval densities (mean number of larvae per dip per habitat) were recorded and the proportion of late instar larvae was calculated as an indicator of larval survival and emergence.

Indoor-resting mosquitoes collected monthly using pyrethrum spray catches from 10 sentinel houses in each valley that were randomly selected from households within 500 m of the valley bottom. The type of household, number of occupants during the night before, and mosquito control methods used were recorded routinely. *An gambiae s.l.* were identified to the species level using PCR, and presence of sporozoites was determined by ELISA of pooled samples of 10 mosquitoes per test.

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Not randomized.
Allocation concealment (selection bias)	High risk	Not randomized.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not blinded; however, the cointervention (ITNs) that were used in study arms had a similar usage in both.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Microscopists were blinded. RDTs and ELISA are objective measurements, but microscopy is objective.
Incomplete outcome data (attrition bias) All outcomes	High risk	Some absences from cross-sectional surveys. Children said to be due to travelling in all cases. Figure 1 in the study paper reported numbers of each survey. Numbers presented appeared to be consistently higher by > 10% in the control group compared to the intervention group.
Selective reporting (reporting bias)	Unclear risk	No protocol was published beforehand. All expected outcomes from a trial such as this were reported.
Baseline characteristics	Low risk	Baseline characteristics appeared similar between the control and intervention arms. Key characteristics such as age and sex were the same.
Contamination	Low risk	Valleys were at least 1 km apart so assumed no contamination risk.
Incorrect analysis	Low risk	Cluster adjustment was carried out. Valleys were treated as the unit of geographic location and included as confounders in their models for analysis.
Baseline outcome measurements All outcomes	Low risk	Baseline outcome measurements appeared similar between the control and intervention arms.

Maheu-Giroux 2013a

Methods

Study design: CBA of a staged programmatic implementation with randomized, cluster sampled household surveys

Unit of allocation: clusters as wards

Larviciding to prevent malaria transmission (Review)

Copyright © 2019 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Maheu-Giroux 2013a (Continued)

Number of units: initially 15 control wards, period 1 (3 intervention, 12 control), period 2 (9 intervention, 6 control), period 3 (15 intervention)

Outcome assessment/surveillance type: 6 rounds of cross-sectional household surveys). A list of TCUs (small administrative units) was assembled for each ward before March 2004 and was regularly updated throughout the study duration. During the first round of the survey, 10 TCUs were randomly sampled from each of the 15 wards. All households located in the sampled TCUs were invited to participate in the survey. From the second round onwards, the TCUs sampled in the first round were followed up longitudinally, and another 10 TCUs per ward were selected for cross-section surveys. Household survey administered and blood films taken.

Length of follow-up: from May 2004 to Dec 2008

Adjustment for clustering: yes

Participants

Number of participants: > 610,000 residents

Population characteristics: used total population

Withdrawal and loss to follow-up: not reported

Interventions

Larvicide:

Active ingredient and dosage: *Bti* (VectoBacH) for open (light-exposed) habitats and *Bs* (VectoLexH) for closed (covered, often highly polluted) habitats. Dosages of 0.04 g/m² for water-dispersible granule formulations and 1 g/m² for corn cob granule formulations of *Bti*. 1 g/m² of *Bs*. The targeting of closed habitats was for *Culex* mosquitoes.

Formulation: water-dispersible granule and corn cob formulations for *Bti* and corn cob formulations for *Bs*

Manufacturer: Valent BioSciences LLC, USA

Quality control of the larvicide: not reported

Duration of the activity of the larvicide: not reported

Method of application: community based but vertically managed intervention implementation. Open habitats (potential to produce *Anopheles* larvae), were treated by the Mosquito Control CORPs, each of whom was assigned to a specific *mtaa* (administrative subunits) or portions of an *mtaa*. Closed habitat were treated by an additional team of CORPs.

Frequency of application: open habitats were treated every week with *Bti*. Closed habitats treated every 3 months.

Buffer size between clusters: not reported.

Cointerventions: all existing interventions such as ITNs, house screening, ceiling boards, repellents, coils, and spray.

Outcomes

Infection prevalence in all ages

EIR

Adult mosquito density

Location profile

Study location: Dar es Salaam, Tanzania

Malaria endemicity: climate was tropical humid with 2 rainy seasons – the long rains during the months of April and May and the short rains of October and November. Malaria transmission was year-round with peaks in incidence after the 2 rainy seasons.

EIR: 1.28 (all *Anopheles*)

Maheu-Giroux 2013a (Continued)

Population proximity/density: not reported

Plasmodium species: *Plasmodium falciparum* accounted for > 90% of cases

Vector profile

Primary (and secondary) vector species: *An gambiae s.l.* and *An funestus s.l.* *An coustani* has a low contribution to transmission.

Vector behaviour (nature, stability, adult habitat, peak biting times, exophilic/endophilic, exophagic/endophagic, anthropophilic/zoophilic): exophagic

Phenotypic resistance profile: not reported

Genotypic resistance profile: not reported

Method of mosquito collection: HLC performed in all clusters. In each of the 67 *mtaa*, 4 different, well-distributed sampling locations were chosen non-randomly to maximize coverage of surveillance, resulting in a total of 268 routinely maintained surveillance sites. HLC was conducted once every 4 weeks, overnight. In order to estimate the total true exposure experienced both indoors and outdoors by residents, directly measured outdoor mosquito densities were multiplied by the coefficient of the estimated total true human exposure divided by the estimated total outdoor biting rate obtained from detailed studies of mosquito–human interactions. These coefficients were derived from an in-depth behavioural survey of both mosquitoes and humans which was conducted during the main rainy season of April to June 2006. Captured mosquitoes were identified to genus morphologically and to subspecies by PCR. ELISA was used to detect infection of sporozoites.

Notes

The entomological outcomes were extracted from an earlier published paper, before all clusters received the intervention (3/15), related to the same study (Geissbühler 2009).

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Intervention was not randomly allocated; however, participants selected for outcome assessment were randomly selected.
Allocation concealment (selection bias)	High risk	Intervention was not randomly allocated; however, participants selected for outcome assessment were randomly selected.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Not blinded; however, cointerventions used in study arms had a similar usage in both. Also analysis was adjusted, taking into account as possible covariates such as cointervention usage.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Not blinded but outcome data collected independently of those implementing control.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Study authors stated loss to follow-up was minimal; however, no numbers were reported.
Selective reporting (reporting bias)	Unclear risk	No protocol was published beforehand. All expected outcomes from a trial such were reported.
Baseline characteristics	Unclear risk	Baseline characteristics appeared similar between the control and intervention arms which we took as their third survey round. Key characteristics such as age and sex were the same.

Maheu-Giroux 2013a (Continued)

Contamination	Unclear risk	No mention of a buffer zone or population migration (both in terms of participants and mosquitoes). Potential for bias but unclear from what the study authors report.
Incorrect analysis	Low risk	Cluster adjustment was carried out. TCU and household was taken into account with their multivariate model.
Baseline outcome measurements All outcomes	Low risk	Baseline outcome measurements appeared similar between the control and intervention arms.

Majambere 2010

Methods	<p>Study design: non-randomized cross-over trial</p> <p>Unit of allocation: zones</p> <p>Number of units: 4 zones approximately 12 × 8 km in area and divided into 3 parallel 4-km wide bands (subzones) perpendicular to the river.</p> <p>Outcome assessment/surveillance type: cross-sectional surveys (start and end of transmission season in 2 years (June and December 2006, then June and December 2007) and passive case detection. Census of residents, including children aged 6 months to 10 years, was carried out in 50 study villages during the dry season in 2006. Children were selected from random lists, with the total in each village proportional to village size.</p> <p>Length of follow-up: baseline entomological data, but not clinical data, were collected during July–November 2005. In 2006 and 2007, entomological and clinical data collection started in May and ended in November. A cross-over design was used for the application of larvicide. From June to November 2006, larvicide was applied to all accessible aquatic habitats in zones 1 and 3 at weekly intervals and zones 2 and 4 served as controls. From May to November 2007, larvicide intervention was applied to zones 2 and 4 and zones 1 and 3 served as controls.</p> <p>Adjustment for clustering: no</p>
Participants	<p>Number of participants: 2039 total children at the first cross-sectional survey. 1862 in the final survey.</p> <p>Population characteristics: children aged 0.5–10 years old</p> <p>Withdrawal and loss to follow-up: from enrolment and first survey to the second survey, there was a high number of participants who were absent (98 in zone 1, 65 in zone 2, 76 in zone 3, and 76 in zone 4). A new cohort of children was used from survey 3 onwards which included previous children that were still under 10 years if age and a selection of new children which replaced either children over 10 years of age or any that had left the study site. From this new cohort, a total of 184 were absent across all zones (33 in zone 1, 50 in zone 2, 47 in zone 3, and 54 in zone 4).</p>
Interventions	<p>Larvicide:</p> <p>Active ingredient and dosage: commercial strains of <i>Bti</i> at 0.2 kg/hectare for water-dispersible granules and at 5 kg/hectare for corn granules</p> <p>Formulation: water-dispersible granule and corn granules</p> <p>Manufacturer: Valent BioSciences LLC, USA</p> <p>Quality control of the larvicide: field applicators were recruited from communities supervised by 1 field supervisor in each zone and trained for 1 month before larviciding. Larval surveys were carried out continuously by the zone supervisor. In 2005, during the baseline period, all aquatic habitats in each zone were visited and the presence or absence of anopheline and culicine larvae recorded as described else-</p>

Larviciding to prevent malaria transmission (Review)

Majambere 2010 (Continued)

where. Each habitat was visited monthly. During the intervention years (2006 and 2007) random larval spot checks were implemented throughout the season to estimate the proportion of habitats containing early and late instar larvae to determine the effectiveness of larvicide application.

Of the total number of habitats identified in each zone during baseline (1076), 40 habitats were randomly (computer-generated) selected every day for each zone respectively by the programme manager (S.M.) and the habitat identification number, including global positioning system co-ordinates, forwarded to the field supervisor for habitat inspection as described above. Selection of sites was stratified according to subzone and the timetable of larvicide application to ensure that inspection of sites took place 1–2 days after the habitat was treated with larvicide and that an equal number of sites were visited weekly in all three subzones in each zone. In addition, 10 sentinel habitats per zone were randomly selected after the first round of complete habitat surveys in 2005 and larval densities measured weekly in these.

At each site visit, purposive dipping was used to sample larvae (10 dips per site), which were categorized as early (first and second instars) stages and late (third and fourth) stages.

Duration of the activity of the larvicide: not reported.

Method of application: the water-dispersible granules were applied as liquid with knapsack compression sprayers (15-L capacity diaphragm knapsack sprayers, Solo 475; Solo Kleinmotoren GmbH, Sindelfingen, Germany) in areas with low vegetation coverage. The corn granules were applied by hand from buckets held with a strap around the waist or neck or motorized knapsack granuleblowers (13-L capacity motorized sprayers; MD 150DX-13; Maruyama, Tokyo, Japan) when aquatic habitats were covered by vegetation and difficult to access.

Frequency of application: weekly

Buffer size between clusters: study villages were recruited from the central band of each zone. The study authors assumed that when larvicide was applied to an entire study zone, the 2 × 4-km bands, either side of the central band, would be sufficiently wide to minimize mosquito movement from untreated sites outside the study zone into the central band, where the study villages were located.

Cointerventions: existing ITNs

Compliance: use increased during study range in 2006 was 6.1% to 38.3%, range in 2007 37.2% to 81.4%.

Outcomes	<p>Incidence of malaria cases per 100 child-years defined as a history of fever within the last 48 hours or axillary temperature ≥ 37.5 °C later confirmed with the presence of <i>Plasmodium falciparum</i> identified microscopically.</p> <p>Prevalence of <i>P falciparum</i> infection</p> <p>Mean haemoglobin level</p> <p>Prevalence of splenomegaly</p> <p>Prevalence of gametocytaemia</p> <p>Seasonal EIR</p> <p>Number of female adult <i>An gambiae s.l.</i></p> <p>Sporozoite rate</p>
----------	---

Location profile	<p>Study location: floodplains of river Gambia, east of Farafenni, The Gambia. Study was carried out in 4 separate areas (referred to as zones 1–4), 2 on the north banks and 2 on the south banks of the Gambia River. Flat open Sudan savannah broadly consisting of farmlands, sparse woodland, and the extensive alluvial floodplains of the river.</p> <p>Malaria endemicity: seasonal transmission</p> <p>EIR: not reported</p>
------------------	--

Larviciding to prevent malaria transmission (Review)

Majambere 2010 (Continued)

Population proximity/density: % of villages in each zone < 1 km from the floodplain was reported. In zones 1 and 2, this was < 20% on average. In zones 3 and 4, this was > 80%.

Plasmodium species: *Plasmodium falciparum*

Vector profile

Primary (and secondary) vector species: *An gambiae s.s.*, *An melas*, and *An arabiensis*

Vector behaviour (nature, stability, adult habitat, peak biting times, exophilic/endophilic, exophagic/endophagic, anthropophilic/zoophilic): not reported

Aquatic habitats (type, stability and extent (number and size), proximity of aquatic habitats to human habitation): flood plains of River Gambia

Phenotypic resistance profile: not reported

Genotypic resistance profile: not reported

Method of mosquito collection: adult vector surveys were implemented in 39 villages (10 in zone 1, 11 in zone 2, 9 in zone 3, and 9 in zone 4) at 2-week intervals from July to November in 2005 and for the duration of larviciding in the intervention years. Each zone had 15 traps divided between the villages with 1–3 sentinel houses per village proportional to village size. Within randomly selected compounds, all houses with open eaves, a thatched roof, no ceiling, and where a single man slept were numbered and 1 was selected randomly. Mosquitoes were sampled using miniature CDC light traps (Model 512; John W. Hock Company, Gainesville, FL) positioned 1 m above the floor at the foot end of the bed where a person slept under an untreated bed net. Traps were set at 7:00 p.m. and collected at 7:00 a.m. the following morning. If the occupant moved house, the trap was moved to the nearest similar house in the same village. If the occupant did not spend the night in the selected room or the trap was faulty, the data were excluded from the analysis.

Mosquitoes were identified to the level of species by microscopy and the numbers of *An gambiae s.l.* females recorded. The presence of sporozoites was identified using ELIZA. In 2005 and 2006, a 1% random sample of the *An gambiae s.l.* females, stratified by zone and sampling period, was typed to the species by PCR.

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Intervention was not randomly allocated; however, participants selected for outcome assessment were randomly selected.
Allocation concealment (selection bias)	High risk	Intervention was not randomly allocated; however, participants selected for outcome assessment were randomly selected.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Residents were aware of ongoing interventions but this was unlikely to have impacted results. There was a large difference in net use in zone 1 compared to the other zones; however, this was measured at baseline and net use increased at a similar rate throughout all zones after the intervention was introduced.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Reading of blood films was blinded and RDTs are objective assessments.
Incomplete outcome data (attrition bias) All outcomes	High risk	> 10% loss in study groups. See above under 'Withdrawal and loss to follow-up'

Majambere 2010 (Continued)

Selective reporting (reporting bias)	Unclear risk	No protocol was published beforehand. All expected outcomes from a trial such were reported.
Baseline characteristics	High risk	Large differences in key baseline characteristics (sex ratio, ethnicity, net use)
Contamination	Low risk	Study authors do not specifically mention contamination. However, the intervention is expected to have a very short-lasting effect so would not carry over to the following year when the cross-over of intervention happened.
Incorrect analysis	Unclear risk	Suggested no paired analysis was done, therefore, it should have an inappropriate weighting in a meta-analysis. However, this is not so much a risk of bias issue and this study was not meta-analyzed.
Baseline outcome measurements All outcomes	High risk	Taking the prevalence of <i>Plasmodium falciparum</i> infections, zone 1 had a much higher prevalence (38.4%) compared to the others (9.5–16.8%)
Other bias	Low risk	<p>Suitability of a cross-over design: low risk. Malaria can be highly seasonal but study authors reported rainfall to be consistent throughout the years of the trial period.</p> <p>Whether only first-period data are available: low risk. Multiple periods of data reported.</p> <p>Comparability of results with those from parallel-group trials: low risk.</p>

Yapabandara 2001

Methods	<p>Study design: cRCT</p> <p>Unit of allocation: clusters (villages)</p> <p>Number of units: 8 villages divided equally into 2 arms. On the basis of 1 year's preintervention data the villages were stratified into 4 with high levels of malaria transmission and 4 with lower transmission. Within each strata 2 villages randomly selected for intervention.</p> <p>Outcome assessment/surveillance type: passive case detection. Also 2 mass blood surveys were carried out in July and December during the pre- and postintervention years. Blood films were taken, regardless of the presence/ absence of fever, from all the residents of the 8 villages.</p> <p>Length of follow-up: January 1994 to December 1995</p> <p>Adjustment for clustering: yes</p>
Participants	<p>Number of participants: 4/8 villages had populations < 500 while the other 4 had populations of 600–1100.</p> <p>Population characteristics: not reported.</p> <p>Withdrawal and loss to follow-up: not reported.</p>
Interventions	<p>Larvicide:</p> <p>Active ingredient and dosage: pyriproxyfen, S31183 (Adeal 0.5% G) applied at a rate of 0.01 mg active ingredient/L (2 g granules/m³)</p> <p>Formulation: not reported.</p> <p>Manufacturer: not reported.</p>

Larviciding to prevent malaria transmission (Review)

Yapabandara 2001 (Continued)

Quality control of the larvicide: community engagement to encourage community to inform about new gem pits so that they could be rapidly treated.

Duration of the activity of the larvicide: not reported but assays were conducted to determine if residual activity was present.

Method of application: not reported.

Frequency of application: 3 applications – 1 in December 1994, 1 between June and July 1995 in the postmonsoon season when river bed pools were formed, and 1 at end of November 1995.

Buffer size between clusters: not reported.

Outcomes

Malaria incidence defined as fever/history of fever and parasites detected by blood film.

Infection prevalence (slide positivity rates)

Number of anophelines

Location profile
Study location:

This study was carried out in Kaluganga, which is part of Elahera gem-mining area situated in Matale District (7°40N, 80°50E) in the dry central zone of Sri Lanka. A cluster of 8 villages with a total area of 23 km² was selected for this study. The numbers of gem pits per village ranged from 311 to 3622. The villages were surrounded by thick jungle. The area was a settlement scheme, which was established about 30 years before the trial was conducted around the rivers, Aban ganga and Kalu ganga.

Malaria endemicity: not reported.

EIR: not reported.

Population proximity/density: treated gem pits and pools up to 1.5 km from villages.

Plasmodium species: *Plasmodium falciparum* and *Plasmodium vivax*

Vector profile

Primary (and secondary) vector species: *An culicifacies* (*An subpictus* and *An varuna*)

Vector behaviour (nature, stability, adult habitat, peak biting times, exophilic/endophilic, exophagic/endophagic, anthropophilic/zoophilic): not reported

Aquatic habitats (type, stability and extent (number and size), proximity of aquatic habitats to human habitation):

Shallow pits dug by gem miners that filled with water. Breeding of *An culicifacies* was almost entirely in gem pits but some breeding of *An subpictus* and most of *An varuna* was in other sites such as river bed pools and slow-moving river margins

Phenotypic resistance profile: not reported

Genotypic resistance profile: not reported

Method of mosquito collection: Anopheline populations in the study area were estimated by 7 sampling methods: window exit trap collection; pyrethrum spray sheet collection; indoor HCs; all night or for the first part of the night up to midnight; cattle-baited hut collection and cattle-baited net trap collection; and light trap collection. The locations chosen for applying these methods were near the centres of each village to try to avoid interference by immigration of mosquitoes from neighbouring villages. Data reported were only from cattle-baited huts, partial night landing catches, and all night landing catches.

Notes
Risk of bias
Larviciding to prevent malaria transmission (Review)

Yapabandara 2001 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Stated that they randomized, but unclear how.
Allocation concealment (selection bias)	Unclear risk	Stated that they randomized, but unclear how.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinding was not possible for the intervention; however, unlikely to affect the outcome.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Not reported whether blinding was used. Unclear whether slide readers were blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No cohort established. Movement in and out of study area not documented.
Selective reporting (reporting bias)	Unclear risk	No protocol was published beforehand. Not all the stated entomological outcomes described in the methods were reported.
Incorrect analysis	High risk	Inappropriate analysis, no adjustment for clustering
Other bias	Unclear risk	<p>Recruitment bias: low risk – randomized study where they had selected clusters based on malaria cases before the intervention was introduced to ensure this was equal in both arms. Mass blood surveys and census attempted to include the entire population.</p> <p>Baseline imbalance: low risk – baseline characteristics appeared similar</p> <p>Loss of clusters: low risk – no clusters were lost</p> <p>Comparability with RCTs randomizing participants: low risk – larviciding is expected to have a community wide effect and should be implemented at a community level.</p>

Abbreviations: *An*: *Anopheles*; *Bti*: *Bacillus thuringiensis israeliensis*; *Bs*: *Bacillus sphaericus*; EIR: entomological inoculation rate; ELISA: enzyme-linked immunosorbent assay; HLC: human landing catches; ITN: insecticide-treated nets; OR: odds ratio; PCR: polymerase chain reaction; RCT: randomized controlled trial; RDT: rapid diagnostic test; TCU: Ten-Cell Unit.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Abdalmagid 2012	No relevant outcomes
Afrane 2016	No relevant outcomes
Ansari 2005	No relevant outcomes
Balaraman 1983	No relevant outcomes
Balaraman 1987	No relevant outcomes

Larviciding to prevent malaria transmission (Review)

Study	Reason for exclusion
Bertram 1950	No relevant outcomes
Bhalwar 1995	No relevant outcomes
Biswas 1997	No relevant outcomes
Bond 2004	No relevant outcomes
Brescia 1947	Study design did not match inclusion criteria
Castro 2002	Study design did not match inclusion criteria
Castro 2009	Intervention did not match inclusion criteria
Chaki 2011	Study design did not match inclusion criteria
Chen 1985	Study design did not match inclusion criteria
Chen 1988	Study design did not match inclusion criteria
Claborn 2002	No relevant outcomes
Derua 2017	No relevant outcomes
Djènonatin 2014	No relevant outcomes
El Safi 1986	No relevant outcomes
Farashiani 2000	Full text not available
Fillinger 2003	No relevant outcomes
Galardo 2013	No relevant outcomes
Giurcă 1978	Duplicate but in a different journal
Haq 2004	No relevant outcomes
Houten 1980	Full text not available
Imbahale 2012	Intervention did not match inclusion criteria
Johnson 1947	No relevant outcomes
Julvez 1987a	Study design did not match inclusion criteria
Julvez 1987b	Duplicate but in a different journal
Kanda 1995	Study design did not match inclusion criteria
Karanja 1994	No relevant outcomes
Konradsen 1999	Study design did not match inclusion criteria
Kramer 2014	Protocol

Study	Reason for exclusion
Kumar 1994	Intervention did not match inclusion criteria
Kumar 1998	Intervention did not match inclusion criteria
Kumar 2013	Intervention did not match inclusion criteria
Kusumawathie 2008	No relevant outcomes
Ladoni 1986	Full text not available
Lee 1990	No relevant outcomes
Liu 2009	Study design did not match inclusion criteria
Lunin 1979	No relevant outcomes
Mahdi 1967	No relevant outcomes
Maheu-Giroux 2013b	No relevant outcomes
Maheu-Giroux 2013c	Duplicate but in a different journal
Maheu-Giroux 2014	No relevant outcomes
Marina 2014	No relevant outcomes
McCann 2017a	Protocol
McCann 2017b	Intervention did not match inclusion criteria
Meng 1996	Study design did not match inclusion criteria
Minakawa 2007	Protocol (study not performed)
Mossadegh 1973	Study design did not match inclusion criteria
Msellemu 2016	Study design did not match inclusion criteria
Müller 1984	Study design did not match inclusion criteria
Obopile 2018	Study design did not match inclusion criteria
Ouedraogo 2017	Study design did not match inclusion criteria
Parvez 2003	No relevant outcomes
Perich 1990	No relevant outcomes
Prabhu 2011	No relevant outcomes
Pridantseva 1980	No relevant outcomes
Ranjbar 2012	Study design did not match inclusion criteria
Rettich 1973	Conference abstract

Larviciding to prevent malaria transmission (Review)

Study	Reason for exclusion
Rifaat 1974	Full text not available
Sharma 1983	No relevant outcomes
Sharma 1989	Intervention did not match inclusion criteria
Sharma 2003	No relevant outcomes
Shililu 2003	No relevant outcomes
Shimada 2007	Study design did not match inclusion criteria
Skovmand 1997	No relevant outcomes
Skovmand 1999	No relevant outcomes
Skovmand 2011	Intervention did not match inclusion criteria
Some 1994	Study design did not match inclusion criteria
Song 2013	Study design did not match inclusion criteria
Srivastava 1996	No relevant outcomes
Tchicaya 2009	No relevant outcomes
Tchicaya 2010	Conference abstract
Teng 2005	No relevant outcomes
Tâcu 1977	Study design did not match inclusion criteria
Usenbaev 2006	Study design did not match inclusion criteria
Vasuki 1992	Study design did not match inclusion criteria
Wang 1983	Study design did not match inclusion criteria
Xu 1980	Study design did not match inclusion criteria
Xu 1983	Study design did not match inclusion criteria
Xu 2004	Study design did not match inclusion criteria
Yapabandara 2002	No relevant outcomes
Yapabandara 2004	Intervention did not match inclusion criteria
Yapabandara 2005	No relevant outcomes
Zhou 2010	Study design did not match inclusion criteria
Zhou 2016	Protocol
Zohdy 1982	Study design did not match inclusion criteria

Larviciding to prevent malaria transmission (Review)

Characteristics of studies awaiting assessment [ordered by study ID]

Fuseini 2017

Methods	Unavailable
Participants	Unavailable
Interventions	Unavailable
Outcomes	Unavailable
Notes	

Javadian 1974

Methods	<p>Study design: CBA</p> <p>Unit of allocation: clusters (dehestans which are administrative units above villages)</p> <p>Number of units: 4 dehestans per arm</p> <p>Outcome assessment/surveillance type: active case detection</p> <p>Length of follow-up: 1 year preintervention and 3 years postintervention. Overall period of April 1965 to the end of 1968</p> <p>Adjustment for clustering: unclear as not reported</p>
Participants	<p>Number of participants: total population varied throughout study period.</p> <p>1965: 40,794 in control, 28,999 in intervention</p> <p>1966: 41,514 in control, 27,446 in intervention</p> <p>1967: 46,226 in control, 43,663 in intervention</p> <p>1968: 46,757 in control, 32,649 in intervention</p> <p>Population characteristics: total population was used. Population denominator not stable, swelled by migration during date palm harvest season which coincides with peak transmission in Aug/Sept.</p> <p>Withdrawal and loss to follow-up: not reported</p>
Interventions	<p>Larvicide:</p> <p>Active ingredient and dosage: the text suggested it is petroleum oil</p> <p>Formulation: not reported</p> <p>Manufacturer: not reported</p> <p>Quality control of the larvicide: not reported</p> <p>Duration of the activity of the larvicide: not reported</p> <p>Method of application: not reported</p> <p>Frequency of application: not reported</p>

Larviciding to prevent malaria transmission (Review)

Javadian 1974 (Continued)

Buffer size between clusters: not reported

Cointerventions: IRS

Active ingredient and dosage: DDT 2 g/m² and malathion 2 g/m²

Formulation: not reported

Frequency of spraying: 2 rounds per year of DDT until 1967. After that, DDT was used for the first round and malathion was used for the second each year.

Coverage: not reported

Outcomes

Malaria incidence

Indoor resting density of *An stephensi*

Notes
Zhou 2013
Methods

Study design: cRCT assessing a combination of vector control interventions. The study spanned 2 years (2010 to 2011), but for this review, only the data from 2011 were included as this was when larviciding was used.

Unit of allocation: clusters

Number of units: 3 study sites which each had 3 paired clusters, making a total of 9 paired clusters. Each of these pair would then be randomly assigned the control (ITNs) or the intervention (ITNs + larviciding). Furthermore, within each cluster, an area was targeted with IRS.

Outcome assessment/surveillance type: active case detection in cohort of 350 participants per cluster in 2010 and 450 per cluster in 2011.

Cross-sectional surveys were done during February and March 2010 before the IRS application and 2011 before the IRS and larvicide application. Another survey was conducted post intervention in May 2010, May 2011, and July 2011. Blood smears were taken from randomly selected participants of different ages within each cluster: approximately 150 in 2010 and 250 in 2011.

Only 12/18 clusters were monitored in 2011.

Length of follow-up: around 2 years: February 2010 to July 2011

Adjustment for clustering: yes

Participants

Number of participants: in 2010, numbers ranged from 2884 to 2906 for the cross-sectional survey. In 2011, numbers ranged from 4323 to 5139. For the cohort, there were 6248 participants in 2010 and 5574 in 2011.

Population characteristics: not reported but no particular age group was targeted.

Withdrawal and loss to follow-up: not reported.

Interventions
Larvicide:

Active ingredient and dosage: commercial strains of *Bti* (VectoMax)

Formulation: corn granules

Manufacturer: Valent BioSciences LLC, USA

Zhou 2013 (Continued)

Quality control of the larvicide: aquatic habitats were searched thoroughly by a team of technicians accompanied by field assistants from local villages.

Duration of the activity of the larvicide: not reported.

Method of application: not reported.

Frequency of application: first round of application was completed in February/March 2011 and the second in March/April 2011, 4 weeks after the first round.

Buffer size between clusters: 250 m buffer zone between control and intervention clusters; however, there was no buffer between the IRS targeted zones and the non-targeted zones.

Cointervention: ITNs

Active ingredient and dosage: not reported.

Method of distribution: Global Fund supported mass distribution in June to September 2006 (targeting < 5 s and pregnant women) in all study areas. A second round was accomplished during May to July 2011 (targeting everyone at risk).

Coverage: increased from 40.7% (range 34.3 to 47.8%) in 2010 to 93.0% (range 81.6 to 100%) in 2011.

Compliance: not reported

Outcomes	Malaria incidence defined as fever/history and plasmodium parasites detected by smear
	Parasite prevalence
	Indoor resting density of mosquitoes
Notes	

Abbreviations: *An*: *Anopheles*; *Bti*: *Bacillus thuringiensis israeliensis*; CBA: controlled before-and-after; cRCT: cluster-randomized controlled trial; DDT: dichlorodiphenyltrichloroethane; IRS: indoor residual spraying.

DATA AND ANALYSES


Comparison 1. Larviciding versus no larviciding

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Malaria incidence (cluster-randomized controlled trial (cRCT))	1		Rate Ratio (Fixed, 95% CI)	Subtotals only
1.1 Sensitivity analysis with estimated mean cluster population 675 and estimated intracluster correlation coefficient (ICC) 0.01	1		Rate Ratio (Fixed, 95% CI)	0.24 [0.14, 0.40]
1.2 Sensitivity analysis with estimated mean cluster population 675 and estimated ICC 0.05	1		Rate Ratio (Fixed, 95% CI)	0.24 [0.08, 0.70]
1.3 Sensitivity analysis with estimated mean cluster population 675 and estimated ICC 0.1	1		Rate Ratio (Fixed, 95% CI)	0.24 [0.05, 1.08]


Larviciding to prevent malaria transmission (Review)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2 Malaria incidence with subgrouping by extent of aquatic habitat (non-randomized study (NRS))	2		Risk Ratio (Random, 95% CI)	1.17 [0.60, 2.26]
2.1 Habitats < 1 km ²	1		Risk Ratio (Random, 95% CI)	0.62 [0.38, 1.01]
2.2 Habitats > 1 km ²	1		Risk Ratio (Random, 95% CI)	1.58 [0.94, 2.65]
3 Parasite prevalence (CRCTs)	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
3.1 Sensitivity analysis with estimated mean cluster population 675 and estimated ICC 0.01	1	763	Risk Ratio (M-H, Fixed, 95% CI)	0.08 [0.02, 0.46]
3.2 Sensitivity analysis with estimated mean cluster population 675 and estimated ICC 0.05	1	169	Risk Ratio (M-H, Fixed, 95% CI)	0.19 [0.02, 1.68]
3.3 Sensitivity analysis with estimated mean cluster population 675 and estimated ICC 0.1	1	87	Risk Ratio (M-H, Fixed, 95% CI)	0.26 [0.03, 2.42]
4 Parasite prevalence (NRS)	3		Risk Ratio (Random, 95% CI)	Subtotals only
4.1 Adjusted data	2		Risk Ratio (Random, 95% CI)	0.79 [0.71, 0.89]
4.2 Sensitivity analysis including Majambere 2010 with estimated ICC 0.01	3		Risk Ratio (Random, 95% CI)	0.88 [0.66, 1.16]
4.3 Sensitivity analysis including Majambere 2010 with estimated ICC 0.05	3		Risk Ratio (Random, 95% CI)	0.80 [0.71, 0.89]
4.4 Sensitivity analysis including Majambere 2010 with estimated ICC 0.1	3		Risk Ratio (Random, 95% CI)	0.80 [0.71, 0.89]
4.5 Sensitivity analysis excluding Majambere 2010 northern zones due to large baseline imbalance; estimated ICC 0.01	3		Risk Ratio (Random, 95% CI)	0.79 [0.71, 0.89]
5 Parasite prevalence with subgrouping by extent of aquatic habitat	3		Risk Ratio (Random, 95% CI)	0.88 [0.66, 1.16]
5.1 Habitats < 1 km ²	2		Risk Ratio (Random, 95% CI)	0.79 [0.71, 0.89]
5.2 Habitats > 1 km ²	1		Risk Ratio (Random, 95% CI)	1.15 [0.41, 3.20]
6 Mean haemoglobin level	1	3586	Mean Difference (IV, Fixed, 95% CI)	-0.13 [-0.40, 0.13]
6.1 Sensitivity analysis with estimated ICC 0.01	1	3586	Mean Difference (IV, Fixed, 95% CI)	-0.13 [-0.40, 0.13]


Analysis 1.1. Comparison 1 Larviciding versus no larviciding, Outcome 1 Malaria incidence (cluster-randomized controlled trial (cRCT)).




Analysis 1.2. Comparison 1 Larviciding versus no larviciding, Outcome 2 Malaria incidence with subgrouping by extent of aquatic habitat (non-randomized study (NRS)).




Analysis 1.3. Comparison 1 Larviciding versus no larviciding, Outcome 3 Parasite prevalence (cRCTs).





Analysis 1.4. Comparison 1 Larviciding versus no larviciding, Outcome 4 Parasite prevalence (NRS).






Analysis 1.5. Comparison 1 Larviciding versus no larviciding, Outcome 5 Parasite prevalence with subgrouping by extent of aquatic habitat.





Analysis 1.6. Comparison 1 Larviciding versus no larviciding, Outcome 6 Mean haemoglobin level.



ADDITIONAL TABLES

Table 1. Characteristics of larviciding

Study	Active ingredient, formulation, dose, and manufacturer	Frequency of application	Targeted aquatic habitats	Who carried out the larviciding	Vector species
Fillinger 2009	Commercial strains of <i>Bs</i> (water-dispersible, Valent BioSciences LLC)	Weekly intervals for first 6 months of the study	All water bodies	Project staff	<i>An gambiae s.l.</i> and <i>An funestus s.l.</i>

Table 1. Characteristics of larviciding (Continued)

	Commercial strains of <i>Bti</i> (water-dispersible, Valent BioSciences LLC)	Weekly intervals for remainder of the study			
Maheu-Giroux 2013a	Commercial strains of <i>Bs</i> (0.04 g/m ² , water-dispersible, Valent BioSciences LLC)	Weekly intervals	All open light-exposed water bodies	Community-owned resource person	<i>An gambiae</i> s.s., <i>An funestus</i> s.l., and <i>An coustani</i>
	Commercial strains of <i>Bti</i> (0.1 g/m ² , corn cob Valent BioSciences LLC)	Once every 3 months	All closed, covered, often highly polluted water bodies		
Majambere 2010	Commercial strains of <i>Bti</i> (0.2 kg/hectare, water-dispersible, Valent BioSciences LLC)	Weekly intervals	Areas of low vegetation across the Gambia river	Project staff using knapsack compression sprayers	<i>An gambiae</i> s.s., <i>An melas</i> , and <i>An arabiensis</i>
	Commercial strains of <i>Bti</i> (0.5 kg/hectare, corn cob, Valent BioSciences LLC)		Areas of high vegetation across the Gambia river	Project staff by hand	
Yapabandara 2001	Pyriproxyfen, S31183 (Adeal 0.5% G) applied at a rate of 0.01 mg active ingredient/L (2 g of granules/m ³)	3 applications: December 1994, June–July 1995, end of November 1995	Gem mining pits	Project staff	<i>An culicifacies</i> , <i>An subpictus</i> , and <i>An aruna</i>

 Abbreviations: *An*: *Anopheles*; *Bs*: *Bacillus sphaericus*; *Bti*: *Bacillus thuringiensis israeliensis*.

Table 2. Entomological inoculation rate from included studies

Study	Study arm	Mean numbers (95% CI)
Fillinger 2009 ^a	No larviciding	1.68 (1.16 to 2.43)
	Larviciding	0.39 (0.16 to 0.79)
Maheu-Giroux 2013a ^a	No larviciding	1.28
	Larviciding	0.683 (0.491 to 0.952)
Majambere 2010 (2006 data)	No larviciding North Zone	0
	Larviciding North Zone	0
	No larviciding South Zone	3.13
	Larviciding South Zone	5.82
Majambere 2010 (2007 data)	No larviciding North Zone	2.24
	Larviciding North Zone	2.32
	No larviciding South Zone	17.00

Larviciding to prevent malaria transmission (Review)

Table 2. Entomological inoculation rate from included studies (Continued)

Larviciding South Zone 3.91

Abbreviations: CI: confidence interval.

^aThere is a statistically significant difference between the study arms ($P < 0.05$).

APPENDICES

Appendix 1. Detailed search strategy

Search set	CIDG SR ^a	CENTRAL	MEDLINE	Embase	LILACS	CABS Abstracts
1	Mosquito*	Malaria [ti, ab, Mesh]	Malaria [ti, ab, Mesh]	Malaria [ti, ab, Emtree]	Mosquito\$	Mosquito*
2	Anopheles	Anopheles [Mesh]	Anopheles [ti, ab, Mesh]	Anopheles ti, ab, Emtree	Anopheles	Anopheles
3	malaria	Mosquito* ti, ab	Mosquito* ti, ab	Mosquito* ti, ab	malaria	malaria
4	1 or 2 or 3	Mosquito control [Mesh]	Mosquito control [Mesh]	Mosquito control ti, ab	1 or 2 or 3	1 or 2 or 3
5	Larvicid*	1 or 2 or 3 or 4	1 or 2 or 3 or 4	1 or 2 or 3 or 4	Larvicid\$ or larval or larva or larvae ti, ab	Larvicid* or larval or larva or larvae ti, ab
6	4 and 5	Larvicid* or larval or larva or larvae ti, ab	Larvicid* or larval or larva or larvae ti, ab	Larvicid* or larval or larva or larvae ti, ab	4 and 5	Bacillus thuringiensis
7	—	"Larval control" ti, ab	"Larval control" ti, ab	"Larval control" ti, ab	—	Bacillus sphericus
8	—	6 or 7	Bacillus thuringiensis [ti, ab, Mesh]	Bacillus thuringiensis ti, ab	—	Paris green
9	—	5 and 8	Bacillus sphericus ti, ab	Bacillus sphericus ti, ab	—	Temefos
10	—	—	Paris green ti, ab, sn	Paris green ti, ab	—	Pyriproxyfen or methoprene OR fenthion OR abate OR "surface oils" OR "surface films" OR chlorpyrifos OR pirimiphos-methyl OR di-

(Continued)

						flubenzuron OR novaluron OR spinosad
11	—	—	Temefos ti, ab, sn	Temefos ti, ab	—	Insect growth regulator*
12	—	—	(Pyriproxyfen or methoprene OR fenthion OR abate OR "surface oils" OR "surface films" OR chlorpyrifos OR pirimiphos-methyl OR diflubenzuron OR novaluron OR spinosad) ti, ab	(Pyriproxyfen or methoprene OR fenthion OR abate OR "surface oils" OR "surface films" OR chlorpyrifos OR pirimiphos-methyl OR diflubenzuron OR novaluron OR spinosad) ti, ab	—	Biological pest control
13	—	—	Juvenile hormones [Mesh]	Insect growth regulator* ti, ab	—	5-12/OR
14	—	—	Insect growth regulator* ti, ab	Biological pest control [Emtree]	—	4 AND 13
15	—	—	Pest Control, Biological [Mesh]	Larvicidal agent [Emtree]	—	—
16	—	—	6-15/OR	6-15/OR	—	—
17	—	—	5 AND 16	5 AND 16	—	—
18	—	—	—	—	—	—

^aCochrane Infectious Diseases Group Specialized Register.

WHAT'S NEW

Date	Event	Description
6 September 2019	Amended	Corrected link to Appendix 1 in review text

CONTRIBUTIONS OF AUTHORS

All authors contributed to the protocol design, wrote the protocol, and approved the final version.

LC and AW screened articles, extracted data from the included studies, and conducted the analysis.

SM acted as arbitrator.

All authors sought to interpret the data, and prepared and interpreted the GRADE summaries.

LC and AW wrote the full text and SM edit modified this. All authors read and approved the final manuscript.

DECLARATIONS OF INTEREST

LC has no known conflicts of interest.

SM is an independent consultant, and occasionally advises governments on malaria control. His main sources of consultancy work are from a research project funded by the Innovative Vector Control Consortium (IVCC). The title of the review is related to the use of insecticide

Larviciding to prevent malaria transmission (Review)

Copyright © 2019 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

applications for malaria vector control. The IVCC as an organization has a programme of working with industry on the development of novel insecticides and other vector control tools. He is a co-chair of the Larval Source Management workstream of the Roll Back Malaria's Vector Control Working Group. SM was a primary investigator and author of one study reviewed. He was not involved in the screening, data extraction, risk of bias assessment, and analysis for this particular study.

AW sits on the Innovative Vector Control Consortium (IVCC) External Scientific Advisory Committee 1 (ESAC1), which provides the IVCC management team with independent scientific advice on all projects proposed to the IVCC relating to the development of new products for malaria vector control.

SOURCES OF SUPPORT

Internal sources

- Liverpool School of Tropical Medicine, UK.

External sources

- Department for International Development, UK.

Project number 300342-104

- World Health Organization (WHO), Switzerland.

WHO Global Malaria Programme Agreement for Performance of Work (APW) Grant 2017 (number 709319)

- Partnership for Increasing the Impact of Vector Control (PIIVeC), UK.

Provided support to LC. PIIVeC is funded by the Medical Research Council of the UK (grant number MR/P027873/1) through the Global Challenges Research Fund.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We changed the title from 'Larviciding to control malaria' in the protocol ([Choi 2017](#)), to 'Larviciding to prevent malaria transmission'.

We amended the study inclusion criteria to allow inclusion of non-randomized cross-over trials.

In the published protocol, [Choi 2017](#), we stated that we would not present results from cluster-randomized controlled trials (cRCTs) that were not adjusted for clustering. In the review we presented unadjusted measures of effect and we have taken this into account in the GRADE assessments.

In the protocol we stated, "If there was considerable heterogeneity i.e. an I^2 statistic value of 75% to 100% or inconsistency in the direction of the effect, or both, then we did not perform a meta-analysis". We performed a meta-analysis in [Analysis 1.2](#) where heterogeneity was $I^2 = 82%$. We accounted for the heterogeneity in the GRADE assessments, and this is reflected in the 'Summary of findings' tables.

We added Silas Majambere as an author.

INDEX TERMS

Medical Subject Headings (MeSH)

Culicidae; Disease Reservoirs [*parasitology]; Ecosystem; Insecticides [*pharmacology]; Interrupted Time Series Analysis; Larva [drug effects]; Malaria [*prevention & control]; Mosquito Control [*methods]; Randomized Controlled Trials as Topic [methods]

MeSH check words

Animals; Humans