

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

disorders that have been most frequently associated with chronic kidney diseases and chronic non-transmitted diseases (diabetes mellitus, high blood pressure, etc.) Depression is an affective disorder which is associated with the emergence of the feeling of loss experienced by many patients with kidney the pandemic in Tarija.Patients suspectified to the feeling of loss experienced by many patients with kidney diseases, underwent immunochromatooperphi viscosticity in partients advoction states of the produces associated with the pandemic claim to the produce of the produ modality for peritoneal dialysis to hemurchisty ejaharateristic continuous the contaminate from renal modality for peritoneal dialysis to hemurchisty ejaharateristic continuous from the first the first to peritone from the peritone from the first to the first to prevent and take care formal health. The outcome of this modern from the first to the first study. 7 patients were excluded for notice meeting with the control of the study in which the study. with a period less than 3 months undergrang and the months are period to the months and the months and the months are period to the months are per who had changed the treatment modality in our interest in the second of the complete will general information on each were excluded too. Of the 63 hemodality is patients in datase was completed will general information on each were excluded too. Of the 63 hemodality sits patients in a patients in the patients of the patients who presented cover in the patients who presented cover in the patients who presented cover in the patients who were in the patients who were the patients of the patients who were in the patients who were the patients who were in the patients who were the patie miological discharge (Graph 1).Of the 6 patients who developed COVIDed 9tic is was assowere women and 3 were men, 4 patients with previous hope alone, some degree of suicidal ideation where women and 3 were men, 4 patients with previous hope alone, some degree of suicidal ideation who were dialyzed in the room with management and 2 patients

without liver infections who were dial					COV-19 negative N= 40 (91%)	COV-19 positive N= 4 (9%)	Total N= 44
patients evolved asympt	tomatic	during	Age	Average	51.75 (DS=14.8)	57.5 (DS= 10.08)	51.65 (DS=14.46)
			Gender	96	22% V / 78% M	100% M	20% V/80% M
Graphic 1	zh, bu						
Prevalence and mortality of COVID 19 in HD Tarija depa	ırtamental center	í	Mixed-race	%	98%	100%	98%
■ 1 ■ 2		1 one	Black	56	2%	0	2%
		'sis cat	Marital status				
		SIS Cat		%	52%	50%	52%
DEADS		ıed in	Single	96	43%	25%	41%
17 %	ica in		Widowed	56	5%	25%	7%
Healthy		Tabl	Grade of education				
Patients Presented COVID 19	■1 ■2	cs of s	Never assisted Primary	% %	2 50	75	2 52
91% RECOVERY				70 %	35	25	34
83%			College	96	13	0	12
			Reason for HD	70	15	0	12
		OVID no	P1	96	42%	75%	45%
			Hypertension	%	17%	25%	16%
Conclusions: The prevalence of patients infected by	SARS-CoV-2	MAL 57	GN	96	30%	0	30%
9% with a mortality of 17% (1 patient), validating		CONT.	PKD	%	8%	0	7%
control and presention measures become a great	Mean.	51.08	Other	96	3%	0	2%
massive spread (Selver Fation) prone to acquire this	ir@ections disc		Months in HD	average	32.7 months	14.5 months	31.9 months
No conflict of interest	magetious dis	36(42)	Px without depression	56	63	25	59
No conflict of interest Ethnicity			Px with depression	96	37	75	41
POS-553 Latin	%	98	Mild	96	53	0	44
POS-553 Latin	70	30	Moderade	%	20	33	28
PRESENCE AND SEVERITY OF COVID-19	%		Severe	56	27	67	28
PANDEMIC-ASSOCIATED DEPRESSION IN			Suicidal Ideation	96			
PATIENTS UNDERGOING HEMODIALYSIS		<u> </u>	Yes	96	15	0	14
THE TARIJA DEPARTMENTAL NEPHROLOG	A.%	35	No	%	85	100	86
		25	Interpersonality				
AND HEMODIALLY STS SCENTER 1 GROEVIA	%	25	Living in solitude (LIS)	%	25	0	23
ZAMORA, N*GMMuz, J1, Molina, A2, Miranda, A	3%	26	Living with family	96	75	100	77
¹ Hospital Regiopeir§an Juan de Dios, Nephrology, Tai			(LIS) with depression (LIS) without	% %	60 40	0	60 40
Bank Departmental Reference, Hematology, Tarija, Bo	ivia ³ Hospital	San	depression	79	40	0	40
Gabriel, Psychia tale Paz, Bolivia	%	9 c	onclusions: The	orevalenc	e of depression	Sompared to	other cohorts

Conclusions: The Operation of the patients who made a mindroduction: The Operation of the patients who made any individual, but a fatches in a very special way those who stiffer from both organic and mental diseases, such situations like the kind we usually live generally aggravate symptoms easily by presenting feelings of loneliness, fear, uncertainty or sadness. Anxiety and depression are the psychiatric