

ERRATUM

Open Access



Erratum to: Do epinephrine auto-injectors have an unsuitable needle length in children and adolescents at risk for anaphylaxis from food allergy?

Sten Dreborg¹, Xia Wen², Laura Kim³, Gina Tsai⁶, Immaculate Nevis⁴, Ryan Potts⁵, Jack Chiu⁶, Arunmozhi Dominic⁷ and Harold Kim^{6,7*}

Erratum to: *Allergy Asthma Clin Immunol* (2016) 12:11 DOI 10.1186/s13223-016-0110-8

After publication of the article [1] it was brought to our attention that the incorrect length of needle was used for one of the epinephrine auto-injectors (EAI), Jext[®]. The length of the needle of the Jext[®] 0.15 mg is not 15.7 mm as mentioned in the paper, but 13 mm. After correcting the needle length for Jext[®], the results of the study do not change aside for those involving Jext[®]. The results of the primary outcome variable for Jext[®] are now the same as the other high pressure EAIs (HPEAIs) and do not lead to intra-osseous injections 38% but 11% between 15 and 30 kg. This is the same risk as the Epipen[®] and the Auvi-Q[®].

Author details

¹ Women's and Children's Health, University of Uppsala, Uppsala, Sweden. ² Faculty of Science, McGill University, Montreal, Canada. ³ Faculty of Medicine, University British Columbia, Vancouver, Canada. ⁴ Goodman School of Business, Brock University, St. Catharines, Canada. ⁵ Farncombe Family Digestive Health Unit, McMaster University, Hamilton, Canada. ⁶ Department of Medicine, Western University, London, Canada. ⁷ Department of Medicine, McMaster University, Hamilton, ON, Canada.

The online version of the original article can be found under doi:[10.1186/s13223-016-0110-8](https://doi.org/10.1186/s13223-016-0110-8).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 27 June 2017 Accepted: 27 June 2017

Published online: 07 July 2017

Reference

1. Dreborg S, Wen X, Kim L, Tsai G, Nevis I, Potts R, Chiu J, Dominic A, Kim H. Do epinephrine auto-injectors have an unsuitable needle length in children and adolescents at risk for anaphylaxis from food allergy? *Allergy Asthma Clin Immunol*. 2016;12:1. doi:[10.1186/s13223-016-0110-8](https://doi.org/10.1186/s13223-016-0110-8).

*Correspondence: hkimkw@gmail.com

⁷ Department of Medicine, McMaster University, Hamilton, ON, Canada
Full list of author information is available at the end of the article