

Research

Open Access

Iloprost-induced desensitization of the prostacyclin receptor in isolated rabbit lungs

Ralph T Schermuly*¹, Soni S Pullamsetti¹, Susanne C Breitenbach¹, Norbert Weissmann¹, Hossein A Ghofrani¹, Friedrich Grimminger¹, Sigrid M Nilius², Karsten Schrör², Jutta Meger-Kirchrath², Werner Seeger¹ and Frank Rose¹

Address: ¹University of Giessen Lung Center (UGLC), Medical Clinic II/V, Justus-Liebig-University Giessen, 35392 Giessen, Germany and ²Institut fuer Pharmakologie und Klinische Pharmakologie, Heinrich-Heine-Universitaet Duesseldorf, 40225 Duesseldorf, Germany

Email: Ralph T Schermuly* - Ralph.Schermuly@innere.med.uni-giessen.de; Soni S Pullamsetti - Soni.Pullamsetti@innere.med.uni-giessen.de; Susanne C Breitenbach - Christiane.Breitenbach@innere.med.uni-giessen.de; Norbert Weissmann - Norbert.Weissmann@innere.med.uni-giessen.de; Hossein A Ghofrani - Ardeschir.Ghofrani@innere.med.uni-giessen.de; Friedrich Grimminger - Friedrich.Grimminger@innere.med.uni-giessen.de; Sigrid M Nilius - Nilius@uni-duesseldorf.de; Karsten Schrör - kschroer@uni-duesseldorf.de; Jutta M Kirchrath - meyerj@uni-duesseldorf.de; Werner Seeger - Werner.Seeger@innere.med.uni-giessen.de; Frank Rose - Frank.Rose@innere.med.uni-giessen.de

* Corresponding author

Published: 26 January 2007

Received: 24 February 2006

Respiratory Research 2007, **8**:4 doi:10.1186/1465-9921-8-4

Accepted: 26 January 2007

This article is available from: <http://respiratory-research.com/content/8/1/4>

© 2007 Schermuly et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The rapid desensitization of the human prostacyclin (IP) in response to agonist binding has been shown in cell culture. Phosphorylation of the IP receptor by protein kinase C (PKC) has been suggested to be involved in this process.

Methods and results: In this study we investigated the vasodilatory effects of iloprost, a stable prostacyclin analogue, in perfused rabbit lungs. Continuous infusion of the thromboxane mimetic U46619 was employed to establish stable pulmonary hypertension. A complete loss of the vasodilatory response to iloprost was observed in experiments with continuous iloprost perfusion, maintaining the intravascular concentration of this prostanoid over a 180 min period. When lungs under chronic iloprost infusion were acutely challenged with inhaled iloprost, a corresponding complete loss of vasoreactivity was observed. This desensitization was not dependent on upregulation of cAMP-specific phosphodiesterases or changes in adenylate cyclase activity, as suggested by unaltered dose-response curves to agents directly affecting these enzymes. Application of a prostaglandin E1 receptor antagonist 6-isopropoxy-9-oxoxanthene-2-carboxylic acid (AH 6809) or the PKC inhibitor bisindolylmaleimide I (BIM) enhanced the vasodilatory response to infused iloprost and partially prevented tachyphylaxis.

Conclusion: A three-hour infusion of iloprost in pulmonary hypertensive rabbit lungs results in complete loss of the lung vasodilatory response to this prostanoid. This rapid desensitization is apparently not linked to changes in adenylate cyclase and phosphodiesterase activation, but may involve PKC function and co-stimulation of the EPI receptor in addition to the IP receptor by this prostacyclin analogue.

Background

Prostacyclin (PGI₂) is the major product of cyclooxygenases (COX) in the vascular endothelium and mediates potent anti-platelet, vasodilator, and anti-inflammatory actions by a prostacyclin receptor (IP) [1]. This receptor is a member of the G protein-coupled receptor (GPCR) superfamily and is coupled to adenylate cyclase (AC) and phospholipase C (PLC) [2-4]. The prostanoid receptors are classified into DP, IP, EP (EP 1-4), FP and TP receptors [5-7] with different affinities for agonists and different roles in signal transduction. The IP, EP2, EP4 and DP receptors are coupled to stimulation of adenylate cyclase, while the TP, EP1 and FP receptors are coupled to Ca²⁺ mobilization. The EP3 receptor is an alternative spliced gene, with at least 8 isoforms identified so far. Depending on the subtype, this receptor can be negatively or positively coupled to Gs [8,9]. Agonist (PGI₂ or PGI₂ mimetics such as cicaprost, iloprost, carbacyclin, and prostaglandin E1)-binding to the IP receptor leads to activation of protein kinase A (PKA) by cyclic adenosine monophosphate (cAMP) [10]. Although, the above mentioned agonist specificities for IP receptors vary and none of them were highly selective for IP receptors alone. The affinity of cicaprost for human IP receptors is merely 3-fold higher than for the prostaglandin E2 EP4 subtype of receptor, which also couples to Gs [11], and only 17-fold higher than for the EP3 receptor in the mouse [12]. Iloprost is equipotent at both human and mouse IP and EP1 receptors, and carbacyclin and prostaglandin E1 show even greater affinity for EP3 than for IP receptors [11,12].

Disturbances to PGI₂ synthesis [13,14], as well as polymorphisms of the PGI₂-synthase (PGIS) [15], have been linked to severe pulmonary hypertension. Substitution of PGI₂, either by overexpression of the PGIS [16] in an experimental model or by application of PGI₂ [17,18] or its analogues iloprost [19,20] or beraprost [21,22] in patients decreases pulmonary artery pressure. However, tolerance of the lung vasodilatory response to continuously infused PGI₂ rapidly develops in patients with severe pulmonary hypertension, and dose adjustments have to be made [17,23]. In COPD patients with pulmonary hypertension, the pulmonary vasodilatory response to continuously infused PGI₂ was found to dissipate within 24 h [24]. Marked tolerance to the anti-mitogenic actions of PGI₂ developed within 24 h in coronary artery smooth muscle cells [25]. Basic studies demonstrated that desensitization of the IP receptor occurs within minutes after exposure to agonists and is due to agonist-induced receptor phosphorylation, mainly mediated by PKC [26], with subsequent sequestration of the intact receptor and removal from the cell surface [26,27]. In addition, there is evidence for changes in adenylate cyclase and phosphodiesterase (PDE) activation occurring in response to IP receptor stimulation, which may contribute to loss of the

vasodilatory response to PGI₂ and its analogues [28-30]. In the present study, we employed the model of pulmonary hypertensive perfused rabbit lungs to investigate the dose-response relationship and features of tolerance development upon continuous iloprost infusion in the intact lung vasculature.

Materials and methods

Materials

Sterile Krebs-Henseleit buffer (KHB) was obtained from Serag-Wiessner (Naila, Germany). The thromboxane-A₂ mimetic U46619 was supplied by Paesel-Lorei (Frankfurt, Germany) and iloprost by Schering (Berlin, Germany). All other chemicals were purchased from Merck (Darmstadt, Germany).

Isolated lung model

The perfused rabbit lung model has previously been described in detail [31]. Briefly, rabbits of either sex weighing 2.6 to 2.9 kg were anticoagulated with heparin (1000 U/kg) and anaesthetized with intravenous ketamine/xylazine. Tracheostomy was performed and the animals were ventilated with room air. After mid-sternal thoracotomy, catheters were placed into the pulmonary artery and the left atrium, and perfusion with KHB was started. For washout of blood, perfusate was initially not recirculated and the lungs were removed without interruption of ventilation and perfusion. The lungs were placed in a temperature-equilibrated chamber at 37.5°C, freely suspended from a force transducer for monitoring of organ weight. In a recirculating system the flow was slowly increased to 120 ml/min (total volume 350 ml). Left atrial pressure was set at 2 mmHg in all experiments. In parallel with onset of artificial perfusion, an air mixture of 80.5% N₂, 15% O₂ and 4.5% CO₂ was used for ventilation. Tidal volume (11 ml/kg) and frequency (10 - 13 breaths/min) were adapted to maintain the pH of the recirculating buffer in the range between 7.35 and 7.37. A positive end-expiratory pressure of 1 mmHg was used throughout. The PO₂ and PCO₂ values in the post-lung buffer fluid ranged between 100 and 120 mmHg and 38 and 43 mmHg, respectively. Pressures in the pulmonary artery, the left atrium and the trachea were registered by means of small diameter tubing threaded into the perfusion catheters and the trachea and connected to pressure transducers (zero referenced at the hilum). The whole system was heated to 37.5°C.

Lungs included in the study had 1) a homogeneous white appearance with no signs of hemostasis, edema or atelectasis; 2) pulmonary artery and ventilation pressures in the normal range; and were 3) isogravimetric (lung weight gain < 0.2 g/h) during an initial steady state period of at least 45-60 min.

Experimental protocols**Isolated lung experiments**

As described previously [32], a sustained increase of P_{PA} from ≈ 6 to ≈ 24 mmHg was achieved by continuous infusion of the thromboxane-mimetic U46619 with a dose range of 70–160 pmol·kg⁻¹·min⁻¹. Individual titration was performed. This level of pulmonary hypertension was then maintained for at least 320 min with variations in P_{PA} of less than 2 mmHg. The following experimental groups were employed:

Control lungs (n = 4): No interventions were undertaken.

U46619-infused lungs (n = 4): After termination of the steady state period, U46619 was continuously infused for 320 min to provoke an increase in P_{PA} to ≈ 24 mmHg.

Iloprost infusion (n = 4): U46619 was administered as described. 20 min after onset of U46619 infusion, iloprost was applied by a bolus injection of 200 ng, followed by a continuous infusion of 33 ng/kg/h to achieve stable buffer concentrations in the range of 350 pg/ml.

Intravenous PGI₂ application (n = 4): U-46619 was administered as described. 20 min after onset of U-46619 infusion, PGI₂ was continuously applied at a dose of 50 ng·kg⁻¹·min⁻¹ until termination of the experiments.

Rolipram i.v. (n = 4): After adjusting stable pulmonary hypertension, the PDE4 inhibitor rolipram was applied in increasing doses (dose range 0.001–10 μM).

Iloprost infusion and rolipram i.v. (n = 4): Iloprost was applied by a bolus injection of 200 ng, followed by a continuous infusion of 33 ng/kg/h. 240 min after starting the iloprost infusion, rolipram was applied in increasing doses (dose range 0.001–10 μM).

Motapizone i.v. (n = 4): After adjusting stable pulmonary hypertension, the PDE3 inhibitor motapizone was applied in increasing doses (dose range 0.01–100 μM).

Iloprost infusion and motapizone i.v. (n = 4): Iloprost was applied by a bolus injection of 200 ng, followed by a continuous infusion of 33 ng/kg/h. 240 min after starting the iloprost infusion, motapizone was applied in increasing doses (dose range 0.01–100 μM).

Iloprost inhalation (n = 4): When stable pulmonary hypertension was achieved, iloprost was nebulized (deposited dose 75 ng) by means of an ultrasonic nebulizer as described previously [33,34].

Iloprost infusion and iloprost inhalation (n = 4): U46619 was continuously infused and iloprost was applied by a bolus

injection of 200 ng, followed by a continuous infusion of 33 ng/kg/h. 240 min after starting the iloprost infusion, iloprost was nebulized (deposited dose 75 ng).

Forskolin i.v. (n = 4): After adjusting stable pulmonary hypertension, the adenylate cyclase stimulator forskolin was applied in increasing doses (dose range 10–100 μM).

Iloprost infusion and forskolin i.v. (n = 4): Iloprost was applied by a bolus injection of 200 ng, followed by a continuous infusion of 33 ng/kg/h. 240 min after starting the iloprost infusion, forskolin was applied in increasing doses (dose range 10–100 μM).

Bisindolylmaleimide I (BIM) infusion (n = 4): BIM (1 μM) was applied by a bolus injection after stable adjustment of pulmonary hypertension with U46619. As described previously, stable buffer levels of BIM are achieved by a single application in the setup of isolated perfused rabbit lungs [35].

Iloprost infusion in presence of BIM (n = 4): U46619 was administered as described. After stable adjustment of pulmonary hypertension, BIM (1 μM) was applied by a bolus injection. Iloprost was bolus injected and infused as described above.

6-isopropoxy-9-oxoxanthene-2-carboxylic acid (AH 6809) infusion (n = 4): AH 6809 (3 μM) was applied by a bolus injection after stable adjustment of pulmonary hypertension with U46619. As described previously, AH6809 demonstrated long acting EP1 blocking activity in fluid perfused small intestine segments after a single application [36].

Iloprost infusion in presence of AH 6809 (n = 4): U46619 was administered as described. 45 min after onset of U46619 infusion, AH 6809 (3 μM) was applied as a bolus injection and the bolus injection plus infusion of iloprost was started as in the preceding groups.

Cell culture

Rabbit pulmonary smooth muscle cells (PSMC) were prepared as described. Primary SMC were isolated from rabbit pulmonary artery by carefully preparing <1 mm³ pieces of media, devoid of adventitial tissue as assessed by microscopic control. The pieces of media were placed into 12-well cell culture plates with 500 μl culture medium. The isolated pulmonary smooth muscle cells (PSMC) identity was verified by characteristic appearance in phase-contrast microscopy, indirect immunofluorescent antibody staining for smooth muscle-specific isoforms of α-actin and myosin (at least 95% of cells stained positive), and lack of staining for von Willebrand factor and vimentin, indicating that the cultures did not contain significant

numbers of endothelial cells or adventitial fibroblasts [37]. Smooth muscle cells were grown in Dulbecco's modified Eagle's high glucose medium supplemented with 10% fetal bovine serum, 100 µg/ml streptomycin and 100 units/ml penicillin.

Desensitization assays and cAMP measurement

Smooth muscle cells were grown to 90% confluence in 24-well plates, as previously described. For desensitization assays, cells were treated with iloprost for the indicated times. Then, cells were washed three times with HBSS and preincubated in HBSS containing 1 mg/ml BSA, 10 mM HEPES (pH 7.3) and 1 mM IBMX for 10 min at 37°C. Afterwards cells were again stimulated with iloprost (100 nM) or forskolin (10 µM) for 10 min and cAMP measurements were performed as described below. Reactions were stopped by aspiration and addition of ice-cold 96% ethanol. Dried samples were overlaid with 300 µl RIA-buffer (150 mM NaCl, 8 mM Na₂HPO₄, 2 mM NaH₂PO₄, pH 7.4) and frozen overnight at -80°C. cAMP in the supernatant was determined by radioimmunoassay [Steiner et al., 1972]. Protein determination was performed according to the method of Bradford. Cells treated with vehicle alone (no ligand) or forskolin (50 µM) were served as negative and positive controls respectively for the cAMP measurement.

Data analysis

Values are means ± SEM. Student's t-test was performed for comparison of two values. For multiple comparisons, ANOVA was followed by post hoc test (Bonferroni/Dunn method). Statistical significance was considered at $p < 0.05$.

Results

Baseline conditions

After termination of the steady state period, all lungs displayed P_{PA} values in the range between 5 and 7 mmHg.

U46619-elicited pulmonary hypertension

Continuous infusion of U46619 provoked an increase in P_{PA} to 23.4 ± 0.9 mmHg within 25 min, with subsequent plateau of the pulmonary hypertension. This level of pulmonary hypertension was then maintained for at least 300 min with variations in P_{PA} of less than 2 mmHg. Total weight gain at the end of the experiments was 6.5 ± 2.7 g.

U46619-elicited pulmonary hypertension and infusion of iloprost or PGI₂

As in the preceding group, U46619 infusion for establishing stable pulmonary hypertension was performed. Bolus injection of 200 ng iloprost was followed by an infusion of 33 ng iloprost per hour. This procedure resulted in a significant vasodilatory response with P_{PA} values decreasing by a mean of 3.3 mmHg. Similarly, the continuous

infusion of a more selective IP receptor ligand, PGI₂ also resulted in significant vasodilatory responses (Fig 1). Stable iloprost buffer concentrations in the range of 350–380 pg/ml were documented over the entire perfusion period. A slow increase in P_{PA} toward pre-infusion values was noted to commence after 100–120 min, and 200 min after beginning of the iloprost infusion, the vasodilatory effect was virtually fully lost. The total weight gain of the lungs was 5.3 ± 3.2 g, and thus did not differ from the groups with mono U46619 infusion.

Dose-effect curves of PDE-inhibitors with or without infusion of iloprost

As shown in Fig. 2A, the intravascular administered PDE3 inhibitor motapizone (dose range 0.01–100 µM) effected a dose-dependent reduction of the elevated P_{PA} values in lungs with U46619-elicited pulmonary hypertension. There was no shift in the dose response curve when applying identical doses of motapizone 240 min after onset of iloprost infusion. Similar experiments were performed with the PDE4 inhibitor rolipram, which dose-dependently reduced pulmonary artery pressure in U46619 pre-constricted lungs, with identical dose-effect curves in the absence and presence of iloprost infusion (Fig. 2B). Total weight gain was 5.7 ± 2.9 g and 5.3 ± 1.9 g (iloprost infusion) in the motapizone treated lungs and 4.6 ± 2.2 g and 5.0 ± 2.6 g (iloprost infusion) in the rolipram treated lungs, respectively.

Nebulization of iloprost

Inhalation of aerosolized iloprost (total dose 75 ng inhaled over 10 min) resulted in a significant reduction of the U46619-induced pulmonary hypertension, with PAP values decreasing by a mean of 6.3 mmHg (Fig. 3). In separate experiments, iloprost was infused in advance for 240 min, and then nebulization was started. There was virtually no response to inhaled iloprost after this preceding iloprost infusion period. The total weight gain of the lungs was 6.7 ± 2.9 g at the end of experiments.

Dose-effect curves of forskolin

As shown in Fig. 4, the adenylate cyclase activator forskolin (dose range 10–300 µM) reduced the U46619-elicited pulmonary hypertension in a dose dependant manner. After iloprost infusion for 240 min, an identical dose response curve was observed. Total weight gain was 5.6 ± 2.1 g at the end of the experiments.

Infusion of iloprost: effects of BIM

As in the preceding groups, U46619 infusion for establishing stable pulmonary hypertension was performed. Subsequently, the PKC inhibitor BIM at a dose of 1 µM was applied alone or in combination with iloprost. In combination group, after 5 min of BIM application, bolus injection of 200 ng iloprost was followed by an infusion




Figure 1
Prostacyclin receptor (IP) desensitization in U46619-induced pulmonary hypertension in intact rabbit lungs in response to PGI₂ or iloprost infusion. Pulmonary hypertension was adjusted by continuous infusion of the thromboxane mimetic U46619. After a bolus injection of 200 ng iloprost followed by an infusion of 33 ng iloprost per hour (stable buffer levels of ~350 pg/ml) or continuous infusion of PGI₂ (50 ng/min) is administered and measured their influence on U46619-elicited pulmonary hypertension.

of 33 ng iloprost per hour. This procedure resulted in a significant vasodilatory response with P_{PA} values decreasing by a mean of 4.3 mmHg (Fig. 5A). However, the sole BIM application demonstrated no effects on U46619-elicited pulmonary hypertension (Fig. 5A). The calculated area under the curve (AUC) of the pressure curve was significantly increased as compared to sole iloprost infusion (Fig. 6). The loss of the vasodilatory response to iloprost was somewhat retarded, but not prevented, as compared to single iloprost application. The total weight gain of the lungs was 5.8 ± 2.5 g.


Infusion of iloprost: effects of AH 6809

After adjusting stable pulmonary hypertension, the EP1 receptor antagonist AH 6809 at a dose of 3 μ M was

applied alone or in combination with iloprost, without effecting pulmonary artery pressure. In combination group, after 5 min of AH 6809 application, a bolus injection of 200 ng iloprost was followed by an infusion of 33 ng iloprost per hour. The maximum vasodilatory response to iloprost was 11.4 mmHg (Fig. 5B) and the area under the curve (AUC) of the pressure curve was 305 mmHg \times min ($p < 0.05$ versus sole iloprost infusion) (Fig. 6). The total weight gain of the lungs was 4.8 ± 2.0 g. In contrary, the AH 6809 application alone had no effect on U46619-induced tension (Fig. 5B).

Measurement of cyclic AMP

We assayed the desensitization kinetics of iloprost and forskolin in rabbit pulmonary smooth muscle cells (Fig.

**Figure 2**

Dose effect curve of the PDE3 inhibitor motapizone or PDE4 inhibitor rolipram on U46619-elevated pulmonary artery pressure in untreated and desensitized (post iloprost infusion) lungs. Motapizone and rolipram dose-dependently reduced the elevated pulmonary artery pressure. There was no change in the dose response curve when applying identical doses of motapizone 240 min after iloprost infusion. (A) The decrease in pulmonary artery pressure (Δ PAP, as decrease of U46619-induced pulmonary hypertension) in response to motapizone or (B) rolipram (concentrations related to the recirculating perfusate) is given. (mean \pm SEM of four independent experiments each; SEM bars are not included when smaller than symbol). *: $p < 0.05$ as compared to U46619-elevated pulmonary artery pressure.

7). Incubation of cells with iloprost led to a time-dependent reduction of iloprost induced cAMP formation to 88% (1 h), 71% (2 h), 76% (3 h), 66% (5 h), 54% (7 h) and 26% (24 h) of control. Incubation of cells with iloprost and stimulation of cells with forskolin led to a reduction of cAMP formation to 83% (1 h), 84% (2 h), 91% (3 h), 83% (5 h), 79% (7 h) and 75% (24 h) of control.

Discussion

The desensitization of the IP receptor after incubation of platelets [38,39] or IP receptor overexpressing cells [26,27,40,41] with PGI₂ or its analogues in vitro is a well-known phenomenon. Tolerance development of the lung vasodilatory response to PGI₂ has also been observed in

patients with severe pulmonary hypertension being chronically treated by intravenous infusion of the prostanoid [23,24]. In the present study we show that such desensitization occurs within 180 min in perfused rabbit lungs.

Desensitization is not dependant on increased PDE activity

Phosphodiesterases (PDE) are enzymes that hydrolyze cyclic AMP and cyclic GMP, the second messengers of PGI₂, and NO [42,43]. The characterization of the various PDEs currently known has profited from the employment of selective PDE inhibitors. Concerning the lung vasculature, the presence of the PDE isoenzymes 1, 3, 4 and 5 in the cytosolic and particulate phases (homogenized




Figure 3

Effect of inhaled iloprost on U46619-elevated pulmonary artery pressure in untreated and desensitized (post iloprost infusion) lungs. Inhalation of aerosolized iloprost for 10 min (deposited dose 75 ng) resulted in a significant reduction of the U46619-induced pulmonary hypertension, with PAP values decreasing by a mean of 6.3 mmHg. The vasodilatory potency of inhaled iloprost was completely lost in the desensitization group (iloprost infusion for 240 min). *: $p < 0.05$ as compared to U46619-elevated pulmonary artery pressure.

human pulmonary artery tissue) has been demonstrated [44]. There is strong evidence of PDE activation and upregulation in response to intracellular elevation of cAMP. It has been shown recently that cyclic AMP upregulates isoforms of PDE4 in airway smooth muscle cells [45], monocytes [46], endothelial cells [47] and Jurkat T-cells [48]. In addition, short term activation (phosphorylation) of PDE4 via PKA was demonstrated by treatment of cells with forskolin, a stimulator of adenylate cyclase [49]. Furthermore, treatment of rats with 7-oxo-prostacyclin resulted in a significant increase in PDE4 activity, which attenuated the elevation of isoprenaline-induced cAMP levels as well as the contractile force development in a Langedorff-heart preparation [50]. Similar results were obtained for PDE3 which has been shown to be regulated on the level of activity [48] and expression [32] by

increased cAMP levels. In the current study, we employed motapizone and rolipram, selective inhibitors of PDE3 and PDE4, respectively and performed dose response studies on vasodilation in normal and desensitized (iloprost-treated) isolated rabbit lungs. As reported earlier, these PDE inhibitors are potent pulmonary vasodilators [51] and we found no shift in the dose response curve in desensitized lungs. Thus, there is no evidence of increased PDE activity in iloprost-treated isolated rabbit lungs.

Desensitization is not dependant on adenylate cyclase activity

We employed forskolin as direct activator of adenylate cyclase and performed dose response studies on pulmonary vasodilation in isolated rabbit lungs. The response to forskolin in the desensitized (iloprost-treated) organs




Figure 4

Dose effect curve of the adenylate cyclase activator forskolin on U46619-elevated pulmonary artery pressure in untreated and desensitized (post iloprost infusion) lungs. Forskolin dose-dependently reduced the elevated pulmonary artery pressure. There was no change in the dose response curve when applying identical doses of forskolin 240 min after iloprost infusion. The decrease in pulmonary artery pressure (ΔPAP , as decrease of U46619-induced pulmonary hypertension) in response to forskolin (concentrations related to the recirculating perfusate) is given (mean \pm SEM of 4 independent experiments each; SEM bars are not included when smaller than symbol). *: $p < 0.05$ as compared to U46619-elevated pulmonary artery pressure.

remained intact (and seemed to be slightly increased), indicating no regulation of adenylate cyclase. Similar results were obtained in rats chronically treated with albuterol in which acetylcholine (ACh)-induced bronchoconstriction was reversible by forskolin, but not by albuterol itself, suggesting that the intracellular site of desensitization is upstream of adenylate cyclase [27].

Moreover, the cAMP production of iloprost-pretreated rabbit pulmonary smooth muscle cells in response to forskolin was also not significantly reduced, except in 24 hours of iloprost-pretreated group (to 75% of control). This is quite contrary to iloprost responses, where a significantly reduced cAMP levels (to 25% of control) was observed, thereby excluding the down-regulation of ade-

nylate cyclase as a possible reason for reduced cAMP formation after continuous stimulation of cells with iloprost. In this context, desensitization of IP receptors by iloprost may possibly explained by its direct influence on the receptor, a mechanism upstream of adenylate cyclases. This involves rapid receptor phosphorylation, which results in uncoupling of receptor-G protein interactions and subsequent receptor internalization. These events result in a diminished response to agonist [26]. This may also be due to preferential reduction of domain-bound pool of the cognate Gproteins $\text{Gs}\alpha\text{L}$ and $\text{Gs}\alpha\text{S}$ [52].

On the other hand, iloprost induced vasorelaxation responses may also be via cAMP-independent manner. Consistent with the observation, PGI_2 analogues iloprost




Figure 5
Influence of iloprost infusion and its combination with AH 6809 and BIM on U46619-elicited pulmonary hypertension. Pulmonary artery pressure (PAP) is given (mean \pm SEM of 4 independent experiments each, SEM bars are missing when enclosed by the symbol). (A) The prostaglandin E_1 receptor antagonist AH 6809 alone or in combination with iloprost (pre-applied followed by an iloprost infusion). (B) The PKC inhibitor BIM alone or in combination with iloprost (pre-applied followed by an iloprost infusion).

were shown to activate MaxiK and K_{IR} channel and subsequent vasorelaxation through a cAMP-independent, G(s)-protein mediated mechanism in vascular smooth muscle cells [53,54]. Concomitantly in guinea-pig aorta, cyclic AMP-dependent and -independent pathways were recently proposed to underlie the relaxation induced by the PGI_2 analogues, iloprost and Beraprost [55-57]. In contrast, in rat tail artery, it is thought that channel phosphorylation by cyclic AMP-dependent PKA is responsible for the MaxiK channel-mediated vascular relaxation due to IP receptor stimulation by iloprost [58,59]. This is well in line with investigations in human fibroblasts, where cells were challenged at any time point of iloprost pre-treatment with forskolin and fully maintained their cAMP response [30]. In contrast, Sobolewski et al., demonstrated in rat pulmonary smooth muscle cells (PSMCs)

which were exposed to cicaprost, another PGI_2 analogue, a significant reduction in forskolin-induced cAMP production [60] as well as a decrease in adenylate cyclase isoform 2 and 5/6 expression after one hour of exposure to this prostanoid. However, both isoforms were expressed at control levels when PSMCs were incubated for longer time periods with cicaprost. It is possible that the agonists employed, as well as species differences are responsible for these conflicting data.

Iloprost influences the EPI receptor in addition to the IP receptor

It is well known that iloprost, the clinically approved compound for treatment of thromboangiitis obliterans [20] and pulmonary arterial hypertension [5], is not highly selective for the IP receptor but also activates the




Figure 6
Influence of iloprost infusion and its combination with AH 6809 and BIM on the area under the curve of the U46619-elicited pulmonary hypertension. Area under the curve (AUC) is given (mean ± SEM of 4 independent experiments each). The AUC was calculated by standard techniques and is given as mmHg × min. **: $p < 0.01$; ***: $p < 0.001$ as compared to AUC of sole iloprost infusion.


EP1receptor which is coupled to Gq and signals via IP_3 generation and increased intracellular calcium (Ca^{2+}) [61]. To investigate the effects of possible involvement of the EP1receptor we employed the EP1receptor antagonist AH 6809 and found a dramatic increase in the efficacy of intravenous iloprost. In contrary, the AH 6809 application alone had no effect on U46619-induced tension. Thus, the agonistic effects on EP1receptor explain the differences between iloprost and selective IP agonists in inhibition of smooth muscle cell response [26,40].

PKC inhibition and IP receptor desensitization

There is evidence that the initial desensitization process is dependent on phosphorylation of the IP receptor by PKC [41,62]. After phosphorylation, the IP receptor is sequestered to clathrin-coated vesicles and by a dynamin-dependant process internalized into endosomes [41]. This

receptor sequestration is, unlike desensitization, PKC-independent [27] and a time of approximately 18 h is needed for a resensitization in human fibroblasts [27,30]. However, several studies including our work employed inhibitors of PKC (GF-109203X, BIM) and could not or could only partially prevent receptor desensitization, suggesting a minor role of PKC in the desensitization process.

In conclusion, this study demonstrates for the first time in an intact lung model that the PGI_2 analogue iloprost caused desensitization of the IP receptor within 2 to 3 hours, resulting in complete loss of responsiveness after this time period. Desensitization was not due to increased PDE activity or loss of adenylate cyclase activity, but may involve PKC function and co-stimulation of the EP1 receptor in addition to the IP receptor by this PGI_2 analogue.

**Figure 7**

Prostacyclin receptor (IP) desensitization in rabbit pulmonary smooth muscle cells. Cells were continuously stimulated with 100 nM iloprost for the indicated preincubation times. cAMP response to iloprost (100 nM) or forskolin (10 μ M) was assayed in these cells and is given as a percent of the non-desensitized control. Values are expressed as mean \pm SEM of three independent experiments performed in triplicate. *: $p < 0.05$; ***: $p < 0.001$ as compared to non-desensitized control.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (SFB 547), Project C6.

References

- Vane JR, Botting RM: **Pharmacodynamic profile of prostacyclin.** *Am J Cardiol* 1995, **75**:3A-10A.
- Boie Y, Rushmore TH, Darmon-Goodwin A, Grygorczyk R, Slipetz DM, Metters KM, Abramovitz M: **Cloning and expression of a cDNA for the human prostanoid IP receptor.** *J Biol Chem* 1994, **269**:12173-12178.
- Coleman RA, Smith WL, Narumiya S: **International Union of Pharmacology classification of prostanoid receptors: properties, distribution, and structure of the receptors and their subtypes.** *Pharmacol Rev* 1994, **46**:205-229.
- Namba T, Oida H, Sugimoto Y, Kakizuka A, Negishi M, Ichikawa A, Narumiya S: **cDNA cloning of a mouse prostacyclin receptor. Multiple signaling pathways and expression in thymic medulla.** *J Biol Chem* 1994, **269**:9986-9992.
- Breyer RM, Bagdassarian CK, Myers SA, Breyer MD: **Prostanoid receptors: subtypes and signaling.** *Annu Rev Pharmacol Toxicol* 2001, **41**:661-690.
- Negishi M, Sugimoto Y, Ichikawa A: **Molecular mechanisms of diverse actions of prostanoid receptors.** *Biochim Biophys Acta* 1995, **1259**:109-119.
- Narumiya S, FitzGerald GA: **Genetic and pharmacological analysis of prostanoid receptor function.** *J Clin Invest* 2001, **108**:25-30.
- Narumiya S, Sugimoto Y, Ushikubi F: **Prostanoid receptors: structures, properties, and functions.** *Physiol Rev* 1999, **79**:1193-1226.
- Hata AN, Breyer RM: **Pharmacology and signaling of prostaglandin receptors: multiple roles in inflammation and immune modulation.** *Pharmacol Ther* 2004, **103**:147-166.
- Wise H, Jones RL: **Focus on prostacyclin and its novel mimetics.** *Trends Pharmacol Sci* 1996, **17**:17-21.
- Abramovitz M, Adam M, Boie Y, Carriere M, Denis D, Godbout C, Lamontagne S, Rochette C, Sawyer N, Tremblay NM, Belley M, Gallant M, Dufresne C, Gareau Y, Ruel R, Juteau H, Labelle M, Ouimet N, Metters KM: **The utilization of recombinant prostanoid receptors to determine the affinities and selectivities of prostaglandins and related analogs.** *Biochim Biophys Acta* 2000, **1483**:285-293.
- Kiryama M, Ushikubi F, Kobayashi T, Hirata M, Sugimoto Y, Narumiya S: **Ligand binding specificities of the eight types and subtypes of the mouse prostanoid receptors expressed in Chinese hamster ovary cells.** *Br J Pharmacol* 1997, **122**:217-224.
- Christman BW, McPherson CD, Newman JH, King GA, Bernard GR, Groves BM, Loyd JE: **An imbalance between the excretion of thromboxane and prostacyclin metabolites in pulmonary hypertension.** *N Engl J Med* 1992, **327**:70-75.
- Tuder RM, Cool CD, Geraci MW, Wang J, Abman SH, Wright L, Badesch D, Voelkel NF: **Prostacyclin synthase expression is decreased in lungs from patients with severe pulmonary hypertension.** *Am J Respir Crit Care Med* 1999, **159**:1925-1932.
- Iwai N, Katsuya T, Ishikawa K, Mannami T, Ogata J, Higaki J, Ogihara T, Tanabe T, Baba S: **Human prostacyclin synthase gene and hypertension: the Suita Study.** *Circulation* 1999, **100**:2231-2236.
- Geraci MW, Gao B, Shepherd DC, Moore MD, Westcott JY, Fagan KA, Alger LA, Tuder RM, Voelkel NF: **Pulmonary prostacyclin synthase overexpression in transgenic mice protects against development of hypoxic pulmonary hypertension.** *J Clin Invest* 1999, **103**:1509-1515.
- Barst RJ, Rubin LJ, Long WA, McGoan MD, Rich S, Badesch DB, Groves BM, Tapson VF, Bourge RC, Brundage BH: **A comparison of continuous intravenous epoprostenol (prostacyclin) with conventional therapy for primary pulmonary hypertension. The Primary Pulmonary Hypertension Study Group [see comments].** *N Engl J Med* 1996, **334**:296-302.
- Rubin LJ, Mendoza J, Hood M, McGoan M, Barst R, Williams WB, Diehl JH, Crow J, Long W: **Treatment of primary pulmonary hypertension with continuous intravenous prostacyclin (epoprostenol). Results of a randomized trial.** *Ann Intern Med* 1990, **112**:485-491.
- Olschewski H, Walrath D, Schermuly R, Ghofrani A, Grimminger F, Seeger W: **Aerosolized prostacyclin and iloprost in severe pulmonary hypertension.** *Ann Intern Med* 1996, **124**:820-824.
- Olschewski H, Simonneau G, Galie N, Higenbottam T, Naeije R, Rubin LJ, Nikkho S, Speich R, Hoepfer MM, Behr J, Winkler J, Sitbon O, Popov W, Ghofrani HA, Manes A, Kiely DG, Ewert R, Meyer A, Corris PA, Delcroix M, Gomez-Sanchez M, Siedentop H, Seeger W: **Inhaled iloprost for severe pulmonary hypertension.** *N Engl J Med* 2002, **347**:322-329.
- Barst RJ, McGoan M, McLaughlin V, Tapson V, Rich S, Rubin L, Wasserman K, Oudiz R, Shapiro S, Robbins IM, Channick R, Badesch D, Rayburn BK, Flinchbaugh R, Sigman J, Arneson C, Jeffs R: **Beraprost therapy for pulmonary arterial hypertension.** *J Am Coll Cardiol* 2003, **41**:2119-2125.
- Galie N, Humbert M, Vachiery JL, Vizza CD, Kneussl M, Manes A, Sitbon O, Torbicki A, Delcroix M, Naeije R, Hoepfer M, Chouat A, Morand S, Besse B, Simonneau G: **Effects of beraprost sodium, an oral prostacyclin analogue, in patients with pulmonary arterial hypertension: a randomized, double-blind, placebo-controlled trial.** *J Am Coll Cardiol* 2002, **39**:1496-1502.
- McLaughlin VV, Genthner DE, Panella MM, Rich S: **Reduction in pulmonary vascular resistance with long-term epoprostenol (prostacyclin) therapy in primary pulmonary hypertension [see comments].** *N Engl J Med* 1998, **338**:273-277.
- Archer SL, Mike D, Crow J, Long W, Weir EK: **A placebo-controlled trial of prostacyclin in acute respiratory failure in COPD.** *Chest* 1996, **109**:750-755.
- Zucker TP, Bonisch D, Hasse A, Grosser T, Weber AA, Schror K: **Tolerance development to antimitogenic actions of prostacyclin but not of prostaglandin E1 in coronary artery smooth muscle cells.** *Eur J Pharmacol* 1998, **345**:213-220.
- Smyth EM, Nestor PV, FitzGerald GA: **Agonist-dependent phosphorylation of an epitope-tagged human prostacyclin receptor.** *J Biol Chem* 1996, **271**:33698-33704.
- Nilius SM, Hasse A, Kuger P, Schror K, Meyer-Kirchdrath J: **Agonist-induced long-term desensitization of the human prostacyclin receptor.** *FEBS Lett* 2000, **484**:211-216.
- Le J I, Shepherd M, Van HG, Houslay MD, Hall IP: **Cyclic AMP-dependent transcriptional up-regulation of phosphodiesterase 4D5 in human airway smooth muscle cells. Identification and characterization of a novel PDE4D5 promoter.** *J Biol Chem* 2002, **277**:35980-35989.
- Mehats C, Tanguy G, Dallot E, Cabrol D, Ferre F, Leroy MJ: **Is up-regulation of phosphodiesterase 4 activity by PGE2 involved in the desensitization of beta-mimetics in late pregnancy human myometrium?** *J Clin Endocrinol Metab* 2001, **86**:5358-5365.
- Sobolewski A, Jourdan KB, Upton PD, Long L, Morrell NW: **Mechanism of cicaprost-induced desensitization in rat pulmonary artery smooth muscle cells involves a PKA-mediated inhibition of adenylyl cyclase.** *Am J Physiol Lung Cell Mol Physiol* 2004, **287**:L352-L359.
- Seeger W, Walrath D, Grimminger F, Rosseau S, Schutte H, Kramer HJ, Ermert L, Kiss L: **Adult respiratory distress syndrome: model systems using isolated perfused rabbit lungs.** *Methods Enzymol* 1994, **233**:549-584.
- Schermuly RT, Roehl A, Weissmann N, Ghofrani HA, Schudt C, Tenor H, Grimminger F, Seeger W, Walrath D: **Subthreshold doses of specific phosphodiesterase type 3 and 4 inhibitors enhance the pulmonary vasodilatory response to nebulized prostacyclin with improvement in gas exchange.** *J Pharmacol Exp Ther* 2000, **292**:512-520.
- Schermuly RT, Krupnik E, Tenor H, Schudt C, Weissmann N, Rose F, Grimminger F, Seeger W, Walrath D, Ghofrani HA: **Coaerosolization of phosphodiesterase inhibitors markedly enhances the pulmonary vasodilatory response to inhaled iloprost in experimental pulmonary hypertension. Maintenance of lung selectivity.** *Am J Respir Crit Care Med* 2001, **164**:1694-1700.
- Schermuly RT, Leuchte H, Ghofrani HA, Weissmann N, Rose F, Kohstall M, Olschewski H, Schudt C, Grimminger F, Seeger W, Walrath D: **Zardaverine and aerosolized iloprost in a model of acute respiratory failure.** *Eur Respir J* 2003, **22**:342-347.
- Weissmann N, Voswinckel R, Hardebusch T, Rosseau S, Ghofrani HA, Schermuly R, Seeger W, Grimminger F: **Evidence for a role of protein kinase C in hypoxic pulmonary vasoconstriction.** *Am J Physiol* 1999, **276**:L90-L95.

36. Shahbazian A, Heinemann A, Peskar BA, Holzer P: **Differential peristaltic motor effects of prostanoid (DP, EP, IP, TP) and leukotriene receptor agonists in the guinea-pig isolated small intestine.** *Br J Pharmacol* 2002, **137**:1047-1054.
37. Rose F, Grimminger F, Appel J, Heller M, Pies V, Weissmann N, Fink L, Schmidt S, Krick S, Camenisch G, Gassmann M, Seeger W, Hanze J: **Hypoxic pulmonary artery fibroblasts trigger proliferation of vascular smooth muscle cells: role of hypoxia-inducible transcription factors.** *FASEB J* 2002, **16**:1660-1661.
38. Darius H, Binz C, Veit K, Fisch A, Meyer J: **Platelet receptor desensitization induced by elevated prostacyclin levels causes platelet-endothelial cell adhesion.** *J Am Coll Cardiol* 1995, **26**:800-806.
39. Fisch A, Tobusch K, Veit K, Meyer J, Darius H: **Prostacyclin receptor desensitization is a reversible phenomenon in human platelets.** *Circulation* 1997, **96**:756-760.
40. Smyth EM, Li WH, FitzGerald GA: **Phosphorylation of the prostacyclin receptor during homologous desensitization. A critical role for protein kinase c.** *J Biol Chem* 1998, **273**:23258-23266.
41. Smyth EM, Austin SC, Reilly MP, FitzGerald GA: **Internalization and sequestration of the human prostacyclin receptor.** *J Biol Chem* 2000, **275**:32037-32045.
42. Manganiello VC, Murata T, Taira M, Belfrage P, Degerman E: **Diversity in cyclic nucleotide phosphodiesterase isoenzyme families.** *Arch Biochem Biophys* 1995, **322**:1-13.
43. Torphy TJ: **Phosphodiesterase isozymes: molecular targets for novel antiasthma agents.** *Am J Respir Crit Care Med* 1998, **157**:351-370.
44. Rabe KF, Tenor H, Dent G, Schudt C, Nakashima M, Magnussen H: **Identification of PDE isozymes in human pulmonary artery and effect of selective PDE inhibitors.** *Am J Physiol* 1994, **266**:L536-L543.
45. Lee R, Wolda S, Moon E, Esselstyn J, Hertel C, Lerner A: **PDE7A is expressed in human B-lymphocytes and is up-regulated by elevation of intracellular cAMP.** *Cell Signal* 2002, **14**:277-284.
46. Torphy TJ, Zhou HL, Foley JJ, Sarau HM, Manning CD, Barnette MS: **Salbutamol up-regulates PDE4 activity and induces a heterologous desensitization of U937 cells to prostaglandin E2. Implications for the therapeutic use of beta-adrenoceptor agonists.** *J Biol Chem* 1995, **270**:23598-23604.
47. Zhu B, Kelly J, Vemavarapu L, Thompson WJ, Strada SJ: **Activation and induction of cyclic AMP phosphodiesterase (PDE4) in rat pulmonary microvascular endothelial cells.** *Biochem Pharmacol* 2004, **68**:479-491.
48. Seybold J, Newton R, Wright L, Finney PA, Suttorp N, Barnes PJ, Adcock IM, Giembycz MA: **Induction of phosphodiesterases 3B, 4A4, 4D1, 4D2, and 4D3 in Jurkat T-cells and in human peripheral blood T-lymphocytes by 8-bromo-cAMP and Gs-coupled receptor agonists. Potential role in beta2-adrenoreceptor desensitization.** *J Biol Chem* 1998, **273**:20575-20588.
49. Murthy KS, Zhou H, Makhlof GM: **PKA-dependent activation of PDE3A and PDE4 and inhibition of adenylyl cyclase V/VI in smooth muscle.** *Am J Physiol Cell Physiol* 2002, **282**:C508-C517.
50. Borchert G, Bartel S, Beyerdorfer I, Kuttner I, Szekeres L, Krause EG: **Long lasting anti-adrenergic effect of 7-oxo-prostacyclin in the heart: a cycloheximide sensitive increase of phosphodiesterase isoform I and IV activities.** *Mol Cell Biochem* 1994, **132**:57-67.
51. Finney PA, Belvisi MG, Donnelly LE, Chuang TT, Mak JC, Scorer C, Barnes PJ, Adcock IM, Giembycz MA: **Albuterol-induced down-regulation of G α accounts for pulmonary beta(2)-adrenoceptor desensitization in vivo.** *J Clin Invest* 2000, **106**:125-135.
52. Moravcova Z, Rudajev V, Stohr J, Novotny J, Cerny J, Parenti M, Miligan G, Svoboda P: **Long-term agonist stimulation of IP prostanoid receptor depletes the cognate G(s) α protein in membrane domains but does not change the receptor level.** *Biochim Biophys Acta* 2004, **1691**:51-65.
53. Tanaka Y, Yamaki F, Koike K, Toro L: **New insights into the intracellular mechanisms by which PGI2 analogues elicit vascular relaxation: cyclic AMP-independent, Gs-protein mediated-activation of MaxiK channel.** *Curr Med Chem Cardiovasc Hematol Agents* 2004, **2**:257-265.
54. Orie NN, Fry CH, Clapp LH: **Evidence that inward rectifier K $^{+}$ channels mediate relaxation by the PGI2 receptor agonist cicaprost via a cyclic AMP-independent mechanism.** *Cardiovasc Res* 2006, **69**:107-115.
55. Yamaki F, Kaga M, Horinouchi T, Tanaka H, Koike K, Shigenobu K, Toro L, Tanaka Y: **MaxiK channel-mediated relaxation of guinea-pig aorta following stimulation of IP receptor with beraprost via cyclic AMP-dependent and -independent mechanisms.** *Naunyn Schmiedebergs Arch Pharmacol* 2001, **364**:538-550.
56. Clapp LH, Turcato S, Hall S, Baloch M: **Evidence that Ca $^{2+}$ -activated K $^{+}$ channels play a major role in mediating the vascular effects of iloprost and cicaprost.** *Eur J Pharmacol* 1998, **356**:215-224.
57. Turcato S, Clapp LH: **Effects of the adenylyl cyclase inhibitor SQ22536 on iloprost-induced vasorelaxation and cyclic AMP elevation in isolated guinea-pig aorta.** *Br J Pharmacol* 1999, **126**:845-847.
58. Schubert R, Serebryakov VN, Engel H, Hopp HH: **Iloprost activates KCa channels of vascular smooth muscle cells: role of cAMP-dependent protein kinase.** *Am J Physiol* 1996, **271**:C1203-C1211.
59. Schubert R, Serebryakov VN, Mewes H, Hopp HH: **Iloprost dilates rat small arteries: role of K(ATP)- and K(Ca)-channel activation by cAMP-dependent protein kinase.** *Am J Physiol* 1997, **272**:H1147-H1156.
60. Fiessinger JN, Schafer M: **Trial of iloprost versus aspirin treatment for critical limb ischaemia of thromboangiitis obliterans. The TAO Study.** *Lancet* 1990, **335**:555-557.
61. Clapp LH, Finney P, Turcato S, Tran S, Rubin LJ, Tinker A: **Differential effects of stable prostacyclin analogs on smooth muscle proliferation and cyclic AMP generation in human pulmonary artery.** *Am J Respir Cell Mol Biol* 2002, **26**:194-201.
62. Smyth EM, Austin SC, FitzGerald GA: **Activation-dependent internalization of the human prostacyclin receptor.** *Adv Exp Med Biol* 2002, **507**:295-301.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

