



POSTER PRESENTATION

Open Access

Onset of action of loratadine/montelukast combination in subjects with seasonal allergic rhinitis in the environmental exposure unit

JH Day^{1*}, MP Briscoe¹, JD Ratz¹, AK Ellis¹, M Danzig², R Yao²

From Canadian Society of Allergy and Clinical Immunology Annual Scientific Meeting 2009
Halifax, Canada. 22-25 October 2009

Background

Onset of action is recognized as an important pharmacologic property of allergic rhinitis medications and can be reliably determined under the controlled conditions of the Environmental Exposure Unit (EEU).

Objective

To evaluate the onset of action of loratadine/montelukast (10 mg/10 mg) versus placebo in subjects with ragweed-induced seasonal allergic rhinitis (SAR).

Methods

A single-center, double-blind, parallel-group study of ragweed-sensitive allergic rhinitis subjects (N=310), performed in the EEU. Subjects were exposed to ragweed pollen in the EEU and symptoms were recorded at 30, 60, 90, and 120 minutes prior to a single dose of loratadine/montelukast or placebo. After dosing, symptoms were recorded for 4 hours - at 15-minute intervals for the first 2 hours and 30-minute intervals for the final 2 hours. The primary endpoint was the time to onset of action for loratadine/montelukast, defined as the first time point at which the mean change from baseline in total symptom score (TSS) for loratadine/montelukast became and remained significantly better than placebo. Secondary endpoints included nasal congestion scores and peak nasal inspiratory flow (PNIF).

Results

The onset of action of loratadine/montelukast for TSS was 1 hour 15 minutes ($p=0.005$ versus placebo). Loratadine/montelukast reduced nasal congestion as indicated

by significant improvements in both the nasal congestion score ($p=0.011$) and PNIF measurements ($p=0.007$) within 1 hour 15 minutes post dose. The incidence of treatment-emergent adverse events was similar between groups.

Conclusion

The onset of action following treatment with loratadine/montelukast was 1 hour 15 minutes for TSS, as well as for nasal congestion. Loratadine/montelukast was well tolerated.

Acknowledgements

Funding for this study was provided by Schering-Plough/Merck Pharmaceuticals.

Author details

¹Department of Medicine, Queen's University and Division of Allergy & Immunology, Kingston General Hospital, Kingston, ON, Canada . ²Schering-Plough Research Institute, Kenilworth, NJ, USA.

Published: 12 May 2010

doi:10.1186/1710-1492-6-S1-P29

Cite this article as: Day et al.: Onset of action of loratadine/montelukast combination in subjects with seasonal allergic rhinitis in the environmental exposure unit. *Allergy, Asthma & Clinical Immunology* 2010 **6**(Suppl 1):P29.

* Correspondence: dayj@kgh.kari.net

¹Department of Medicine, Queen's University and Division of Allergy & Immunology, Kingston General Hospital, Kingston, ON, Canada