CORRECTION Open Access

Correction to: *Pseudomonas aeruginosa* prioritizes detoxification of hydrogen peroxide over nitric oxide

Darshan M. Sivaloganathan¹ and Mark P. Brynildsen^{2*}

Correction to: BMC Res Notes (2021) 14:120

https://doi.org/10.1186/s13104-021-05534-7

Following the publication of the original article [1], the authors brought to our attention that an error was introduced in Figure 1 during the implementation of their corrections: The error bars that were present in panel d of Figure 1 were accidentally removed.

The correct Figure 1 is shown here below and has now been updated in the original article.

Author details

¹Program in Quantitative and Computational Biology, Princeton University, Princeton, NJ, USA. ²Department of Chemical and Biological Engineering, Princeton University, Princeton, NJ, USA.

Published online: 12 July 2021

Reference

 Sivaloganathan DM, Brynildsen MP. Pseudomonas aeruginosa prioritizes detoxification of hydrogen peroxide over nitric oxide. BMC Res Notes. 2021;14:120. https://doi.org/10.1186/s13104-021-05534-7.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at https://doi.org/10.1186/s13104-021-05534-7.

Full list of author information is available at the end of the article



© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*}Correspondence: mbrynild@princeton.edu

 $^{^{\}rm 2}$ Department of Chemical and Biological Engineering, Princeton University, Princeton, NJ, USA




Fig. 1 H_2O_2 clearance is prioritized over that of NO. *P. aeruginosa* cultures were grown to exponential phase and inoculated, at an OD₆₀₀ of 0.025, into a bioreactor containing either 0, 10, or 20 μ M H_2O_2 . Immediately after addition of cells, cultures were treated with either 50 μ M DPTA NONOate or the same volume of the DPTA NONOate solvent. **a** NO concentrations in the bioreactor were continuously measured. **b** Cumulative cellular NO consumption was assessed using a kinetic model with a black-box cellular compartment. **c** H_2O_2 concentrations were measured at 10-min intervals. **d** Culturability of *P. aeruginosa* in the presence of 50 μ M DPTA and 50 μ M DPTA + 20 μ M H_2O_2 were assessed at the beginning and 1 h after treatment. All data represents the mean of three replicates, with error bars representing the standard error of the mean