

Clinic and care: associations with adolescent antiretroviral therapy adherence in a prospective cohort in South Africa

Lucie Cluver^{a,b,*}, Yulia Shenderovich^{a,c,*}, Elona Toska^{d,e,a},
William E. Rudgard^a, Siyanai Zhou^d, Mark Orkin^{a,f},
Roxanna Haghghat^a, Angelique N. Chetty^{a,d}, Caroline Kuo^g,
Alice Armstrong^h and Lorraine Sherrⁱ

Objective: Adolescent antiretroviral treatment (ART) adherence remains critically low. We lack research testing protective factors across both clinic and care environments.

Design: A prospective cohort of adolescents living with HIV (sample $n = 969$, 55% girls, baseline mean age 13.6) in the Eastern Cape Province in South Africa were interviewed at baseline and 18-month follow-up (2014–2015, 2015–2016). We traced all adolescents ever initiated on treatment in 52 government health facilities (90% uptake, 93% 18-month retention, 1.2% mortality).

Methods: Clinical records were collected; standardized questionnaires were administered by trained data collectors in adolescents' language of choice. Probit within-between regressions and average adjusted probability calculations were used to examine associations of caregiving and clinic factors with adherence, controlling for household structure, socioeconomic and HIV factors.

Results: Past-week ART adherence was 66% (baseline), 65% (follow-up), validated against viral load in subsample. Within-individual changes in three factors were associated with improved adherence: no physical and emotional violence (12.1 percentage points increase in adjusted probability of adherence, $P < 0.001$), improvement in perceived healthcare confidentiality (7.1 percentage points, $P < 0.04$) and shorter travel time to the clinic (13.7 percentage points, $P < 0.02$). In combination, improvement in violence prevention, travel time and confidentiality were associated with 81% probability of ART adherence, compared with 47% with a worsening in all three.

Conclusion: Adolescents living with HIV need to be safe at home and feel safe from stigma in an accessible clinic. This will require active collaboration between health and child protection systems, and utilization of effective violence prevention interventions.

Copyright © 2021 The Author(s). Published by Wolters Kluwer Health, Inc.

AIDS 2021, **35**:1263–1271

Keywords: adherence, adolescence, antiretroviral therapy, confidentiality, decentralization, violence

^aCentre for Evidence-Based Intervention, Department of Social Policy and Intervention, University of Oxford, Oxford, UK,

^bDepartment of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa, ^cCentre for the Development and Evaluation of Complex Interventions for Public Health Improvement (DECIPHer), School of Social Sciences, Cardiff University Cardiff, UK, ^dCentre for Social Science Research, ^eDepartment of Sociology, University of Cape Town, Cape Town, ^fMedical Research Council Development Pathways to Health Research Unit, School of Clinical Medicine, University of the Witwatersrand, Johannesburg, South Africa, ^gSchool of Public Health, Brown University, Providence, Rhode Island, USA, ^hUNICEF Eastern and Southern Africa Regional Office, Nairobi, Kenya, and ⁱHealth Psychology Unit, Institute of Global Health, University College London, London, UK.

Correspondence to Lucie Cluver, DPhil (PhD), Department of Social Policy and Intervention, Barnett House, 32 Wellington Square, Oxford OX1 2ER, UK.

E-mail: lucie.cluver@spi.ox.ac.uk.

* Lucie Cluver and Yulia Shenderovich contributed equally to this study.

Received: 9 November 2020; revised: 2 March 2021; accepted: 8 March 2021.

DOI:10.1097/QAD.0000000000002882

Introduction

Adolescents have the lowest antiretroviral therapy (ART) adherence rate among all age groups [1], leading to elevated morbidity, viral resistance, onwards infection [2,3] and mortality [4]. Retention in HIV care is further challenged in the context of COVID-19 [5].

Eighty-five percent of adolescents living with HIV live in sub-Saharan Africa. Systematic reviews have found limited evidence for effective interventions to increase adolescent ART adherence [6,7]. Evidence shows that healthcare factors are important [8], and interventions to date have primarily focused on health systems: supporting young people to take their medication and increasing the availability, affordability and acceptability of HIV care [9,10].

Social relationships in healthcare settings and relationships in the community have also been identified as key for adolescents [11]. Confidentiality and trust may be both indicators of and predictors of quality of care and healthcare outcomes [12]. New evidence in the region also shows positive impacts of interventions, including peer supporters, support groups and counselling [9,13].

But we may be missing an additional social pathway. Developmental psychology shows that adolescent behaviour is closely linked to family and home relationships [14]. In particular, caregiving or parenting – which refers to any biological or nonbiological primary caregiver – has lasting effects including and beyond health-seeking behaviours [15]. Nurturing and supportive caregiving increases adolescent capacity to manage risks [16], while violent caregiving increases risk behaviours and mental health distress [17]. In qualitative research, adolescents living with HIV have identified both healthcare and family influences on their medication-taking [18–21]. The importance of caregiving has been heightened during the COVID-19 epidemic, with revolving lockdowns, closures of schools and community-based services, and restricted peer-mixing [22].

Despite this, we lack longitudinal studies that test whether caregiving is linked with adolescent adherence in sub-Saharan Africa. There are strong indications from cross-sectional studies: in Uganda, family and caregiver social support were associated with higher adolescent ART adherence, but support from peers and teachers were not [23]. Caregiver violence was associated with lower adherence in cross-sectional studies in South Africa, Malawi and Zambia [24–26].

It is essential to examine both healthcare provision and family care factors in longitudinal research to identify further ways to support adherence. We conducted a community-based prospective cohort of adolescents living with HIV (aged 10–19 years at baseline) from 2014 to 2017 in 180 communities of South Africa's Eastern Cape. We use

baseline and 18-month follow-up data from 969 adolescents to examine associations of clinic and care factors with past-week ART adherence, accounting for household structure, socioeconomic and HIV-related factors, shown in prior research to affect adherence [27–29].

Materials and methods

The study took place in the Eastern Cape province of South Africa, characterized by high morbidity, low human development and poor infrastructure. We conducted standardized interviews and extracted clinical records for 1046 adolescents living with HIV at baseline (2014–2015), with 979 reinterviewed at 18-month follow-up (2016–2017). Nine hundred and sixty-nine had complete data on key variables. The study included both adolescents engaged in clinical care and those who were lost to follow-up in clinical care, and thus, it is the region's first large-scale community-traced cohort of this group.

In a health district including rural, urban and peri-urban settlements, we identified all 52 community health centres, primary clinics and hospitals that provided ART to adolescents. In each facility, all patient files (paper and electronic) were reviewed to identify all those who had ever initiated ART and were aged 10–19 years. Throughout 180 communities, adolescents were interviewed at a location of their choice. At 18 months after baseline, all adolescents who had consented to be reapproached were asked for consent for follow-up. Reflecting high mobility, 18% of participants have moved households between study waves, and by follow-up participants lived in six provinces: Eastern Cape, Gauteng, KwaZulu-Natal, Free State, Western Cape and North-West.

Ethical approvals were given by the University of Cape Town (CSSR 2013/4), Oxford University (CUREC2/12–21), provincial Departments of Health and Education, and healthcare facilities. All adolescents and their primary caregivers gave written informed consent at both time points in their language of choice (Xhosa or English), read aloud in cases of low literacy. The study did not provide financial incentives, but adolescents did receive a snack, a certificate of participation, and a small gift pack including soap and pencils. These were recommended by our adolescent advisory group [30] and provided regardless of interview completion.

Clinical records were extracted in healthcare facilities (see Supplementary Materials Box S1, <http://links.lww.com/QAD/C74>), and trained local researchers supported participants in completing tablet-based questionnaires lasting 60–90 min. Adolescents chose their language of participation. Questionnaires were codesigned with an adolescent advisory group; the South African National

Departments of Health, Social Development, Basic Education and National AIDS Council; UNICEF; PEPFAR-USAID, and local NGOs. Prepiloting was conducted locally with 25 adolescents living with HIV.

To avoid stigma or unintended disclosure of HIV status, the research focus was presented as general adolescent social and health needs, and 456 neighbouring adolescents were also interviewed (not included in these analyses). Confidentiality was maintained except in cases of risk of harm. For rape, abuse, suicidality or untreated severe illness [e.g. symptomatic tuberculosis (TB)], researchers made immediate health and social service referrals with follow-up support ($n = 157$ referrals over 3 years for 104 adolescents).

Procedures

Full questionnaires are available at <http://www.young-carers.org.za/youthpulse>. All variables in these analyses were measured and defined in the same way at baseline and follow-up.

Outcome

ART adherence was measured using adapted items from the Patient Medication Adherence Questionnaire and measures developed in Botswana [31,32]. ART adherence was defined as past 7 days adherence more than 95% (including weekdays and weekend), based on currently taking ART and not having missed any doses in the past seven days [33]. We validated self-reported adherence against viral loads available in clinical records, using the viral load measurement closest to the interview date. Eight percent of adolescents' clinical records did not include any viral load measurements, and about 60% of adolescents with an available viral load had a measurement from the two years before or after the questionnaire date [34]. Thus, our validation focused on adolescents whose clinical records included viral load measurements within this period, excluding measurements in a 30-day range around ART initiation ($n = 650$ adolescents at baseline and $n = 598$ at follow-up).

Explanatory variables

Clinic factors: *Medication stock-outs* were measured as past-year inability to access ART due to clinic stock-outs. *Confidentiality* was measured as feeling that their information would be kept safe and confidential at the clinic most or all of the time in the past year. *Wait time in the clinic* above 1 h and *travel to the clinic* above 1 h were measured as past-year experiences in respect to the main clinic the adolescent attends [27–29].

Caregiving factors: *Past-year physical or emotional abuse victimization* by caregivers or other adults were measured using UNICEF Measures for National-level Monitoring of Orphans and Vulnerable Children [35] (12 items) and defined as having experienced at least one type of violence, such as being hit with a hard item. *Past-week witnessing domestic violence between adults in the home* (physical or verbal)

also used these UNICEF measures [35] (two items). *Good parent/caregiver monitoring and supervision* (nine items, e.g., having rules for when adolescents come home) and *positive caregiving* (six items, e.g. positive reinforcement) were measured over the past 2 months, using subscales of the Alabama Parenting Questionnaire [36], and defined as not having experienced any poor monitoring and supervision, and as always experiencing positive caregiving. *Good communication* between primary caregivers and adolescents was measured over the past 2 months using adapted Child-Parent Communication Apprehension Scale for use with young adults, asking about ease and openness of communication and defined as agreeing or strongly agreeing to all items (five items) [37]. All caregiving and clinic factors were dichotomized to facilitate interpretability.

Control variables

Household structure factors included *orphanhood* (maternal or paternal) measured using items adapted from the 2011 South African census [38]; *number of changes of caregiver* experienced; *household size*, and *relationship of primary caregiver to child*, that is biological parent or not. Socioeconomic factors included adolescent *age*, *sex*, *urban/rural location* and living in *formal or informal housing*, using census-based items [39]. *Household poverty* was measured as access to eight highest socially perceived necessities for children in the nationally representative South African Social Attitudes Survey (e.g. enough food) [40]. HIV-related factors were measured using clinical records, checked against self-report where possible, and included *mode of HIV infection* (vertical/horizontal) and *recent HIV infection* (<2 years before baseline).

Statistical analyses

First, we validated self-reported ART adherence against undetectable viral load (<50 copies/ml) from clinical records using multivariable logistic regression, controlling for age, sex, rural/urban, orphanhood, informal housing, mode of infection and health status. Second, we examined frequencies of adherence, hypothesized explanatory and control variables. Third, we compared participants who were followed up and included in this analysis, and those who were not, to check for potential differences using *t*-tests for continuous variables and Chi-squared tests for dichotomous variables.

Fourth, we examined associations of clinic and care factors with past-week adherence. We used a within-between regression model, also known as a hybrid model, which allows us to look at within-person variation as well as compute average adjusted probabilities of the outcome [41–43]. For each explanatory variable, we used a person's average value and time-specific deviation from this average [see Eq. (1)].

$$P(\text{Adherence}_{it} = 1 | x_{it}) = \Phi(\beta_0 + \beta_1(\bar{x}_i) + \beta_2(x_{it} - \bar{x}_i) + v_{i0} + \varepsilon_{it}), \quad (1)$$

where Φ is the cumulative normal distribution, adherence is the time-varying dependent variable, β_0 represents the overall intercept, x_{it} is a time-varying explanatory variable for person i at time t , \bar{x}_i is the average of the explanatory variable for person i across both time points, β_1 represents the average between-person effect and β_2 represents the average within-person effect, ν_{i0} – a random person-level intercept, assumed to be normally distributed, and ε_{it} – the residuals.

This provides a ‘between-person’ coefficient (estimated by \bar{x}_i), the difference between individuals and a ‘within-person’ coefficient (estimated by $x_{it} - \bar{x}_i$), which examines the changes within individual’s levels of the explanatory variable over time. With two time points, the within-person estimate is equivalent to a first difference model [44]. The within-person estimate is the focus of our analyses as it allows to account for all time-invariant confounders [44]. Participant sex, ART initiation and mode of infection were modelled as time-invariant, while all other control variables were modelled as time-variant.

Analyses were conducted in Stata 14.2 (code available at <https://osf.io/znse9/>). We used a probit model (*xtprobit* command) with robust clustered standard errors at the individual level. To aid interpretation of the relationships between the explanatory variables and adherence, we estimated average adjusted probabilities [45] of the clinic and caregiving factors that were statistically significant in the multivariable regression ($P < 0.05$), using *margins* and *lincom* commands [46].

Results

Adherence rates and viral load

Self-reported ART adherence was low, with 66% of adolescents at baseline and 65% at follow-up reporting adherence in the past week. Only 45% reported past-week adherence at both time points. Self-reported adherence was associated with undetectable viral load (<50 copies/ml) at both time points, controlling for age,

sex, rural/urban, orphanhood, informal housing, mode of infection and health status [see Table 1, baseline odds ratio (OR) 1.46, 95% confidence interval (95% CI) 1.02–2.08, $P = 0.037$, $n = 650$; follow-up OR 1.45 95% CI 1.01–2.07, $P = 0.045$, $n = 598$].

Sample characteristics

Descriptive data are summarized in Table 2. The sample was 55% female with mean age 13.6 years (SD = 2.9) at baseline. Around two-thirds of adolescents were orphaned (maternal, paternal or double), and less than half lived with a biological parent. At both baseline and follow-up, around one-third of adolescents reported good caregiver supervision and good communication. Eleven percent reported past-week physical or verbal violence between adults in the home. Thirty percent (baseline) and 38% (follow-up) reported experiencing physical or emotional abuse. Eleven percent (baseline) and 5% (follow-up) reported travel time to clinic more than 1 h, and about half of the sample at baseline and follow-up reported wait time at the clinic more than 1 h. Five percent at each time-point reported medication stock-outs, and 24% (at baseline) and 36% (at follow-up) did not feel that their information would mostly be kept confidential at the clinic.

Sixty-seven (6%) adolescents recruited at baseline were lost to follow-up, including 12 who died between timepoints. Ten (1%) had incomplete data. We compare 969 participants included (complete cases) and 77 participants not included (Table S1, <http://links.lww.com/QAD/C74>). There were no differences on most baseline characteristics, but participants lost to follow up were more likely to be nonadherent, older, horizontally infected and live with a biological parent.

Associations of caregiving and healthcare factors with adherence

Key results of the within-between probit regression are presented in Table 3 (single-variable regression results in Table S2, <http://links.lww.com/QAD/C74> and full output of multivariable regression in Table S3, <http://links.lww.com/QAD/C74>).

Table 1. Multivariable logistic regressions testing associations between self-reported antiretroviral therapy adherence and undetectable viral load.

Explanatory variable	Baseline viral load ($n = 650$)				Follow-up viral load ($n = 598$)			
	AOR	Lower 95% CI	Higher 95% CI	P	AOR	Lower 95% CI	Higher 95% CI	P
Self-reported adherence	1.457*	1.024	2.075	0.037	1.447*	1.009	2.074	0.045
Age	0.928*	0.867	0.994	0.032	0.971	0.903	1.045	0.433
Sex (girl)	1.117	0.801	1.557	0.516	0.935	0.665	1.315	0.700
Rural area	0.948	0.654	1.374	0.777	0.618*	0.425	0.899	0.012
Orphan	0.661*	0.464	0.941	0.022	0.809	0.567	1.154	0.242
Informal housing	0.707	0.471	1.061	0.094	1.151	0.720	1.841	0.557
Horizontal infection	0.877	0.534	1.441	0.604	0.829	0.491	1.399	0.482
Poor health	0.459*	0.230	0.917	0.027	0.518	0.225	1.194	0.123

95% CI, 95% confidence interval; AOR, adjusted odds ratio.

* $P < 0.05$.

Table 2. Descriptive characteristics of the analytic sample (n = 969).

Variables	Baseline n (%)	Follow-up n (%)
ART adherence	642 (66)	628 (65)
Healthcare factors		
No medication stock-outs	918 (95)	920 (95)
Confidentiality	740 (76)	625 (64)
Travel to the clinic below 1 h	859 (89)	923 (95)
Wait time in the clinic below 1 h	503 (52)	450 (46)
Caregiving factors		
Good monitoring and supervision	332 (34)	364 (38)
Good caregiver-teen communication	258 (27)	331 (34)
No emotional or physical violence in the past year	680 (70)	601 (62)
Not witnessing arguments or fights between adults in the home in the past week	859 (89)	861 (89)
Household structure		
Orphan	578 (60)	592 (61)
Number of caregivers with whom the teen has ever lived	1.89 (1.04)	2.07 (1.25)
Biological parent is primary caregiver	423 (44)	395 (41)
Household size	6.78 (2.94)	6.21 (3.87)
Socioeconomic factors		
Sex (girl)		532 (55)
Age (years)	13.58 (2.87)	15.10 (2.87)
Rural area	258 (27)	238 (25)
Informal housing	180 (19)	141 (15)
Can afford 8 basic necessities	310 (32)	215 (22)
HIV factors		
Mode of infection (horizontal)		201 (21)
Recent ART initiation (under 2 years prior to baseline)		256 (26)

Table 3. Multivariable probit regressions testing associations of adolescent clinic and care experiences and their adherence (n = 969 people, 1928 observations).

	Coefficient	Lower 95% CI	Higher 95% CI	P
Healthcare factors – within-person variation				
No medication stock outs	0.335	-0.116	0.786	0.146
Perceived confidentiality at the clinic	0.240*	0.042	0.438	0.018
Travel to the clinic below 1 h	0.451*	0.105	0.796	0.011
Wait time in the clinic below 1 h	-0.170	-0.352	0.012	0.068
Family care factors – within-person variation				
Good monitoring and supervision	-0.082	-0.287	0.124	0.438
Good caregiver-teen communication	-0.003	-0.196	0.189	0.972
No emotional or physical violence victimization	0.378***	0.183	0.574	<0.001
Not witnessing any arguments or fights between adults at home	0.191	-0.083	0.464	0.171

Control factors are orphanhood, caregiver being a biological parent, household size, number of previous caregivers, adolescent age, rural residence, informal housing, poverty, study wave, participant sex, mode of HIV infection and recent ART initiation. 95% CI, confidence interval. All coefficients are presented in supplementary Table S3, <http://links.lww.com/QAD/C74>.

* $P < 0.05$.

*** $P < 0.001$.

Table 4. Adjusted average predicted probabilities of adherence at various levels of the key explanatory variables, based on regressions reported in Table 3.

Explanatory variables	Adjusted probability of adherence	Lower 95% CI	Higher 95% CI	Difference of adjusted probabilities of adherence	Lower 95% CI	Higher 95% CI	P
Perceived confidentiality at the clinic							
Decrease in confidentiality	0.621	0.580	0.661	ref			
Increase in confidentiality	0.692	0.654	0.730	0.071*	0.006	0.137	0.0325
Travel to the clinic below 1 h							
Increase in travel time	0.586	0.523	0.649	ref			
Decrease in travel time	0.723	0.668	0.777	0.137*	0.027	0.2460	0.0143
No emotional or physical violence victimization							
Increase in violence	0.595	0.555	0.635	ref			
Decrease in violence	0.716	0.680	0.752	0.121***	0.058	0.183	0.0001

Average adjusted probabilities are reported at the observed values of other variables in the model (average marginal effects). 95% CI, confidence interval.

* $P < 0.05$.

*** $P < 0.001$.

To aid the interpretation of the regression coefficients' magnitude, Table 4 demonstrates adjusted probabilities of adherence at each value of the within-person explanatory variables that were statistically significant in the regression analysis, and the differences between the probabilities of adherence at the different values of these explanatory variables.

Within-person variation in two clinic factors was associated with adherence. Improvement in confidentiality in the clinic setting was associated with a greater likelihood of adolescents becoming adherent (7.1 percentage points, $P=0.0325$). Travel time to the clinic reducing to under 1 h was also associated with higher likelihood of becoming adherent (13.7 percentage points difference, $P=0.0143$).

Within-person variation in one caregiving factor was associated with adherence. Ending exposure to violence was associated with a greater likelihood of adolescents becoming adherent (12.1 percentage points difference, $P=0.0001$).

Predicted probability of adherence for those experiencing a reduction in violence and travel time as well as improved confidentiality was estimated, based on the regression model, to be 81%, compared with 47% for those experiencing a deterioration on these three factors, thus a 34 percentage points difference (95% CI 21–46, $P < 0.001$).

Discussion

The high rates of adolescent nonadherence in this study reflect similar trends in national and regional data, and are well below UNAIDS targets [47]. It is essential that we understand and mitigate barriers to ART adherence for young people. These are the first known findings from a longitudinal study identifying associations of both care and clinic factors to adolescent adherence, advancing the evidence by using a prospective cohort and including variables across different settings of the adolescents' lives.

The study finds three factors were associated with better adherence. Nonexposure to violence was associated with better adherence. Associations of violence with ART adherence were as strong or stronger than healthcare factors. This is particularly relevant in the context of South Africa, which has high rates of violence against children [48]. There is also evidence that social protection such as cash transfers or other forms of economic strengthening can help to reduce family violence [49], and existing cash transfer policies within South Africa may be an essential component of violence reduction. Other effective interventions to reduce child abuse do exist, but are not yet scaled to population levels in any high HIV-prevalence country [50]. Evidence-based

noncommercialized parenting/caregiving programmes have been tested in randomized trials in South Africa, South Sudan and Burundi. They show reduced violence, and improved parenting, reduced parental substance use and mental health distress that are strong predictors of violence [51–53], with similar engagement and outcomes amongst HIV/AIDS-affected and nonaffected families [54,55]. Parenting/caregiver programmes have also been shown to be cost-effective [56] with delivery costs of around \$20 per family.

HIV and child protection services are rarely explicitly linked. Overcoming this may require increased collaboration between these systems, and across health and social development agencies. There are existing entry points. Health facilities and peer supporters could ask brief screening questions to identify violence exposure, such as those included in the WHO-recommended HEADSS and HEADSS+ questions [57,58], although caregiver presence may present challenges in reporting. Anecdotally, HIV peer supporter services already report high levels of violence disclosure. Governments, UNICEF, PEPFAR-USAID and the International Rescue Committee deliver parenting/caregiving programmes in the region, and these could provide platforms for reaching more adolescents living with HIV. Community-based services ([9,59,60]) may provide excellent opportunities to co-deliver effective violence prevention, screening, and support. In the COVID-19 pandemic, violence prevention and adherence support services are being adapted to remote and digital delivery modes [61,62], and this may provide opportunities for combining with HIV services.

Reductions in travel time to clinics and improvements in perceived confidentiality within clinics were also associated with improved adolescent adherence. This suggests potential positive implications for sub-Saharan Africa's decentralization from hospital-based ART provision to primary care, which often reduces patient travel burden. Systematic reviews suggest – from limited data – that decentralization may support adolescent retention in care [63,34]. WHO Global Standards for Quality Healthcare Services for Adolescents have identified insufficient respect for adolescent confidentiality and privacy as a major barrier for service access [64]. Confidentiality is important in building trust between adolescents living with HIV and their healthcare providers, and this has new implications as we grapple with providing remote support in the COVID context.

This study has limitations. First, we need to be cautious in determining causality. A strength of the within-person estimates is that they account for unmeasured stable person-level characteristics, such as adolescent early childhood experiences, as we examine changes within the same individuals [65,66]. However, this estimate assumes no unobserved time-varying confounders (for

instance, a policy change that affects both violence and adherence) [41]. We control for wave of data collection to help account for period effects. The within-person estimate relies on the adolescents who experienced changes in their exposures and in their adherence, reducing statistical power to detect associations. We assume that our outcome does not affect the explanatory variables, but we should also consider risks of reverse causality. It is possible that nonadherence could increase the travel time to HIV services if an adolescent was subsequently up-referred to a more specialized and distant facility; however, the reverse could also be true of down-referral to improve accessibility. Our findings suggest that caregiver violence is associated with adolescent ART-nonadherence. We also considered the reverse possibility that adolescent ART nonadherence could contribute to caregiver violence, but epidemiological evidence suggests that adolescent behaviour is not a primary contributor to the social and structural factors driving family violence [67]. Notably, two-thirds of adolescents living with HIV who participated in recent qualitative research in Zambia felt that family violence was negatively impacting their adherence, clinic attendance or virologic results [68].

Second, we use self-reported adherence as the outcome, due to very limited clinical recording of treatment outcomes. We validated adherence against viral load measures but the low rates of viral load testing in this context meant that only up to 650 measures could be compared. Third, adolescents who reported nonadherence at baseline were more likely to be lost to follow-up. Fourth, our measure of violence focused on victimization by adults at home or school (e.g. at a boarding school or orphanage) but in this sample almost all children lived in households. Fifth, we were not able to measure treatment stockouts for adolescents who were disengaged from HIV care. We did not include clinic in which adolescents received care as a factor in the models as inter-facility mobility and attendance of multiple healthcare facilities concurrently were common in this cohort: 31% had clinical records from at least two facilities within the study area.

The study also has several important strengths. It provides evidence from a sample of adolescents initiated on ART through government services in over 50 clinics in the South African Eastern Cape province. Affected by the legacy of apartheid, the Eastern Cape has high rates of poverty, unemployment, with over a quarter of the province's households food insecure [69]. Eastern Cape also has poor healthcare infrastructure and high rates of HIV prevalence (at 15.3% in 2017 [70]). Therefore, findings from this study setting may be generalizable to other countries in sub-Saharan Africa as well as other resource-constrained settings.

Our study uses standardized questionnaires and repeated measures, with clinical record data. It traces and interviews adolescents as a community sample, allowing

inclusion of adolescents who have disengaged from care or move between healthcare settings. It is the only known longitudinal study to provide evidence of associations between caregiving factors and adherence amongst adolescents living with HIV in Sub-Saharan Africa, and it is also unique in allowing the simultaneous examination of clinic and care factors, while controlling for HIV, social and economic factors. Further research on the multiple drivers of adolescent adherence is needed to build on these results and inform more effective services. It will also be essential to build longer-term evidence to improve lifetime adherence. Our future research includes a third wave of data collection, and using linked National Health Laboratory Service data as these adolescents become adults.

The HIV epidemic has taught us a hard lesson: that our capacity to prevent and treat a fatal virus is intertwined with the complexity of life experiences. For this highly vulnerable group of adolescents living with HIV, we need to respond both within and beyond the clinic system. Ensuring access to effective violence prevention and response services is an essential component for adolescent survival and success.

Acknowledgements

We would like to thank the young people and their families who shared their experiences, and the fieldwork team who have shown such care, kindness and tenacity over the years. With particular thanks to Laurie Gulaid at UNICEF ESAR Office for conceptual thinking, and to Dr Tobias Rüttenauer, for valuable advice on methods.

This study was funded by the UK Research and Innovation Global Challenges Research Fund (UKRI GCRF) Accelerate Hub [ES/S008101/1]; UNICEF Eastern and Southern Africa Office, Nuffield Foundation [CPF/41513]; Evidence for HIV Prevention in Southern Africa, a UKAID programme managed by Mott MacDonald; Janssen Pharmaceutica NV; and the International AIDS Society through the CIPHER grant (155-Hod; 2018/625-TOS); Claude Leon Foundation (08 559/C); Oak Foundation (R46194/AA001); the John Fell Fund (103/757 and 161/033); the University of Oxford's Economic and Social Research Council Impact Acceleration Account (IAA-MT13-003; 1602-KEA-189; K1311-KEA-004); the Leverhulme Trust (PLP-2014-095); Research England; the European Research Council (ERC) under the European Union's (EU) Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement 313421, the EU's Horizon 2020 research and innovation programme/ERC grant agreement 737476); the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID

Concordat agreement, and by the Department of Health Social Care through its National Institutes of Health Research (MR/R022372/1), and the Oak Foundation (OFIL-20-057).

Author contributions: E.T. and L.C. designed and oversaw the data collection. Y.S. and L.C. led the statistical analysis and data interpretation with contributions from E.T., S.Z., W.E.R., A. A. and M.O. All authors contributed to the writing of the manuscript and approved its final version.

Conflicts of interest

There are no conflicts of interest.

References

- Lowenthal E, Bakeera-Kitaka S, Marukutira T, Chapman J, Goldrath K, Ferrand R. **Perinatally acquired HIV infection in adolescents from sub-Saharan Africa: a review of emerging challenges.** *Lancet Infect Dis* 2014; **14**:627–639.
- Pillay D. **The emergence and epidemiology of resistance in the nucleoside-experienced HIV-infected population.** *Antivir Ther* 2001; **6 Suppl 3**:15–24.
- UNICEF. *For every child, end AIDS: Seventh Stocktaking Report.* New York: UNICEF; 2016.
- UNAIDS. *Global HIV and AIDS statistics 2019.* Geneva: UN-AIDS; 2019.
- UNICEF. *Prioritizing the continuity of services for adolescents living with HIV during the COVID-19 pandemic.* New York: UNICEF; 2020. Available at <https://www.corecommitments.unicef.org/kp/prioritizing-the-continuity-of-services-for-adolescents-living-with-hiv-during-the-covid-19-pandemic.url>.
- Casale M, Carlqvist A, Cluver L. **Recent interventions to improve retention in HIV care and adherence to antiretroviral treatment among adolescents and youth: a systematic review.** *AIDS Patient Care STDS* 2019; **33**:237–252.
- MacPherson P, Munthali C, Ferguson J, Armstrong A, Kranzer K, Ferrand R, et al. **Service delivery interventions to improve adolescents' linkage, retention and adherence to antiretroviral therapy and HIV care.** *Trop Med Int Health* 2015; **20**:1015–1032.
- Cluver L, Pantelic M, Toska E, Orkin M, Casale M, Bungane N, et al. **STACKing the odds for adolescent survival: health service factors associated with full retention in care and adherence amongst adolescents living with HIV in South Africa.** *J Int AIDS Soc* 2018; **21**:e25176.
- Mavhu W, Willis N, Mufuka J, Bernays S, Tshuma M, Manganah C, et al. **Effect of a differentiated service delivery model on virological failure in adolescents with HIV in Zimbabwe (Zvan-diri): a cluster-randomised controlled trial.** *Lancet Glob Health* 2020; **8**:e264–e275.
- Fatti G, Jackson D, Goga AE, Shaikh N, Eley B, Nachega JB, Grimwood A. **The effectiveness and cost-effectiveness of community-based support for adolescents receiving antiretroviral treatment: an operational research study in South Africa.** *J Int AIDS Soc* 2018; **21 (Suppl 1)**:e25041.
- Zuma T, Seeley J, Mdulsi S, Chimbindi N, McGrath N, Floyd S, et al. **Young people's experiences of sexual and reproductive health interventions in rural KwaZulu-Natal, South Africa.** *Int J Adolesc Youth* 2020; **25**:1058–1075.
- Hall MA, Dugan E, Zheng B, Mishra AK. **Trust in physicians and medical institutions: what is it, can it be measured, and does it matter?** *Milbank Q* 2001; **79**:613–639v.
- Ferrand RA, Simms V, Dauya E, Bandason T, McHugh G, Mujuru H, et al. **The effect of community-based support for caregivers on the risk of virological failure in children and adolescents with HIV in Harare, Zimbabwe (ZENITH): an open-label, randomised controlled trial.** *Lancet Child Adolesc Health* 2017; **1**:175–183.
- Ungar M, Ghazinour M, Richter J. **Annual research review: what is resilience within the social ecology of human development?** *J Child Psychol Psychiatry* 2013; **54**:348–366.
- Nemeroff CB. **Paradise lost: the neurobiological and clinical consequences of child abuse and neglect.** *Neuron* 2016; **89**:892–909.
- Bornstein M. *Handbook of parenting.* Washington, DC: Routledge; 2019.
- Anda R, Felitti V, Bremner J, Walker J, Whitfield C, Perry B, et al. **The enduring effects of abuse and related adverse experiences in childhood.** *Eur Arch Psychiatry Clin Neurosci* 2006; **256**:174–186.
- Denison JA, Banda H, Dennis AC, Packer C, Nyambe N, Stalter RM, et al. **The sky is the limit': adhering to antiretroviral therapy and HIV self-management from the perspectives of adolescents living with HIV and their adult caregivers.** *J Int AIDS Soc* 2015; **18**:19358.
- Hodes R. **The stuff that dreams are made of: using participatory research to explore interlinkages in HIV-positive adolescents' aspirations for development.** *J Int AIDS Soc* 2018; **21**:e25057.
- MacCarthy S, Saya U, Samba C, Birungi J, Okoboi S, Linnemayr S. **How am I going to live? Exploring barriers to ART adherence among adolescents and young adults living with HIV in Uganda.** *BMC Public Health* 2018; **18**:1158.
- Bernays S, Seeley J, Rhodes T, Mupambireyi Z. **What am I 'living' with? Growing up with HIV in Uganda and Zimbabwe.** *Sociol Health Illn* 2015; **37**:270–283.
- Orben A, Tomova L, Blakemore SJ. **The effects of social deprivation on adolescent development and mental health.** *Lancet Child Adolesc Health* 2020; **4**:634–640.
- Damulira C, Mukasa MN, Byansi W, Nabunya P, Kivumbi A, Namatovu P, et al. **Examining the relationship of social support and family cohesion on ART adherence among HIV-positive adolescents in southern Uganda: baseline findings.** *Vulnerable Child Youth Stud* 2019; **14**:181–190.
- Cluver L, Meinck F, Toska E, Orkin FM, Hodes R, Sherr L. **Multitype violence exposures and adolescent antiretroviral nonadherence in South Africa.** *AIDS* 2018; **32**:975–983.
- Kim MH, Mazenga AC, Yu X, Ahmed S, Paul ME, Kazembe PN, et al. **High self-reported nonadherence to antiretroviral therapy amongst adolescents living with HIV in Malawi: barriers and associated factors.** *J Int AIDS Soc* 2017; **20**:21437.
- Merrill KG, Campbell JC, Decker MR, McGready J, Burke VM, Mwansa JK, et al. **Past-year violence victimization is associated with viral load failure among HIV-positive adolescents and young adults.** *AIDS Behav* 2020. Available at <https://doi.org/10.1007/s10461-020-02958-3>.
- Ojikutu B, Higgins-Biddle M, Greeson D, Phelps BR, Amzel A, Okechukwu E, et al. **The association between quality of HIV care, loss to follow-up and mortality in pediatric and adolescent patients receiving antiretroviral therapy in Nigeria.** *PLoS One* 2014; **9**:e100039.
- Okoboi S, Ssali L, Yansaneh AI, Bakanda C, Birungi J, Nantume S, et al. **Factors associated with long-term antiretroviral therapy attrition among adolescents in rural Uganda: a retrospective study.** *J Int AIDS Soc* 2016; **19 (5 Suppl 4)**:20841.
- Reif LK, Bertrand R, Benedict C, Lamb MR, Rouzier V, Verdier R, et al. **Impact of a youth-friendly HIV clinic: 10 years of adolescent outcomes in Port-au-Prince, Haiti.** *J Int AIDS Soc* 2016; **19**:20859.
- Oliveras C, Cluver L, Sharma A, Caswell G, Wanjiku L, Bernays S, et al. **'Nothing about us without RIGHTS': Meaningful engagement of children and youth: from research prioritization to clinical trials, implementation science and policy.** *J AIDS* 2018; **78**:S27–S31.
- Duong M, Piroth L, Grappin M, Forte F, Peytavin G, Buisson M, et al. **Evaluation of the Patient Medication Adherence Questionnaire as a tool for self-reported adherence assessment in HIV-infected patients on antiretroviral regimens.** *HIV Clin Trials* 2001; **2**:128–135.
- Lowenthal E, Haruna B, Mmapula L, Keofentse M, Tshume O, Anabwani G. **Disclosure of HIV status to HIV-infected children in a large African treatment center: lessons learned in Botswana.** *Child Youth Serv Rev* 2014; **45**:143–149.
- Paterson DL, Swindells S, Mohr J, Brester M, Vergis EN, Squier C, et al. **Adherence to protease inhibitor therapy and outcomes in patients with HIV infection.** *Ann Intern Med* 2000; **133**:21–30.

34. Haghight R, Toska E, Cluver L, Gulaid L, Mark D, Bains A. **Transition pathways out of pediatric care and associated HIV outcomes for adolescents living with HIV in South Africa.** *J Acquir Immune Defic Syndr* 2019; **82**:166–174.
35. Snider L, Dawes A. *Psychosocial vulnerability and resilience measures for national-level monitoring of orphans and other vulnerable children: recommendations for revision of the UNICEF Psychological Indicator.* Cape Town: UNICEF; 2006.
36. Frick P. *The Alabama Parenting Questionnaire.* Alabama: University of Alabama; 1991.
37. Lucchetti A, Powers W, Love D. The empirical development of the child-parent communication apprehension scale for use with young adults. *J Fam.* 2002; **2**:109–131. http://www.tandfonline.com/doi/abs/10.1207/S15327698JFC0203_1.
38. Yihun BA, Kibret GD, Leshargie CT. **Correction: Incidence and predictors of treatment failure among children on first-line antiretroviral therapy in Amhara Region Referral Hospitals, northwest Ethiopia 2018: a retrospective study.** *PLoS One* 2019; **14**:e0217901.
39. Statistics South Africa. Census 2011: household questionnaire. Pretoria; 2011.
40. Pillay U, Roberts B, Rule S. *South African social attitudes: changing times, diverse voices.* Cape Town: HSRC Press; 2006.
41. Firebaugh G, Warner C, Massoglia M. **Fixed effects, random effects, and hybrid models for causal analysis.** In *Handbook of causal analysis for social research.* 2013:113–132. Dordrecht, The Netherlands: Springer Science + Business Media. Available at <https://doi.org/10.1007/978-94-007-6094-3>
42. Dieleman JL, Templin T. **Random-effects, fixed-effects and the within-between specification for clustered data in observational health studies: a simulation study.** *PLoS One* 2014; **9**:e110257.
43. Bell A, Fairbrother M, Jones K. **Fixed and random effects models: making an informed choice.** *Qual Quan* 2019; **53**:1051–1074.
44. Allison PD. *Fixed effects regression models,* vol. 160. Los Angeles, USA: Sage Publications; 2009.
45. Bland J, Cook A. **Random effects probit and logit: understanding predictions and marginal effects.** *Appl Econ Lett* 2019; **26**:116–123.
46. Williams R. **Using the margins command to estimate and interpret adjusted predictions and marginal effects.** *Stata J* 2012; **12**:308–331.
47. UNAIDS. *90-90-90: an ambitious treatment target to help end the AIDS epidemic.* Geneva: UNAIDS; 2014, Available at https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=ahUKEwievbXujrjvAhWIZhUIHdXICZsQFjAEegQIARAD&url=https%3A%2F%2Fwww.unaids.org%2Ffiles%2Fdefault%2Ffiles%2Fsub_landing%2F90-90-90_en.pdf&usq=AOvVaw0-iEYrTVs2RTj-NUFb0Mos. [Accessed 17 March 2021]
48. Burton P, Ward CL, Artz L. *The optimus study on child abuse, violence and neglect in South Africa.* Cape Town: The Centre for Justice and Crime Prevention; 2015.
49. World Health Organisation. *INSPIRE: seven strategies for ending violence against children.* Geneva: World Health Organisation; 2016.
50. WHO. *Global status report on preventing violence against children.* Geneva: WHO; 2020.
51. Cluver L, Meinck F, Steinert J, Shenderovich Y, Doubt J, Romero R, et al. **Parenting for lifelong health: a pragmatic cluster randomised controlled trial of a noncommercialised parenting programme for adolescents and their families in South Africa.** *BMJ Global Health* 2018; **3**:e000539.
52. Ward CL, Wessels IM, Lachman JM, Hutchings J, Cluver LD, Kassinjee R, et al. **Parenting for lifelong health for young children: a randomized controlled trial of a parenting program in South Africa to prevent harsh parenting and child conduct problems.** *J Child Psychol Psychiatry* 2020; **61**:503–512.
53. Annan J, Bunderviet T, Seban J, Costigan J. *Urwaruka Rushasha: a randomised impact evaluation of village savings and loans associations and family-based interventions in Burundi.* Washington, DC: International Rescue Committee and USAID; 2013.
54. Shenderovich Y, Eisner M, Cluver L, Doubt J, Berezin M, Majokweni S, et al. **What affects attendance and engagement in a parenting program in South Africa?** *Prev Sci* 2018; **19**:977–986.
55. Shenderovich Y, Cluver L, Eisner M, Murray AL. **Moderators of treatment effects in a child maltreatment prevention programme in South Africa.** *Child Abuse Negl* 2020; **106**:104519.
56. Redfern A, Cluver LD, Casale M, Steinert JI. **Cost and cost-effectiveness of a parenting programme to prevent violence against adolescents in South Africa.** *BMJ Glob Health* 2019; **4**:e001147.
57. Goldenring J, Cohen E. **Getting into adolescent heads.** *Contemp Pediatr* 1988; **5**:75–90.
58. World Health Organisation, UNAIDS, UNICEF, UNESCO, Women U. *Global accelerated action for the health of adolescents (AA-HA!).* Geneva: WHO; 2017.
59. Mark D, Armstrong A, Andrade C, Penazzato M, Hatane L, Taing L, et al. **HIV treatment and care services for adolescents: a situational analysis of 218 facilities in 23 sub-Saharan African countries.** *J Int AIDS Soc* 2017; **20** (Suppl 3):21591.
60. Grimwood A, Fatti G, Mothibi E, Malahlela M, Shea J, Eley B. **Community adherence support improves programme retention in children on antiretroviral treatment: a multicentre cohort study in South Africa.** *J Int AIDS Soc* 2012; **15**:17381.
61. Cluver L, Lachman JM, Sherr L, Wessels I, Krug E, Rakotomalala S, et al. **Parenting in a time of COVID-19.** *Lancet* 2020; **395**:e64.
62. Perks B, Cluver L. **The parenting vaccine.** *Nat Hum Behav* 2020; **4**:985. doi:10.1038/s41562-020-0932-8.
63. Haghight R, Steinert J, Cluver L. **The effects of decentralising antiretroviral therapy care delivery on health outcomes for adolescents and young adults in low- and middle-income countries: a systematic review.** *Glob Health Action* 2019; **12**:1668596.
64. WHO, UNAIDS. *Global standards for quality health-care services for adolescents: a guide to implement a standards-driven approach to improve the quality of healthcare services for adolescents. Volume 1: Standards and criteria.* Geneva: WHO; 2015.
65. Johnson D. **Two-wave panel analysis: comparing statistical methods for studying the effects of transitions.** *J Marriage Fam* 2005; **67**:1061–1075.
66. Vaisey S, Miles A. **What you can: and can't: do with three-wave panel data.** *Sociol Meth Res* 2017; **46**:44–67.
67. Akmatov MK. **Child abuse in 28 developing and transitional countries: results from the Multiple Indicator Cluster Surveys.** *Int J Epidemiol* 2011; **40**:219–227.
68. Merrill KG, Campbell JC, Kennedy CE, Burke VM, Miti S, Frimpong C, et al. **'So hurt and broken': a qualitative study of experiences of violence and HIV outcomes among Zambian youth living with HIV.** *Global Public Health* 2021:1–13.
69. Statistics SA. Provincial profile: Eastern Cape <http://cs2016.stats-sa.gov.za/wp-content/uploads/2018/07/EasternCape.pdf>. Pretoria: Statistics South Africa; 2018.
70. Simbayi L, Zuma K, Zungu N, Moyo S, Marinda E, Jooste S, et al. *South African National HIV prevalence, incidence, behaviour and communication survey, 2017.* http://www.hsrc.ac.za/en/departments/hsc/National_HIV_Survey. Cape Town: HSRC Press; 2019.