

REVIEW

Open Access

# Phytochemical and pharmacological progress on the genus *Syringa*

Guozhu Su<sup>1,2</sup>, Yuan Cao<sup>1,2</sup>, Chun Li<sup>1</sup>, Xuelong Yu<sup>1,2</sup>, Xiaoli Gao<sup>1</sup>, Pengfei Tu<sup>1</sup> and Xingyun Chai<sup>1\*</sup>

## Abstract

Genus *Syringa*, belonging to the Oleaceae family, consists of more than 40 plant species worldwide, of which 22 species, including 18 endemic species, are found in China. Most *Syringa* plants are used in making ornaments and traditional medicines, whereas some are employed for construction or economic use. Previous studies have shown that extracts of *Syringa* plants mainly contain iridoids, lignans, and phenylethanoids that have antitumor, antihypertensive, anti-oxidant, and anti-inflammatory activities. This study reviews phytochemical and pharmacological progress on *Syringa* in the recent 20 years and discusses the future research prospects to provide a reference in further promotion and application of the genus.

**Keywords:** *Syringa*, Oleaceae, Iridoid, Lignan, Phenylethanoid, Bioactivities, Review

## Introduction

Plants belonging to the family Oleaceae, which consists of 27 genera and 400 species worldwide, have important applications in the daily life of people living in developing countries. Plants of many well-known genera, including *Forsythia*, *Syringa*, and *Osmanthus*, have been widely used for medicinal and industrial purposes. For instance, the stems and roots of *S. pinnatifolia* var. *alashanensis* is the major composition of a traditional formula 'Ba wei chen-xiang' powder that is used for treatment of asthma, cardiopalmus, and angina [1].

Most *Syringa* plants are deciduous shrubs and arbors and include more than 40 species distributed around Europe and Asia [2]. At present, 22 species are found in China, of which 18 are endemic species that are mainly distributed in the southwestern part of Sichuan, Yunnan, Tibet, and other Northwestern regions. Many *Syringa* species, such as *S. chinensis*, *S. meyeri*, and *S. pekinensis*, are used for making ornaments. Flowers of *S. oblata* and *S. reticulata* var. *mandshurica* are an ideal source of aroma oils or nectar. Some *Syringa* plants are also used for construction purposes or for manufacturing furniture [1].

Previous phytochemical studies on *Syringa* species have revealed the presence of more than 140 secondary

metabolites, including iridoids, lignans, phenylethanoids, their glycosides, minor organic acids, and essential oils [3,4]. Modern pharmacological studies have shown the bioactivities of these metabolites, such as antitumor, antihypertensive, anti-oxidant, anti-inflammatory activities, and so on [5]. However, a systematic review of these studies has not been performed to date. This review summarizes the phytochemical and pharmacological progress on *Syringa* to date by focusing on its chemical classification, structural features, and biological and pharmacological applications to provide information for further research on this genus.

## Chemical constituents

Previous studies have reported that extracts of *Syringa* plants contain iridoids (1–46), lignans (47–80), phenylpropanoids (81–105), phenylethanoids (106–121), and other compounds (122–142). The structures of these compounds are shown in Figures 1, 2, and 3 and related information are listed in Tables 1, 2, and 3.


## Iridoids

Iridoids are one of the most important natural compounds that are widely distributed in various plant families such as Plantaginaceae, Rubiaceae, and Scrophulariaceae [6]. Iridoids are extensively present in almost all *Syringa* species and have antitumor, antihypertensive, anti-inflammatory, anti-oxidant, and antifungal activities. In addition, iridoids

\* Correspondence: xingyunchai@yeah.net

<sup>1</sup>Modern Research Center for Traditional Chinese Medicine, Beijing University of Chinese Medicine, 11 North 3rd Ring Road, Chaoyang District, Beijing 100029, P. R. China

Full list of author information is available at the end of the article




**Figure 1** The structures of iridoids from the genus *Syringa*.

play an important role in defense mechanism of ants [7]. Among all the iridoids reported in this genus, secoiridoids are the most abundant and have been shown to have anti-tumor activity. To date, 46 iridoids (1–46) have been described, including secoiridoids (1–30 and 40–44), eight typical iridoids (32–39), and three minor dimers (31, 45, and 46). Most iridoids exist as glycosides and are mainly produced by the glycosylation of glucose and galactose. *Syringa* iridoids are generally substituted by various acid fragments and phenolic moieties such as 1-*O*-cinnamoyl- $\beta$ -D-glucopyranosyl, *p*-hydroxyphenethyl, 3, 4-dihydroxy-phenethyl, and caffeic acid, which contribute to their low polarity. *Syringa* iridoids have antitumor (33 and 40) [8,9], antihypertensive (4), and anti-oxidant (4 and 31) activities [10].

## Lignans

Lignans are another major compounds in this genus, particularly in *S. komarowii* [27], *S. pubescens* [3], *S. reticulata* [10], *S. velutina* [28], *S. patula* [5], *S. vulgaris* [29], *S. pinatifolia* var. *alashanensis* [30,31], and *S. reticulata* var. *mandshurica* [32]. *Syringa* species have 34 lignans and their glycosides (47–80), including monoepoxylic lignans (47–60, 62) and their dimers (63 and 64), neolignans (61, 73–74), cyclolignans (65 and 66), simple lignans (67–72), and bisepoxylic lignans (75–80). Lignans also exhibit many bioactivities. For example, compound 50 has anti-oxidant activity [10]; compounds 57 and 58 have antifungal activities [32]; and compound 75 has significant cytotoxic, antihypertensive, anti-inflammatory, and anti-oxidant activities [5].



**Figure 2** The structures of lignans from the genus *Syringa*.

### Other compounds


Phenylethanoids (**81–105**), phenylpropanoids and their analogues (**106–121**), flavonoids (**122–128**), sesquiterpenes (**129** and **130**), and other minor compounds have been described in *Syringa* plants. Of these, phenylethanoids are predominant, particularly in *S. reticulata* [10,12,35], *S. vulgaris* [29], *S. pubescens* [3], *S. oblata* var. *alba* [36], *S. reticulata* var. *mandshurica* [35], *S. afghanica* [13], and *S. komarowii* [27]. Sesquiterpenes (**129** and **130**) are present in the stems of *S. pinnatifolia* var. *alashanensis* [37]. These miscellaneous compounds have cytotoxic, anti-inflammatory, antihypertensive, anti-oxidant, and antifungal properties.

Besides the abovementioned compounds, *Syringa* plants contain essential oils that form the most important constituents not only because of their economic utility but also because of their potential medicinal value as antimicrobial, antipyretic, and antiviral agents. Multiple analytical techniques such as headspace solid-phase microextraction, gas chromatography–mass spectrometry (GC–MS), GC–MS coupled with heuristic evolving latent projections, moving subwindow searching, nuclear magnetic

resonance spectroscopy, and X-ray single-crystal diffraction analysis have been used to identify essential oils from fresh flowers of *S. oblata* var. *alba*. For instance, 39 volatile oil constituents were identified, including four characteristic isomers of lilac alcohols (lilac alcohols A–D) and lilac aldehydes A–D [38]. Ninety-five components, including 15 terpenes, 14 oxygenated terpenes, 10 aromatic compounds, and 13 *n*-alkanes were quantitatively analyzed from *S. oblata* buds [39]. Forty-nine components were described from essential oil of *S. pubescens* flowers, most of which are monoterpenes and sesquiterpenes [40]. Thirty-four volatile oil components, accounting for around 64.7% (zerumbone) of the oil, were identified from roots and barks of *S. pinnatifolia* var. *alashanensis* [4]. These data imply that *Syringa* plants could be considerably different from each other in terms of their essential oil components.

### Pharmacological activities

Various crude extracts and isolated compounds from *Syringa* plants have shown significant antitumor, antihypertensive, anti-inflammatory, anti-oxidant, and antifungal activities.



**Figure 3** The structures of other type of compounds from the genus *Syringa*.

**Table 1 Iridoids from the genus *Syringa***

No	Compound	Part of plants	Source	Reference
1	Isoligustroside	leaves	<i>S. vulgaris</i>	[11]
2	Isooleuropein	leaves	<i>S. vulgaris</i>	[11]
3	Oleoside 11-methyl ester	flowers, leaves and floral buds	<i>S. pubescens</i> <i>S. patula</i>	[3,5]
4	Oleuropein	flowers, leaves, barks and floral buds	<i>S. pubescens</i> <i>S. reticulata</i> , <i>S. dilatata</i> , <i>S. velutina</i> , <i>S. afghanica</i> , <i>S. oblata</i> var. <i>alba</i> , <i>S. patula</i>	[3,5,8,10,12-14]
5	Neooleuropein	leaves	<i>S. vulgaris</i>	[15]
6	8(E)-Ligstroside	flowers, leaves and barks	<i>S. pubescens</i> , <i>S. reticulata</i> , <i>S. dilatata</i> , <i>S. afghanica</i>	[3,8,10,13]
7	8(E)-Nüzhenide	leaves	<i>S. reticulata</i>	[16]
8	Safghanoside A	leaves	<i>S. afghanica</i>	[13]
9	Safghanoside B	leaves	<i>S. afghanica</i>	[13]
10	Safghanoside C	leaves	<i>S. afghanica</i>	[13]
11	Safghanoside D	leaves	<i>S. afghanica</i>	[13]
12	Safghanoside E	leaves	<i>S. afghanica</i>	[13]
13	Safghanoside F	leaves	<i>S. afghanica</i>	[13]
14	Formoside	leaves	<i>S. afghanica</i>	[13]
15	Fraxiformoside	leaves	<i>S. afghanica</i>	[13]
16	2"- <i>epi</i> -framoside	leaves	<i>S. afghanica</i>	[13]
17	1'''- <i>O</i> - $\beta$ -D-glucosylformoside	leaves	<i>S. afghanica</i>	[13]
18	1'''- <i>O</i> - $\beta$ -D-glucosylfraxiformoside	leaves	<i>S. afghanica</i>	[13]
19	Lilacoside	barks and leaves	<i>S. vulgaris</i>	[17,18]
20	Fliederoside	barks and leaves	<i>S. vulgaris</i>	[17,18]
21	8(Z)-Ligstroside	leaves	<i>S. reticulata</i>	[16]
22	8(Z)-Nüzhenide	leaves	<i>S. reticulata</i>	[16]
23	Oleoside dimethyl ester	leaves	<i>S. afghanica</i>	[13]
24	10-Hydroxyoleuropein	flowers and leaves	<i>S. pubescens</i>	[3]
25	10-Hydroxyoleoside dimehyl ester	flowers and leaves	<i>S. pubescens</i>	[3]
26	Secologanoside 7-methyl ester	leaves	<i>S. reticulata</i>	[19]
27	Grandifloroside 11-methyl ester	flowers and leaves	<i>S. pubescens</i>	[3]
28	8-Epikingiside	barks	<i>S. vulgaris</i>	[20]
29	Syrveoside A	leaves	<i>S. velutina</i>	[21]
30	Syrveoside B	leaves	<i>S. velutina</i>	[21]
31	Jaspolyoside	barks	<i>S. reticulata</i>	[10]
32	Syringopicroside	leaves	<i>S. dilatata</i> , <i>S. vulgaris</i> , <i>S. oblata</i> , <i>S. reticulata</i>	[8,16,19,22,23]
33	Syringopicroside B	leaves	<i>S. vulgaris</i>	[9]
34	3'- <i>O</i> - $\beta$ -D-glucopyranosylsyring-opicroside	leaves	<i>S. reticulata</i>	[16]
35	4'- <i>O</i> - $\beta$ -D-glucopyranosylsyring-opicroside	leaves	<i>S. reticulata</i>	[16]
36	6'- <i>O</i> - $\alpha$ -D-glucopyranosylsyring-opicroside	leaves	<i>S. reticulata</i>	[16]
37	6'- <i>O</i> - $\alpha$ -D-galactopyranosylsyring-opicroside	leaves	<i>S. reticulata</i>	[19]
38	Syringopicrogenin C	seeds	<i>S. oblata</i>	[24]
39	Syringopicrogenin A	seeds and crust	<i>S. oblata</i>	[24,25]

**Table 1 Iridoids from the genus *Syringa* (Continued)**

40	Isooleoacteoside	leaves	<i>S. vulgaris</i>	[9]
41	Oleoacteoside	leaves	<i>S. reticulata</i>	[9,26]
42	Oleoechinacoside	leaves	<i>S. reticulata</i>	[9,26]
43	Reticuloside	barks	<i>S. reticulata</i>	[10]
44	Jasminoside	whole plant	<i>S. komarowii</i>	[27]
45	Safghanoside H	leaves	<i>S. afghanica</i>	[13]
46	Safghanoside G	leaves	<i>S. afghanica</i>	[13]

### Antitumor activity

Cytotoxic activities of crude extracts and chemicals obtained from *Syringa* plants have been extensively evaluated against various tumor cell lines. Aqueous extracts from the flowers and leaves of *S. pubescens* inhibited the growth of L2215 (hepatitis B virus) cells, with a 50% inhibitory concentration ( $IC_{50}$ ) value of 78  $\mu\text{g}/\text{mL}$  [51]. Hydrolysis of isoligustroside (**1**) and isooleuropein (**2**) were assayed using a disease-oriented panel of 39 human cancer cell lines. The results showed that the hydrolysis product of compound **2** had moderate cytotoxic activity against lung cancer cell lines DMS273 [ $\log GI_{50} = 5.19$  (6.4  $\mu\text{M}$ )] and DMS114 [ $\log GI_{50} = 5.06$  (8.7  $\mu\text{M}$ )]. Preliminary analysis of structure–activity relationship suggested that C-5'-OH plays an important role in this cytotoxic activity [11]. Isooleoacteoside (**40**) showed weak cytotoxicity against LOX-IMVI melanoma cell line, with  $GI_{50}$  value of 16  $\mu\text{M}$ , and syringopicroside B (**33**) showed weak cytotoxic activity against NCI-H522 lung cancer cell line, with  $GI_{50}$  value of 13  $\mu\text{M}$  [9]. MTT assay used to assess the cytotoxicities of syringaresinol (**78**) and oleoside 11-methyl ester (**3**) showed that compound **78** had a strong dose-dependent effect on HepG2 cell line, with an  $IC_{50}$  value of 94.6  $\mu\text{M}$ , and compound **3** has a dose–response curve of low slope, with a high  $IC_{50}$  value of 186.5  $\mu\text{M}$ , compared with positive controls dexamethasone ( $IC_{50}$  14.2  $\mu\text{M}$ ) and paclitaxel ( $IC_{50}$  700 nM). However, compound **78** was cytotoxic even at the lowest concentration of 29.9  $\mu\text{M}$ .  $\beta$ -Amyrin acetate (**139**) showed weak cytotoxicity against A2780 human ovarian cancer and HepG2 cell lines [5]. Oleuropein (**4**) and 2-(3, 4-dihydroxy)-phenylethyl- $\beta$ -D-glucopyranoside (**83**) showed evident cytotoxicities against P-388, L-1210, SNU-5, and HL-60 cell lines, with  $IC_{50}$  values varying from 8.5 to 139.8  $\mu\text{M}$  [12]. Verbascoside (**86**) showed moderate cytotoxic activity against SNB-75 (brain cancer) and SNB-78 cell lines, with  $GI_{50}$  values of 7.4 and 7.7  $\mu\text{M}$ , respectively [9]. A pharmacokinetic study showed that compound **86** interacted with the catalytic domain of PKC and acted as a competitive inhibitor of adenosine triphosphate ( $K_i = 22 \mu\text{M}$ ) and non-competitive inhibitor of phosphate acceptor (histone III). Because **83** is one part of

**86** in its molecular structure, the cytotoxic effect could be attributed to 3, 4-dihydroxyphenylethoxy moiety, which may act as a competitive inhibitor to the catalytic domain of PKC. Therefore, **83** is a potentially essential skeleton of most cytotoxic phenylethanoid glycosides [12].

### Hypotensive activity

Syringin (**110**) and kaempferol-3-*O*-rutinoside (**125**) showed antihypertensive activity. Intravenous injection of 10 mg/kg of compound **86** significantly decreased systolic, diastolic, and mean arterial blood pressure in Pentothal-anesthetized rats. Moreover, the depressor effect of compound **86** was independent of muscarinic and histaminergic receptors because it did not block the effect of atropine (an antimuscarinic agent) and chlorpheniramine/cimetidine (antihistaminergic agents) [36]. *In vitro* studies showed that oleuropein (**4**) significantly lowered blood pressure. It is interesting to note that antihypertensive effect of compound **4** (33% at 30 mg/kg dose) on the blood pressure of anesthetized rats was similar to that of compound **86** (39.04%  $\pm$  2.38% at 10 mg/kg dose) [14,36], which is probably because of the similarity in their structures, with both possessing the same aromatic fragment having two hydroxy groups.

### Anti-inflammatory activity

Iridoid glycosides (IGs) exerted obvious anti-inflammatory effects on ulcerative colitis *in vivo* by inhibiting relative proinflammatory cytokines [53]. IGs significantly ameliorated macroscopic damages and histological changes, reduced the activity of myeloperoxidase, and strongly inhibited epithelial cell apoptosis. Moreover, IGs markedly decreased the levels of tumor necrosis factor- $\alpha$ , interleukin-8, cyclooxygenase-2, and transforming growth factor- $\beta$ 1 in colonic tissues in a dose-dependent manner. Moreover, effects of IGs (160 and 240 mg/kg) were superior to those of positive control salicylazosulfapyridine (150 mg/kg). Furthermore, IGs significantly blocked NF- $\kappa$ B signaling by inhibiting inflammatory bowel phosphorylation/degradation and inhibitor kappa B kinase  $\beta$  activity; downregulated protein and mRNA expressions of Fas/FasL, Bax, and caspase-3; and activated Bcl-2 in intestinal epithelial cells



**Table 2 Lignans from the genus *Syringa***

No	Name	Part of plant	Source	Reference
47	(-)-Olivil	whole plant	<i>S. komarowii</i>	[27]
48	Olivil 4-O- $\beta$ -D-glucopyranoside	barks	<i>S. reticulata</i> , <i>S. patula</i>	[10]
49	Olivil 4''-O- $\beta$ -D-glucopyranoside	barks	<i>S. reticulata</i>	[27]
50	Armandiside	barks	<i>S. reticulata</i>	[10]
51	Syripinnalignan A	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[31]
52	Syripinnalignan B	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[31]
53	(8R, 8'R, 9S)-4-4'-dihydroxy-3, 3', 9-trimethoxy-9-9'-epoxylignan	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30]
54	(8R, 8'R, 9R)-4-4'-dihydroxy-3, 3', 9-trimethoxy-9-9'-epoxylignan	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30]
55	(8S, 8'S, 9R)-4-4'-dihydroxy-3, 3', 9-trimethoxy-9-9'-epoxylignan	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30]
56	(8S, 8'S, 9S)-4-4'-dihydroxy-3, 3', 9-trimethoxy-9-9'-epoxylignan	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30]
57	Mandshuricol A	leaves	<i>S. reticulata</i> var. <i>mandshurica</i>	[32]
58	Mandshuricol B	leaves	<i>S. reticulata</i> var. <i>mandshurica</i>	[32]
59	(+)-Lariciresinol	seeds crust	<i>S. oblata</i>	[25]
60	(+)-Lariciresinol 4-O- $\beta$ -D-glucopyranoside	barks	<i>S. vulgaris</i>	[29]
61	Balanophonin	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30]
62	(+)-Lariciresinol 4'-O- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranoside	leaves	<i>S. reticulata</i>	[19]
63	Syripinnalignan A	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[33]
64	Syripinnalignan B	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[33]
65	Cycloolivil 6-O- $\beta$ -D-glucoside	barks	<i>S. reticulata</i>	[10]
66	(+)-Cycloolivil	whole plant	<i>S. komarowii</i>	[27]
67	(-)-Secoisolariciresinol	stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30,31]
68	PiperphilippininVI	roots and stems	<i>S. pinnatifolia</i>	[30]
69	Dihydrocubebin	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30]
70	Syripinnalignan C	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[34]
71	Syripinnalignan D	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[34]
72	Syripinnalignan E	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[34]
73	(7S, 8R)-Guaiacylglycerol-8-O-4'-sinapyl ether 9'-O- $\beta$ -D-glucopyranoside	leaves	<i>S. velutina</i>	[28]
74	(7S, 8R)-Syringylglycerol-8-O-4'-sinapyl ether 9'-O- $\beta$ -D-glucopyranoside	leaves	<i>S. velutina</i>	[28]
75	Pinoresinol-4-O- $\beta$ -monoglycoside	barks	<i>S. reticulata</i>	[10]
76	Syringaresinol-4-O-bis- $\beta$ -D-monoglycoside	barks	<i>S. reticulata</i>	[10]
77	Syringaresinol-4, 4''-O-bis- $\beta$ -D-glucoside	barks	<i>S. reticulata</i>	[10]
78	Syringaresinol	floral buds, flowers and leaves	<i>S. patula</i> , <i>S. pubescens</i>	[3,5]
79	(+)-Medioresinol-4-O-glucoside	floral buds	<i>S. patula</i>	[5]
80	(-)-Pinoresinol	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30]

[53,54].  $\beta$ -Amyrin acetate (**139**) and syringaresinol (**78**) at a dose of 20  $\mu$ g/mL evidently inhibited lipopolysaccharide-induced nitric oxide (NO) production, with inhibition rates of 49.97% and 33.21%, respectively [5].

#### Liver-protective and cholagogic effects

Crude extract of *Syringa* species, interferon (IFN), and an injection of "Gan-Yan-Ling" were compared to

evaluate their liver-protective effects on the survival rates of HepG2.215 cells and secretion of hepatitis B surface antigen (HBsAg) and HBeAg. The results indicated that all the three assayed drugs may suppress the secretion of HBsAg and HBeAg from HepG2.215 cells in a dose-dependent manner, with the effect of crude extract of *Syringa* being intermediate those of IFN and Gan-Yan-Ling. Therefore, extracts of *Syringa* plant

**Table 3 Other type of compounds from the genus *Syringa***

No	Name	Part of plant	Source	Reference
81	Isosyringalide	leaves	<i>S. reticulata</i>	[41]
82	Forsythiaside	barks	<i>S. vulgaris</i>	[29]
83	2-(3, 4-dihydroxy)-phenylethyl- $\beta$ -D-glucopyranoside	barks	<i>S. reticulata</i>	[10,12]
84	<i>cis</i> -Echinacoside	leaves	<i>S. reticulata</i>	[35]
85	Isoverbascoside	leaves	<i>S. pubescens</i>	[3]
86	Verbascoside	leaves	<i>S. pubescens</i> , <i>S. oblata</i> var. <i>alba</i> , <i>S. vulgaris</i>	[3,9,29,14,36]
87	Echinacoside	barks, leaves and flowers	<i>S. pubescens</i> , <i>S. reticulata</i> <i>S. vulgaris</i>	[3,29,42]
88	Forsythoside B	leaves	<i>S. reticulata</i> var. <i>mandshurica</i>	[35]
89	Salidroside	barks	<i>S. reticulata</i>	[10]
90	3'-O- $\beta$ -D-glucopyranosylsalidroside	leaves	<i>S. reticulata</i> var. <i>mandshurica</i>	[35]
91	2-(3, 4-dihydroxyphenyl) ethanol	leaves	<i>S. pubescens</i>	[3]
92	Osmanthuside F	leaves	<i>S. reticulata</i>	[35]
93	(S)-(+)-2-(3, 4-dihydroxyphenyl)-2-ethoxyethanol	leaves	<i>S. reticulata</i> var. <i>mandshurica</i>	[43]
94	(S)-(+)-2-(3, 4-dihydroxyphenyl)-2-acetoxyethanol	leaves	<i>S. reticulata</i> var. <i>mandshurica</i>	[43]
95	Decaffeoylacteoside	leaves	<i>S. reticulata</i>	[35]
96	Syringalide B	leaves	<i>S. reticulata</i>	[41]
97	Poliumoside	leaves	<i>S. afghanica</i>	[13]
98	2-(4-hydroxyphenyl)-ethyl behenate	whole plant	<i>S. komarowii</i>	[27]
99	2-(4-hydroxyphenyl)-ethyl tricosanoate	whole plant	<i>S. komarowii</i>	[27]
100	2-(4-hydroxyphenyl)-ethyl lignocerate	whole plant	<i>S. komarowii</i>	[27]
101	2-(4-hydroxyphenyl)-ethyl pentacosanoate	whole plant	<i>S. komarowii</i>	[27]
102	2-(4-hydroxyphenyl)-ethyl hexacosanoate	whole plant	<i>S. komarowii</i>	[27]
103	Bongardol	whole plant	<i>S. komarowii</i>	[27]
104	2-(4-hydroxyphenyl)-ethyl 1-dodecyloctadecanoate	whole plant	<i>S. komarowii</i>	[27]
105	2-(4-hydroxyphenyl)-ethyl dotriacontanoate	whole plant	<i>S. komarowii</i>	[27]
106	Coniferin	barks	<i>S. vulgaris</i>	[29]
107	Coniferylaldehyde	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[44]
108	Coniferylaldehyde glucoside	barks	<i>S. reticulata</i>	[10]
109	Sinapaldehyde glucoside	barks	<i>S. reticulata</i>	[10]
110	Syringin	barks	<i>S. vulgaris</i> , <i>S. reticulata</i>	[10,45,46]
111	Isosyringinioside	barks	<i>S. reticulata</i>	[10]
112	Eugenol	foral buds	<i>S. patula</i>	[5]
113	Larixnaphthano	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[30]
114	Cinnamic acid	leaves, roots and stems	<i>S. afghanica</i> , <i>S. pinnatifolia</i> var. <i>alashanensi</i> , <i>S. reticulata</i>	[44,47]
115	Caffeic acid	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[44]
116	Ferulic acid	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[44]
117	7-Methoxycoumarin	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[44]
118	Esculetine	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[44]
119	Umbelliferone	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[44]
120	O-[[ $\beta$ -D-xylopyranosyl (1-6) $\beta$ -D-glucopyranosyl]-7-hydroxycoumarin	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[44]
121	Syringfghanoside	leaves	<i>S. afghanica</i>	[13]
122	Astragalinalin	bark	<i>S. vulgaris</i>	[48]



**Table 3 Other type of compounds from the genus *Syringa* (Continued)**

123	Kaempferol-3, 7- $\alpha$ -L-dirhamnoside	flowers and leaves	<i>S. pubescens</i>	[3]
124	Kaempferol-3- $\beta$ -D-glucoside-7- $\alpha$ -L-dirhamnoside	flowers and leaves	<i>S. pubescens</i>	[3]
125	Kaempferol-3-O-rutinoside	flowers	<i>S. vulgaris</i>	[49]
126	Luteolin	leaves	<i>S. afghanica</i>	[13]
127	Rutin	leaves	<i>S. vulgaris</i>	[49,50]
128	Rhoifolin	leaves	<i>S. afghanica</i>	[13]
129	Guai-9-en-4 $\beta$ -ol	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[37]
130	14, 15-dinorguai-1, 11-dien-9, 10-dione	roots and stems	<i>S. pinnatifolia</i> var. <i>alashanensis</i>	[37]
131	Momorcerebroside I	whole plant	<i>S. komarowii</i>	[27]
132	Phytolacca cerebroside	whole plant	<i>S. komarowii</i>	[27]
133	Pubescenside A	flowers and leaves	<i>S. pubescens</i>	[51]
134	Stigmastane-3 $\beta$ , 6 $\alpha$ -diol 3-O-tetradecanoate	whole plant	<i>S. komarowii</i>	[27]
135	Stigmastane-3 $\beta$ , 6 $\alpha$ -diol 3-O-palmitate	whole plant	<i>S. komarowii</i>	[27]
136	Stigmastane-3 $\beta$ , 6 $\alpha$ -diol 3-O-stearate	whole plant	<i>S. komarowii</i>	[27]
137	$\beta$ -sitosterol	foral buds and whole plant	<i>S. patula</i> , <i>S. komarowii</i>	[5,27]
138	Daucosterol	whole plant	<i>S. komarowii</i>	[27]
139	$\beta$ -Amyrin acetate	foral buds	<i>S. patula</i>	[5]
140	Jasminidin	leaves	<i>S. vulgaris</i>	[52]
141	Jasminin	leaves	<i>S. vulgaris</i>	[52]
142	Nortropin	foral buds	<i>S. patula</i>	[5]

could be used to develop effective and less toxic anti-hepatitis B medicines [55].

Aqueous extracts of *S. reticulata* var. *mandshurica* significantly decreased the levels of alanine transaminase and aspartate transaminase and the concentration of malondialdehyde in the serum but increased the activity of superoxide dismutase (SOD) in the liver. These extracts showed protective effects on acute liver injury induced by CCl<sub>4</sub> in mice [56]. In addition, the essential oils of *Syringa* exerted protective effects on the liver and cholecyst [39].

#### Antifungal activity

Phenylpropanoids such as verbascoside (**86**) and forsythiaside (**82**) exhibit significant antimicrobial activity [29]. Compounds **93** and **94** at 1- mM concentration inhibited the radial growth of *Phytophthora capsici* after 6 days of incubation, with inhibition rates 59.1% and 72.5%, respectively [43]. Two sesquiterpenes, guai-9-en-4 $\beta$ -ol (**129**) and 4, 15-dinorguai-1, 11-dien-9, 10-dione (**130**), have antibacterial and antifungal properties. Compound **129** was active against *Bacillus coagulans* [inhibition zone (IZ) = 15.34 mm] and *Aspergillus niger* (IZ = 13.20 mm) while compound **130** significantly inhibited *Escherichia coli* (IZ = 15.34 mm) and *Fusarium oxysporum* (IZ = 15.32 mm) [37].

Compound **3** showed effective antimicrobial activity against *Lactobacillus pentosus* (IZ = 1 mm), and compound

**139** inhibited the growth of *Candida* species at concentrations of 30–250  $\mu$ g/mL [5].

#### Antioxidant activity

A 70% EtOH extract of *S. reticulata* barks showed potent superoxide anion and DPPH free radical scavenging activities, with EC<sub>50</sub> values of 5.88 and 38.10  $\mu$ g/mL, respectively [10].

Among the compounds isolated from the bark of *S. reticulata*, six (**4**, **31**, **50**, **77**, **83**, and **111**) showed significant superoxide anion scavenging activity, with EC<sub>50</sub> values of 2.57, 4.97, 10.64, 15.98, 4.97, and 14.14  $\mu$ g/mL, respectively. Compound **4** also interacted with the stable free radical DPPH, with an IC<sub>50</sub> value of 40.4  $\mu$ M [8,10]. These different anti-oxidant activities are closely related to their structural features. Presence of 2-(3, 4-dihydroxyphenyl)-ethoxy moiety might be important for a higher activity because the most potent compounds (EC<sub>50</sub> = 2.57–4.97  $\mu$ M), including the two secoiridoid glycosides (**31** and **4**) and a phenylethanoid glycoside (**83**), possess the same structural features. Comparison of the structures of compounds **4** and **83** with those of 8(Z)-ligstroside (**21**) and salidroside (**89**) showed that presence of *ortho*-coupling hydroxyl group at C-2 might be responsible for their different activities. It has been previously reported that 1, 2-dihydroxybenzene moiety is crucial to its DPPH scavenging activity [10].

Syringaresinol (**78**) showed a strong scavenging activity against DPPH, with  $EC_{50}$  value as low as 12.5  $\mu\text{g/mL}$ , which might be responsible for its strong inhibition of NO production [5].

Eugenol (**112**) inhibited the catalytic activity of  $\text{H}_2\text{O}_2/\text{Ca}^{2+}$  human erythrocyte membrane lipid peroxidation at a concentration of 200  $\mu\text{mol/L}$ , with an inhibition rate of 62%, and completely suppressed the catalytic activity of dibenzoyl peroxide/ $\text{Ca}^{2+}$  human erythrocyte membrane lipid peroxidation at a concentration of 100  $\mu\text{mol/L}$ . Compound **112** exerted its effect in a non-competitive manner by reacting with  $\text{Ca}^{2+}$  and inhibiting the formation of hydroxyl radicals, thus, protecting the cell membrane lipid from oxidation [2].

#### Inhibition of platelet aggregation

Aqueous extract of *S. aramaticum* significantly inhibited adenosine diphosphate (ADP) and collagen-induced platelet aggregation, with inhibition rates of 37.4% and 69.7%, respectively [57]. Mandshuricols A (**57**) and B (**58**) showed antagonistic activities on platelet-activating factor (PAF) in [3H]PAF receptor binding assay, with  $IC_{50}$  values of  $4.8 \times 10^{-5}$  and  $3.5 \times 10^{-5}$  M, respectively [32].

#### Others

Essential oils from the stems and roots of *S. pinnatifolia* var. *alashanensis* (SPEO) reduced the deviation of ST segment; decreased the levels of lactate dehydrogenase, creatine kinase, and troponin T; and increased the activity of SOD. These protective effects were further confirmed by histopathological examination [58]. Treatment with both 8 and 32 mg/kg SPEO prolonged the survival of mice under hypoxia conditions, showing a remarkable protective effect against  $\text{H}_2\text{O}_2$ -induced death in cultured rat myocytes. Moreover, 5, 2.5 and 1.25  $\mu\text{g/mL}$  doses of SPEO inhibited ADP-induced rat platelet aggregation by 47.4%, 37.0%, and 32.9%, respectively [58], implying that SPEO exerted protective effects against myocardial ischemia.

Oral and intraperitoneal administration of 0.2–0.4 g of leaf extract of *S. vulgaris* in cats or rabbits exerted an antipyretic effect that was equal to the effect of 0.1–0.3 g of aminopyrine administered orally or intraperitoneally. However, leaf extracts of *S. vulgaris* are considerably more toxic than aminopyrine, with their toxic dosages being 0.4 and 1.2 g/kg, respectively [59]. *In vitro* evaluation of leaf extract of *S. aramaticum* showed its antiviral activity against herpes simplex virus at concentrations 1.25%–2.5%. The protective effect was more obvious when controlling the amount of virus attacks at 9.2–92 tissue culture infective dose (TCID<sub>50</sub>), suggesting that *S. aramaticum* effectively killed the virus without any harmful side effects [60–62].

Studies have reported that leaf extracts of *S. aramaticum* could be used for treating hemorrhoids [63].

Eugenol (**112**) inhibited the metabolism of arachidonic acid. Extracts of *S. reticulata* var. *mandshurica* have been used for treating bronchitis, and one of its constituents 2-(3, 4-dihydroxyphenyl) ethanol (**91**) significantly inhibited the production of phlegm [2].

#### Review and conclusions

This review describes phytochemical and pharmacological progress on the genus *Syringa* in the recent 20 years and discusses the future research prospects.

*Syringa* plants are used not only as traditional medicines to treat rheumatoid arthritis, asthma, cardiopalmus, and angina pectoris by natives in China but also for making ornaments, volatile oils, food additives, and bactericides worldwide, particularly in developing countries. Previous phytochemical studies on crude extracts from various species of this genus have identified iridoids, lignans, phenylpropanoids, and phenylethanoids having antitumor, anti-hypertensive, anti-oxidant, and anti-inflammatory activities. Iridoids, lignans, and phenylethanoids are the most predominant compounds in *Syringa* plants that probably contribute independently or synergistically to their main biological activities.

To the best of our knowledge, 46 iridoid representatives have been reported in *Syringa* plants, with high concentrations present in the leaves of *S. vulgaris*, *S. pubescens*, *S. afghanica*, *S. reticulata*, and *S. velutina* and barks of *S. vulgaris* and *S. reticulata* and low concentrations present in the flowers (*S. pubescens*), seeds, and seeds crust (*S. oblata*). This difference may be associated with their ecological roles, because iridoids are produced mainly to fight predators and/or microbes. Moreover, high concentrations of lignans in the stems and roots can be attributed to the rigidity of these plants. This may be the reason for the absence of iridoids in *S. pinnatifolia* var. *alashanensis* because materials used for chemical investigation included peeled stems and roots. Anti-inflammatory effects of extracts from these plants are mainly responsible for their applications in traditional medicine. However, only preliminary work has been performed on most isolated compounds, such as *in vitro* cytotoxicity screening (**1**, **2**, **78**, and **139**). Limited studies have been performed on the *in vivo* effects of these compounds; thus, providing opportunities for further detailed research. It is particularly worthy to mention that China has an abundant resource of *Syringa*, with many endemic species. For instance, *S. pinnatifolia* var. *alashanensis* is a well-known Mongolian medicine traditionally used for myocardial ischemia in clinical practice. However, no substantial evidence is available on its bioactive ingredients and mechanisms of action underlying this effect. Therefore, it deserves further phytochemical and pharmacological studies.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

SG, CY, LC, GX, and YX have all been involved in preparing this review. SG, TP and CX are responsible for writing, checking and revising the manuscript. All authors read, discussed and approved final version of the manuscript.

### Acknowledgments

This paper is financially supported by the National Natural Science Foundation of China (No.81473426).

### Author details

<sup>1</sup>Modern Research Center for Traditional Chinese Medicine, Beijing University of Chinese Medicine, 11 North 3rd Ring Road, Chaoyang District, Beijing 100029, P. R. China. <sup>2</sup>School of Chinese Materia Medica, Beijing University of Chinese Medicine, 6 Wangjing Southern Middle Ring Road, Chaoyang District, Beijing 100102, P. R. China.

Received: 26 April 2014 Accepted: 7 January 2015

Published online: 27 January 2015

### References

- Editorial Committee of Chinese Material Medica. Chinese Material Medica-Mongolian volume. Shanghai Sci Technol Press. 1992;61:2–50.
- Zhang JF, Zhang SJ. An overview of the genus *Syringa*: phytochemical and pharmacological aspects. *Nat Sci J Hainan Univ.* 2007;2:201–5.
- Deng RX, Yuan H, Liu P, Yin WP, Wang XS, Zhao TZ. Chemical constituents from *Syringa pubescens* Turcz. *Biochem Syst Ecol.* 2010;38:813–5.
- Zhang GB, Liu LJ, Wang MK. Chemical constituents of volatile oils from *Syringa pinnatifolia*. *Chin Pharm J.* 1994;5:271.
- SamyK ED, GamalEldeen AM. Cytotoxic and anti-inflammatory activities of some constituents from the floral buds of *Syringa patula*. *Pharm Biol.* 2009;47:872–7.
- Dinda B, Debnath S, Harigaya Y. Naturally occurring iridoids. a review, part 1. *Chem Pharm Bull.* 2007;55:159–222.
- Ghisalberti EL. Biological and pharmacological activity of naturally occurring iridoids and secoiridoids. *Phytomedicine.* 1998;5:147–63.
- Oh H, Ko EK, Kim DH, Jang KK, Park SE, Lee HS, et al. Secoiridoid glucosides with free radical scavenging activity from the leaves of *Syringa dilatata*. *Phytother Res.* 2003;17:417–9.
- Masafumi K, Yasunri Y, Nariyasu M. Glycoside from the leaves of *Syringa vulgaris* and their growth inhibitory activity against human cancer cell lines. *Japan Soc Pharm.* 2010;64:104–5.
- Bi XY, Li W, Sasaki T, Li Q, Mitsuhashi N, Asada Y, et al. Secoiridoid glucosides and related compounds from *Syringa reticulata* and their anti-oxidant activities. *Bioorg Med Chem Lett.* 2011;21:6426–9.
- Kikuchi M, Yaoita Y, Mano N, Kikuchi M. Structure and cytotoxic activity of enzymatic hydrolysis products of secoiridoid glucosides, isoligustroside and isooleuropein. *Chem Biodivers.* 2011;8:651–7.
- Park HJ, Lee MS, Lee KT, Sohn IC, Han YN, Miyamoto K. Studies on constituents with cytotoxic activity from the stem bark of *Syringa velutina*. *Chem Pharm Bull.* 1999;47:1029–31.
- Yk T, Okazaki N, Tanahashi T, Nagakura N, Nishi T. Secoiridoid and iridoid glucosides from *Syringa afghanica*. *Phytochemistry.* 2002;59:779–87.
- Nenadis N, Vervoort J, Boeren S, Tsimidou MZ. *Syringa oblata* Lindl var. *alba* as a source of oleuropein and related compounds. *J Sci Food Agr.* 2007;87:160–6.
- Kikuch IM, Yamauchi Y, Yanase C. Structures of new secoiridoid glucosides from the leaves of *Syringa vulgaris* Linn. *Yakugakuzasshi.* 1987;107:245–8.
- Machida K, Kaneko A, Hosogai T, Kakuda R, Yaoita Y, Kikuchi M. Studies on the constituents of *Syringa* species. X. five new iridoid glycosides from the leaves of *Syringa reticulata* (Blume) Hara. *Chem Pharm Bull.* 2002;50:493–7.
- Sticher O, Ahmad M, Salama O, Winkler T. Two new secoiridoid glucosides from *Syringa vulgaris*. *Planta Med.* 1982;45:151.
- Kikuch IM, Yamauchi Y, Takahashi Y. Structures of new secoiridoid glycosides from the leaves of *Syringa vulgaris* Linn. *Yakugakuzasshi.* 1988;108:355–60.
- Machida K, Unagami E, Ojima H, Kikuchi M. Studies on the constituents of *Syringa* species. XII. new glycosides from the leaves of *Syringa reticulata* (Blume) Hara. *Chem Pharm Bull.* 2003;51:883–4.
- Garcia J, Lavaitte S, Gey C. 8-Epikingside and its vanillate ester, isolated from *Gentiana pyrenaica*. *Phytochemistry.* 1989;28:2199–201.
- Feng XS, Qu Y, Wu ZHXL, Zhang DS, Gao HY, Wu LJ. Two new secoiridoid glucosides from *Syringa velutina* Kom. *Chinese Chem Lett.* 2009;20:702–5.
- Asaka Y, Kamikawa T, Tokoroyama T, Kubota T. The structure and absolute configuration of syringopicroside. a new iridoid glucoside from *Syringa vulgaris*. *Tetrahedron.* 1970;26:2365–70.
- Zhou YM, Gao HY, Wu B, Li SH, Wu LJ. Structure determination of syringopicroside from the leaves of *Syringa oblata* Lindl. using 2D NMR spectral methods. *Asian J Tradit Med.* 2006;1:31–3.
- Zhang SJ, Guo HQ, Han J, Zhao M, Wang JJ. Chemical constituents from seeds of *Syringa oblata*. *Chin Tradit Herbal Drugs.* 2011;42:1894–900.
- Wang JL, Zhang GF, Dong LW, Zhao M, Zhang SJ. Chemical constituents from seed crust of *Syringa oblata*. *Chin Tradit Herbal Drugs.* 2010;41:1598–601.
- Kikuch IM, Yamauchi Y, Takahashi Y. Studies on constituents of *Syringa* species. VIII. Isolation and structures of phenylpropanoid glycosides from the leaves of *Syringa reticulata* (Blume) Hara. *Yakugakuzasshi.* 1989;109:366–71.
- Luo Y, Liu Y, Qi H, Wu Z, Zhang G. Steryl esters and phenylethanol esters from *Syringa komarowii*. *Steroids.* 2006;71:700–5.
- Feng XS, Qu Y, Xu L, Wang DC, Wu LJ, Meng FH, et al. Two new neolignans from *Syringa velutina* Kom. *Molecules.* 2009;14:953–8.
- Kurkin VA. Phenylpropanoids from medicinal plants: distribution, classification, structural analysis, and biological activity. *Chem Nat Compd.* 2003;39:123–53.
- Zeng XJ, Wang GC, Wu X, Li YL. Chemical constituents from *Syringa pinnatifolia*. *Chin Tradit Herbal Drugs.* 2013;7:1721–5.
- Ao WLJ, Bao XH, Wu XL, Wang QH. Lignans from *Syringa pinnatifolia* var. *alashanensis*. *J Asian Nat Prod Res.* 2012;14:396–400.
- Xu QM, Li Q, Liu YL, Feng YL, Yang SL, Li XR. New lignans from *Syringa reticulata* var. *mandshurica*. *Chem Nat Compd.* 2010;46:366–9.
- Wang QH, Ao WLJ, Wu QS, Ying X. Two new neolignans from the stems of *Syringa pinnatifolia* var. *alashanensis*. *Helv Chim Acta.* 2012;95:1680–5.
- Ao WLJ, Wang QH, Wang XL, Dai YNT. Extraction and structure elucidation of three lignans from peeled dried roots and stems of *Syringa pinnatifolia*. *Chin Tradit Herbal Drugs.* 2013;7:791–3.
- Machida K, Ohkawa N, Ohsawa A, Kikuchi M. Two new phenolic glycosides from *Syringa reticulata*. *J Nat Med.* 2009;63:192–4.
- Ahmad M, Rizwan GH, Aftab K, Ahmad VU, Gilani AH, Ahmad SP. Acteoside: a new antihypertensive drug. *Phytother Res.* 1995;9:525–7.
- Ao WLJ, Wang QH, Si Q, Mu D, Sa RTY, Dai NYT, et al. The structural elucidation and antimicrobial activities of two new sesquiterpenes from *Syringa pinnatifolia*. *Chin J Nat Med.* 2012;10:477–80.
- Li ZG, Lee MR, Shen DL. Analysis of volatile compounds emitted from fresh *Syringa oblata* flowers in different florescence by headspace solid-phase microextraction-gas chromatography-mass spectrometry. *Anal Chim Acta.* 2006;1:43–9.
- Chen XZ, Liang YZ, Fang HZ, Li XN. Temperature-programmed retention indices for gas chromatography-mass spectroscopy analysis of plant essential oils. *J Chromatogr A.* 2005;1096:76–85.
- Yu AN, Yang XZ. Chemical composition of the essential oil of fresh wild *Syringa pubescens* flowers from China. *Adv MaterRes (Durnten-Zurich, Switzerland).* 2012;22:581–2.
- Kikuch IM, Yamauchi Y. Studies the constituents of *Syringa* species. I. Isolation and structures of iridoids and secoiridoids from the leaves of *Syringa reticulata* (Blume) Hara. *Yakugakuzasshi.* 1987;107:350–4.
- Ahmad M, Salama O, Sticher O. Isolation of echinacoside from the white flowers of *Syringa vulgaris*. *J Chem Soc Pakistan.* 1987;9:451–4.
- Xu QM, Liu YL, Li XR, Feng YL, Yang SL. Two new phenylglycol derivatives isolated from *Syringa reticulata* var. *mandshurica* and their antifungal activities. *Chem Pharm Bull.* 2009;57:863–6.
- Ao WLJ, Wang QH, Dai YNT. Chemical detection on the stems of *Syringa pinnatifolia* var. *alashanensis*. *Chin Hosp Pharm J.* 2013;33:1544–6.
- Kurkin VA. Phenylpropanoids as the biologically active compounds of the medicinal plants and phytopharmaceuticals. *Adv Biol Chem.* 2013;3:26–8.
- Ahmad M, Aftab K. Hypotensive action of syringin from *Syringa vulgaris*. *Phytother Res.* 1995;9:452–4.
- Kikuch IM, Yamauchi Y. Studies on constituents of *Syringa* species I. Isolation and structure of iridoids and secoiridoids from the leaves of *Syringa reticulata* (Blume) Hara. *Yakugakuzasshi.* 1987;107:23–7.
- Kurkin VA, Zapesochynaya GG, Ginenko NA, Zolotarev BM. Phenolic compounds from the bark of *Syringa vulgaris*. *Chem Nat Compd.* 1989;4:581–3.

49. Ahmad M, Salama O. Flavonoids from violet flowers of *Syringa vulgaris*. *Pakistan J Sci Ind Res*. 1987;30:150–1.
50. Ahmad M, Salama O. Isolation of rutin from the fresh leaves of *Syringa vulgaris*. *J Pharm*. 1985;4:9–11.
51. Yin WP, Zhao TZ, Zhang HY. A novel oligosaccharide ester from *Syringa pubescens*. *J Asian Nat Prod Res*. 2008;10:95–100.
52. Helmut R. Jasminidin, ein neues monoterpenalkaloid aus *Syringa vulgaris*. *Phytochemistry*. 1978;17:1069–70.
53. Liu X, Wang JM. Iridoid glycosides fraction of *Folium Syringae* leaves modulate NF- $\kappa$ B signal pathway and intestinal epithelial cells apoptosis in experimental colitis. *Plos One*. 2011;6:e24740.
54. Liu X, Wang JM. Anti-inflammatory effects of iridoid glycosides fraction of *Folium Syringae* leaves on TNBS-induced colitis in rats. *J Ethnopharmacol*. 2011;133:780–7.
55. Gao SQ, Niu JQ, Wang F, Wang F, Liu XD, Jin ZX, et al. Comparative studies on the resistance of hepatitis B by *Syringa* extract, IFN and ganyanling in HepG2.2.15 cells. *Chin J Cell Mol Immunol*. 2003;19:385–6.
56. Yao DL, Han ZG, Zhou W, Ju GM, Li G. Protective effects of the *Syringa reticulata* var. *mandshurica* extract against acute liver injury in mice. *J Med Sci Yan B Univ*. 2009;32:29–31.
57. Li JX. The *Syringa aramaticum* modern pharmacological research. *P J Pract Tradit Chin Med*. 2002;18:54.
58. Yan Y, WLJ O, Zhao XJ, Ye XC, Zhang C, Hao JJ, et al. Effect of essential oil of *Syringa pinnatifolia* var. *alashanensis* on ischemia of myocardium, hypoxia and platelet aggregation. *J Ethnopharmacol*. 2010;131:248–55.
59. Balint G, Eperjessy ET, Thuranszky K. Hypothermic action of the leaf extract of *Syringa vulgaris*. *Acta Physiolo Acad Sci Hung*. 1965;28:399–406.
60. Yang WW, Xing MY. The preparations of *Syringa oblata* leaves change in immunoassay after the treatment on herpes simplex keratitis. *Chin J Pract Ophthalmol*. 1990;8:252–61.
61. Xing MY. The preparations of *Syringa oblata* leaves change in tears' PH after the treatment on herpes simplex keratitis. *Chin J Ophthalmol*. 1992;10:312–21.
62. Xing MY, Li B. The preparations of *Syringa oblata* leaves treat epidemic hemorrhagic conjunctivitis. *Chin J Ophthalmol*. 1996;14:301.
63. Li SJ, Xu LY. The preparations of *Syringa oblata* leaves treat hemorrhoids with 18 cases. *China Acad J*. 2001;1:57.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

*“Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge.”*

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
<http://www.chemistrycentral.com/manuscript/>



**ChemistryCentral**