

RESEARCH ARTICLE

Factors Associated with Low-Level Viraemia and Virological Failure: Results from the Austrian HIV Cohort Study

Gisela Leierer^{1,2*}, Katharina Grabmeier-Pfistershammer³, Andrea Steuer⁴, Maria Geit⁵, Mario Sarcletti¹, Bernhard Haas⁶, Manfred Kanatschnig⁷, Michaela Rappold^{1,2}, Robert Zangerle¹, Bruno Ledergerber⁸, Ninon Taylor^{9*}, Austrian HIV Cohort Study Group[¶]

1 Department of Dermatology and Venereology, Medical University of Innsbruck, Innsbruck, Austria, **2** Austrian HIV Cohort Study, Innsbruck, Austria, **3** Department of Dermatology, Medical University of Vienna, Vienna, Austria, **4** Otto-Wagner Hospital, Department of Pulmonology, Vienna, Austria, **5** Department of Dermatology, General Hospital Linz, Linz, Austria, **6** Department of Internal Medicine, General Hospital Graz-West, Graz, Austria, **7** 1st Medical Department, General Hospital Klagenfurt, Klagenfurt, Austria, **8** Division of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Zurich, Switzerland, **9** Department of Internal Medicine III with Hematology, Medical Oncology, Hemostaseology, Infectious Diseases, Rheumatology, Oncologic Center, Laboratory of Immunological and Molecular Cancer Research, Paracelsus Medical University, Salzburg, Austria

¶ Membership of the Austrian HIV Cohort Study Group is provided in the Acknowledgments.

* n.taylor@salk.at (NT); gisela.leierer@tirol-kliniken.at (GL)



OPEN ACCESS

Citation: Leierer G, Grabmeier-Pfistershammer K, Steuer A, Geit M, Sarcletti M, Haas B, et al. (2015) Factors Associated with Low-Level Viraemia and Virological Failure: Results from the Austrian HIV Cohort Study. PLoS ONE 10(11): e0142923. doi:10.1371/journal.pone.0142923

Editor: Paul Richard Harrigan, University of British Columbia, CANADA

Received: April 16, 2015

Accepted: October 28, 2015

Published: November 13, 2015

Copyright: © 2015 Leierer et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the Austrian Agency for Health and Food Safety (AGES). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Background

In human immunodeficiency virus treatment adequate virological suppression is warranted, nevertheless for some patients it remains a challenge. We investigated factors associated with low-level viraemia (LLV) and virological failure (VF) under combined antiretroviral therapy (cART).

Materials and Methods

We analysed patients receiving standard regimens between 1st July 2012 and 1st July 2013 with at least one viral load (VL) measurement below the quantification limit (BLQ) in their treatment history. After a minimum of 6 months of unmodified cART, the next single VL measurement within 6 months was analysed. VF was defined as HIV RNA levels ≥ 200 copies/mL and all other quantifiable measurements were classified as LLV. Factors associated with LLV and VF compared to BLQ were identified by logistic regression models.

Results

Of 2276 participants, 1972 (86.6%) were BLQ, 222 (9.8%) showed LLV and 82 (3.6%) had VF. A higher risk for LLV and VF was shown in patients with cART interruptions and in patients with boosted PI therapy. The risk for LLV and VF was lower in patients from centres using the Abbott compared to the Roche assay to measure VL. A higher risk for LLV but not for VF was found in patients with a higher VL before cART [for >99.999 copies/mL: aOR

(95% CI): 4.19 (2.07–8.49); for 10.000–99.999 copies/mL: aOR (95% CI): 2.52 (1.23–5.19)] and shorter cART duration [for <9 months: aOR (95% CI): 2.59 (1.38–4.86)]. A higher risk for VF but not for LLV was found in younger patients [for <30 years: aOR (95% CI): 2.76 (1.03–7.35); for 30–50 years: aOR (95% CI): 2.70 (1.26–5.79)], people originating from high prevalence countries [aOR (95% CI): 2.20 (1.09–4.42)] and in male injecting drug users [aOR (95% CI): 2.72 (1.38–5.34)].

Conclusions

For both VF and LLV, factors associated with adherence play a prominent role. Furthermore, performance characteristics of the diagnostic assay used for VL quantification should also be taken into consideration.

Introduction

The advent of combined antiretroviral therapy (cART) resulted in a significant reduction in morbidity and mortality for persons infected with the human immunodeficiency virus type 1 (HIV-1) [1,2]. Suppression of plasma HIV-1 RNA through antiretroviral therapy is the goal of HIV treatment. Adequate virological suppression also hinders the development of resistance mutations. Current guidelines define virological suppression below the limit of quantification of commercial assays which is usually between <20 to 75 copies/mL [3]. On the other hand, the threshold for virological failure (VF) remains controversial ranging from 50 to 200 copies/mL [3,4]. These discrepancies reflect the fact that optimal virological suppression remains often an ideal goal but some patients may show persistent low-level viraemia (LLV) and the best clinical practice in such cases, especially concerning LLV between 20–200 copies/mL, is not resolved. Further, the threshold where detectable viral load (VL) in plasma comes from ongoing replication is not known and remains indistinguishable from circulating virus due to the release of HIV from latently infected cells.

The objective of this study was to investigate factors associated with LLV defined with a cut-off <200 copies/mL and VF in a cohort of HIV-1 positive patients.

Methods

Study Design and Data Collection

The Austrian HIV Cohort Study (AHIVCOS) is an open, multicentre, prospective, observational cohort study of HIV-infected individuals followed at seven HIV treatment centres in Austria. The study was initiated in 2001 and patients are enrolled actively and prospectively covering approximately 80% of all treated HIV-infected patients in Austria. Data collection includes sociodemographic information, complete history of cART as well as laboratory results with VL and CD4 cell count measurements, usually repeated every 3–6 months. The laboratory analyses were performed in each single centre and for HIV-1 VL assessment, five centres used the Roche Cobas AmpliPrep/Cobas TaqMan 2.0 assay (Taqman, Roche Diagnostics, Mannheim, Germany), referred to as the Roche assay in this manuscript, while the remaining two centres used the Abbott RealTime HIV-1 assay (Abbott RT, Abbott Diagnostics, Wiesbaden, Germany), referred to as the Abbott assay, during the period of time analysed.

Approval for this study was obtained from the local ethical committees of all participating centres: ethics committee of the Vienna Medical University (No. 898/ 2010), of the Salzburg

Federal Government (No. 1159/ 2010), of the Graz Medical University (No. 21-431/ 2010), of the Innsbruck Medical University (No 283/4.4/ 2009), of the Upper Austria Federal Government (No C-3-10/ 2010) and of the Carinthian Federal State (No A-13-11/2011). Written informed consent was given by the patients for their information to be stored in the hospital database and used for research.

Statistical Analysis

Patient selection. A retrospective analysis was conducted on individuals receiving unmodified cART for more than 6 months between 1st July 2012 and 1st July 2013 enrolled in AHIVCOS (N = 4289). Patients who were not on stable cART due to either interruptions or switches were excluded (N = 290). During these 6 months, patients had to receive cART with 2 nucleoside reverse-transcriptase inhibitors (NRTIs) and either a non-nucleoside reverse-transcriptase inhibitor (NNRTI) or a boosted protease inhibitor (PI) or an integrase inhibitor (INSTI). Individuals with other cART regimens than those mentioned above, were excluded (N = 489). A further exclusion criterion was a lacking VL below the respective limit of quantification in their treatment history. Thus, another 461 individuals were excluded from analyses (35 tested with the Abbott assay and 426 with the Taqman assay).

In total, 3049 patients fulfilled the inclusion criteria. 773 patients did not have any VL measurement between 6 and 12 months after initiation of the respective cART regimen and were too excluded. 2276 patients remained and their first VL measurement between 6 and 12 months after initiation of the respective cART regimen was evaluated.

Definition of LLV and VF. All quantifiable HIV VL measurements below <200 copies/mL were classified as LLV. For further analyses, LLV was divided into two categories, namely HIV RNA levels of 51–199 copies/mL and HIV RNA levels of ≤ 50 copies/mL. VF was defined as HIV RNA levels ≥ 200 copies/mL.

Statistical methods. Groups were compared using χ^2 test or Fisher's exact test where appropriate. To investigate the association between various demographic and clinical parameters with LLV as well as VF compared to BLQ, two separate univariable and multivariable logistic regression models were performed. As reference category for both analyses, patients who were BLQ at baseline were used (N = 1972). For the first analysis, to find factors associated with LLV, 82 patients with VF were excluded. 222 (10.1%) of the remaining 2194 patients had LLV below 200 copies/mL. For the second analysis, to find factors associated with VF, 222 patients who had LLV below 200 copies/mL were excluded. 82 (4.0%) of the remaining 2054 patients had a VF. Variables considered for univariable analyses were age at VL measurement, HIV transmission category, nationality, CD4 count before cART, ever cART interruptions, type of VL assay used, diabetes, cART regimen, VL before cART and cART duration. Multivariable models were adjusted for demographic variables age at VL measurement and HIV transmission category whether or not significant in univariable analyses. Further adjustment was made for variables with $p < 0.05$ in univariable analyses. Ever cART interruptions were defined as interruptions prior to 6 months of stable cART of the respective cART regimen. A cART discontinuation for at least 8 days after having started cART was regarded as cART interruption.

In addition, different sensitivity analyses were performed for different criteria in patient selection. Another sensitivity analysis was conducted in order to analyse patients' characteristics stratified by cART regimen. Further, we additionally did an analysis for the two outcomes LLV and VF excluding the centres measuring VL with the Abbott assay. P-values of < 0.05 were considered to indicate statistical significance.

All analyses were conducted using Stata software, version 13.1 (StataCorp, College Station, TX, USA).

Results

Patient characteristics stratified by VL

The flowchart of the patient selection is shown in [Fig 1](#) and the characteristics of the 2276 included participants, stratified by HIV RNA levels are listed in [Table 1](#). For all the patients included in the study, the median age was 43.8 years (interquartile range IQR: 36.6–51.6) and the majority were men (72.1%). cART was PI-based in 42.9% (976/2276) of the patients, with darunavir being mostly used. NNRTI- and INSTI-based therapies were used in 57.1% (1300/2276) of the cases. The median time on cART was 68.4 months (IQR: 32.0–129.8). Seventy eight percent of the participants had their HIV RNA measured by the Roche assay and 22% had their HIV RNA measured using the Abbott assay. 773 did not have a VL measurement within 12 months of respective cART initiation and were excluded. 377 (16.6%) patients had no prior cART. 1972 of the 2276 included patients were BLQ (86.4%), while 222 (9.8%) showed a LLV and 82 (3.6%) had VF. Stratification by VL of sociodemographic and laboratory analyses showed a higher frequency of male injecting drug users (12.6 and 22.0 versus 8.8%) and of patients with cART interruptions (32.0 and 48.8 versus 24.1%) in the LLV and VF group compared to patients who were BLQ. PI-based regimens were more frequent (65.9 and 51.4 versus 41.0%). Patients whose HIV RNA was measured by the Roche assay rather than by the Abbott assay had more quantifiable VL measurements ≥ 200 copies/mL and below 200 copies/mL compared to individuals who were BLQ (96.3 and 90.5 versus 75.9%). A higher VL before cART ($>99,999$ copies/mL: 50.0% versus 36.9%) was more likely to be found in patients with LLV. Patients with VF were younger (<30 years: 12.2 versus 8.5%; 30–50 years: 78.1 versus 64.8%) and more likely originating from high prevalence countries (18.3 versus 8.8%). Concerning VL cases, the majority (35.4%) of the VF cases were below 1000 copies/mL and only 6.1% reached very high VLs ($>99,000$ copies/mL).

Correlates of LLV

As shown in [Table 2](#), a reduction in risk for LLV was found in female heterosexuals compared to men who have sex with men [adjusted Odds Ratio, aOR (95% CI): 0.62 (0.41–0.95)] as well as in patients from 2 centres which used the Abbott assay as compared to the other centres measuring VL by the Roche assay [aOR (95% CI): 0.33 (0.21–0.51)]. Interrupted cART [aOR (95% CI): 1.69 (1.22–2.34)] versus uninterrupted cART, 2 NRTIs with a boosted PI as cART regimen [aOR (95% CI): 1.54 (1.15–2.06)] versus 2 NRTIs with a NNRTI or INSTI, a higher VL before cART [for $>99,999$ copies/mL: aOR (95% CI): 4.19 (2.07–8.49); for 10,000–99,999 copies/mL: aOR (95% CI): 2.52 (1.23–5.19); compared to $\leq 9,999$ copies/mL] and a shorter cART duration [<9 months: aOR (95% CI): 2.59 (1.38–4.86) compared to >18 months] increased the risk of LLV.

Correlates of VF

A reduction in risk of VF was found in patients from 2 centres using the Abbott assay as compared to the other centres measuring VL by the Roche assay [aOR (95% CI): 0.09 (0.03–0.30)]. Younger age [for <30 years: aOR (95% CI): 2.76 (1.03–7.35); for 30–50 years: aOR (95% CI): 2.70 (1.26–5.79); compared to >50 years], male intravenous drug users [aOR (95% CI): 2.72 (1.38–5.34)] versus men who have sex with men, originating from high prevalence countries [aOR (95% CI): 2.20 (1.09–4.42)], interrupted cART [aOR (95% CI): 2.93 (1.84–4.67)] compared to uninterrupted cART and 2 NRTIs with a boosted PI as cART regimen [aOR (95% CI): 2.36 (1.45–3.83)] versus 2 NRTIs with a NNRTI or INSTI increased the risk of VF ([Table 2](#)).




Fig 1. Flowchart of patient selection.

doi:10.1371/journal.pone.0142923.g001

Table 1. Characteristics of patients stratified by HIV RNA levels below the limit of quantification and low-level viraemia defined as quantifiable HIV RNA levels <200 copies/mL and virological failure defined as HIV RNA levels ≥200 copies/mL.

No. of patients	All patients N = 2276 N (%)	BLQ N = 1972 N (%)	LLV <200 N = 222 N (%)	VF ≥200 N = 82 N (%)	P value ¹	P value ²
Age at VL measurement					0.648	0.002
<30 years	201 (8.8)	168 (8.5)	23 (10.4)	10 (12.2)		
30–50 years	1481 (65.1)	1277 (64.8)	140 (63.1)	64 (78.1)		
>50 years	594 (26.1)	527 (26.7)	59 (26.6)	8 (9.8)		
HIV transmission category					0.034	0.001
Male injecting drug user	220 (9.7)	174 (8.8)	28 (12.6)	18 (22.0)		
Female injecting drug user	95 (4.2)	85 (4.3)	5 (2.3)	5 (6.1)		
Male heterosexual	477 (21.0)	421 (21.4)	45 (20.3)	11 (13.4)		
Female heterosexual	507 (22.3)	448 (22.7)	35 (15.8)	24 (29.3)		
Other	98 (4.3)	82 (4.2)	13 (5.9)	3 (3.7)		
Men who have sex with men	879 (38.6)	762 (38.6)	96 (43.2)	21 (25.6)		
Nationality					0.448	0.004
High prevalence country	212 (9.3)	174 (8.8)	23 (10.4)	15 (18.3)		
Low prevalence country	2064 (90.7)	1798 (91.2)	199 (89.6)	67 (81.7)		
CD4 count before cART					0.349	0.770
Missing	346 (15.2)	295 (15.0)	39 (17.6)	12 (14.6)		
<50 cells/μL	233 (10.2)	207 (10.5)	21 (9.5)	5 (6.1)		
50–199 cells/μL	465 (20.4)	399 (20.2)	48 (21.6)	18 (22.0)		
200–349 cells/μL	716 (31.5)	615 (31.2)	75 (33.8)	26 (31.7)		
≥350 cells/μL	516 (22.7)	456 (23.1)	39 (17.6)	21 (25.6)		
Ever cART interruptions³					0.010	<0.001
≥1	587 (25.8)	476 (24.1)	71 (32.0)	40 (48.8)		
None	1689 (74.2)	1496 (75.9)	151 (68.0)	42 (51.2)		
Assay used					<0.001	<0.001
Abbott RealTime (2 centres)	501 (22.0)	475 (24.1)	23 (10.4)	3 (3.7)		
Roche TaqMan 2.0 (5 centres)	1775 (78.0)	1497 (75.9)	199 (89.5)	79 (96.3)		
Ever diabetes⁴					0.246	0.767
Yes	89 (3.9)	75 (3.8)	12 (5.4)	2 (2.4)		
No	2187 (96.1)	1897 (96.2)	210 (94.6)	80 (97.6)		
cART regimen					0.003	<0.001
2 NRTIs + PI/r	976 (42.9)	808 (41.0)	114 (51.4)	54 (65.9)		
2 NRTIs + NNRTI/INSTI	1300 (57.1)	1164 (59.0)	108 (48.7)	28 (34.2)		
VL before cART					<0.001	0.746
Missing	406 (17.8)	353 (17.9)	39 (17.6)	14 (17.1)		
>99.999 copies/mL	867 (38.1)	728 (36.9)	111 (50.0)	28 (34.2)		
10.000–99.999 copies/mL	753 (33.1)	658 (33.4)	63 (28.4)	32 (39.0)		
≤9.999 copies/mL	250 (11.0)	233 (11.8)	9 (4.1)	8 (9.8)		
cART duration⁵					0.056	0.781
<9 months	86 (3.8)	69 (3.5)	15 (6.8)	2 (2.4)		
9–18 months	183 (8.0)	158 (8.0)	17 (7.7)	8 (9.8)		
>18 months	2007 (88.2)	1745 (88.5)	190 (85.6)	72 (87.8)		
First-line cART⁶					0.949	0.091
Yes	377 (16.6)	332 (16.8)	37 (16.7)	8 (9.8)		

(Continued)

Table 1. (Continued)

No. of patients	All patients N = 2276 N (%)	BLQ N = 1972 N (%)	LLV <200 N = 222 N (%)	VF ≥200 N = 82 N (%)	P value ¹	P value ²
No	1899 (83.4)	1640 (83.2)	185 (83.3)	74 (90.2)		

Abbreviations: LLV, low-level viraemia; VF, virological failure; VL, viral load; NRTIs, nucleoside reverse transcriptase inhibitors; NNRTI, non-nucleoside reverse transcriptase inhibitor; PI/r, boosted protease inhibitor; INSTI, integrase inhibitor; BLQ, below the limit of quantification; cART, combination antiretroviral therapy;

¹ Comparison between quantifiable HIV RNA levels <200 copies/mL (LLV) and HIV RNA levels below the limit of quantification (BLQ).

² Comparison between HIV RNA levels ≥200 copies/mL (VF) and HIV RNA levels below the limit of quantification (BLQ).

³ Interruptions prior to 6 months stable cART of the respective cART regimen.

⁴ Diabetes mellitus prior to 6 months stable cART of the respective cART regimen.

⁵ cART duration until 6 months stable cART of the respective cART regimen.

⁶ Whether the respective cART regimen is a first-line cART or not.

doi:10.1371/journal.pone.0142923.t001

Sensitivity analyses

We performed several sensitivity analyses with different criteria concerning patient selection. One analysis was performed without considering a VL measurement BLQ in treatment history in order to exclude patients with very short cART durations. Another analysis was conducted regarding a shorter recruitment period of 6 months instead of 12 months and one analysis including patients receiving unmodified cART for >9 months instead of >6 months. All additional analyses did, however, not reveal any substantial differences in ORs compared to our primary analysis (data not shown). Stratification of the patients by regimen showed that VL and age at baseline, HIV transmission category, nationality, CD4 cell count before cART, prior cART interruptions, assay used and VL before cART differ significantly (see [S1 Table](#)). In addition, sensitivity analyses were conducted for the two outcomes (LLV <200 copies/mL and VF ≥200 copies/mL) excluding the samples analysed with the Abbott assay. The exclusion of patients whose VL had been measured by the Abbott assay did not reveal any major changes in HRs compared to the original models (see [S2 Table](#)).

Discussion

This study of well-defined patients on stable cART over a period of more than 6 months gives insights into the different factors associated with LLV and VF. As expected, adherence-associated factors play a predominant role. Accordingly, cART interruptions constitute an independent risk factor for both LLV and VF, while younger age, the category of male injecting drug users and originating from high prevalence regions are associated with VF alone. Additionally, PI-based regimens appear as independent markers for both groups. Several studies have already shown that NNRTI-based combinations may result in a better suppression of plasma viral load than PI-based cART [5–7]. On the other hand, we have to acknowledge a potential clinician induced bias for favorably prescribing PIs in case of suspected adherence issues. We performed a supplementary stratification of our study patients by regimen and we observed that patients receiving PIs were found significantly more frequently in the group of intravenous drug users and in the group originating from high prevalence regions (see [S1 Table](#)). In our study, for first-line therapy, NNRTI-based regimens were the most commonly used (N = 226), followed by PIs (N = 116). INSTIs were less represented due to recent release during the observation period (N = 35). However, the majority of our patients were on 2nd or more lines of

Table 2. Univariable and multivariable logistic regression results: Association between different factors and low-level viraemia as well as virological failure compared to HIV RNA levels below the limit of quantification.

Outcome No. of patients included No. of outcomes	LLV <200 N = 2194 N = 222				VF ≥200 N = 2054 N = 82			
	Univariable		Multivariable		Univariable		Multivariable	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Age at VL measurement								
<30 years	1.22	(0.73–2.04)	1.03	(0.60–1.79)	3.92	(1.52–10.10)	2.76	(1.03–7.35)
30–50 years	0.98	(0.71–1.35)	0.97	(0.69–1.35)	3.30	(1.57–6.93)	2.70	(1.26–5.79)
>50 years	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)
HIV transmission category								
Male injecting drug user	1.28	(0.81–2.01)	1.11	(0.69–1.79)	3.75	(1.96–7.20)	2.72	(1.38–5.34)
Female injecting drug user	0.47	(0.18–1.18)	0.43	(0.17–1.12)	2.13	(0.78–5.81)	1.98	(0.70–5.60)
Male heterosexual	0.85	(0.58–1.23)	0.85	(0.58–1.25)	0.95	(0.45–1.99)	0.79	(0.36–1.72)
Female heterosexual	0.62	(0.41–0.93)	0.62	(0.41–0.95)	1.94	(1.07–3.53)	1.09	(0.55–2.16)
Other	1.26	(0.68–2.34)	1.31	(0.69–2.48)	1.33	(0.39–4.55)	1.30	(0.37–4.56)
Men who have sex with men	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)
Nationality								
High prevalence country	1.19	(0.75–1.89)			2.31	(1.29–4.14)	2.20	(1.09–4.42)
Low prevalence country	1.00	(Reference)			1.00	(Reference)	1.00	(Reference)
CD4 count before cART								
Missing	1.55	(0.97–2.47)			0.88	(0.43–1.82)		
<50 cells/μL	1.19	(0.68–2.07)			0.52	(0.20–1.41)		
50–199 cells/μL	1.41	(0.90–2.19)			0.80	(0.51–1.86)		
200–349 cells/μL	1.43	(0.95–2.14)			0.92	(0.51–1.65)		
≥350 cells/μL	1.00	(Reference)			1.00	(Reference)		
Ever cART interruptions¹								
≥1	1.48	(1.09–2.00)	1.69	(1.22–2.34)	2.99	(1.92–4.67)	2.93	(1.84–4.67)
None	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)
Assay used								
Abbott RealTime (2 centres)	0.36	(0.23–0.57)	0.33	(0.21–0.51)	0.12	(0.04–0.38)	0.09	(0.03–0.30)
Roche TaqMan 2.0 (5 centres)	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)
Ever diabetes²								
Yes	1.45	(0.77–2.70)			0.63	(0.15–2.62)		
No	1.00	(Reference)			1.00	(Reference)		
cART regimen								
2 NRTIs + PI/r	1.52	(1.15–2.01)	1.54	(1.15–2.06)	2.78	(1.74–4.42)	2.36	(1.45–3.83)
2 NRTIs + NNRTI/INSTI	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)
VL before cART								
Missing	2.86	(1.36–6.02)	2.70	(1.27–5.76)	1.16	(0.48–2.80)		
>99,999 copies/mL	3.95	(1.97–7.91)	4.19	(2.07–8.49)	1.12	(0.50–2.49)		
10,000–99,999 copies/mL	2.48	(1.21–5.06)	2.52	(1.23–5.19)	1.42	(0.64–3.12)		
≤9,999 copies/mL	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)		
cART duration³								
<9 months	2.00	(1.12–3.56)	2.59	(1.38–4.86)	0.70	(0.17–2.92)		
9–18 months	0.99	(0.59–1.67)	1.00	(0.58–1.73)	1.23	(0.58–2.59)		
>18 months	1.00	(Reference)	1.00	(Reference)	1.00	(Reference)		

(Continued)

Table 2. (Continued)

Outcome	LLV <200				VF ≥200			
	N = 2194 N = 222				N = 2054 N = 82			
	Univariable		Multivariable		Univariable		Multivariable	
No. of patients included No. of outcomes	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
First-line cART⁴								
Yes	0.99	(0.68–1.43)			0.53	(0.26–1.12)		
No	1.00	(Reference)			1.00	(Reference)		

Abbreviations: LLV, low-level viraemia; VF, virological failure; VL, viral load; CI, confidence interval; OR, odds ratio; NRTIs, nucleoside reverse transcriptase inhibitors; NNRTI, non-nucleoside reverse transcriptase inhibitor; PI/r, boosted protease inhibitor; INSTI, integrase inhibitor; cART, combination antiretroviral therapy;

¹ Interruptions prior to 6 months stable cART of the respective cART regimen.

² Diabetes mellitus prior to 6 months stable cART of the respective cART regimen.

³ cART duration until 6 months stable cART of the respective cART regimen.

⁴ Whether the respective cART regimen is a first-line cART or not.

doi:10.1371/journal.pone.0142923.t002

therapy and PIs such as NNRTIs were both equally represented (N = 855 and N = 860, respectively), followed again by INSTIs (N = 184).

Further, the association found between high baseline VL and the risk of developing LLV is in line with the hypothesis that the size of the cellular compartment infected before cART initiation determines the level of residual viraemia [6] and is in accordance with other studies [5,8]. Interestingly, the CD4 count before cART start, especially in the very low range of below 50 cells/μl was comparable in all our 3 groups, BLQ, LLV and VF. This observation argues against the concept that LLV may be more frequent in patients presenting at a late stage of infection.

For our group of patients on stable suppressive therapy, we found a significant risk reduction for LLV and VF according to the assay used for VL quantification. This finding is interesting since both the Abbott as well as the Roche assays are widely used in clinical settings to measure HIV RNA and differ significantly regarding their extraction system, primers and probe design. Although the overall correlation between the Roche and Abbott assay in the higher range of quantification is good [9,10], significant discrepancies between these assays have been described at LLV [11–13]. In an international comparative analysis by Swenson et al., the concordance between four commercial assays including the Roche and Abbott assay was determined and it was shown that the inter-assay concordance at a threshold of 200 copies/mL was much higher than at the clinically relevant threshold of 50 copies/mL [14]. The published literature is controversial concerning the impact of LLV as a predictor of progressive viral rebound [15–19]. Findings by Ribaud et al. from a large retrospective analysis found the same predictive value for virological rebound at an HIV RNA of <200 copies/mL as compared to a threshold of <50 copies/mL [18]. These discrepancies may be partly explained due to different definitions of LLV and VF but also due to different quantification assays used. The latter is most problematic since treatment decisions are based on the definitions of VF and virological suppression, all within the lower range of viraemia. Consecutively, if quantification assays are not concordant in such settings, a univocal threshold of VF becomes impossible to establish. Many factors may contribute to such assay discordance, ranging from preanalytic issues, such as handling procedures, to viral blips or different primer target regions [13,20–25]. In our analysis, the Roche assay was predominantly used (78% versus 22% Abbott assay) limiting a

comparative analysis. Additionally, due to the multicentre study design, we cannot rule out different local practices as potential further bias.

Finally, concerned by the high number of patients excluded from our study due to missing VL assessment within 6 to 12 months ($N = 773$), we compared patients' characteristics of the two groups (data not shown). Individuals excluded differed significantly from included patients in some factors associated with LLV and/or VF. Excluded patients were significantly older, males were more frequently infected through injecting drug use and females through heterosexual contacts. HIV RNA of excluded individuals was more frequently measured by the Roche assay. On the other hand, VL and CD4 count before cART initiation were significantly lower and excluded patients had a shorter overall therapy duration, more cART interruptions and were more frequently on first-line therapy.

In conclusion, the strengths of our study include its large sample size with similar national standards of care and our results support that adherence associated issues play a central part in LLV and VF development. The association between PI-based regimens and risk of LLV and VF remains biased due to confounding by indication. Our data seems to support the concept that performance characteristics of the diagnostic assay used has to be taken into account for evaluation of LLV and especially in cases where clinical decisions such as therapy switches are pending. The present study does not yet include clinical follow-up and predictivity. Despite its observational design, it provides implications for patient management and forms the basis for future outcome studies. We suggest that similar studies with clinical follow-up including different test systems may be able to evaluate the need for guidelines taking into account different performance characteristics of each assay to establish adequate HIV RNA cutoffs.

Supporting Information

S1 Table. Characteristics of patients stratified by regimen.

(DOCX)

S2 Table. Univariable and multivariable logistic regression results: Association between different factors and low-level viraemia as well as virological failure compared to HIV RNA levels below the limit of quantification. Exclusion of the samples in which the Abbott assay was used.

(DOCX)

Acknowledgments

We are grateful to Hannes Oberkofler for fruitful discussion and we thank Peter Taylor for revising the manuscript. The authors are grateful to all clinicians, data managers and research nurses in participating HIV treatment centres listed below. Furthermore, a special thank you goes to the company DI Heinz Appoyer (now called network vita) who was entrusted with the development of the HIV Patient Management System (HIP). All authors are members of the Austrian HIV Cohort Study Group.

The members of the Austrian HIV Cohort Study Group are: HIV treatment centres: LKH Innsbruck (coordinating centre): Martin Gisinger, Maria Kitchen, Elisabeth Rieser, Brigitte Rühr, Mario Sarcletti, Robert Zangerle. LKH Salzburg: Alexander Egle, Richard Greil, Michaela Schachner, Ninon Taylor. AKH Linz: Jörg Berg, Maria Geit, Angela Öllinger. AKH Vienna: Regina Aichwalder, Katharina Grabmeier-Pfistershammer, Armin Rieger, Veronique Touzeau. Otto-Wagner Hospital Vienna: Piotr Cichon, Manfred Gartner, Brigitte Schmied, Andrea Steuer. LKH Graz-West: Bernhard Haas, Andreas Kapper, Elmar Wallner. LKH Klagenfurt: Silvana Achatz, Manfred Kanatschnig, Georg Schober.

Data management group: Heinz Appoyer (IT-related), Gisela Leierer (AHIVCOS), Michaela Rappold (AHIVCOS), Stefanie Strickner (AHIVCOS).

Author Contributions

Conceived and designed the experiments: GL NT MR BL. Performed the experiments: GL NT. Analyzed the data: GL NT RZ BL. Contributed reagents/materials/analysis tools: GL KG AS MG MS BH MK MR RZ NT. Wrote the paper: GL NT BL MR KG.

References

1. Life expectancy of individuals on combination antiretroviral therapy in high-income countries: a collaborative analysis of 14 cohort studies. (2008) *Lancet* 372: 293–299. doi: [10.1016/S0140-6736\(08\)61113-7](https://doi.org/10.1016/S0140-6736(08)61113-7) PMID: [18657708](https://pubmed.ncbi.nlm.nih.gov/18657708/)
2. Palella FJ Jr., Delaney KM, Moorman AC, Loveless MO, Fuhrer J, Satten GA, et al. (1998) Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV Outpatient Study Investigators. *N Engl J Med* 338: 853–860. PMID: [9516219](https://pubmed.ncbi.nlm.nih.gov/9516219/)
3. Panel on Antiretroviral Guidelines for adults and adolescents for the use of ART in HIV -1 infected Adults and Adolescents. Department of Health and Human Services. Available at <http://aidsinfo.nih.gov/guidelines>. Accessed 26th August 2015.
4. European AIDS Clinical Society version 7.1, November 2014. EACS, Brussels, Belgium. available at <http://www.eacsociety.org/guidelines/eacs-guidelines/eacs-guidelines.html>. Accessed 26th August 2015.
5. Vancoillie L, Demecheleer E, Callens S, Vogelaers D, Vandekerckhove L, Mortier V, et al. (2014) Markers associated with persisting low-level viraemia under antiretroviral therapy in HIV-1 infection. *J Antimicrob Chemother* 69: 1098–1103. doi: [10.1093/jac/dkt484](https://doi.org/10.1093/jac/dkt484) PMID: [24335486](https://pubmed.ncbi.nlm.nih.gov/24335486/)
6. Maldarelli F, Palmer S, King MS, Wiegand A, Polis MA, Mican J, et al. (2007) ART suppresses plasma HIV-1 RNA to a stable set point predicted by pretherapy viremia. *PLoS Pathog* 3: e46. PMID: [17411338](https://pubmed.ncbi.nlm.nih.gov/17411338/)
7. Palmisano L, Giuliano M, Nicastrì E, Pirillo MF, Andreotti M, Galluzzo CM, et al. (2005) Residual viraemia in subjects with chronic HIV infection and viral load < 50 copies/ml: the impact of highly active antiretroviral therapy. *Aids* 19: 1843–1847. PMID: [16227792](https://pubmed.ncbi.nlm.nih.gov/16227792/)
8. Martin-Blondel G, Saune K, Vu Hai V, Marchou B, Delobel P, Izopet J, et al. (2012) Factors associated with a strictly undetectable viral load in HIV-1-infected patients. *HIV Med* 13: 568–573. doi: [10.1111/j.1468-1293.2012.01012.x](https://doi.org/10.1111/j.1468-1293.2012.01012.x) PMID: [22435457](https://pubmed.ncbi.nlm.nih.gov/22435457/)
9. Sire JM, Vray M, Merzouk M, Plantier JC, Pavie J, Maylin S, et al. (2011) Comparative RNA quantification of HIV-1 group M and non-M with the Roche Cobas AmpliPrep/Cobas TaqMan HIV-1 v2.0 and Abbott Real-Time HIV-1 PCR assays. *J Acquir Immune Defic Syndr* 56: 239–243. doi: [10.1097/QAI.0b013e3182099891](https://doi.org/10.1097/QAI.0b013e3182099891) PMID: [21164353](https://pubmed.ncbi.nlm.nih.gov/21164353/)
10. Taylor N, Schmid I, Egle A, Greil R, Patsch W, Oberkofler H (2009) Initial evaluation of the Roche COBAS TaqMan HIV-1 v2.0 assay for determining viral load in HIV-infected individuals. *Antivir Ther* 14: 1189–1193. doi: [10.3851/IMP1427](https://doi.org/10.3851/IMP1427) PMID: [20032549](https://pubmed.ncbi.nlm.nih.gov/20032549/)
11. Amendola A, Marsella P, Bloisi M, Forbici F, Angeletti C, Capobianchi MR (2014) Ability of two commercially available assays (Abbott RealTime HIV-1 and Roche Cobas AmpliPrep/Cobas TaqMan HIV-1 Version 2.0) to quantify low HIV-1 RNA Levels (<1,000 copies/milliliter): comparison with clinical samples and NIBSC working reagent for nucleic acid testing assays. *J Clin Microbiol* 52: 2019–2026. doi: [10.1128/JCM.00288-14](https://doi.org/10.1128/JCM.00288-14) PMID: [24671791](https://pubmed.ncbi.nlm.nih.gov/24671791/)
12. Naeth G, Ehret R, Wiesmann F, Braun P, Knechten H, Berger A (2013) Comparison of HIV-1 viral load assay performance in immunological stable patients with low or undetectable viremia. *Med Microbiol Immunol* 202: 67–75. doi: [10.1007/s00430-012-0249-y](https://doi.org/10.1007/s00430-012-0249-y) PMID: [22699843](https://pubmed.ncbi.nlm.nih.gov/22699843/)
13. Taylor N, Grabmeier-Pfistershammer K, Egle A, Greil R, Rieger A, Ledergerber B, et al. (2013) Cobas ampliPrep/cobas TaqMan HIV-1 v2.0 assay: consequences at the cohort level. *PLoS One* 8: e74024. doi: [10.1371/journal.pone.0074024](https://doi.org/10.1371/journal.pone.0074024) PMID: [24023696](https://pubmed.ncbi.nlm.nih.gov/24023696/)
14. Swenson LC, Cobb B, Geretti AM, Harrigan PR, Poljak M, Seguin-Devaux C, et al. (2014) Comparative performances of HIV-1 RNA load assays at low viral load levels: results of an international collaboration. *J Clin Microbiol* 52: 517–523. doi: [10.1128/JCM.02461-13](https://doi.org/10.1128/JCM.02461-13) PMID: [24478482](https://pubmed.ncbi.nlm.nih.gov/24478482/)
15. Doyle T, Smith C, Vitiello P, Cambiano V, Johnson M, Owen A, et al. (2012) Plasma HIV-1 RNA detection below 50 copies/ml and risk of virologic rebound in patients receiving highly active antiretroviral therapy. *Clin Infect Dis* 54: 724–732. doi: [10.1093/cid/cir936](https://doi.org/10.1093/cid/cir936) PMID: [22238167](https://pubmed.ncbi.nlm.nih.gov/22238167/)

16. Eron JJ, Cooper DA, Steigbigel RT, Clotet B, Gatell JM, Kumar PN, et al. (2013) Efficacy and safety of raltegravir for treatment of HIV for 5 years in the BENCHMRK studies: final results of two randomised, placebo-controlled trials. *Lancet Infect Dis* 13: 587–596. doi: [10.1016/S1473-3099\(13\)70093-8](https://doi.org/10.1016/S1473-3099(13)70093-8) PMID: [23664333](https://pubmed.ncbi.nlm.nih.gov/23664333/)
17. Laprise C, de Pokomandy A, Baril JG, Dufresne S, Trottier H (2013) Virologic failure following persistent low-level viremia in a cohort of HIV-positive patients: results from 12 years of observation. *Clin Infect Dis* 57: 1489–1496. doi: [10.1093/cid/cit529](https://doi.org/10.1093/cid/cit529) PMID: [23946221](https://pubmed.ncbi.nlm.nih.gov/23946221/)
18. Ribaudo H, Lennox J, Currier J (2009) Virological failure endpoint definition in clinical trials: Is using HIV-1 RNA threshold <200 copies/mL better than 50 copies/mL? An analysis of the ACTG studies. Paper presented at: Conference on Retroviruses and Opportunistic Infections; Montreal, Canada.
19. Taiwo B, Gallien S, Aga E, Ribaudo H, Haubrich R, Kuritzkes DR, et al. (2011) Antiretroviral drug resistance in HIV-1-infected patients experiencing persistent low-level viremia during first-line therapy. *J Infect Dis* 204: 515–520. doi: [10.1093/infdis/jir353](https://doi.org/10.1093/infdis/jir353) PMID: [21791652](https://pubmed.ncbi.nlm.nih.gov/21791652/)
20. Djedaini M, Peraldi P, Drici MD, Darini C, Saint-Marc P, Dani C, et al. (2009) Lopinavir co-induces insulin resistance and ER stress in human adipocytes. *Biochem Biophys Res Commun* 386: 96–100. doi: [10.1016/j.bbrc.2009.05.148](https://doi.org/10.1016/j.bbrc.2009.05.148) PMID: [19501568](https://pubmed.ncbi.nlm.nih.gov/19501568/)
21. Gatanaga H, Tsukada K, Honda H, Tanuma J, Yazaki H, Watanabe T, et al. (2009) Detection of HIV type 1 load by the Roche Cobas TaqMan assay in patients with viral loads previously undetectable by the Roche Cobas Amplicor Monitor. *Clin Infect Dis* 48: 260–262. doi: [10.1086/595707](https://doi.org/10.1086/595707) PMID: [19113986](https://pubmed.ncbi.nlm.nih.gov/19113986/)
22. Griffith BP, Mayo DR (2006) Increased levels of HIV RNA detected in samples with viral loads close to the detection limit collected in Plasma Preparation Tubes (PPT). *J Clin Virol* 35: 197–200. PMID: [16253548](https://pubmed.ncbi.nlm.nih.gov/16253548/)
23. Lima V, Harrigan R, Montaner JS (2009) Increased reporting of detectable plasma HIV-1 RNA levels at the critical threshold of 50 copies per milliliter with the Taqman assay in comparison to the Amplicor assay. *J Acquir Immune Defic Syndr* 51: 3–6. doi: [10.1097/QAI.0b013e31819e721b](https://doi.org/10.1097/QAI.0b013e31819e721b) PMID: [19247185](https://pubmed.ncbi.nlm.nih.gov/19247185/)
24. Portman MD, Lacey CJ (2012) Apparent low-level HIV RNA viraemia related to sample processing time. *HIV Med* 13: 578–579. doi: [10.1111/j.1468-1293.2012.01016.x](https://doi.org/10.1111/j.1468-1293.2012.01016.x) PMID: [22963641](https://pubmed.ncbi.nlm.nih.gov/22963641/)
25. Ruelle J, Debaisieux L, Vancutsem E, De Bel A, Delforge ML, Piérard D, et al. (2012) HIV-1 low-level viraemia assessed with 3 commercial real-time PCR assays show high variability. *BMC Infect Dis* 12: 100. PMID: [22530816](https://pubmed.ncbi.nlm.nih.gov/22530816/)