

ARTICLE

Open Access

GADD45 α sensitizes cervical cancer cells to radiotherapy via increasing cytoplasmic APE1 level

Qing Li¹, Xi Wei², Zhi-Wei Zhou³, Shu-Nan Wang⁴, Hua Jin⁵, Kui-Jun Chen⁶, Jia Luo¹, Kenneth D. Westover³, Jian-Min Wang⁶, Dong Wang¹, Cheng-Xiong Xu¹ and Jin-Lu Shan¹

Abstract

Radioresistance remains a major clinical challenge in cervical cancer therapy. However, the mechanism for the development of radioresistance in cervical cancer is unclear. Herein, we determined that growth arrest and DNA-damage-inducible protein 45 α (GADD45 α) is decreased in radioresistant cervical cancer compared to radiosensitive cancer both in vitro and in vivo. In addition, silencing GADD45 α prevents cervical cancer cells from undergoing radiation-induced DNA damage, cell cycle arrest, and apoptosis. More importantly, our data show that the overexpression of GADD45 α significantly enhances the radiosensitivity of radioresistant cervical cancer cells. These data show that GADD45 α decreases the cytoplasmic distribution of APE1, thereby enhancing the radiosensitivity of cervical cancer cells. Furthermore, we show that GADD45 α inhibits the production of nitric oxide (NO), a nuclear APE1 export stimulator, by suppressing both endothelial NO synthase (eNOS) and inducible NO synthase (iNOS) in cervical cancer cells. In conclusion, our findings suggest that decreased GADD45 α expression significantly contributes to the development of radioresistance and that ectopic expression of GADD45 α sensitizes cervical cancer cells to radiotherapy. GADD45 α inhibits the NO-regulated cytoplasmic localization of APE1 through inhibiting eNOS and iNOS, thereby enhancing the radiosensitivity of cervical cancer cells.

Introduction

Cervical cancer is the fourth most common malignant disease¹ and one of the major causes of cancer-related death among females worldwide². Clinically, radiotherapy is one of the most commonly used treatments for cervical cancer as it significantly reduces the risk of cervical cancer relapse³. Over 60% of patients with cervical cancer undergo radiotherapy⁴; however, some cervical cancers

develop resistance to radiotherapy, which can significantly compromise clinical outcome. Unfortunately, the mechanism for acquiring and developing radioresistance in cervical cancer remains unclear.

Mechanistically, radiotherapy causes cell cycle arrest and tumor cell death by inducing DNA damage⁵. Thus, aberrant DNA repair is one mechanism whereby cancer cells may become radioresistant. Growth arrest and DNA-damage-inducible protein 45 α (GADD45 α) is a radiation-inducible gene⁶ that is involved in DNA repair^{7, 8}. The effects of GADD45 α on cancer cell radiosensitivity have been investigated in several cancer types, but its role in radioresistance remains inconclusive. Lu et al.⁹ and Hur et al.¹⁰ showed that the inactivation of GADD45 α sensitized epithelial cancer cells and hepatoma cells, respectively, to radiation treatment, whereas Zhang et al.¹¹ and

Correspondence: C-X. Xu (xuchengxiong@hanmail.net) or J-L. Shan (lulu7476@sina.com)

¹Cancer Center, Daping Hospital and Research Institute of Surgery, Third Military Medical University, Chongqing 400042, China

²Department of Diagnostic and Therapeutic Ultrasonography, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center of Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin, China

Full list of author information is available at the end of the article.


These authors contributed equally: Qing Li, Xi Wei, Zhi-Wei Zhou.

Edited by A Stephanou

© The Author(s) 2018



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



Asuthkar et al.¹² reported that the overexpression of GADD45 α enhanced the sensitivity of squamous cell carcinoma of the tongue and medulloblastoma cells, respectively, to radiation treatment. Klopp et al.¹³ demonstrated a decrease in GADD45 α expression in recurrent cervical squamous cell carcinoma patients. Notably, our group previously found that GADD45 α expression was decreased in radioresistant cervical cancer cells¹⁴. Taken together, these findings implicate GADD45 α in the development of radioresistance; however, the function and mechanism whereby GADD45 α regulates cervical cancer radiosensitivity remains elusive.

Apurinic/apyrimidinic endonuclease 1 (APE1) is a multifunctional protein involved in DNA repair and gene transcription during the adaptive cellular response to oxidative stress, and APE1 reportedly contributes to the development of therapeutic resistance, tumor aggressiveness, and metastasis¹⁵. The elevated expression or activity of APE1 is associated with increased resistance to radiation in several cancers, including cervical cancer^{16–19}. In addition, inhibition or silencing of APE1 dramatically enhances cancer cell sensitivity to radiotherapy in prostate cancer²⁰, colorectal cancer²¹, non-small-cell lung cancer²², pancreatic cancer²³, and hepatocellular


carcinoma²⁴, suggesting an association between APE1 and radiosensitivity across different cancer types. Recent studies have shown that GADD45 α regulates APE1 activity in cancer cells through direct interaction^{25, 26}.

Given these findings, we propose that GADD45 α regulates APE1 and that reduction of GADD45 α contributes to the development of radioresistance in cervical cancer. In this work, we demonstrate that GADD45 α levels are inversely correlated with radioresistance in cervical cancer patients. Our data indicate that GADD45 α sensitizes tumors to radiotherapy by enhancing radiation-induced cell cycle arrest and apoptosis in cervical cancer cells. In addition, our data illustrate that GADD45 α enhances the radiosensitivity of cervical cancer cells through the suppression of cytoplasmic APE1 levels via the inhibition of nitric oxide (NO) production.

Results

HeLa-XR is a radioresistant cervical cancer cell line

First, we confirmed that the X-ray-resistant HeLa cell line (HeLa-XR) is indeed resistant to radiation treatment. As shown in Fig. 1a, a clonogenic assay revealed that HeLa-XR cells exhibited a higher survival fraction compared to parental HeLa cells when treated with the same



dose of irradiation (IR). Consistent with the clonogenic assay, a comet assay also illustrated that HeLa-XR cells exhibited reduced DNA damage compared to HeLa cells when treated with the same dose of IR (Fig. 1b). Furthermore, we compared IR treatment-induced cell apoptosis and cell cycle arrest between HeLa-XR and HeLa cells by flow cytometry. As shown in Fig. 1c, d, 6 Gy IR treatment-induced apoptosis and G_2/M cell cycle arrest in HeLa cells compared controls (0 Gy IR treatment), but HeLa-XR cells did not exhibit these effects. These findings confirm that HeLa-XR is a radioresistant cervical cancer cell line.


Decreased expression of GADD45 α contributes to radioresistance in cervical cancer

Our previous microarray analysis indicated that levels of the DNA damage repair gene GADD45 α were significantly reduced in X-ray-resistant HeLa cells¹⁴. We confirmed these results using Quantitative Real-time Polymerase Chain Reaction, RT-qPCR and western blot analysis in cultured cells and cervical cancer patient specimens. Our data showed that the expression of GADD45 α was significantly decreased in radioresistant HeLa-XR cervical cancer cells at both the mRNA and protein level compared to parental HeLa cells (Fig. 2a, b). In agreement with the in vitro experimental results, clinical specimens also presented with significantly reduced GADD45 α expression compared to specimens from radiotherapy-sensitive patients (Fig. 2c), suggesting that decreased GADD45 α expression correlates with radioresistance in cervical cancer. Next, we investigated the impact of GADD45 α on the radiosensitivity of cervical cancer cells. As expected, our data showed that silencing

GADD45 α (Fig. 3a) dramatically protected HeLa cells from IR-induced cell death (Fig. 3b), DNA damage (Fig. 3c), apoptosis (Fig. 3d), and G_2/M phase cell cycle arrest (Fig. 3e). Consistent with these results, silencing of GADD45 α in the GADD45 α -high expressing cervical cancer cell lines CasKi and SiHa (Supplementary Fig. 1) also protected these two cell lines from IR-induced cell death, DNA damage, apoptosis, and cell cycle arrest (Supplementary Figs. 1 and 2). Altogether, these data clearly demonstrate that the suppression of GADD45 α significantly contributes to the development of radioresistance in cervical cancer cells.

Ectopic overexpression of GADD45 α overcomes radiosensitivity in radioresistant cervical cancer cells

Our observation that silencing GADD45 α contributes to the development of radioresistance in cervical cancer cells prompted us to investigate whether the overexpression of GADD45 α could enhance the sensitivity of radioresistant cervical cancer cells to radiotherapy. To address this, GADD45 α was overexpressed in HeLa-XR cells (Fig. 4a), and cells were subsequently treated with 6 Gy IR and subjected to clonogenic, cell viability, apoptosis, cell cycle, and comet assays. As expected, clonogenic (Fig. 4b) and cell viability (Fig. 4c) assays demonstrated that the overexpression of GADD45 α significantly enhanced IR-induced cell growth inhibition and cell death compared to vector controls in HeLa-XR cells. Consistent with these results, the comet and flow cytometry analyses showed that the combination of GADD45 α overexpression and IR treatment significantly induced DNA damage (Fig. 4d) and apoptosis (Fig. 4e) in HeLa-XR cells compared to IR treatment alone. In addition, cell cycle




analysis illustrated that the overexpression of GADD45 α with IR exposure considerably increased G₂/M cell cycle arrest in HeLa-XR cells compared to IR treatment alone (Fig. 4f). Furthermore, we confirmed these in vitro results in vivo using xenograft animal models. As shown in Fig. 5, a single treatment of IR or GADD45 α overexpression alone did not significantly inhibit tumor growth, cell proliferation, or induce cell apoptosis, compared to vector controls in the HeLa-XR xenograft model. In contrast, a combination of GADD45 α overexpression (Fig. 5a) and IR treatment dramatically suppressed tumor growth (Fig. 5b), with significant inhibition of cell proliferation (Fig. 5c) and induction of cell apoptosis (Fig. 5d) compared to both controls and a single treatment of IR or GADD45 α overexpression alone in the HeLa-XR xenograft model. These findings suggest that the overexpression of GADD45 α augments radiosensitivity in radioresistant cervical cancer cells via enhancing IR-induced apoptosis and cell cycle inhibition.

GADD45 α reduces cytoplasmic APE1 levels in cervical cancer cells

To investigate the mechanism by which GADD45 α enhances radiosensitivity, we examined the effects of GADD45 α on APE1 expression as APE1 is implicated in

cancer radioresistance²⁷ and is involved in GADD45 α function²⁵. Western blot data showed no difference in total APE1 expression levels between HeLa and HeLa-XR cells (Fig. 6a). However, we observed a significant difference in the subcellular distribution of APE1 between HeLa and HeLa-XR cells. Our data showed that radioresistant HeLa-XR cells exhibited higher levels of cytoplasmic APE1 and lower levels of nuclear APE1 than did HeLa cells (Fig. 6a). Similarly, immunofluorescence (IF) showed high levels of cytosolic APE1 in HeLa-XR cells compared to HeLa cells (Fig. 6b). Next, we tested whether GADD45 α is involved in the regulation of APE1 subcellular distribution. As expected, western blot and IF data show that silencing of GADD45 α did not affect total APE1 expression, but resulted in a significant increase in cytoplasmic APE1 and a significant decrease in nuclear APE1 compared to vector controls in HeLa cells (Fig. 6c, d). In contrast, the overexpression of GADD45 α decreased cytoplasmic APE1 levels and increased nuclear APE1 compared to control HeLa-XR cells but did not affect total APE1 expression levels (Fig. 6e, f). Similar to our in vitro data, the clinical data indicated that GADD45 α expression is inversely correlated with cytoplasmic APE1 levels and resistance to radiotherapy (Fig. 6g). Collectively, these findings suggest that



GADD45 α mediates APE1 subcellular distribution in cervical cancer cells.


GADD45 α reduces cytoplasmic APE1 levels through inhibition of NO production in cervical cancer cells

To investigate the mechanism of GADD45 α -mediated regulation of APE1 subcellular distribution, we examined the effects of GADD45 α on the production of NO and the expression of inducible NO synthase (iNOS) and endothelial NO synthase (eNOS). As shown in Fig. 7a, levels of NO, iNOS, and eNOS were significantly increased in HeLa-XR cells compared to parental HeLa cells. Next, we investigated whether GADD45 α is involved in the regulation of NOS expression and NO production. Our data demonstrate that silencing GADD45 α significantly increases NO production and eNOS and iNOS expression in HeLa cells (Fig. 7b). In addition, silencing GADD45 α increases IR treatment-induced production of NO, iNOS, and eNOS more significantly than in vector controls (Fig. 7b). In contrast, the overexpression of GADD45 α significantly suppressed

NO production and eNOS and iNOS expression in HeLa-XR cells compared to vector controls. In addition, the overexpression of GADD45 α significantly inhibited IR-induced NO production and eNOS and iNOS expression in HeLa-XR cells compared to vector controls (Fig. 7c). On the other hand, inhibiting NO production using a NOS inhibitor dramatically suppressed GADD45 α -induced cytoplasmic APE1 accumulation (Fig. 7d). Taken together, these findings indicate that GADD45 α reduces cytoplasmic APE1 through the inhibition of NO production by reducing both iNOS and eNOS expression in cervical cancer cells.

Discussion

Radioresistance correlates with poor clinical outcome in cervical cancer patients. Unfortunately, there is no available treatment specific to cervical cancer patients with radioresistance. Herein, we identified that the expression of GADD45 α was significantly reduced in radioresistant cervical cancer cells and specimens from radioresistant cervical cancer patients compared to radiosensitive




cervical cancer cells and specimens from radiosensitive cervical cancer patients, respectively. In addition, our in vitro and in vivo experimental data show that silencing GADD45α increases the radiosensitivity of radioresistant cervical cancer cells to IR treatment via the inhibition of IR-induced apoptosis and cell cycle arrest. More importantly, our findings demonstrate that ectopic overexpression of GADD45α significantly enhances radiosensitivity in radioresistant cervical cancer cells. These results suggest that the suppression of GADD45α expression is partially responsible for the development of radioresistance and that the ectopic expression of GADD45α is a potential strategy to overcome radioresistance in cervical cancer.

Next, we have examined the mechanism by which GADD45α regulates cervical cancer cell radiosensitivity. A positive correlation between increased total APE1 expression and radioresistance has been previously identified in numerous cancer studies. Recent studies have shown that not only expression level but also dysregulation of the subcellular localization of APE1 is associated with therapeutic resistance and cancer progression. Clinical studies show that cytoplasmic localization of APE1 correlates with increased aggressiveness and poor prognosis in several

cancer types, including lung cancer²⁸, ovarian cancer²⁹, breast cancer³⁰, and hepatocellular carcinoma³¹. More importantly, Qing et al.³² reported that patients with greater cytoplasmic APE1 expression experienced a shorter survival time compared to those expressing more nuclear APE1 among cervical patients treated with 252Cf radiation. Herein, our data demonstrate that decreased GADD45α results in cervical cancer cell radioresistance, leading to increased cytoplasmic APE1. In contrast, the overexpression of GADD45α enhances radiosensitivity and decreases cytoplasmic APE1 levels in radioresistant cervical cancer cells. Overall, these findings suggest that GADD45α enhances the radiosensitivity of cervical cancer cells by increasing cytoplasmic APE1 localization.

Finally, we identified a novel mechanism by which GADD45α regulates the subcellular localization of APE1 in cervical cancer cells. NO is an important factor in the regulation of APE1 subcellular localization. Qu et al.³³ reported that increased NO stimulates the nuclear export of APE1, resulting in cytoplasmic APE1 accumulation. Here, our data show that the overexpression of GADD45α dramatically suppresses radiation-induced NO by significantly inhibiting both eNOS and iNOS expression in cervical cancer cells. In addition, the



repression of GADD45 α -induced cytoplasmic APE1 accumulation was reduced by treatment with an NOS inhibitor, suggesting that GADD45 α hinders nuclear APE1 export by inhibiting NO production and increasing cytoplasmic APE1 levels in cervical cancer cells. Moreover, via the regulation of NO production, GADD45 α mediates the subcellular localization of APE1 by interacting with APE1 in the nucleus. According to Jung et al.²⁵, APE1 nuclear localization was observed in GADD45 α wild-type cells, whereas APE1's distribution became primarily cytoplasmic in GADD45 α -deficient cells. In addition, their study showed that GADD45 α contributes to the nuclear interaction between APE1 and proliferating cell nuclear antigen (PCNA). Consistent with these results, Kim et al.²⁶ also demonstrated that GADD45 α plays a critical role in modulating the interaction of PCNA and APE1. Collectively, our findings combined with reports from other groups suggest that GADD45 α represses the nuclear export of APE1 to the


cytoplasm via inhibiting NO production and stimulating APE1's interaction with PCNA in cervical cancer cells.

In summary (Fig. 8), this study demonstrates that suppressing GADD45 α contributes to development of radioresistance and that overexpressing GADD45 α can overcome radioresistance in cervical cancer cells. Furthermore, we demonstrated that GADD45 α enhances radiosensitivity by decreasing the cytoplasmic distribution of APE1 in cervical cancer cells. Additionally, we illustrated that GADD45 α inhibits APE1 cytoplasmic distribution by inhibiting NO production via the reduction of eNOS and iNOS expression. Further studies are still needed to elucidate the mechanism whereby GADD45 α mediates NOS regulation.

Materials and Methods

Cell lines and cell culture

The HeLa cell line was obtained from American Type Culture Collection (ATCC, Manassas, VA), and X-ray-



resistant HeLa cells (HeLa-XR) were developed using HeLa cells as previously described¹⁴. HeLa and HeLa-XR cell lines were maintained in Dulbecco’s modified Eagle’s medium with 10% fetal bovine serum (HyClone, Logan, UT) at 37 °C and 5% CO₂.

Plasmids and transfection

The GADD45a plasmid and shRNA constructs were obtained from Western Biomedical Technology (Chongqing, China). Cells were transfected with indicated plasmids using Lipofectamine 2000 according to manufacturer’s instructions (Invitrogen, Carlsbad, CA).


Cell viability and clonogenic assay

To determine cell viability, cells were plated in 96-well plates at a density of 5000 cells per well. Twelve hours after seeding, cells were exposed to indicated

doses of IR. After 24 h of IR exposure, cell viability was determined using a CCK-8 kit (Dojindo Laboratories, Kumamoto, Japan) according to the manufacturer’s instructions. For the clonogenic assays, cells were seeded in 60 mm² dishes (1000 cells per plate). Twelve hours after seeding, cells were exposed to indicated doses of IR and were allowed to grow until visible colonies appeared. Colonies were subsequently stained with 0.01% crystal violet (Sigma, St. Louis, MO) and counted under a microscope.

Apoptosis and cell cycle analysis

Cells were exposed to the indicated doses of IR. After 24 h of IR treatment, cells were subjected to cell cycle and apoptosis analysis. Apoptotic cells were identified by flow cytometry using the Annexin V-FITC kit (Calbiochem, Shanghai, China) according to the manufacturer’s



instructions. Briefly, cells were washed twice with cold PBS and resuspended in binding buffer. Next, Annexin V and fluorescein isothiocyanate, FITC were added. Cells were gently mixed followed by incubation for 15 min at room temperature in the dark and then subjected to flow cytometry analysis. For cell cycle analysis, cells were washed twice with cold PBS and fixed in 70% ethanol overnight at -20°C . Next, the cells were treated with a DNA staining solution, and cell cycle analysis was performed with a Fluorescence activated Cell Sorting, FACS flow cytometer. Apoptotic cells in the tissue were determined using the In Situ Cell Death Detection Kit (Roche, Mannheim, Germany) according to the manufacturer's instructions.

Comet analysis

Cells were exposed to the indicated doses of IR. After 4 h of IR, cells were collected and diluted to 4×10^4 cells per ml. Cells were then mixed with 0.5% low melting point agarose and placed into a horizontal electrophoresis tank, and a fresh preparation of cold electrophoresis buffer was added. Next, cells were incubated for 30 min to allow for the formation of DNA chains and then electrophoresed for 25 min (20 V, 300 mA). The glass was washed with a neutral buffer three times (5 min each time) after completion of electrophoresis. The dry glass was stained with 10–15 μl of ethidium bromide (20 $\mu\text{g}/\text{ml}$) and observed by fluorescence microscopy. Comet was analyzed using Comet 5.5 software.

Real-time quantitative reverse transcription PCR analysis

Total RNA was isolated from cells and fresh tissues with TRIzol reagent (Invitrogen) according to the manufacturer's protocol. Reverse transcript and real-time PCR assays were performed using a High-Capacity cDNA Reverse Transcription Kit and the QuantiTect SYBR Green PCR kit (Qiagen, Germantown, MD), respectively. The primer sequences for PCR assays were as follows: GADD45 α forward, 5'-AGUCGCUACAUGGAUCAAUT T-3'; reverse, 5'-AUUGAUCCAUGUAGCGACUTT-3'; GAPDH forward, 5'-GCAGGGGGGAGCCAAAAGGGT-3'; and reverse, 5'-TGGGTGGCAGTGATGGCATGG-3'.

Immunohistochemistry, immunofluorescence, and western blot

Immunohistochemistry (IHC), IF, and western blot assays were performed as previously described^{34, 35}. Antibodies against APE1, actin, Ki-67, and GADD45 α , and secondary antibodies conjugated to horseradish peroxidase were purchased from Abcam (Cambridge, MA).

Detection of nitric oxide, inducible nitric oxide synthase and endothelial nitric oxide synthase

NO, iNOS, and eNOS were measured in cell lysates using an ELISA Kit according to the manufacturer's instructions. The Human iNOS ELISA Kit and Human eNOS ELISA Kit were purchased from Wuhan Huamei Biotech Co. (Wuhan, Hubei, China). The NO detection ELISA Kit was obtained from Beyotime Biotech (Shanghai, China).

Human specimens and animal experiments

Human samples were collected by biopsy from patients before and after radiotherapy. Initial samples were prepared to measure gene expression, and follow-up samples (after radiotherapy) underwent an evaluation to investigate the effectiveness of the radiotherapy. The effectiveness of radiotherapy was evaluated according to methods previously described by Kitahara et al.³⁶, and samples were divided into radiosensitive and radioresistant groups based on the results of the evaluation. This study was approved by the Ethics Committee of the Third Military Medical University, and clinical samples were obtained from the Daping Hospital and Research Institute of Surgery at the Third Military Medical University.

Animal experiments were performed using 5-week-old female BALB/c nude mice (Shanghai Laboratory Animal Center, Shanghai, China). Twenty mice were divided into two groups ($n = 10$ per group), and HeLa-XR cells stably overexpressing GADD45 α or the empty vector were subcutaneously injected into the left anterior axillae of nude mice (1.5×10^7 cells per mouse). One week following inoculation with the tumor cells, each group was further divided into two groups ($n = 5$ per group), with one group

irradiated with 8 Gy every 5 days, three times. Tumor volume was measured using MRI (7.0-T MRI, Bruker Biospec 70/20USR, Germany) every week. On day 35 after inoculation, all mice were killed, and tumors were excised. All animal experiments were approved by the Animal Care Committee of the Third Military Medical University.

Statistical analyses

All *p* values <0.05 were considered significant. All assays were independently performed at least three times. All values are presented as the mean \pm SD. Differences between groups were determined using Student's *t* test or one-way analysis of variance using the SAS statistical software package version 6.12 (SAS Institute, Cary, NC).

Acknowledgements

This study was supported by the National Natural Science Foundation of China (81402155).

Author details

¹Cancer Center, Daping Hospital and Research Institute of Surgery, Third Military Medical University, Chongqing 400042, China. ²Department of Diagnostic and Therapeutic Ultrasonography, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center of Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin, China. ³Department of Radiation Oncology and Biochemistry, University of Texas Southwestern Medical Center, Dallas, TX, USA. ⁴Department of Radiology, Daping Hospital and Research Institute of Surgery, Third Military Medical University, Chongqing 400042, China. ⁵Department of Thoracic surgery, Daping Hospital and Research Institute of Surgery, Third Military Medical University, Chongqing 400042, China. ⁶State Key Laboratory of Trauma, Burn and Combined Injury, Daping Hospital and Research Institute of Surgery, Third Military Medical University, Chongqing 400042, China

Conflict of interest

The authors declare that they have no conflict of interest.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41419-018-0452-x>).

Received: 24 October 2017 Revised: 26 February 2018 Accepted: 28 February 2018

Published online: 09 May 2018

References

- Ferlay, J. S. I. et al. *GLOBOCAN 2012v1.1, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11* (International Agency for Research on Cancer, Lyon, 2015).
- Cheng, H. Inhibiting CD146 by its monoclonal antibody AA98 improves radiosensitivity of cervical cancer cells. *Med Sci. Monit.* **22**, 3328–3333 (2016).
- Barney, B. M. et al. Intraoperative electron beam radiotherapy (IOERT) in the management of locally advanced or recurrent cervical cancer. *Radiat. Oncol.* **8**, 80 (2013).
- Powell, M. E. Modern radiotherapy and cervical cancer. *Int J. Gynecol. Cancer* **20**, S49–S51 (2010).
- Chen, Y. et al. IL-6 signaling promotes DNA repair and prevents apoptosis in CD133+ stem-like cells of lung cancer after radiation. *Radiat. Oncol.* **10**, 227 (2015).
- Papathanasiou, M. A. et al. Induction by ionizing radiation of the gadd45 gene in cultured human cells: lack of mediation by protein kinase C. *Mol. Cell Biol.* **11**, 1009–1016 (1991).
- Rosemary Siafakas, A. & Richardson, D. R. Growth arrest and DNA damage-45 alpha (GADD45alpha). *Int. J. Biochem. Cell Biol.* **41**, 986–989 (2009).
- Hollander, M. C. & Fornace, A. J. Jr. Genomic instability, centrosome amplification, cell cycle checkpoints and Gadd45a. *Oncogene* **21**, 6228–6233 (2002).
- Lu, X. et al. Inactivation of gadd45a sensitizes epithelial cancer cells to ionizing radiation in vivo resulting in prolonged survival. *Cancer Res.* **68**, 3579–3583 (2008).
- Hur, J. M. et al. Gliotoxin enhances radiotherapy via inhibition of radiation-induced GADD45a, p38, and NFkappaB activation. *J. Cell. Biochem.* **104**, 2174–2184 (2008).
- Zhang, X. Y. et al. Over-expression of Gadd45a enhances radiotherapy efficacy in human Tca8113 cell line. *Acta Pharmacol. Sin.* **32**, 253–258 (2011).
- Asuthkar, S. et al. Gadd45a sensitizes medulloblastoma cells to irradiation and suppresses MMP-9-mediated EMT. *Neuro Oncol.* **13**, 1059–1073 (2011).
- Klopp, A. H. et al. Gene expression changes in cervical squamous cell carcinoma after initiation of chemoradiation and correlation with clinical outcome. *Int. J. Radiat. Oncol. Biol. Phys.* **71**, 226–236 (2008).
- Qing, Y. et al. Microarray analysis of DNA damage repair gene expression profiles in cervical cancer cells radioresistant to 252Cf neutron and X-rays. *BMC Cancer* **10**, 71 (2010).
- Wu, H. H. et al. Cytoplasmic Ape1 expression elevated by p53 aberration may predict survival and relapse in resected non-small cell lung cancer. *Ann. Surg. Oncol.* **20**, S336–S347 (2013).
- Bobola, M. S., Blank, A., Berger, M. S., Stevens, B. A. & Silber, J. R. Apurinic/apyrimidinic endonuclease activity is elevated in human adult gliomas. *Clin. Cancer Res.* **7**, 3510–3518 (2001).
- Sak, S. C., Harnden, P., Johnston, C. F., Paul, A. B. & Kiltie, A. E. APE1 and XRCC1 protein expression levels predict cancer-specific survival following radical radiotherapy in bladder cancer. *Clin. Cancer Res.* **11**, 6205–6211 (2005).
- Bobola, M. S. et al. Apurinic/apyrimidinic endonuclease is inversely associated with response to radiotherapy in pediatric ependymoma. *Int. J. Cancer* **129**, 2370–2379 (2011).
- Herring, C. J. et al. Levels of the DNA repair enzyme human apurinic/apyrimidinic endonuclease (APE1, APEX, Ref-1) are associated with the intrinsic radiosensitivity of cervical cancers. *Br. J. Cancer* **78**, 1128–1133 (1998).
- Raffoul, J. J. et al. Down-regulation of apurinic/apyrimidinic endonuclease 1/redox factor-1 expression by soy isoflavones enhances prostate cancer radiotherapy in vitro and in vivo. *Cancer Res.* **67**, 2141–2149 (2007).
- Xiang, D. B. et al. Chimeric adenoviral vector Ad5/F35-mediated APE1 siRNA enhances sensitivity of human colorectal cancer cells to radiotherapy in vitro and in vivo. *Cancer Gene Ther.* **15**, 625–635 (2008).
- Singh-Gupta, V. et al. Soy isoflavones augment radiation effect by inhibiting APE1/Ref-1 DNA repair activity in non-small cell lung cancer. *J. Thorac. Oncol.* **6**, 688–698 (2011).
- Chen, S. M., Xiong, G. S., Wu, S. M. & Mo, J. Z. Downregulation of apurinic/apyrimidinic endonuclease 1/redox factor-1 enhances the sensitivity of human pancreatic cancer cells to radiotherapy in vitro. *Cancer Biother. Radiopharm.* **28**, 169–176 (2013).
- Cun, Y. P. et al. Silencing of APE1 enhances sensitivity of human hepatocellular carcinoma cells to radiotherapy in vitro and in a xenograft model. *PLoS ONE* **8**(2) e55313 (2013).
- Jung, H. J. et al. Base excision DNA repair defect in Gadd45a-deficient cells. *Oncogene* **26**, 7517–7525 (2007).
- Kim, H. L., Kim, S. U. & Seo, Y. R. A novel role for Gadd45alpha in base excision repair: modulation of APE1 activity by the direct interaction of Gadd45alpha with PCNA. *Biochem. Biophys. Res. Commun.* **434**, 185–190 (2013).
- Cun, Y. et al. Silencing of APE1 enhances sensitivity of human hepatocellular carcinoma cells to radiotherapy in vitro and in a xenograft model. *PLoS ONE* **8**, e55313 (2013).
- Wu, H. H. et al. Cytoplasmic Ape1 expression elevated by p53 aberration may predict survival and relapse in resected non-small cell lung cancer. *Ann. Surg. Oncol.* **20**, S336–S347 (2013).
- Sheng, Q. S. et al. Prognostic significance of APE1 cytoplasmic localization in human epithelial ovarian cancer. *Med. Oncol.* **29**, 1265–1271 (2012).
- Puglisi, F. et al. Prognostic role of Ape/Ref-1 subcellular expression in stage I-III breast carcinomas. *Oncol. Rep.* **9**, 11–17 (2002).
- Di Maso, V. et al. Subcellular localization of APE1/Ref-1 in human hepatocellular carcinoma: possible prognostic significance. *Mol. Med.* **13**, 89–96 (2007).

32. Qing, Y. et al. The expression of APE1 and its correlation with prognostic significance after 252Cf radiotherapy in cervical cancer. *Sichuan Da Xue Xue Bao Yi Xue Ban.* **40**, 125–128 (2009).
33. Qu, J., Liu, G. H., Huang, B. & Chen, C. Nitric oxide controls nuclear export of APE1/Ref-1 through S-nitrosation of cysteines 93 and 310. *Nucleic Acids Res.* **35**, 2522–2532 (2007).
34. Xu, C. X. et al. Poly(ester amine)-mediated, aerosol-delivered Akt1 small interfering RNA suppresses lung tumorigenesis. *Am. J. Resp. Crit. Care* **178**, 60–73 (2008).
35. Xu, C. X. et al. Chondroitin sulfate extracted from the *Styela clava* tunic suppresses TNF-alpha-induced expression of inflammatory factors, VCAM-1 and iNOS by blocking Akt/NF-kappaB signal in JB6 cells. *Cancer Lett.* **264**, 93–100 (2008).
36. Kitahara, O., Katagiri, T., Tsunoda, T., Harima, Y. & Nakamura, Y. Classification of sensitivity or resistance of cervical cancers to ionizing radiation according to expression profiles of 62 genes selected by cDNA microarray analysis. *Neoplasia* **4**, 295–303 (2002).