


Knowledge about the Developmental Origins of Health and Disease is independently associated with variation in diet quality during pregnancy

Luseadra McKerracher^{1,2}  | Tina Moffat¹ | Mary Barker³ | Meghan McConnell^{4,5} |
Stephanie A. Atkinson⁶ | Beth Murray-Davis⁷ | Sarah D. McDonald^{8,9,10} |
Deborah M. Sloboda^{2,6,7,10}

¹Department of Anthropology, McMaster University, Hamilton, Ontario, Canada

²Department of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, Ontario, Canada

³MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK

⁴Department of Innovation in Medical Education, University of Ottawa, Ottawa, Ontario, Canada

⁵Department of Anesthesiology and Pain Medicine, University of Ottawa, Ottawa, Ontario, Canada

⁶Department of Pediatrics, McMaster University, Hamilton, Ontario, Canada

⁷Department of Obstetrics and Gynecology, McMaster University, Hamilton, Ontario, Canada

⁸Department of Radiology, McMaster University, Hamilton, Ontario, Canada

⁹Department of Health Research Health Research Methods, Evidence & Impact, McMaster University, Hamilton, Ontario, Canada

¹⁰Farncombe Family Digestive Diseases Research Institute, McMaster University, Hamilton, Ontario, Canada

Correspondence

Dr Luseadra McKerracher, Department of Anthropology and Department of Biochemistry and Biomedical Sciences, McMaster University, 1280 Main Street West, CNH 524, Hamilton, Ontario L8S 4L8, Canada. Email: mckerrl@mcmaster.ca

Professor Deborah M. Sloboda, Department of Biochemistry and Biomedical Sciences, McMaster University, 1280 Main Street West, HSC 4H30A, Hamilton, Ontario L8S 4K1, Canada. Email: sloboda@mcmaster.ca

Funding information

Canadian Institutes of Health Research, Grant/Award Number: 146333; Women's College Hospital Xchange Challenge; McMaster University, Grant/Award Number: Provost's Award; Canadian Foundation for Innovation; Canada Research Chairs program; Canadian Institute for Health Research, Grant/Award Number: 146333

Abstract

Environmental factors affecting development through embryogenesis, pregnancy, and infancy impact health through all subsequent stages of life. Known as the Developmental Origins of Health and Disease (DOHaD) hypothesis, this concept is widely accepted among health and social scientists. However, it is unclear whether DOHaD-based ideas are reaching the general public and/or influencing behaviour. This study thus investigated whether and under what circumstances pregnant people in Canada are familiar with DOHaD, and if DOHaD familiarity relates to eating behaviour. Survey responses from pregnant people from Hamilton, Canada, were used to assess respondents' knowledge of DOHaD (hereafter, DOHaD_{KNOWLEDGE}) compared with their knowledge of more general pregnancy health recommendations (Pregnancy Guideline_{KNOWLEDGE}). The survey also characterized respondents' pregnancy diet quality and sociodemographic profiles. We fit two multiple, linear, mixed regression models to the data, one with DOHaD_{KNOWLEDGE} score as the dependent variable and the other with diet quality score as the dependent. In both models, responses were clustered by respondents' neighbourhoods. Complete, internally consistent responses were available for 330 study-eligible respondents. Relative to Pregnancy Guideline_{KNOWLEDGE}, respondents had lower, more variable DOHaD_{KNOWLEDGE}

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Maternal & Child Nutrition* published by John Wiley & Sons, Ltd.

scores. Additionally, higher DOHaD_{KNOWLEDGE} was associated with higher socio-economic position, older age, and lower parity, independent of Pregnancy Guideline_{KNOWLEDGE}. Diet quality during pregnancy was positively associated with DOHaD_{KNOWLEDGE}, adjusting for sociodemographic factors. A subset of relatively high socio-economic position respondents was familiar with DOHaD. Greater familiarity with DOHaD was associated with better pregnancy diet quality, hinting that translating DOHaD knowledge to pregnant people may motivate improved pregnancy nutrition and thus later-life health for developing babies.

KEYWORDS

developmental origins, diet quality, health inequities, knowledge translation, pregnancy, nutrition

1 | INTRODUCTION

Non-communicable diseases (NCDs) related to obesity and metabolic function (e.g., cardiovascular disease and type 2 diabetes mellitus) represent leading causes of illness and death globally (World Health Organization, 2017). Although these diseases manifest mainly in adulthood, abundant evidence indicates that the main risk factors for metabolism-related NCDs are established during the peri-conceptual and foetal stages of development (Barker et al., 2017; Fleming et al., 2018; Stephenson et al., 2018). The idea that early life—especially prenatal—environments shape developmental trajectories, which then affect later-life disease risk (Barker & Osmond, 1986), is now most widely referred to as the Developmental Origins of Health and Disease (DOHaD) hypothesis (Barnes, Heaton, Coates, & Packer, 2016).

The DOHaD hypothesis offers a powerful set of explanations accounting for much of the epidemiology of NCDs related to metabolic function (Godfrey et al., 2017; Hanson & Gluckman, 2015). Furthermore, models derived from the DOHaD hypothesis indicate that strategies for reducing the burden of metabolic disease should focus on intervening in the earliest stages of life (Garmendia, Corvalan, & Uauy, 2014; Godfrey et al., 2017; Hanson & Gluckman, 2015; Low, Gluckman, & Hanson, 2018; Stephenson et al., 2018; Victora et al., 2016). As such, over the last decade, there has been an increasing push to translate key findings from DOHaD-related research into social and public health policy and guidelines as well as into clinical practice (Anon., 2011; Barker et al., 2017; Mckerracher, Moffat, Barker, Williams, & Sloboda, 2019).

Indeed, influential stakeholders and institutions in several nations with strong DOHaD-oriented research communities are beginning to direct efforts towards DOHaD knowledge translation (Baird, Cooper, Margetts, Barker, & Inskip, 2009; Barker et al., 2017; Bay, Mora, Sloboda, & Morton, 2012, 2017b; McMullan et al., 2017; MacNab & Mukisa, 2017; Mckerracher et al., 2019; Oyamada, Lim, Dixon, Wall, & Bay, 2018; Woods-Townsend et al., 2018; Perreault et al., 2016). What remains unclear is the extent to which DOHaD findings are effectively reaching populations and organizations best positioned to

Key messages

- It is now well established that early life environmental exposures, especially maternal diet during pregnancy, impact children's development and health across the life span. As such, identification of motivational factors that influence diets during pregnancy is needed.
- Understanding long-term impacts on children of diet and health during pregnancy may help motivate pregnant people to consume macronutrient and micronutrient dense foods.
- Socio-economic and related demographic factors are associated with both quality of pregnancy diet and understanding of the long-term impacts of pregnancy diet.
- To support healthy eating during pregnancy, policy should focus simultaneously on providing income supports for pregnant people unable to afford to prioritize eating high-quality diets and promoting understanding of the idea that prioritizing high-quality nutrition during pregnancy has outsized and long-term effects on children's health.

make use of DOHaD-related research findings. In particular, we do not yet know the extent to which pregnant people and their clinical, social, and local political supports are receiving the idea that early life environments impact metabolic profile and function over the entire lifespan (McMullan et al., 2017). Understanding the current state of DOHaD knowledge translation and its potential impacts is critical to developing next steps, both on the ground and in public policy arenas.

To begin to address this gap in our understanding, this study had three aims: (1) to measure familiarity with key concepts underpinning the DOHaD hypothesis in a sample of pregnant people in Canada, (2) to assess the extent to which familiarity with DOHaD concepts varied within the sample, and (3) to investigate whether having some

familiarity with DOHaD concepts might influence behaviour, particularly regarding diet quality during pregnancy.

We hypothesized that, because DOHaD knowledge translation efforts are relatively new in general and in Canada in particular (McKerracher et al., 2019), people outside of the research community would be largely unfamiliar with DOHaD-related concepts. We predicted that pregnant Canadians would have lower familiarity with DOHaD-related concepts relative to their familiarity with general pregnancy health guidelines as issued by Health Canada. The core of these latter guidelines has been in place since 1995 (Haertsch, Campbell, & Sanson-Fisher, 1999) and is widely communicated by public health practitioners and media channels (Chalmers, Dzakupasu, Heaman, & Kaczorowski, 2008).

We further hypothesized that pregnant people's understanding of DOHaD-related concepts would vary and that this variation would be patterned sociodemographically. Health knowledge and level of trust in health workers and systems differs substantially among communities and sociodemographic groups in Canada (Glouberman & Millar, 2003; Jenkins & Keating, 1999), as elsewhere (Chinn, 2011; Sørensen et al., 2012). Specifically, people of higher socio-economic position (SEP; i.e., higher income, greater material wealth, and more years of formal education) tend to be more knowledgeable about their own health and more trusting of health care institutions (Sørensen et al., 2015; Willems, De Maesschalck, Deveugele, Derese, & De Maeseneer, 2005). As such, these people generally have greater health literacy (ability to understand health information and to navigate health systems) and a stronger sense of self-efficacy in relation to health care (Nutbeam, 2008; Smith, Dixon, Trevena, Nutbeam, & McCaffery, 2009). Additionally, older people with greater parenting experience have had more opportunities to accrue knowledge and understanding (Bornstein, Cote, Haynes, Hahn, & Park, 2010). Multi-year experience with navigating and understanding a given health system is also associated with improved health knowledge and literacy (Kalich, Heinemann, & Ghahari, 2016; Simich, 2009). Thus, we anticipated that familiarity with DOHaD concepts would be positively associated with respondents' SEP, age, and parity and negatively associated with respondents' statuses as newcomers to Canada, holding constant familiarity with general pregnancy health guidelines.

The last hypothesis we tested was that pregnant respondents would be more likely to prioritize eating nutritious foods if they understood that what they ate during pregnancy could impact their children's health not only perinatally but through subsequent stages of life. Our test prediction was that self-reported quality of diet during pregnancy would be positively associated with familiarity with DOHaD-related concepts, independent of other factors known/expected to influence pregnancy diet (age, SEP, parity, newcomer status, and knowledge of general pregnancy health recommendations).

2 | MATERIALS AND METHODS

2.1 | Setting and participants

Participation in this study was restricted to people who were pregnant and living in Hamilton, Canada, at the time of data collection (between

December 2015 and September 2018). Hamilton is a suitable site to evaluate our hypotheses for three main reasons. First, it is sociodemographically diverse, with known, marked variation in access to knowledge, support, and financial resources (Dean & Elliott, 2012; DeLuca, Buist, & Johnston, 2012; Latham & Moffat, 2007; Moffat & Galloway, 2008). Second, the city has a high prevalence of obesity and metabolism-related NCDs for an urban centre of its size (Navaneelan & Janz, 2014). Lastly, Hamilton is characterized by well-documented, systematic health inequities among neighbourhoods, including in early life indicators of later-life NCD risks such as low birthweight (DeLuca et al., 2012).

We advertised our anonymous survey, described further below, at community health centres, midwifery clinics, prenatal classes, hospitals, and other community organizations across Hamilton. We used a targeted approach when advertising our survey to ensure that pregnant people from diverse backgrounds were appropriately represented in our sample. That is, we promoted the study intensively at community health centres with prenatal programming where attendees were relatively likely to come from low-income, low-educational attainment households (Denny & Grady, 2007) and/or to identify as non-White and/or as newcomers to Canada (Durant et al., 2007; Oh et al., 2015).

2.2 | Study instrument

We developed on-line (LimeSurvey©) and print versions of a 140-item questionnaire in English. It was piloted with 78 pregnant people between December of 2015 and August of 2016 to ensure reliability of items and range of response selection across items. It was slightly modified, then relaunched, and rerun between June 2017 and September 2018; following the relaunch, a print version was also translated and made available in Arabic, the most common language spoken at a main study promotion location. The questionnaire included questions grouped into the following four domains: (1) the extent to which respondents were familiar with core concepts and evidence related to the DOHaD hypothesis (hereafter, DOHaD_{KNOWLEDGE}), (2) the extent to which respondents were familiar and in agreement with Health Canada's general guidelines for health during pregnancy (hereafter, Pregnancy Guideline_{KNOWLEDGE}), (3) respondents' self-reported diet quality during pregnancy, and (4) respondents' sociodemographic characteristics.

2.3 | Measures

To assess DOHaD_{KNOWLEDGE}, we aggregated scores concerning the level at which respondents agreed, using a 5-point Likert scale (0 = *strongly disagree* to 4 = *strongly agree*), with each of five statements about the long-term impacts of peri-conceptual and prenatal and perinatal parental and/or grandparental health and behaviour on children's health. The five statements are presented in Table 1. The aggregated scores resulted in a 20-point DOHaD_{KNOWLEDGE} scale, which could take a minimum value of 0 (indicating no knowledge/familiarity of concepts and evidence related to the

TABLE 1 Components of the DOHaD_{KNOWLEDGE} scale and the Pregnancy Guideline_{KNOWLEDGE} scale

DOHaD _{KNOWLEDGE} scale	<p>What a woman eats during her pregnancy affects her baby's risk of becoming obese as an adult.</p> <p>What a woman eats during her pregnancy affects her grandchildren's risk of becoming obese.</p> <p>Before pregnancy, both what the mother and the father eat affects the growth and health of their baby.</p> <p>What I ate before pregnancy affects my child's chance of becoming obese as an adult.</p> <p>What I eat while I am breastfeeding affects my child's chance of becoming obese as an adult.</p>
Pregnancy Guideline _{KNOWLEDGE} scale	<p>Smoking during pregnancy will harm my baby.</p> <p>Taking daily prenatal vitamins during my pregnancy is good for my baby's health.</p> <p>Pregnant women should not eat as much as they like because they are "eating for two."</p> <p>Eating nutritious food is my top priority.</p> <p>I would consider ____ to be a healthy birthweight for my baby. (Options: ≤ 1.5 kg = 0 points; 1.5–2.5 kg = 1 point; 2.5–3 kg = 2 points; 3–3.5 kg = 4 points; 3.5–4 kg = 5 points; 4–4.5 kg = 2 points; ≥ 4.5 kg = 0 points)</p>

DOHaD hypothesis) and a maximum value of 20 (indicating very strong knowledge and uptake of such concepts and evidence).

Pregnancy Guideline_{KNOWLEDGE} was assessed in the same way as DOHaD_{KNOWLEDGE}, but using three statements central to Health Canada's general recommendations for having a healthy pregnancy, one statement implicit in Health Canada's recommendations, and one statement based on its prenatal guidelines for health professionals. These Pregnancy Guideline_{KNOWLEDGE} statements are presented in Table 1, alongside those comprising DOHaD_{KNOWLEDGE}. The first four of these Pregnancy Guideline_{KNOWLEDGE} statements were assessed on a 5-point Likert scale. The last was assessed on a 5-point scale based on how close a participant's selected response came to Health Canada's reported 50th percentile for a healthy, full-term birthweight as of 2016 (3.5 kg). As with DOHaD_{KNOWLEDGE}, Pregnancy Guideline_{KNOWLEDGE} statements were aggregated into a 20-point scale, ranging from 0 to 20. To further probe respondents' familiarity with Health Canada's pregnancy guidelines, respondents were also given the opportunity to answer two open-ended questions concerning foods to eat and foods to avoid during pregnancy. These responses were not included in the Pregnancy Guideline_{KNOWLEDGE} scale but are reported briefly in Text S1.

Dietary practices during pregnancy were reported by the respondents using the PrimeScreen© Food Frequency Questionnaire (Rifas-Shiman et al., 2001), which was integrated into our larger questionnaire. Respondents were asked to record how often during their pregnancies they ate foods from each of 20 mutually exclusive food categories. Possible responses ranged from "never" to "more than twice a day." Frequent consumption of foods from each nutrient-dense food category was given a positive score up to a maximum of 4 (with the exception of the category low-fat dairy foods, which could

take a maximum value of 8). Frequent consumption of each nutrient-poor food was given a negative score, with the most negative possible value being -4 . All possible responses to the food frequency questions are presented in Table 2, along with three example food categories and their associated scores. A full list of food categories and scores is available in Text S2 and Table S1. After scoring each individual food category, a total of all food category scores for each respondent was calculated to give a diet quality score, which could take a minimum value of -16 and a maximum of 36.

The sociodemographic characteristics assessed in the survey comprised respondent's age (years), highest completed level of education (ranging from 1 = *had not obtained a high school diploma* to 4 = *had completed post-secondary degree*; further information on this variable available in Text S3 and Table S2), household annual income bracket (1 = $< \$23,000$, 2 = $\$23,000$ – $\$39,999$, 3 = $\$40,000$ – $\$79,999$, and 4 = $> \$80,000$), main source of household income (wages/salaries; other, e.g., social assistance and inheritance), current postal (zip) code, self-identified ethnicity, status as a newcomer or not (temporary residents and permanent residents of Canada who had lived in Canada ≤ 5 years vs. Canadian citizens and permanent residents who had lived in Canada > 5 years), marital status (married/common-law partnership, single, divorced, widowed, and others), and parity. Details on collation of these sociodemographic variables are reported in Section 2.4

2.4 | Data preparation

We began by exploring the data, characterizing the sample sociodemographically, reducing the number of sociodemographic predictors (to avoid multicollinearity and model overfitting in regression analyses, described below), and removing 63 cases for which

TABLE 2 PrimeScreen© Food Frequency Questionnaire (Rifas-Shiman et al., 2001)

Example food category	More than once a day	Almost daily	Two to four times a week	Once a week	Less than once a week	Never
Dark green leafy vegetables (e.g., kale, turnip greens, bokchoy, and Swiss chard)	<input type="checkbox"/> +4	<input type="checkbox"/> +3	<input type="checkbox"/> +2	<input type="checkbox"/> +1	<input type="checkbox"/> 0	<input type="checkbox"/> 0
Fish/seafood	<input type="checkbox"/> +4	<input type="checkbox"/> +3	<input type="checkbox"/> +2	<input type="checkbox"/> +1	<input type="checkbox"/> 0	<input type="checkbox"/> 0
Sugary drinks (e.g., soda, fruit drinks, and Gatorade)	<input type="checkbox"/> -4	<input type="checkbox"/> -3	<input type="checkbox"/> -2	<input type="checkbox"/> -1	<input type="checkbox"/> 0	<input type="checkbox"/> 0

participants discontinued responding within the first 20 survey questions. We also removed one duplicate submission, six submissions for which respondents' postal codes fell outside of Hamilton's boundaries, and four submissions in which there were clear internal inconsistencies in responses.

To decrease the number of sociodemographic variables included in our models, we combined those directly related to SEP (respondent's household income bracket, respondent's educational attainment group, and source of income) into an SEP index (hereafter, SEP score) for use in statistical analyses. SEP, an additive score of educational and income brackets, ranges from two to eight. A score of 2 indicates that, at the time of data collection, the respondent had not completed high school AND lived in a household in which the annual income is <\$23,000; all such respondents also indicated some form of social assistance as a source of income. A score of 8 indicates that the respondent had completed at least one college/university degree AND lived in a household with an annual, employment-based income >\$80,000. A detailed breakdown of all SEP scores is available in Text S3 and Table S3.

Other intercorrelated sociodemographic variables did not lend themselves to straightforward amalgamation, so we selectively dropped out those that were highly collinear with others but for which we did not have theoretical justifications for retaining in our models. Specifically, marital status and self-reported non-White ethnicity were not included because we had not identified them a priori as independent drivers of pregnancy health knowledge or pregnancy diet quality, and they were correlated with respondents' ages, SEP, parities, and/or with status as newcomers to Canada. Further details on the relationships among sociodemographic variables, along with the results of one sensitivity model in which we adjusted for non-White ethnicity instead of newcomer status, are available in Text S3.

Postal code, truncated to its first three digits to represent 25 neighbourhood areas in Hamilton, was treated as a clustering variable, because previous empirical research indicates that sociodemographic variables, behavioural patterns, and health outcomes are strongly geographically patterned in the city (DeLuca et al., 2012). So, treating respondents from the same neighbourhoods as observations independent from one another is inappropriate.

2.5 | Statistical analyses

We carried out four sets of analyses, the first to assess the internal reliability of our DOHaD_{KNOWLEDGE} and Pregnancy Guideline_{KNOWLEDGE} scales and the subsequent ones to evaluate our three test predictions.

To assess the internal consistency of the DOHaD_{KNOWLEDGE} and Pregnancy Guideline_{KNOWLEDGE} scores, we calculated the Cronbach's α (Cronbach & Murphy, 1970) for each scale. This statistic can take values between 0 and 1. A Cronbach's α above \sim .7 indicates that component scale items take similar values for a given respondent, whereas totalled scores vary substantially among participants' responses. Low Cronbach's α values (below .5) may indicate either low intra-item

cohesiveness within respondents or low inter-respondent variation in responses.

To evaluate our first prediction that DOHaD_{KNOWLEDGE} is less likely to have successfully reached pregnant people than Pregnancy Guideline_{KNOWLEDGE}, we compared means (using a Wilcoxon ranked sum test) and coefficients of variation between the two knowledge scales. To assess our second prediction that DOHaD_{KNOWLEDGE} is likely to be sociodemographically patterned, we fit a linear mixed effects regression model using maximum likelihood to the data. Fixed effects in the model were respondent's age, SEP score, status as a newcomer to Canada, and parity. Intercepts were allowed to vary randomly among the 25 postal code (neighbourhood) groups. As SEP score is strongly left skewed, resulting in heteroscedasticity in the model's residuals, we power-transformed this variable for use in a sensitivity analysis, raising the exponent to five, at which point the residual error appears homoscedastic (Text S4 and Figure S1). Third, to assess the prediction that DOHaD_{KNOWLEDGE} would be associated with greater positive attention to health-related behaviours during pregnancy, we modelled diet quality score as a maximum likelihood function of DOHaD_{KNOWLEDGE}, adjusting for respondent's age, SEP score, newcomer status, Pregnancy Guideline_{KNOWLEDGE} score and parity, again clustering the data by postal code group and allowing intercepts to vary randomly among neighbourhoods. No data transformations or other modifications to this model appeared necessary (Text S4 and Figure S2).

Alpha was set at $P = 0.05$ for all statistical analyses. All analyses were carried out and all plots were made in the *stat* (R Core Development Team, 2017), *nlme* (Pinheiro, Bates, DebRoy, Sarkar, & Team, 2018), *psych* (Revelle, 2018), and *ggplot2* (Wickham, 2016) packages of the statistical environment, R.

2.6 | Research ethics approval

The study protocol used to evaluate these predictions was reviewed and approved by the Hamilton Integrated Research Ethics Board, approval #0570.

3 | RESULTS

Complete, internally consistent responses to all questions used to assess DOHaD_{KNOWLEDGE}, Pregnancy Guideline_{KNOWLEDGE}, diet quality, respondent's age, SEP score, parity, and status as a newcomer to Canada were available from 330 respondents, inclusive of the 78 respondents to the pilot. Descriptive statistics for the continuous sociodemographic characteristics, DOHaD_{KNOWLEDGE} score, Pregnancy Guideline_{KNOWLEDGE} score, and diet quality score are presented in the text below and in Table 3. The distribution of SEP scores is also represented in Figure 1.

Participants varied sociodemographically, ranging in age from 17 to 47 (mean = 30.4, $SE = \pm 0.29$), in SEP score from two to eight (mean = 6.4, median = 7, $SE = \pm 0.11$), and in parity from zero to six (mean = 0.8, $SE = \pm 0.06$). Nine percent of respondents were identified

TABLE 3 Descriptive statistics for continuous and ordinal sociodemographic characteristics of sample and for DOHaD_{KNOWLEDGE} score, Pregnancy Guideline_{KNOWLEDGE} score, and diet quality score: N = 330

Characteristic	Min	Median (IQR)	Mean (SE)	Max
DOHaD _{KNOWLEDGE} score	0	10 (6–13)	9.4 (±0.25)	20
Pregnancy Guideline _{KNOWLEDGE} score	7	15 (14–16)	14.5 (±0.10)	20
Diet quality score	–14	12 (5–18)	11.4 (±0.52)	36
Maternal age (years)	17	30 (27–34)	30.5 (±0.29)	47
SEP score	2	7 (5–8)	6.4 (±0.11)	8
Number of previous births	0	0 (0–1)	0.8 (±0.06)	6
Status as newcomer to Canada (0 = has lived in Canada more than 5 years and 1 = has lived in Canada fewer than 5 years)	0	N/A	N/A	1

as newcomers to Canada (i.e., they had immigrated to Canada within the 5 years before participating in the study).

The DOHaD_{KNOWLEDGE} scale was internally reliable, with a Cronbach's α of .82, indicating that its component statements effectively measure the same mental construct (Cronbach & Murphy, 1970). Respondents' DOHaD_{KNOWLEDGE} scores ranged from 0 to 20, with a mean (SE) of 9.4 (±0.25; Table 3).

Pregnancy Guideline_{KNOWLEDGE} scale had a low Cronbach's α (.27). This low value can be attributed to low inter-individual variation in responses to any items (90% of respondents scored 4 or 5 on all five items), rather than to low inter-item agreement. Respondents'

pregnancy Guideline_{KNOWLEDGE} scores took values ranging from 7 to 20. The mean (SE) of this variable was 14.5 (±0.10; Table 3).

The minimum diet quality score was –14 and the maximum was 36. The mean (SE) for this variable was 11.4 (±0.52).

3.1 | Analysis for Hypothesis 1 (comparing DOHaD_{KNOWLEDGE} and Pregnancy Guideline_{KNOWLEDGE} scores)

The mean DOHaD_{KNOWLEDGE} score of 9.4 was ~5 points lower than the sample's mean Pregnancy Guideline_{KNOWLEDGE} score of 14.5 ($P =$




FIGURE 1 Frequency of responses from participants in each socio-economic position (SEP) bracket. SEP scores of 2–4 indicate lower household income (<\$23,000 per year) and low (no completed post-secondary) respondent educational attainment levels. SEP scores of 5–7 indicate higher respondent educational attainment levels (at least some post-secondary) and household income brackets near the city's median (\$40,000–\$79,000). SEP scores of 8 indicate high respondent educational attainment levels (completion of at least one post-secondary degree) and household incomes >\$80,000 per year. Seventy (21%) of respondents have SEP scores of ≤ 4 , 127 (38%) have scores of 5–7, and 133 (40%) have SEP scores of 8.




FIGURE 2 Distributions of respondents' Pregnancy Guideline_{KNOWLEDGE} and DOHaD_{KNOWLEDGE} scores. Ninety-one (27%) of respondents have Pregnancy Guideline_{KNOWLEDGE} scores of 16, the modal score for this variable. Forty (12%) of respondents have DOHaD_{KNOWLEDGE} scores of 10, the modal score for this variable

0.000). The coefficient of variation for DOHaD_{KNOWLEDGE} score was 0.49, indicating substantial variation in respondents' familiarity with DOHaD-related concepts. The coefficient of variation for Pregnancy Guideline_{KNOWLEDGE} was 0.13, indicating a high level of consensus among respondents in whether to agree with the statements this scale comprises. These differences in the spreads and central tendencies between DOHaD_{KNOWLEDGE} and Pregnancy Guideline_{KNOWLEDGE} are summarized in Figure 2. Moreover, respondents' qualitative answers about foods to eat and to avoid during pregnancy aligned with recommendations from Health Canada (Text S1), further supporting a high level of consensus with respect to Pregnancy Guideline_{KNOWLEDGE}.

3.2 | Analysis for Hypothesis 2 (inequities in DOHaD_{KNOWLEDGE})

The results of the DOHaD_{KNOWLEDGE} by sociodemographic factors regression are presented in Table 4. Consistent with our predictions, we found that DOHaD_{KNOWLEDGE} was positively associated with SEP score (Figure 3) and respondent's age, adjusting for status as a newcomer to Canada, Pregnancy Guideline_{KNOWLEDGE} score, and parity and clustering the data by neighbourhood. There was no clear statistical evidence that DOHaD_{KNOWLEDGE} is independently associated with status as a newcomer to Canada or with Pregnancy Guideline_{KNOWLEDGE}.

TABLE 4 Relationships between DOHaD_{KNOWLEDGE} score and maternal age, SEP score, previous childbearing experience, and Pregnancy Guideline_{KNOWLEDGE} score in a full, linear mixed effects model

Fixed effect	Estimate	95% CI	P value
SEP score	0.78	[0.49, 1.07]	0.000***
Maternal age (years)	0.12	[0.02, 0.21]	0.021*
Number of previous births	-0.47	[-0.91, -0.02]	0.040*
Status as a newcomer to Canada	1.44	[-0.05, 3.03]	0.078 [†]
Pregnancy Guideline _{KNOWLEDGE}	0.22	[-0.01, 0.45]	0.062 [†]

Note. Intercepts were allowed to vary randomly among postal code groupings (neighbourhoods). Model summary: AIC = 1892.8, log likelihood = -938.4.

* $P \leq 0.050$, ** $P \leq 0.010$, *** $P \leq 0.001$. [†] $P \leq 0.100$.

DOHaD_{KNOWLEDGE} was negatively associated with parity, adjusting for all other predictors.

Although we report here the results of the model using raw SEP score as a predictor to simplify interpretation of this analysis focused largely on SEP, we acknowledge that the error is likely mis-specified in this model, as the residual plot indicates high levels of heteroscedasticity (Text S4 and Figure S1a). Transforming SEP score




FIGURE 3 Scatterplot of linear relationship between DOHaD_{KNOWLEDGE} score and SEP score. Data are presented as individual scores of DOHaD_{KNOWLEDGE} and SEP. Open circles = respondents without prior pregnancies, closed circles = respondents who have one or two children, and pluses = respondents with three and more children. As SEP score increases, DOHaD_{KNOWLEDGE} score increases. Respondents with three and more children (pluses) are relatively likely to fall below the fit line.

to the fifth exponent eliminates this problem (Text S4 and Figure S1b), and the model in which (SEP score)⁵ is included in lieu of raw SEP score produces results consistent with our main findings (see Text S4 and Table S4).

3.3 | Analysis for Hypothesis 3 (factors associated with diet quality score)

The results of the analysis in which we regressed diet quality score on DOHaD_{KNOWLEDGE} score, Pregnancy Guideline_{KNOWLEDGE} score, and sociodemographic factors are summarized in Table 5. Consistent with

TABLE 5 Relationships between diet quality score and DOHaD_{KNOWLEDGE} score, maternal age, SEP score, previous childbearing experience, and Pregnancy Guideline_{KNOWLEDGE} score in a full, linear mixed effects model

Fixed effect	Estimate	95% CI	P value
DOHaD _{KNOWLEDGE} score	0.35	[0.13, 0.56]	0.003**
SEP score	0.62	[0.00, 1.24]	0.051 [†]
Age	0.31	[0.10, 0.51]	0.003**
Number of previous births	-1.35	[-2.27, -0.43]	0.005**
Status as a newcomer to Canada	-6.46	[-3.17, 1.12]	0.062 [†]
Pregnancy Guideline _{KNOWLEDGE}	0.04	[-0.44, 0.52]	0.869

Note. Intercepts were allowed to vary randomly among postal code groupings (neighbourhoods). Model summary: AIC = 2372.0, log likelihood = -1177.0.

* $P \leq 0.050$. ** $P \leq 0.010$. *** $P \leq 0.001$. [†] $P \leq 0.100$.

expectations, diet quality score was positively associated with DOHaD_{KNOWLEDGE} score (Figure 4). Diet quality score was also positively associated with respondent's age, SEP score, and negatively associated with parity. We highlight also that there was no statistical evidence that diet quality score was independently associated with general Pregnancy Guideline_{KNOWLEDGE} score.

4 | DISCUSSION

We developed and used a novel instrument to show that most respondents were very knowledgeable about general pregnancy health recommendations as laid out by Health Canada (Pregnancy Guideline_{KNOWLEDGE}), but strikingly few respondents, by comparison, were knowledgeable about DOHaD concepts (DOHaD_{KNOWLEDGE}). Further, the data indicate that DOHaD_{KNOWLEDGE} is inequitably distributed with higher DOHaD_{KNOWLEDGE} scores concentrated among respondents of higher SEP, older age, and lower parity. We also show that those respondents who have some understanding of DOHaD were more likely to report a more nutrient-dense diet during pregnancy, above and beyond what well-known diet quality predictors like age, parity, and SEP would predict. We argue that these inequities in DOHaD_{KNOWLEDGE} and in pregnancy diet quality may be partly attributable to politically-, institutionally-, and historically structured differences among sociodemographic subgroups within the sample in the dissemination, accessibility, and/or uptake of DOHaD concepts.




FIGURE 4 Scatterplot of linear relationship between diet quality score and DOHaD_{KNOWLEDGE} score. Data are presented as individual scores of diet quality score and DOHaD_{KNOWLEDGE} score. Open circles = respondents without prior pregnancies, closed circles = respondents who have one or two children, and pluses = respondents with three and more children. As DOHaD_{KNOWLEDGE} score increases, diet quality score increases. Respondents with three and more previous children (pluses) are relatively likely to fall below the fit line.

4.1 | Familiarity with pregnancy health guidelines, but not their long-term implications

Although respondents were clearly familiar with general recommendations about how to live healthfully during pregnancy, they were generally unfamiliar with the idea that prenatal health/nutrition influences the long-term obesity and metabolic disease risks of their children. This may suggest that, as predicted, pregnancy guidelines are being systematically and confidently disseminated and monitored by reputable sources, mainly Health Canada (Chalmers et al., 2008). While Health Canada is educating health professionals about the long-term consequences of excessive gestational weight gain (Health Canada, 2014), it may not yet be systematically disseminating the idea directly to pregnant people that the early life nutritional environment plays a significant role in determining later-life obesity and metabolic disease risks.

Although these results align with our test predictions regarding knowledge translation patterns, we acknowledge that there are other plausible explanations for the discrepancy in scores between the DOHaD_{KNOWLEDGE} and Pregnancy Guideline_{KNOWLEDGE} scales. One such alternative explanation includes how cognitively easy or difficult it is to assimilate concepts measured by each of the scales. The DOHaD hypothesis has been discussed academically for the last 30 years (Barnes et al., 2016), but its translation has been difficult (McKerracher et al., 2019). The core idea that the early life environment shapes how you grow and thus can affect your later-life health is relatively straightforward. However, what exactly is meant by environment, how different kinds of environments might

affect growth and development and in what ways, and how developmental trajectories might influence risks for developing diabetes, obesity, or heart disease decades after exposures are not easily understandable (Barnes et al., 2016; Winett, Wallack, Richardson, Boone-Heinonen, & Messer, 2016; Winett, Wulf, & Wallack, 2016). Furthermore, the DOHaD hypothesis confronts the dominant Western narrative about obesity and metabolic health (Monaghan, 2017; Warin, Turner, Moore, & Davies, 2008). Namely, Westerners most often intuit that individuals, especially adults, are responsible for and cause their own health-related behaviours and health outcomes (Marmot & Bell, 2019; Winett, Wallack, et al., 2016). The DOHaD hypothesis shifts responsibility for health outcomes away from individuals to the multiple layers of the environment in which we developed. These environmental layers include our mothers' bodies, our familial histories, our household economies, our communities, and the political structures in which our communities are embedded (Winett, Wallack, et al., 2016; Winett, Wulf, & Wallack, 2016). Whereas Health Canada's general pregnancy nutrition guidelines are compatible with a cognitively easy, traditional narrative, arguably, the DOHaD concepts are not.

Regardless of which proximate barriers prevent understanding of DOHaD ideas, these data indicate that pregnant people in Hamilton are well informed about what the general dietary and behavioural guidelines for pregnancy health are. But, they do not necessarily understand why they should eat in a particular way, or what the long-term consequences of their prenatal health might be for their children's health.

4.2 | Inequity in who has DOHaD knowledge

Our initial hypothesis that inequities in health literacy are likely to drive inequities in familiarity with DOHaD-related concepts offers a compelling explanation for our findings that DOHaD_{KNOWLEDGE} score varies with SEP and age. This is consistent with the handful of previous studies that have assessed DOHaD_{KNOWLEDGE} translation, which show that people living in lower SEP contexts tend to have lower health knowledge and literacy generally, have little self-efficacy with which to make use of health knowledge, and are more distrustful of health service provision (Barker et al., 2017; Sørensen et al., 2015). Fortunately, the available literature also suggests that these kinds of barriers can be overcome. DOHaD-related concepts have been successfully translated to several hard-to-reach, albeit selected, populations with low levels of health literacy (adolescents in New Zealand and the Cook Islands, adolescents in Uganda, and low SEP women in the United Kingdom; Baird et al., 2014; Bay et al., 2012, 2016a, Bay, Morton, & Vickers, 2016b, Bay, Vickers, Mora, Sloboda, & Morton, 2017a, Bay et al., 2017b; MacNab & Mukisa, 2017; Barker et al., 2017). In these selected cohorts, knowledge translation has required development of hands-on curricula and/or deep engagement and cultivation of meaningful relationships with community partners but is proving fruitful (Grace & Bay, 2011; MacNab & Mukisa, 2017).

The data reported here indicate that DOHaD_{KNOWLEDGE} score is negatively associated with respondents' parity. This is counter to our initial expectations, where we had assumed more pregnancy "experience" would lead to more knowledge. However, this negative association accords with some data from other populations indicating that experienced mothers are less likely to seek out information about prenatal health than first time mothers, relying instead on their own past experiences (Declercq, Sakala, Corry, & Applebaum, 2007). Equally, evidence from other populations suggests that health service providers reduce efforts to engage with pregnant people who they perceive to be "experienced" (Blondin & LoGiudice, 2018; Downs, Savage, & Rauff, 2014). As such, pregnant people who have previously given birth may have had and/or taken fewer opportunities to access and assimilate information about DOHaD.

4.3 | Why pregnancy diet quality might be independently associated with DOHaD knowledge

The significant independent association between pregnancy diet quality and an understanding of DOHaD concepts does not imply a causal relationship between DOHaD_{KNOWLEDGE} and diet quality score, but it might hint at one. It is well established that motivation is central to engaging in positive health behaviours like eating healthfully during pregnancy (Martin, Haskard-Zolnieriek, & DiMatteo, 2010), and health behaviours and health surveillance are more salient to pregnant people than to the general population (Phelan, 2010). Additionally, understanding the relevance of a set of recommendations may contribute to motivating people to follow them (Deci &

Ryan, 2002). However, because numerous other factors were also associated with variation in pregnancy diet quality such as age, parity (Table 4), and main source of household income (data not shown), these other factors likely constrain and/or mediate the relationship between diet quality and DOHaD_{KNOWLEDGE}. Nonetheless, if ongoing qualitative research in the study population provides evidence further supporting hypothesized links between deeper understanding, motivation, and eating behaviour, continuing efforts at DOHaD knowledge translation may pay off in improved diets during pregnancy and, ultimately, better health outcomes and reduced health inequities for mothers and babies.

4.4 | Limitations

We recognize that our study has some limitations that should be considered going forward. Specifically, the sample is biased towards higher SEP respondents (Figure 1). Despite efforts to target hard-to-reach, marginalized segments of Hamilton's population, our highest rates of response came from relatively educated, affluent pregnant Hamiltonians. Nonetheless, 24% of respondents did come from the lowest income bracket. Additionally, responses to other parts of the survey's sociodemographic section as well as our own field observations suggest that we were able to sample not only lower SEP Hamiltonians but also those who identify as Indigenous, as newcomers to Canada, and as racialized minorities, suggesting our analyses reflect much of the city's sociodemographic diversity. Moreover, although caution is necessary, we highlight that, in spite of the sample's noisiness, we found clear signals indicating that we should reject the null hypotheses of no relationships between DOHaD_{KNOWLEDGE} score, diet quality score, and sociodemographic factors.

We also recognize the need for validated instruments to measure DOHaD_{KNOWLEDGE} and Pregnancy Health Guideline_{KNOWLEDGE}. Specifically, both scales comprise predominantly positively phrased statements, and we should aim for an equal number of positively and negatively phrased items in each scale. Our team is thus working towards further developing and validating these scales. At the same time, though, the high internal reliability of the DOHaD_{KNOWLEDGE} scale suggests that a coherent mental construct is being effectively measured (Gliem & Gliem, 2003). We also note that, with the exception of recent work by Oyamada and colleagues in Japan (2018), we are the first to measure DOHaD_{KNOWLEDGE} using multiple items. Our method builds on previous work, which has used a single question/statement to assess familiarity with DOHaD (e.g., Bay et al., 2012, 2017b); it thus represents an important step towards better characterizing how people think about DOHaD-related ideas.

Lastly, as mentioned, earlier work emphasizes that effective DOHaD knowledge translation depends on deep understandings of the needs and priorities of communities and genuine efforts at relationship-building and mutual exchange (Baird et al., 2014; Barker et al., 2017; Bay et al., 2012; Lawrence et al., 2016; MacNab & Mukisa, 2017). Obtaining a deep understanding of pregnancy health and nutrition experiences solely through an anonymous questionnaire is

impossible. As such, the survey used here constitutes only one prong of a research strategy involving engagement with a wide variety of community stakeholders in Hamilton through focus group discussions, stakeholder meetings, and participant observation. In future work, we will situate survey responses in a tapestry of contextual information. Until then, these anonymous data highlight some gaps in respondents' understanding and therefore suggest where and how we might focus our attention when translating knowledge via stakeholder engagement.

5 | CONCLUSIONS

We investigated familiarity of pregnant respondents living in Hamilton, Canada, with concepts related to the DOHaD hypothesis. There were substantial inequities in access to and/or uptake of DOHaD concepts. Furthermore, evidence indicated that having more familiarity with DOHaD-related concepts is associated with relatively high-quality pregnancy diet, suggesting that inequities in DOHaD knowledge may be associated with inequities in a health-related behaviour during a window critical for the development and long-term health of children to be. These findings underscore the potential value of DOHaD knowledge translation efforts. Moreover, deep collaboration with community and political partners can build capacity for supporting pregnancy health and nutrition not only now, but for the generations to come.

ACKNOWLEDGMENTS

We acknowledge the financial support of the Canadian Institutes of Health Research (Grant 146333), the Canada Research Chairs program, the Canadian Foundation for Innovation, McMaster University, and the Women's College Hospital Xchange 15k Challenge funding scheme. Additionally, this research would not have been possible without the support of a number of people and organizations. In particular, we express our gratitude for the practical and logistical support we have received from our partners in Hamilton's Public Health Services and in Hamilton's Ontario Early Years Centres (OEYCs/EarlyON). We would especially like to highlight the integral roles of Jennifer Vickers-Manzin, Dr Vanessa Parlette, Lindsay Zalot, Jackie Cooper-Boyd, Gayle Reece, Michaela Servos, Kathy Pierce, Laura Wilson, and their staff and colleagues, who collectively run a variety of programmes (mainly the Hamilton Prenatal Nutrition Program, the Nurse-Family Partnerships program, and EarlyON drop-in sessions) aimed at improving maternal-child health, well-being, and nutrition in the city of Hamilton. Much of our study promotion was carried out through and with these organizations. We also thank other members of the M2B study team, especially the following research assistants and students: Debbie Kao, Sarah Oresnik, Dianna Williams, and Fei Fei Xia, along with members of the Sloboda Lab for thoughtful comments on earlier presentations of the data reported here. We further acknowledge the contributions of the editor and two anonymous reviewers,

whose comments improved the clarity and quality of this manuscript. Finally, our deepest thanks go to the 404 pregnant people of Hamilton who took time from their busy lives to respond carefully and often creatively to our questionnaire.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

CONTRIBUTIONS

All authors contributed to study design, study promotion/data collection, interpretation of results, and editing and approval of manuscript; LM collated and analysed data and drafted majority of manuscript text; TM, MB, and DS drafted portions of manuscript text.

ORCID

Luseadra McKerracher  <https://orcid.org/0000-0002-9687-6896>

REFERENCES

- Anon (2011). The time to prevent disease begins before conception. *Journal of the World Public Health Nutrition Association*, 2, 195–205.
- Baird, J., Cooper, C., Margetts, B. M., Barker, M., & Inskip, H. (2009). Changing health behaviour of young women from disadvantaged backgrounds: Evidence from systematic reviews. *Proceedings of the Nutrition Society*, 68(2), 195–204. <https://doi.org/10.1017/S0029665109001050>
- Baird, J., Jarman, M., Lawrence, W., Black, C., Davies, J., Tinati, T., ... Inskip, H. (2014). The effect of a behaviour change intervention on the diets and physical activity levels of women attending Sure Start Children's Centres: Results from a complex public health intervention. *BMJ Open*, 4, e005290–e005302. <https://doi.org/10.1136/bmjopen-2014-005290>
- Barker, D. J. P., & Osmond, C. (1986). Infant mortality, childhood nutrition, and ischaemic heart disease. *The Lancet*, 1986-1, 1077–1081.
- Barker, M. E., Baird, J., Tinati, T., Vogel, C., Strömmer, S., Rose, T., ... Lawrence, W. (2017). Translating developmental origins: Improving the health of women and their children using a sustainable approach to behaviour change. *Healthcare*, 5, 17–30. <https://doi.org/10.3390/healthcare5010017>
- Barnes, M. D., Heaton, T. L., Coates, M. C., & Packer, J. M. (2016). Inter-system implications of the Developmental Origins of Health and Disease: Advancing health promotion in the 21st century. *Healthcare*, 4, 1–17.
- Bay, J. L., Fohoko, F., La'Akulu, M., Leota, O., Pulotu, L., Tu'Ipulotu, S., ... Pouvalu, E. H. (2016a). Question in Tongan science classrooms: A pilot study to identify current practice, barriers, and facilitators. *Asia-Pacific Forum on Science Learning and Teaching*, 17(2), 1–25.
- Bay, J. L., Mora, H. A., Sloboda, D. M., & Morton, S. M. (2012). Adolescent understanding of DOHaD concepts: A school-based intervention to support knowledge translation and behaviour change. *Journal of Developmental Origins of Health and Disease*, 3, 469–482. <https://doi.org/10.1017/S2040174412000505>
- Bay, J. L., Morton, S. M., & Vickers, M. H. (2016b). Realizing the potential of adolescence to prevent transgenerational conditioning of non-communicable disease risk: Multi-sectoral design frameworks. *Healthcare*, 4(3), 39. <https://doi.org/10.3390/healthcare4030039>

- Bay, J. L., Vickers, M. H., Mora, H. A., Sloboda, D. M., & Morton, S. M. (2017a). Adolescents as agents of healthful change through scientific literacy development: A school-university partnership program in New Zealand. *International Journal of STEM Education*, 4, 15–35. <https://doi.org/10.1186/s40594-017-0077-0>
- Bay, J. L., Yaqona, D., Barrett-Watson, C., Tairea, K., Herrmann, U., & Vickers, M. H. (2017b). We learnt and now we are teaching our family. *Journal of Developmental Origins of Health and Disease*, 8(suppl1), s152–s152.
- Blondin, J. H., & LoGiudice, J. A. (2018). Pregnant women's knowledge and awareness of nutrition. *Applied Nursing Research*, 39, 167–174. <https://doi.org/10.1016/j.apnr.2017.11.020>
- Bornstein, M. H., Cote, L. R., Haynes, O. M., Hahn, C. S., & Park, Y. (2010). Parenting knowledge: Experiential and sociodemographic factors in European American mothers of young children. *Developmental Psychology*, 46(6), 1677.
- Chalmers, B., Dzakpasu, S., Heaman, M., & Kaczorowski, J. (2008). The Canadian maternity experiences survey: An overview of findings. *Journal of Obstetrics and Gynaecology Canada*, 30(3), 217–228. [https://doi.org/10.1016/S1701-2163\(16\)32758-X](https://doi.org/10.1016/S1701-2163(16)32758-X)
- Chinn, D. (2011). Critical health literacy: A review and critical analysis. *Social Science and Medicine*, 73, 60–67. <https://doi.org/10.1016/j.socscimed.2011.04.004>
- Cronbach, L. J., & Murphy, J. (1970). *Essentials of psychological testing*. New York, NY: Harper and Row.
- Dean, J. A., & Elliott, S. J. (2012). Prioritizing obesity in the city. *Journal of Urban Health*, 89, 196–213. <https://doi.org/10.1007/s11524-011-9620-3>
- Deci, E. L., & Ryan, R. M. (2002). *Handbook of self-determination research*. Rochester: University of Rochester Press.
- Declercq, E. R., Sakala, C., Corry, M. P., & Applebaum, S. (2007). Listening to mothers II: Report of the second national US survey of women's childbearing experiences. *The Journal of Perinatal Education*, 16(4), 9–14. <https://doi.org/10.1624/105812407X244769>
- DeLuca, P. F., Buist, S., & Johnston, N. (2012). The code red project: Engaging communities in health system change in Hamilton, Canada. *Social Indicators Research*, 108, 317–327. <https://doi.org/10.1007/s11205-012-0068-y>
- Denny, C. C., & Grady, C. (2007). Clinical research with economically disadvantaged populations. *Journal of Medical Ethics*, 33(7), 382–385. <https://doi.org/10.1136/jme.2006.017681>
- Downs, D. S., Savage, J. S., & Rauff, E. L. (2014). Falling short of guidelines? Nutrition and weight gain knowledge in pregnancy. *Journal of Women's Health Care*, 3, 184–190.
- Durant, R. W., Davis, R. B., George, D. M. M. S., Williams, I. C., Blumenthal, C., & Corbie-Smith, G. M. (2007). Participation in research studies: Factors associated with failing to meet minority recruitment goals. *Annals of Epidemiology*, 17(8), 634–642. <https://doi.org/10.1016/j.annepidem.2007.02.003>
- Fleming, T. P., Watkins, A. J., Velazquez, M. A., Mathers, J. C., Prentice, A. M., Stephenson, J., ... Godfrey, K. M. (2018). Origins of lifetime health around the time of conception: Causes and consequences. *The Lancet*, 391, 1842–1852. [https://doi.org/10.1016/S0140-6736\(18\)30312-X](https://doi.org/10.1016/S0140-6736(18)30312-X)
- Garmendia, M. L., Corvalan, C., & Uauy, R. (2014). Assessing the public health impact of Developmental Origins of Health and Disease interventions. *Annals of Nutrition and Metabolism*, 64, 226–230. <https://doi.org/10.1159/000365024>
- Gliem, J. A., & Gliem, R. R. (2003). Calculating, interpreting, and reporting Cronbach's alpha reliability coefficient for Likert-type scales. Midwest Research-to-Practice Conference in Adult, Continuing, and Community Education. Columbus: Midwest Research-to-Practice Conference. 1–7.
- Glouberman, S., & Millar, J. (2003). Evolution of the determinants of health, health policy, and health information systems in Canada. *American Journal of Public Health*, 93, 388–392. <https://doi.org/10.2105/AJPH.93.3.388>
- Godfrey, K. M., Reynolds, R. M., Prescott, S. L., Nyirenda, M., Jaddoe, V. W. V., Eriksson, J. G., & Broekman, B. F. P. (2017). Influence of maternal obesity on the long-term health of offspring. *The Lancet: Diabetes and Endocrinology*, 5, 53–64. [https://doi.org/10.1016/S2213-8587\(16\)30107-3](https://doi.org/10.1016/S2213-8587(16)30107-3)
- Grace, B., & Bay, J. (2011). Developing a pedagogy to support science for health literacy. *Asia-Pacific Forum on Science Learning and Teaching*, 12(2), 1–26.
- Haertsch, M., Campbell, E., & Sanson-Fisher, R. (1999). What is recommended for healthy women during pregnancy? A comparison of seven prenatal clinical practice guideline documents. *Birth*, 26(1), 24–30. <https://doi.org/10.1046/j.1523-536x.1999.00024.x>
- Hanson, M. A., & Gluckman, P. D. (2015). Developmental Origins of Health and Disease: Global public health implications. *Best Practice & Research. Clinical Obstetrics & Gynaecology*, 29(1), 24–31. <https://doi.org/10.1016/j.bpobgyn.2014.06.007>
- Health Canada. (2014). Prenatal nutrition guidelines for health professionals: Gestational weight gain. Guidelines, Ottawa: Health Canada.
- Jenkins, J., & Keating, D. (1999). *Risk and resilience in six and ten year old children*. Toronto, ON: University of Toronto Press.
- Kalich, A., Heinemann, L., & Ghahari, S. (2016). A scoping review of immigrant experience of health care access barriers in Canada. *Journal of Immigrant and Minority Health*, 18, 697–709. <https://doi.org/10.1007/s10903-015-0237-6>
- Latham, J., & Moffat, T. (2007). Determinants of variation in food cost and availability in two socio-economically contrasting neighbourhoods of Hamilton, Ontario, Canada. *Health and Place*, 13, 273–287. <https://doi.org/10.1016/j.healthplace.2006.01.006>
- Lawrence, W., Black, C., Tinati, T., Cradock, S., Begum, R., Jarman, M., ... Cooper, C. (2016). Making every contact count: Evaluation of the impact of an intervention to train health and social practitioners in skills to support behaviour change. *Journal of Health Psychology*, 12, 138–151.
- Low, F. M., Gluckman, P. D., & Hanson, M. A. (2018). A lifecourse approach to public health: Why early life matters. In M. van den Bosch, & W. Bird (Eds.), *Oxford textbook of nature in public health: The role of nature in improving the health of a population*. Oxford: Oxford University Press.
- MacNab, A. J., & Mukisa, R. (2017). Priorities for African youth engaging in DOHaD. *Journal of Developmental Origins of Health and Disease*, 9, 15–19.
- Marmot, M., & Bell, R. (2019). Social determinants and non-communicable diseases: A time for integrated action. *British Medical Journal*, 364, 1251.
- Martin, L. R., Haskard-Zolnieriek, K. B., & DiMatteo, M. R. (2010). *Health behavior change and treatment adherence: Evidence-based guidelines for improving healthcare*. Oxford: Oxford University Press.
- McKerracher, L., Moffat, T., Barker, M. E., Williams, D., & Sloboda, D. M. (2019). Translating the Developmental Origins of Health and Disease concept to improve the nutritional environment for our next generations: A call for a reflexive, positive, multi-level approach. *Journal of Developmental Origins of Health and Disease* 10(4), 420–428.
- McMullan, R. L., Fuller, N. R., Celermajer, I. D., Caterson, D. S., Gordon, A., & Hesse, C. (2017). Developmental Origins of Health and Disease: Who knows? Who cares? *Journal of Paediatrics and Child Health*, 53, 613–613. <https://doi.org/10.1111/jpc.13557>
- Moffat, T., & Galloway, T. (2008). Food consumption patterns in elementary school children. *Canadian Journal of Dietetic Practice and Research*, 69, 152–154. <https://doi.org/10.3148/69.3.2008.152>
- Monaghan, L. F. (2017). Re-framing weight-related stigma: From spoiled identity to macro-social structures. *Social Theory and Health*, 15(2), 182–205. <https://doi.org/10.1057/s41285-016-0022-1>

- Navaneelan, T., & Janz T. Adjusting the scales: Obesity in the Canadian after correcting for respondent bias. 2014.
- Nutbeam, D. (2008). The evolving concept of health literacy. *Social Science and Medicine*, 67(12), 2072–2078. <https://doi.org/10.1016/j.socscimed.2008.09.050>
- Oh, S. S., Galanter, J., Thakur, N., Pino-Yanes, M., Barcelo, N. E., White, M. J., ... Burchard, E. G. (2015). Diversity in clinical and biomedical research: A promise yet to be fulfilled. *PLoS Medicine*, 12(12), e1001918. <https://doi.org/10.1371/journal.pmed.1001918>
- Oyamada, M., Lim, A., Dixon, R., Wall, C., & Bay, J. (2018). Development of understanding of DOHaD concepts in students during undergraduate health professional programs in Japan and New Zealand. *Journal of Developmental Origins of Health and Disease*, 9(3), 253–259. <https://doi.org/10.1017/S2040174418000338>
- Perreault, M., Mikail, S., & Atkinson, S. A. (2016). New Health Canada nutrition recommendations for infants birth to 24 months address the importance of early nutrition. *Nutrition Today*, 51(4), 186–190.
- Phelan, S. (2010). Pregnancy: A 'teachable moment' for pregnancy weight control and obesity prevention. *American Journal of Obstetrics and Gynecology*, 202, 135e1–135e8. <https://doi.org/10.1016/j.ajog.2009.06.008>
- Pinheiro, J, Bates, D., DebRoy, S., Sarkar, S., & R Core Team. (2018). nlme: Linear and nonlinear mixed effects models
- R Core Development Team. (2017). R: A language and environment for statistical computing.
- Revelle, W. (2018). psych: Procedures for personality and psychological research.
- Rifas-Shiman, S. L., Willett, W. C., Lobb, R., Kotch, J., Dart, C., & Gillman, M. W. (2001). Primescreen, a brief dietary screen tool: Reproducibility and comparability with both a larger food frequency questionnaire and biomarkers. *Public Health Nutrition*, 4(2), 249–254. <https://doi.org/10.1079/PHN2000061>
- Simich, L. (2009). Health literacy and immigrant populations. Policy Brief, Public Health Agency of Canada, Ottawa: Public Health Agency of Canada and Metropolis of Canada.
- Smith, S. K., Dixon, A., Trevena, L., Nutbeam, D., & McCaffery, K. J. (2009). Exploring patient involvement in healthcare decision making across different education and functional health literacy groups. *Social Science and Medicine*, 69, 1805–1812. <https://doi.org/10.1016/j.socscimed.2009.09.056>
- Sørensen, K., Van den Broucke, S., Fullam, J., Doyle, G., Pelikan, J., Slonska, Z., & Brand, H. (2012). Health literacy and public health: A systematic review and integration of definitions and models. *BMC Public Health*, 12, 80–93. <https://doi.org/10.1186/1471-2458-12-80>
- Sørensen, K., Pelikan, J. M., Röthlin, F., Ganahl, K., Slonska, Z., Doyle, G., ... Falcon, M. (2015). Health literacy in Europe: Comparative results of the European health literacy survey (HLS-EU). *European Journal of Public Health*, 25(6), 1053–1058.
- Stephenson, J., Heslehurst, N., Hall, J., Schoenaker, D. A. J. M., Hutchinson, J., Cade, J. E., ... Mishra, G. D. (2018). Before the beginning: Nutrition and lifestyle in the preconception period and its importance for future health. *The Lancet*, 391, 1830–1841. [https://doi.org/10.1016/S0140-6736\(18\)30311-8](https://doi.org/10.1016/S0140-6736(18)30311-8)
- Victora, C. G., Bahl, R., Barros, A. J. D., França, G. V. A., Horton, S., Krasevec, J., ... Rollins, N. C. (2016). Breastfeeding in the 21st century: Epidemiology, mechanisms, and lifelong effect. *The Lancet*, 387(10017), 475–490. [https://doi.org/10.1016/S0140-6736\(15\)01024-7](https://doi.org/10.1016/S0140-6736(15)01024-7)
- Warin, M., Turner, K., Moore, V., & Davies, M. (2008). Bodies, mothers and identities: Rethinking obesity and the BMI. *Sociology of Health & Illness*, 30(1), 97–111. <https://doi.org/10.1111/j.1467-9566.2007.01029.x>
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. New York, NY: Springer-Verlag. <https://doi.org/10.1007/978-3-319-24277-4>
- Willems, S., De Maesschalck, S., Deveugele, M., Derese, A., & De Maeseneer, J. (2005). Socio-economic status of the patient and doctor-patient communication: Does it make a difference? *Patient Education and Counseling*, 56, 139–146. <https://doi.org/10.1016/j.pec.2004.02.011>
- Winett, L., Wallack, L., Richardson, D., Boone-Heinonen, J., & Messer, L. (2016). A framework to address challenges in communicating the Developmental Origins of Health and Disease. *Current Environmental Health Report*, 3, 169–177. <https://doi.org/10.1007/s40572-016-0102-3>
- Winett, L., Wulf, A. B., & Wallack, L. (2016). Framing strategies to avoid mother blame in communicating the origins of chronic disease. *American Journal of Public Health*, 106, 1369–1373. <https://doi.org/10.2105/AJPH.2016.303239>
- Woods-Townsend, K., Leat, H., Bay, J., Bagust, L., Davey, H., Lovelock, D., ... Hanson, M. (2018). LifeLab Southampton: A programme to engage adolescents with DOHaD concepts as a tool for increasing health literacy in teenagers—A pilot cluster-randomised control trial. *Journal of Developmental Origins of Health and Disease*, 9(5), 475–480.
- World Health Organization. (2017). Non-communicable diseases progress monitor. World Health Organization.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: McKerracher L, Moffat T, Barker M, et al. Knowledge about the Developmental Origins of Health and Disease is independently associated with variation in diet quality during pregnancy. *Matern Child Nutr.* 2020;16:e12891. <https://doi.org/10.1111/mcn.12891>