

RESEARCH ARTICLE

Fencing farm dams to exclude livestock halves methane emissions and improves water quality

Martino E. Malerba¹  | David B. Lindenmayer²  | Ben C. Scheele²  |
Pawel Waryszak¹  | I. Noyan Yilmaz¹  | Lukas Schuster³  | Peter I. Macreadie¹ 

¹Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Melbourne, Victoria, Australia

²Sustainable Farms, Fenner School of Environment and Society, The Australian National University, Canberra, Australian Capital Territory, Australia

³Centre of Geometric Biology, School of Biological Sciences, Monash University, Melbourne, Victoria, Australia

Correspondence

Martino E. Malerba, Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Melbourne, VIC 3125, Australia.

Email: m.malerba@deakin.edu.au

Funding information

Alfred Deakin Postgraduate Research Fellowship; Australian Research Council, Grant/Award Number: DE220100752;

Abstract

Agricultural practices have created tens of millions of small artificial water bodies (“farm dams” or “agricultural ponds”) to provide water for domestic livestock worldwide. Among freshwater ecosystems, farm dams have some of the highest greenhouse gas (GHG) emissions per m² due to fertilizer and manure run-off boosting methane production—an extremely potent GHG. However, management strategies to mitigate the substantial emissions from millions of farm dams remain unexplored. We tested the hypothesis that installing fences to exclude livestock could reduce nutrients, improve water quality, and lower aquatic GHG emissions. We established a large-scale experiment spanning 400 km across south-eastern Australia where we compared unfenced ($N = 33$) and fenced farm dams ($N = 31$) within 17 livestock farms. Fenced farm dams recorded 32% less dissolved nitrogen, 39% less dissolved phosphorus, 22% more dissolved oxygen, and produced 56% less diffusive methane emissions than unfenced dams. We found no effect of farm dam management on diffusive carbon dioxide emissions and on the organic carbon in the soil. Dissolved oxygen was the most important variable explaining changes in carbon fluxes across dams, whereby doubling dissolved oxygen from 5 to 10 mg L⁻¹ led to a 74% decrease in methane fluxes, a 124% decrease in carbon dioxide fluxes, and a 96% decrease in CO₂-eq (CH₄ + CO₂) fluxes. Dams with very high dissolved oxygen (>10 mg L⁻¹) showed a switch from positive to negative CO₂-eq. (CO₂ + CH₄) fluxes (i.e., negative radiative balance), indicating a positive contribution to reduce atmospheric warming. Our results demonstrate that simple management actions can dramatically improve water quality and decrease methane emissions while contributing to more productive and sustainable farming.

KEYWORDS

agricultural reservoirs, dugouts, eutrophication, impoundment, inland water, sustainable farms

1 | INTRODUCTION

Global methane emissions are rising rapidly, nearly tripling from ca. 700 ppb in pre-industrial times to 1900 ppb today (Conrad, 2009;

Dlugokencky, 2022). The accumulation of artificial water bodies has contributed to the growth in atmospheric methane, with aquatic ecosystems now accounting for half of natural and anthropogenic methane emissions (Rosentreter et al., 2021). With farm dams estimated to

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Global Change Biology* published by John Wiley & Sons Ltd.

cover a surface area $>75,000\text{km}^2$ globally (Downing et al., 2006), these artificial systems are now a key part of aquatic ecosystems globally (Malerba et al., 2021; Swartz & Miller, 2021). Therefore, it is likely that farm dams are an important contributor to global carbon cycles—even though this link is often overlooked in national and global carbon inventories. Indeed, the Intergovernmental Panel on Climate Change (IPCC) recently revised their guidelines to promote the inclusion of agricultural ponds in national GHG inventories and tackle this form of anthropogenic carbon emission (IPCC, 2019).

Farm dams (or agricultural ponds) are small, human-made freshwater bodies created for the purpose of storing water for livestock or crop irrigation (Malerba et al., 2022). These systems have some of the highest greenhouse gas (GHG) emissions per m^2 among freshwater ecosystems (Grinham et al., 2018; Ollivier et al., 2018, 2019) due to their much higher nitrogen and phosphorus concentrations than natural ponds (Westgate et al., 2022), creating the perfect conditions for methanogenesis and GHG emissions (Li et al., 2021; Panneer Selvam et al., 2014; Peacock et al., 2021). Importantly, eutrophication appears to have a disproportionate effect on farm dams. That is, a 25% increase in nitrate concentration was observed to double the CO_2 -equivalent carbon flux per m^2 of a farm dam (Ollivier et al., 2018). Hence, understanding how to reduce the emissions of millions of farm dams worldwide has the potential to make a substantial difference in mitigating climate change. Yet, there is no evidence of the effects of management practices on reducing these emissions.

Using fences to exclude livestock from farm dams improves water quality by reducing direct depositions of nutrient-rich manure and urine into the water (Westgate et al., 2022). In addition, fencing a farm dam avoids hooved livestock (ungulates) disturbing soils and promotes higher vegetation cover around the dam, acting as a filter to reduce dissolved nutrients (i.e., “phytoremediation”; reviewed in Pilon-Smits, 2005). A recent study showed that partially or fully fenced farm dams have higher vegetation cover, higher water quality (i.e., lower nutrients, turbidity, and fecal coliforms), and higher macroinvertebrate richness and abundance than unfenced farm dams (Westgate et al., 2022). Moreover, fencing farm dams is often cost-effective, with the benefits for livestock health and weight gain from higher water quality often exceeding the costs of this management intervention (Dobes et al., 2021).

In summary: (1) nutrient pollution drives high GHG emissions from farm dams; (2) excluding livestock from accessing farm dams favors vegetation growth and improves water quality; and (3) higher water quality provides benefits to livestock health, biodiversity, and aesthetic value. Based on these premises, installing fences could reduce aquatic GHG emissions from farm dams while improving agricultural productivity and biodiversity. Previous studies have already shown that excluding livestock and reducing grazing intensity can reduce methane emissions and enhance carbon sequestration and storage of freshwater wetlands (Limpert et al., 2021; Oates et al., 2008; Watkins et al., 2017). Yet, the effects of installing fences (or any other management intervention) on farm dam GHG production remain untested. Similarly, there is little evidence of the benefits of farm dam fencing on water quality (Westgate et al., 2022). Hence, we here addressed two key questions:

1. What are the effects of fencing farm dams on water quality (i.e., total dissolved nitrogen, total dissolved phosphorus, and dissolved oxygen), soil organic carbon, and GHG fluxes of methane and carbon dioxide?
2. What are the mechanisms linking farm dam management to aquatic GHG fluxes?

To answer these questions, we completed a cross-sectional field-based study comparing the effects of fencing farm dams on their water quality and carbon footprint. We surveyed 64 farm dams across 17 farming properties. At each property, we compared farm dams under two management regimes: “unfenced dams” where livestock have free access to the water, and “fenced dams” where water access has been controlled for at least 2 years using either full fencing or partial fencing (with a hardened livestock access point). We predicted that fencing a farm dam would reduce dissolved nutrients, increase dissolved oxygen, and lower GHG emissions. Testing these processes contributes to identifying novel GHG abatement methods to reduce the carbon footprint of farming practices.

2 | METHODS

2.1 | Study area and experimental design

In April 2021, we sampled farm dams across 400 km of the Australian South West Slopes bioregion in south-eastern New South Wales. The study region has a warm temperate climate, with hot dry summers and cool humid winters (the largest city of Albury has an annual mean temperature of 22°C and annual rainfall of 691 mm). Most of the area is dedicated to livestock grazing (especially beef cattle and sheep) and dryland cropping (mainly cereals and oilseed). We surveyed 64 farm dams located in pastures on 17 farming properties. Within each property, we established two experimental treatments: “unfenced” farm dams and “fenced” farm dams. For each experimental treatment within a farming property, we measured between 1 and 5 dams (depending on availability) on the same day. Unfenced farm dams ($N = 33$) received no management intervention to improve their ecological condition. Fenced farm dams ($N = 31$) were either entirely fenced (with a pump delivering water into drinking troughs) or partly fenced (providing water access through a hardened access point) for at least 2 years prior to sampling. We avoided small ($<200\text{m}^2$) farm dams because they were too ephemeral. We measured farm dam areas by tracing the most recent satellite images on Google Earth Pro (version 7.3.4).

2.2 | Aquatic greenhouse gas emissions

We measured diffusive emissions of methane and carbon dioxide at each farm dam using the methods described in Ollivier et al. (2018, 2019). Briefly, a white plastic floating chamber (0.021m^3 volume and 0.14m^2 surface area) was sealed and connected to an Ultraportable Greenhouse Gas Analyzer (UGGA, Los Gatos Research, Model 915-0011) through

two tubes (influx and outflux) on the chamber roof to create a closed circuit. We sampled methane and carbon dioxide (ppm) at 1-second intervals for ca. 5 min (ca. 300 data points per sample). We measured each farm dam three times from different locations along the shore ensuring the starting concentration matched atmospheric levels.

Floating chambers can measure constant fluxes (diffusion) and stochastic releases of gas bubbles (ebullition). Here we focused only on diffusive fluxes. To do so, we excluded any trajectories showing sudden increases in gas concentration due to a gas bubble being released inside the floating chamber. We estimated the linear rate of change of diffusive gas flux from the water surface to the atmosphere (F ; $\text{mg m}^{-2} \text{d}^{-1}$) as:

$$F = \frac{\text{slope} \times \text{volume} \times F_1 \times F_2}{F_3 \times \text{surface}} \quad (1)$$

where *slope* is the linear rate of change in gas concentrations over time within the chamber (ppms^{-1}), *volume* is the chamber volume (0.021 m^3), F_1 is the conversion factor from ppm to $\mu\text{g m}^{-3}$ for methane (655.47), F_2 is the conversion factor from minutes to day (86,400), F_3 is the conversion factor from μg to mg (1000), *surface* is the surface area of the chamber (0.14 m^2 ; Lambert & Fr chet te, 2005). We retained all diffusive rates without applying any filtering method (e.g., R^2 threshold).

2.3 | Sediment carbon stocks

At each dam, we collected two cores (45 mm diameter, 50 mm deep, 79.52 cm^3 volume) from the edge of the pond within the water (wet sediments). We preserved the cores in a freezer until returning to the laboratory. We dried all cores at 60°C until there was no more weight loss (approx. a week) and measured their dry weights. Finally, we ground the cores and determined the organic carbon content by analyzing 10 mg of each sample using a EuroVector MicroElemental CN Analyser (see Gulliver et al., 2020 for details). We quantified each sample's C:N ratio using Acetanilide as standards (71.09% C, 0.5–1 mg input mass; $R^2 > 0.98$). The carbon density of each core was the product of dry biomass density (g cm^{-3}) and carbon content (i.e., % C/100) in units of tons of carbon per hectare (t C ha^{-1}).

2.4 | Water quality and nutrient analysis

At each site, we measured dissolved oxygen ($\text{mg O}_2 \text{ L}^{-1}$), conductivity ($\mu\text{S cm}^{-1}$), and water temperature ($^\circ\text{C}$) using a Hach HQ30D portable Multi Meter. We also filtered 50 ml of water from each farm dam using syringe filters with Filtech 483 Glass fiber filter paper (1.10 μm retention, 25 mm diameter). We froze all filtrated water samples immediately after collection and sent them to ALS Environmental (alsglabal.com, Everton Park QLD 4053 Australia) to analyze total nitrogen following APHA 4500- N_{org} / 4500- NO_3^- (method EK062G; mg N L^{-1}) and total phosphorus following APHA 4500-P (method EK067G; mg P L^{-1}). All analyses followed standard protocols and included quality controls. Finally, we took three pH measurements at each dam using the YSI ProDSS Multiparameter Digital Water Quality Meter (Xylem

Analytics, Yellow Springs, OH 45387 USA), taking measurements at 1.5 m from the water's edge and at 20 cm depth. We rinsed the sensors with demineralized water between samples and sites and always calibrated probes before use.

2.5 | Statistical analyses

First, we used individual linear mixed-effects models to evaluate whether the management regime (categorical variable, either "fenced" or "unfenced") affected total dissolved nitrogen ($\log_{10} \text{ mg N L}^{-1}$), total dissolved phosphorus ($\log_{10} \text{ mg P L}^{-1}$), dissolved oxygen (mg L^{-1}), organic carbon stock ($\log_{10} \text{ t C ha}^{-1}$), and rates of methane emissions ($\log_{10} \text{ mg m}^{-2} \text{ d}^{-1} + 2$), carbon dioxide emissions ($\log_{10} \text{ g m}^{-2} \text{ d}^{-1} + 1.8$), and CO_2 -equivalent emissions (carbon dioxide + methane; $\log_{10} \text{ g m}^{-2} \text{ d}^{-1} + 1.8$). We added two units to methane emissions and 1.8 units to carbon dioxide and CO_2 -eq. emissions to avoid negative values when applying the \log_{10} transformation. We calculated CO_2 -equivalent units by combining methane and carbon dioxide fluxes using the 20-year Sustained-Flux Global Warming Potential (SGWP) metric from Neubauer and Megonigal (2015), where 1 Kg of CH_4 traps as much infrared radiation as 96 Kg of CO_2 . The SGWP calculates the decay rate assuming a sustained gas flux rate over time, and this approach is more realistic for farm dams than the one-time pulse assumed in the Global Warming Potential metric. We did not correct the p-values for multiple statistical testing, yet we ensured that reducing the risk of type I error by adopting more conservative thresholds for statistical significance using the false discovery rate (Benjamini & Hochberg, 1995) did not change any of our conclusions.

Second, we used three linear mixed-effects models to quantify the statistical association of each environmental variable with fluxes of carbon dioxide, methane, and CO_2 -equivalent (carbon dioxide + methane) of a farm dam. In the models, the independent variables were farm dam surface area ($\log_{10} \text{ m}^2$), dissolved oxygen ($\log_{10} \text{ mg L}^{-1}$), pH, conductivity ($\log_{10} \mu\text{S cm}^{-1}$), water temperature ($^\circ\text{C}$), total dissolved nitrogen ($\log_{10} \text{ mg N L}^{-1}$), total dissolved phosphorus ($\log_{10} \text{ mg P L}^{-1}$), and organic carbon stock ($\log_{10} \text{ t C ha}^{-1}$). The initial fully parameterized model included all main effects and a two-way interaction term to account for the potential interplay between total nitrogen and total phosphorus. To avoid bias from multicollinearity between main effects, we ensured a cut-off value of five for the maximum variance inflation factor (VIF) in the model, as recommended by Zuur et al. (2009). As a result, pH and dissolved oxygen could not be included together in the models because they are highly correlated ($r = 0.72$ and $\text{VIF} > 5$). Therefore, we used only dissolved oxygen in the mixed-effects models as this variable is associated with fluxes of both carbon dioxide and methane (whereas pH is only associated with carbon dioxide). Finally, we quantified the importance of each statistically significant explanatory variable by calculating its contribution to the total model prediction power using a permutation approach (Fisher et al., 2019; Niittynen & Luoto, 2018; Virkkala et al., 2021). This analysis consisted of three steps. First, we extracted the predictions from the best-fitting model ($\text{Predictions}_{\text{original}}$). Second, we

created simulated datasets using random permutations of each statistically significant explanatory variable to remove its explanatory power. Third, we re-fitted the model to each simulated dataset, computed model predictions, and quantified the Pearson correlation coefficient between the predictions of the original model ($Predictions_{original}$) and the predictions with the explanatory variable being permuted ($Predictions_{shuffled,v}$), as:

$$Importance_v = 1 - cor(Predictions_{original} - Predictions_{shuffled,v}) \quad (2)$$

Values close to -1 or 1 indicate greater importance of the shuffled variable for the model's explanatory power. We repeated this process 100 times for each variable to calculate the average importance and 95% confidence intervals.

We centered and scaled all variables before fitting the linear mixed-effects models. We also added a random intercept to account for the experimental block design where each of the 17 farming properties contained one or more fenced and unfenced dams. To analyze repeated flux measurements from the same pond, we added a nested random intercept of site within farming property. When standardized residuals showed unequal variances or a systematic trend, we included treatment-specific variance coefficients (function `varIdent`) or other variance functions (functions `varExp` or `varPower`) in the model. We identified the best-fitting model using Akaike information criteria corrected for small sample sizes (AICc; Burnham & Anderson, 2004). We used standard diagnostics to ensure normality, homoscedasticity, and the absence of influential points or outliers.




FIGURE 1 Effects of farm dam fencing on (a) methane fluxes, (b) carbon dioxide fluxes, (c) CO_2 -eq (methane + carbon dioxide) fluxes, and (d) organic carbon in the soil. Black point ranges represent the means \pm 95% confidence intervals from the best-fitting linear models. Grey points are the raw data. All statistics are calculated on a sample size of 64 farm dams across 17 farming properties. We reported percentage changes only on statistically significant effects (see Table S1 for test statistics).

We used the statistical software R version 4.0.3 (R Core Team, 2020) with the packages nlme (Pinheiro et al., 2020) and effects (Fox & Weisberg, 2018, 2019) for the statistical analyses, and dplyr (Wickham et al., 2018), plyr (Wickham, 2011), and ggplot2 (Wickham, 2009) for data manipulation and plotting.

3 | RESULTS

3.1 | Effects of fencing farm dams on greenhouse gas emissions and organic carbon stocks

On average, methane emissions from fenced farm dams ($3.5 \text{ mg m}^{-2} \text{ d}^{-1}$) were 56% lower than unfenced farm dams ($8.05 \text{ mg m}^{-2} \text{ d}^{-1}$; Figure 1a). Conversely, we found no significant difference for carbon dioxide fluxes ($p = .2$; Figure 1b) or for CO_2 -eq fluxes ($p = .08$; Figure 1c). Finally, there was no effect of fencing on the organic carbon stock in the sediments of the farm dams ($p = .42$; Figure 1d). See Table 1 for summary statistics and Table S1 for statistical scores.

3.2 | Effects of fencing farm dams on water quality

Fenced farm dams recorded higher water quality than unfenced ones across all parameters measured here. Specifically, water from fenced farm dams had on average 32% less total dissolved nitrogen (from 2.4 to 1.6 mg L^{-1} ; Figure 2a), 39% less total dissolved phosphorus (from 0.078 to 0.047 mg L^{-1} ; Figure 2b), and 22% more dissolved oxygen than unfenced dams (from 6.32 to 7.74 mg L^{-1} ; Figure 2c). We found no difference in the water temperature (Figure 2d) and water

pH (data not shown) of fenced and unfenced farm dams (see Table 1 for summary statistics and Table S1 for statistical scores).

3.3 | Drivers of greenhouse gas fluxes

Overall, most relationships between greenhouse gas fluxes and environmental variables show a high degree of variability. Yet, the methane flux of a farm dam was statistically associated with dissolved oxygen (Figure 3a), sediment organic carbon stocks (Figure 3b), total dissolved nitrogen (Figure 3c), and total dissolved phosphorus (Figure 3d). In contrast, the carbon dioxide flux of a farm dam only showed a negative association with dissolved oxygen (Figure 3f). The total carbon flux of a farm dam, calculated as CO_2 -eq (methane + carbon dioxide) fluxes, showed statistically significant associations with dissolved oxygen (Figure 3k), sediment organic carbon stocks (Figure 3l), and total dissolved nitrogen (Figure 3m). Conversely, farm dam area, conductivity, and a two-way interaction between dissolved nitrogen and dissolved phosphorus were systematically excluded from the best-fitting models following Akaike information criteria.

Dissolved oxygen was the most important variable for explaining all three greenhouse gas fluxes (see Table S3 for importance scores). Specifically, doubling dissolved oxygen from 5 to 10 mg L^{-1} corresponded to a 74% decrease in methane fluxes (from 6.92 to $1.8 \text{ mg CH}_4 \text{ m}^{-2} \text{ d}^{-1}$; Figure 3a), a 124% decrease in carbon dioxide fluxes (from 2.27 to $-0.56 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$; Figure 3f), and a 96% decrease in CO_2 -eq fluxes (from 3.77 to $0.13 \text{ g CO}_2\text{-eq m}^{-2} \text{ d}^{-1}$; Figure 3k). Farm dams with dissolved oxygen levels higher than ca. 10 mg L^{-1} showed a switch from positive to negative CO_2 -eq fluxes (i.e., negative radiative balance; Figure 4).

TABLE 1 Summary of farm dam properties in this study. Water volume was estimated using the model in Figure 1 of Malerba et al. (2021): $\text{Water Volume} = -3.593 + 1.237 \times \text{Water Area}$. Water depth was estimated using the formula $(\text{Water Volume} \times 1000) / (\text{Water Area} \times 0.4)$ (Agriculture Victoria, 2022)

Variable	Unit	Rep	Min	Mean	Median	Max
Area	m^2	64	227	1978	886	20,796
Water volume (est.)	ML	64	0.21	3.05	1.13	56
Water depth (est.)	m	64	2.31	3.86	3.19	6.73
Longitude		64	146.79	147.63	147.19	149.45
Latitude		64	-36.10	-35.32	-35.86	-33.51
Total Nitrogen	mg N L^{-1}	64	0.40	2.68	2.10	9.20
Total Phosphorus	mg P L^{-1}	64	0.01	0.12	0.06	0.80
Dissolved Oxygen	$\text{mg O}_2 \text{ L}^{-1}$	64	3.16	6.94	6.52	17.60
Water Temperature	$^{\circ}\text{C}$	64	12.83	16.20	15.57	22.13
pH		63	6.50	7.79	7.71	9.52
Conductivity	$\mu\text{S cm}^{-1}$	64	11.89	294.3	227.5	1647
CH_4 diffusion	$\text{g m}^{-2} \text{ d}^{-1}$	63	0.0001	0.0151	0.0034	0.1639
CO_2 diffusion	$\text{g m}^{-2} \text{ d}^{-1}$	64	-1.6995	1.4897	0.7887	13.9746
CO_2 -eq diffusion	$\text{g m}^{-2} \text{ d}^{-1}$	63	-1.2161	2.9364	1.5619	21.3560
Sediment organic C stock	t C ha^{-1}	63	0.56	6.26	4.59	28.84




FIGURE 2 Effects of farm dam fencing on (a) total nitrogen, (b) total phosphorus, (c) dissolved oxygen, and (d) temperature. All statistics are calculated on a sample size of 64 farm dams across 17 farming properties. We reported percentage changes only on statistically significant effects (see Table S1 for test statistics).

Changes in both dissolved oxygen and carbon dioxide were pH related (Figure 5). Dissolved oxygen was positively correlated with the pH ($r = 0.72$; Figure 5a) and negatively correlated with the carbon dioxide flux ($r = -0.82$; Figure 5b), while carbon dioxide flux was negatively correlated with pH ($r = -0.76$; Figure 5c). Conversely, we found no significant correlation between pH and methane fluxes ($p = .39$; data not shown).

4 | DISCUSSION

Farm dams are common in many rural landscapes worldwide and make important contributions to carbon cycles and greenhouse gas (GHG) emissions (Grinham et al., 2018; Ollivier et al., 2018; Peacock et al., 2021). We discovered that simple management practices, such

as fencing off livestock from farm dams, increased water quality and dramatically lowered methane emissions. Fenced farm dams were characterized by 32% less dissolved nitrogen, 39% less dissolved phosphorus, 22% more dissolved oxygen, and 56% lower methane emissions than unfenced dams. Dissolved oxygen was the most important variable explaining changes in carbon fluxes across dams, whereby doubling dissolved oxygen from 5 to 10 mgL⁻¹ led to a 74% decrease in methane fluxes, a 124% decrease in carbon dioxide fluxes, and a 96% decrease in CO₂-eq (CH₄ + CO₂) fluxes. Moreover, farm dams with very high oxygen levels (>10 mgL⁻¹) exhibited a switch from positive to negative CO₂-eq fluxes. Finally, we found a strong negative correlation between the pH of the water and both the dissolved oxygen and fluxes of carbon dioxide.

We found that fencing farm dams, on average, more than halves diffusive methane emissions to 3.56 mg CH₄ m⁻² d⁻¹ compared to 8.16 mg




FIGURE 3 Effects of five environmental parameters (columns) on three carbon fluxes (rows) from farm dams. Points in each panel represent the partial residuals of farm dams after controlling for the effects of the other fixed and random variables in the models. Lines indicate statistically significant effects following the best-fitting mixed effect models ($\pm 95\%$ confidence intervals). Labelled in each panel are the p-values and the importance scores of statistically significant terms (see Table S2 for all test statistics and Table S3 for importance scores).

CH_4 $\text{m}^{-2} \text{d}^{-1}$ of unfenced farm dams. Our fieldwork took place in the bioregion of South Western Slopes in south-eastern Australia, an important agricultural hotspot covering 86,811 km^2 . This region contains an estimated 172 thousands farm dams with a cumulative surface area of 278 km^2 (Malerba et al., 2021), which is equivalent to the surface area of all lakes in the region (277 km^2 ; Crossman & Li, 2015). Assuming our data are representative of average yearly fluxes, we estimated that fencing farm dams in this region would avoid emissions of 468 tonnes CH_4 year $^{-1}$, which corresponds to 44,917 tonnes CO_2 -eq year $^{-1}$ using the 20-year Sustained-Flux Global Warming Potential (SGWP) metric. These are only ballpark estimates, and more data are needed to better estimate the opportunity for avoided emissions using farm dam restoration. Considering that most farm dams have broadly similar properties and serve the same purposes (i.e., collect water for agricultural uses), our results and qualitative mechanisms may also apply to other regions of the world—albeit with different magnitudes. Thus, an important next step is to use a cost-benefit analysis to determine if improving farm dam conditions could be a cost-effective way to help decarbonize agricultural practices at scale (Figure 6).

The range of diffusive carbon fluxes measured here (1 to 164 CH_4 $\text{mg m}^{-2} \text{d}^{-1}$ and -1.7 to 14 CO_2 $\text{g m}^{-2} \text{d}^{-1}$) is comparable to

previously published values for farm dams in Australia, Canada, India, and Sweden (Figure 7; Table S4). Yet, our study (and most others) measured diffusive methane fluxes without accounting for other pathways of methane emissions (e.g., ebullition events; Bastviken et al., 2008; Bastviken et al., 2011). Grinham et al. (2018) quantified both ebullitive and diffusive methane fluxes from Australian irrigation and stock dams and reported higher values than ours (up to 3.6 CH_4 $\text{g m}^{-2} \text{d}^{-1}$; Figure 7). It is possible that the benefits of fencing farm dams on carbon emissions are even higher than our estimates after accounting for multiple types of methane fluxes. However, research is needed to establish if fencing farm dams can influence ebullitive methane fluxes.

The two main findings of this study were: (1) that excluding livestock from farm dams improves water quality, and (2) that higher water quality corresponds to lower methane emissions (Figure 6). For the first finding, fenced farm dams recorded 32% less dissolved nitrogen, 39% less phosphorus, and 22% more dissolved oxygen than unfenced farm dams. Westgate et al. (2022) is the only other study on this topic and showed comparable results to ours, with a 45–50% reduction in total nitrogen and phosphorus in fenced farm dams over unfenced farm dams, together with reduced turbidity and




FIGURE 4 Dashed lines indicate where fluxes equal zero. Also indicated in the figure is the dissolved oxygen concentration (10 mg L^{-1}) associated with a switch from positive to negative CO_2 -eq. fluxes. Variables were linearized using \log_{10} -transformations when fitting the model, but they are presented here on an arithmetic scale (see [Figure 3k](#) for model fits presented in log scales).

lower fecal contamination. The similar results between two field studies from different years (2019 and 2021) and seasons (summer and autumn) suggest that the positive effects of fencing on water quality may be maintained throughout the year.

For the second finding, the higher water quality of fenced farm dams corresponded to 56% lower methane emissions ([Figure 6](#)). We found that total dissolved oxygen was a key driver explaining the reduced methane emissions. The strong negative effect of dissolved oxygen is consistent with our understanding of methanogenesis as a microbiological process requiring anaerobic conditions (Segers, 1998). Similarly, the positive effects of total dissolved nitrogen and sediment organic carbon stocks meet the expectation that freshwater environments rich with nutrients and labile organic materials emit more GHG (Beaulieu et al., 2019; Li et al., 2021; Peacock et al., 2021). Instead, a surprising result was the negative effect of total phosphorus on methane fluxes, particularly since phosphorus is thought to promote methane production rates (Peacock et al., 2019; Peacock et al., 2021). Phosphorus concentration only had a weak negative effect on methane fluxes but not on carbon dioxide or CO_2 -eq fluxes. As shown by Nijman et al. (2022), one explanation could be that a greater phosphorus availability increases the growth and activity of methane-oxidizing bacteria, resulting in a reduction of methane emissions through the oxidation of methane to hydrogen and carbon monoxide. Yet more studies are needed to clarify the effects of phosphorus on methanogenesis in farm dams.

We found that farm dams with very high concentrations of dissolved oxygen exhibited negative CO_2 -eq GHG fluxes (i.e., negative radiative balance), indicating a positive contribution to reduce

atmospheric warming. Most farm dams contribute to climate change by emitting substantial amounts of atmospheric GHG (Holgerson & Raymond, 2016; Ollivier et al., 2018; Peacock et al., 2021). Yet, under certain circumstances, small freshwater systems can remove GHG from the atmosphere and act as a carbon sink (Ollivier et al., 2018; Peacock et al., 2021; Webb, Hayes, et al., 2019; Webb, Leavitt, et al., 2019). While we found negative fluxes in only a minority of cases (13 farm dams out of 64), the effect of oxygen on CO_2 -eq fluxes was very predictable: every farm dam recording oxygen levels $>10 \text{ mg L}^{-1}$ also showed a carbon drawdown (at up to $1.2 \text{ g CO}_2\text{-eq m}^{-2} \text{ d}^{-1}$). These negative fluxes are due to aquatic photosynthesis (i.e., net ecosystem production) sequestering carbon dioxide from the atmosphere at higher rates than CO_2 -eq methane emissions. This finding further emphasizes the importance of farm dam management, even suggesting that increasing oxygen levels could turn farm dams into carbon sinks. Nonetheless, these results are likely to change during the night phase when plant respiration replaces photosynthesis, highlighting the importance of long-term studies on carbon dynamics in farm dams.

There is still considerable uncertainty on the net radiative balance of farm dams, as there is little data on the rates of carbon sequestration and storage in dam sediments. Yet, farm dams appear to have the highest burial rates of organic carbon among freshwater systems, ranging from 148 to $17,000 \text{ g C m}^{-2} \text{ year}^{-1}$ (Downing et al., 2008; Rogers et al., 2022). Therefore, it is possible that farm dams can sequester more carbon in the sediments than what they emit to the atmosphere. Future studies should investigate if fencing farm dams can increase carbon sequestration together with decreasing methane emissions.

Dissolved oxygen was strongly positively correlated with pH and strongly negatively correlated with carbon dioxide, which is evidence that aquatic primary production is the key process regulating dissolved oxygen in the farm dams of this study. Specifically, photosynthetic activity produces oxygen and consumes carbon dioxide, which results in higher pH from faster dissociation of HCO_3^- into CO_2 and OH^- (Zang et al., 2010). Had there been no correlation between pH and dissolved oxygen (as is often the case with aquaculture systems), other factors unrelated to photosynthesis (e.g., decomposition of organic matter) may have been more likely to drive changes in dissolved oxygen (Zang et al., 2010). Importantly, the pH increase from aquatic photosynthesis is likely to further reduce the carbon emissions of a farm dam by moving the carbonate equilibria toward carbonic acid and away from gaseous CO_2 . Specifically, as the system becomes more basic, the carbonate system changes from CO_2 dominated to CO_3^{2-} dominated, with negligible carbon dioxide left at $\text{pH} > 8.5$ (Andersen, 2018; Drever, 1997).

5 | CONCLUSIONS

We discovered that fencing to exclude livestock from farm dams improves water quality (i.e., fewer dissolved nutrients and higher

FIGURE 5 Statistical associations between (a) the dissolved oxygen and the pH of the water, (b) the dissolved oxygen and the CO₂ fluxes, and (c) CO₂ fluxes and the pH of the water. Each point represents the average value recorded from a farm dam. Reported in each panel are the correlation coefficient and the p-value of statistically significant relationships.





FIGURE 6 Effects of fencing farm dams on water quality and methane emissions. Installing fences to exclude livestock from farm dams reduces the direct deposition of nutrient-rich manure and urine into the water, avoids hooved livestock (ungulates) disturbing soil, and promotes higher vegetation cover around the dam. As a result, fenced farm dams have lower dissolved nutrients, higher dissolved oxygen, and lower methane emissions than unfenced dams. Percentages associated with fencing farm dams represent the relative change compared to unfenced dams (Figures 1 and 2).

dissolved oxygen) and reduces diffusive methane emissions. Our data also revealed a threshold in dissolved oxygen at 10 mg L⁻¹ above which farm dams switch from positive to negative CO₂-eq fluxes,

helping mitigate climate change. Considering avoided carbon emissions and additional economic and ecological co-benefits (i.e., higher biodiversity, increased livestock health, and capital value; Dobes




FIGURE 7 Emissions of (a) CH₄ and (b) CO₂ from agricultural ponds from the scientific literature divided by land-use type. Each symbol indicates the arithmetic mean and the range of values. Reported in each panel are the overall average emissions and the average ranges. All studies measured methane diffusion, except for Grinham et al. (2018) who measured both methane diffusion and ebullition.

et al., 2021; Hazell et al., 2001; Lewis-Phillips et al., 2019; Westgate et al., 2022), investing in better farm dam management appears to be a promising strategy for improving farming productivity and environmental sustainability. Nevertheless, carbon cycles in farm dams remain one of the least explored among freshwater systems. Promising avenues for follow-up studies include environmental work to analyze long-term cycles for several carbon pathways (e.g., methane ebullition, plant-mediated methane emissions, rate of carbon sedimentation), economic assessments to determine the best allocation of incentives for sustainable management interventions, and social studies to establish non-market benefits and farmers' willingness to adopt management interventions. This information will help deliver policy recommendations on the cost-effectiveness of investing in farm dam management as a novel carbon abatement strategy, as well as for additional co-benefits.

AUTHOR CONTRIBUTIONS

M.E.M., D.B.L., B.C.S., and P.I.M. designed the research, M.E.M., P.W., and I.N.Y. collected the data, M.E.M. and L.S. analyzed the data, M.E.M. wrote the first draft, and all authors contributed to the final draft.

ACKNOWLEDGMENTS

We thank the anonymous reviewers and the Editor for their many insightful comments and suggestions. This work was supported by the Australian Government through the Australian Research Council (project ID DE220100752) and the Alfred Deakin Fellowship scheme. We thank David Smith, Clare Cane,

Dan Florance, Michelle Young, Suzannah Macbeth, Eleanor Lang, Colleen O'Malley, Angelina Siegrist, Amber Croft, and the team at Sustainable Farms for facilitating fieldwork. We are grateful to landowners who allowed us to work on their farms. We also thank Dr Tertius de Kluiver (Dept. of Industry, Science, Energy and Resources, Australian Government) for insightful comments and suggestions. Open access publishing facilitated by Deakin University, as part of the Wiley - Deakin University agreement via the Council of Australian University Librarians.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

We published a public online repository with all raw data, R codes, analyses, plots, tables, and the results of the literature review on GitHub (<https://github.com/martinomalerba/FarmDamEmissions>) and Dryad (<https://doi.org/10.5061/dryad.g1jwstqt5>).

ORCID

Martino E. Malerba <https://orcid.org/0000-0002-7480-4779>
 David B. Lindenmayer <https://orcid.org/0000-0002-4766-4088>
 Ben C. Scheele <https://orcid.org/0000-0001-7284-629X>
 Pawel Waryszak <https://orcid.org/0000-0002-4245-3150>
 I. Noyan Yilmaz <https://orcid.org/0000-0003-0260-4708>
 Lukas Schuster <https://orcid.org/0000-0003-2691-9085>
 Peter I. Macreadie <https://orcid.org/0000-0001-7362-0882>

REFERENCES

- Agriculture Victoria. (2022). How much water is in my dam? <https://agriculture.vic.gov.au/farm-management/water/farm-water-solutions/how-much-water-is-in-my-dam> (access date 26 April 2022).
- Andersen, C. B. (2018). Understanding carbonate equilibria by measuring alkalinity in experimental and natural systems. *Journal of Geoscience Education*, 50(4), 389–403. <https://doi.org/10.5408/1089-9995-50.4.389>
- Bastviken, D., Cole, J. J., Pace, M. L., & Van de Bogert, M. C. (2008). Fates of methane from different lake habitats: Connecting whole-lake budgets and CH₄ emissions. *Journal of Geophysical Research: Biogeosciences*, 113(G2), 1–13.
- Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M., & Enrich-Prast, A. (2011). Freshwater methane emissions offset the continental carbon sink. *Science*, 331(6013), 50. <https://doi.org/10.1126/science.1196808>
- Beaulieu, J. J., DelSontro, T., & Downing, J. A. (2019). Eutrophication will increase methane emissions from lakes and impoundments during the 21st century. *Nature Communications*, 10(1), 1375. <https://doi.org/10.1038/s41467-019-09100-5>
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*, 57, 289–300.
- Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference - understanding AIC and BIC in model selection. *Sociological Methods and Research*, 33(2), 261–304. <https://doi.org/10.1177/0049124104268644>
- Conrad, R. (2009). The global methane cycle: recent advances in understanding the microbial processes involved. *Environmental Microbiology Reports*, 1(5), 285–292. <https://doi.org/10.1111/j.1758-2229.2009.00038.x>
- Crossman, S., & Li, O. (2015). *Surface hydrology polygons (regional)*. *Geoscience Australia*. <http://pid.geoscience.gov.au/dataset/ga/83134> (data sourced on the 4th Feb 2022 from).
- Dlugokencky, E. (2022). NOAA/GML. gml.noaa.gov/ccgg/trends_ch4/ (access date: 8 Feb 2022).
- Dobes, L., Crane, M., Higgins, T., Van Dijk, A., & Lindenmayer, D. B. (2021). Increased livestock weight gain from improved water quality in farm dams: A cost-benefit analysis. *PLoS One*, 16(8), e0256089. <https://doi.org/10.1371/journal.pone.0256089>
- Downing, J. A., Cole, J. J., Middelburg, J. J., Striegl, R. G., Duarte, C. M., Kortelainen, P., Prairie, K. A., & Laube, K. A. (2008). Sediment organic carbon burial in agriculturally eutrophic impoundments over the last century. *Global Biogeochemical Cycles*, 22(1), 1–10. <https://doi.org/10.1029/2006gb002854>
- Downing, J. A., Prairie, Y. T., Cole, J. J., Duarte, C. M., Tranvik, L. J., Striegl, R. G., McDowell, W. H., Kortelainen, P., Caraco, N. F., Melack, J. M., & Middelburg, J. J. (2006). The global abundance and size distribution of lakes, ponds, and impoundments. *Limnology and Oceanography*, 51(5), 2388–2397. <https://doi.org/10.4319/lo.2006.51.5.2388>
- Drever, J. I. (1997). *The geochemistry of natural waters: Surface and ground-water environments*. Prentice Hall.
- Fisher, A., Rudin, C., & Dominici, F. (2019). All models are wrong, but many are useful: Learning a variable's importance by studying an entire class of prediction models simultaneously. *Journal of Machine Learning Research*, 20(177), 1–81.
- Fox, J., & Weisberg, S. (2018). Visualizing fit and lack of fit in complex regression models with predictor effect plots and partial residuals. *Journal of Statistical Software*, 87(9), 1–27.
- Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (3rd ed.). SAGE Publications, Inc. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/index.html>
- Grinham, A., Albert, S., Deering, N., Dunbabin, M., Bastviken, D., Sherman, B., Lovelock, C. E., & Evans, C. D. (2018). The importance of small artificial water bodies as sources of methane emissions in Queensland, Australia. *Hydrology and Earth System Sciences*, 22(10), 5281–5298.
- Gulliver, A., Carnell, P. E., Trevathan-Tackett, S. M., Duarte de Paula Costa, M., Masqué, P., & Macreadie, P. I. (2020). Estimating the potential blue carbon gains from tidal marsh rehabilitation: A case study from south eastern Australia. *Frontiers in Marine Science*, 7, 403.
- Hazell, D., Cunningham, R., Lindenmayer, D., Mackey, B., & Osborne, W. (2001). Use of farm dams as frog habitat in an Australian agricultural landscape: factors affecting species richness and distribution. *Biological Conservation*, 102(2), 155–169.
- Holgerson, M. A., & Raymond, P. A. (2016). Large contribution to inland water CO₂ and CH₄ emissions from very small ponds. *Nature Geoscience*, 9(3), 222–226. <https://doi.org/10.1038/ngeo2654>
- IPCC. (2019). 2019 refinement to the 2006 IPCC guidelines for national greenhouse gas inventories.
- Lambert, M., & Fréchet, J.-L. (2005). Analytical techniques for measuring fluxes of CO₂ and CH₄ from hydroelectric reservoirs and natural water bodies. In *Greenhouse gas emissions—fluxes and processes* (pp. 37–60). Springer.
- Lewis-Phillips, J., Brooks, S., Sayer, C. D., McCrea, R., Siriwardena, G., & Axmacher, J. C. (2019). Pond management enhances the local abundance and species richness of farmland bird communities. *Agriculture, Ecosystems and Environment*, 273, 130–140. <https://doi.org/10.1016/j.agee.2018.12.015>
- Li, Y., Shang, J., Zhang, C., Zhang, W., Niu, L., Wang, L., & Zhang, H. (2021). The role of freshwater eutrophication in greenhouse gas emissions: A review. *Science of the Total Environment*, 768, 144582. <https://doi.org/10.1016/j.scitotenv.2020.144582>
- Limpert, K. E., Carnell, P. E., & Macreadie, P. I. (2021). Managing agricultural grazing to enhance the carbon sequestration capacity of freshwater wetlands. *Wetlands Ecology and Management*, 29(2), 231–244. <https://doi.org/10.1007/s11273-020-09780-7>
- Malerba, M. E., Wright, N., & Macreadie, P. I. (2021). A continental-scale assessment of density, size, distribution and historical trends of farm dams using deep learning convolutional neural networks. *Remote Sensing*, 13(2), 319.
- Malerba, M. E., Wright, N., & Macreadie, P. I. (2022). Australian farm dams are becoming less reliable water sources under climate change. *Science of the Total Environment*, 829, 154360. <https://doi.org/10.1016/j.scitotenv.2022.154360>
- Neubauer, S. C., & Megonigal, J. P. (2015). Moving beyond global warming potentials to quantify the climatic role of ecosystems. *Ecosystems*, 18(6), 1000–1013.
- Niittynen, P., & Luoto, M. (2018). The importance of snow in species distribution models of arctic vegetation. *Ecography*, 41(6), 1024–1037.
- Nijman, T. P. A., Amado, A. M., Bodelier, P. L. E., & Veraart, A. J. (2022). Relief of phosphate limitation stimulates methane oxidation. *Frontiers in Environmental Science*, 10, 1–8. <https://doi.org/10.3389/fenvs.2022.804512>
- Oates, L. G., Jackson, R. D., & Allen-Diaz, B. (2008). Grazing removal decreases the magnitude of methane and the variability of nitrous oxide emissions from spring-fed wetlands of a California oak savanna. *Wetlands Ecology and Management*, 16(5), 395–404. <https://doi.org/10.1007/s11273-007-9076-0>
- Ollivier, Q. R., Maher, D. T., Pitfield, C., & Macreadie, P. I. (2018). Punching above their weight: Large release of greenhouse gases from small agricultural dams. *Global Change Biology*, 25(2), 721–732. <https://doi.org/10.1111/gcb.14477>
- Ollivier, Q. R., Maher, D. T., Pitfield, C., & Macreadie, P. I. (2019). Winter emissions of CO₂, CH₄, and N₂O from temperate agricultural dams: fluxes, sources, and processes. *Ecosphere*, 10(11), 1–16. <https://doi.org/10.1002/ecs2.2914>
- Panneer Selvam, B., Natchimuthu, S., Arunachalam, L., & Bastviken, D. (2014). Methane and carbon dioxide emissions from inland waters

- in India—Implications for large scale greenhouse gas balances. *Global Change Biology*, 20(11), 3397–3407. <https://doi.org/10.1111/gcb.12575>
- Peacock, M., Audet, J., Bastviken, D., Cook, S., Evans, C. D., Grinham, A., Holgerson, M. A., Högbom, L., Pickard, A. E., Zieliński, P., & Futter, M. N. (2021). Small artificial waterbodies are widespread and persistent emitters of methane and carbon dioxide. *Global Change Biology*, 27, 5109–5123. <https://doi.org/10.1111/gcb.15762>
- Peacock, M., Audet, J., Jordan, S., Smeds, J., & Wallin, M. B. (2019). Greenhouse gas emissions from urban ponds are driven by nutrient status and hydrology. *Ecosphere*, 10(3), e02643.
- Pilon-Smits, E. (2005). Phytoremediation. *Annual Review of Plant Biology*, 56, 15–39.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & Team, R. C. (2020). nlme: Linear and nonlinear mixed effects models. R package version 3.1-150.
- R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rogers, M. N., Williamson, T. J., Knoll, L. B., & Vanni, M. J. (2022). Temporal patterns in sediment, carbon, and nutrient burial in ponds associated with changing agricultural tillage. *Biogeochemistry*, 159, 87–102. <https://doi.org/10.1007/s10533-022-00916-w>
- Rosentreter, J. A., Borges, A. V., Deemer, B. R., Holgerson, M. A., Liu, S., Song, C., Melack, J., Raymond, P. A., Duarte, C. M., Allen, G. H., Olefeldt, D., Poulter, B., Battin, T. I., & Eyre, B. D. (2021). Half of global methane emissions come from highly variable aquatic ecosystem sources. *Nature Geoscience*, 14(4), 225–230. <https://doi.org/10.1038/s41561-021-00715-2>
- Segers, R. (1998). Methane production and methane consumption: A review of processes underlying wetland methane fluxes. *Biogeochemistry*, 41(1), 23–51.
- Swartz, T. M., & Miller, J. R. (2021). The American Pond Belt: An untold story of conservation challenges and opportunities. *Frontiers in Ecology and the Environment*, 19(9), 501–509. <https://doi.org/10.1002/fee.2381>
- Virkkala, A. M., Aalto, J., Rogers, B. M., Tagesson, T., Treat, C. C., Natali, S. M., Watts, J. D., Potter, S., Lehtonen, A., Mauritz, M., Schuur, E. A. G., Kochendorfer, J., Zona, D., Oechel, W., Kobayashi, H., Humphreys, E., Goeckede, M., Iwata, H., Lafleur, P. M., ... Luoto, M. (2021). Statistical upscaling of ecosystem CO₂ fluxes across the terrestrial tundra and boreal domain: Regional patterns and uncertainties. *Global Change Biology*, 27, 4040–4059. <https://doi.org/10.1111/gcb.15659>
- Watkins, S. C., Baldwin, D. S., Waudby, H. P., & Ning, S. E. M. A. (2017). Managing rain-filled wetlands for carbon sequestration: A synthesis. *The Rangeland Journal*, 39(2), 145. <https://doi.org/10.1071/rj16077>
- Webb, J. R., Hayes, N. M., Simpson, G. L., Leavitt, P. R., Baulch, H. M., & Finlay, K. (2019). Widespread nitrous oxide undersaturation in farm waterbodies creates an unexpected greenhouse gas sink. *Proceedings of the National Academy of Sciences of the United States of America*, 116(20), 9814–9819.
- Webb, J. R., Leavitt, P. R., Simpson, G. L., Baulch, H. M., Haig, H. A., Hodder, K. R., & Finlay, K. (2019). Regulation of carbon dioxide and methane in small agricultural reservoirs: optimizing potential for greenhouse gas uptake. *Biogeosciences*, 16(21), 4211–4227. <https://doi.org/10.5194/bg-16-4211-2019>
- Westgate, M. J., Crane, M., Scheele, B. C., Crane, C., O'Malley, C., Siegrist, A., & Lindenmayer, D. B. (2022). Improved management of farm dams increases vegetation cover, water quality, and macroinvertebrate biodiversity. *Ecology and Evolution*, 12(3), e8636.
- Wickham, H. (2009). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York.
- Wickham, H. (2011). plyr: The split-apply-combine strategy for data analysis. *Journal of Statistical Software*, 40(1), 1–29.
- Wickham, H., François, R., Henry, L., & Müller, K. (2018). dplyr: A grammar of data manipulation.
- Zang, C., Huang, S., Wu, M., Du, S., Scholz, M., Gao, F., Lin, C., Guo, Y., & Dong, Y. (2010). Comparison of relationships between pH, dissolved oxygen and chlorophyll a for aquaculture and non-aquaculture waters. *Water, Air, and Soil Pollution*, 219(1–4), 157–174. <https://doi.org/10.1007/s11270-010-0695-3>
- Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects models and extensions in ecology with R*. Springer.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Malerba, M. E., Lindenmayer, D. B., Scheele, B. C., Waryszak, P., Yilmaz, I. N., Schuster, L., & Macreadie, P. I. (2022). Fencing farm dams to exclude livestock halves methane emissions and improves water quality. *Global Change Biology*, 28, 4701–4712. <https://doi.org/10.1111/gcb.16237>