



Cite this: *Med. Chem. Commun.*,
2019, 10, 1165

Synthesis of an α -phosphono- α,α -difluoroacetamide analogue of the diphosphoinositol pentakisphosphate 5-InsP₇†

Andrew M. Riley,^a Huanchen Wang,^b
Stephen B. Shears^b and Barry V. L. Potter^{b*}

Diphosphoinositol phosphates (PP-InsPs) are an evolutionarily ancient group of signalling molecules that are essential to cellular and organismal homeostasis. As the detailed mechanisms of PP-InsP signalling begin to emerge, synthetic analogues of PP-InsPs containing stabilised mimics of the labile diphosphate group can provide valuable investigational tools. We synthesised 5-PCF₂Am-InsP₅ (**1**), a novel fluorinated phosphonate analogue of 5-PP-InsP₅, and obtained an X-ray crystal structure of **1** in complex with diphosphoinositol pentakisphosphate kinase 2 (PPIP5K2). 5-PCF₂Am-InsP₅ binds to the kinase domain of PPIP5K2 in a similar orientation to that of the natural substrate 5-PP-InsP₅ and the PCF₂Am structure can mimic many aspects of the diphosphate group in 5-PP-InsP₅. We propose that **1**, the structural and electronic properties of which are in some ways complementary to those of existing phosphonoacetate and methylenebisphosphonate analogues of 5-PP-InsP₅, may be a useful addition to the expanding array of chemical tools for the investigation of signalling by PP-InsPs. The PCF₂Am group may also deserve attention for wider application as a diphosphate mimic.

Received 15th March 2019,
Accepted 8th May 2019

DOI: 10.1039/c9md00163h

rsc.li/medchemcomm

Introduction

The diphosphoinositol phosphates (PP-InsPs, “inositol pyrophosphates”) are of fundamental importance to all eukaryotes, with pivotal roles in cellular and organismic metabolic homeostasis (Fig. 1).¹ The acute clustering of monophosphate and diphosphate groups around the hexahydroxycyclohexane ring of *myo*-inositol (Ins) endows the PP-InsPs with the most concentrated three-dimensional array of phosphate groups found in Nature.¹ The PP-InsPs are formed through the enzymatic phosphorylation of *myo*-inositol hexakisphosphate (InsP₆, Fig. 1) by inositol hexakisphosphate kinases (IP6Ks) and diphosphoinositol pentakisphosphate kinases (PPIP5Ks).² Among the PP-InsPs, 5-diphospho-*myo*-inositol pentakisphosphate (5-PP-InsP₅, also known as “5-InsP₇”) is both the most abundant and the most intensively studied member of this signalling family.

An understanding of the molecular actions of PP-InsPs by their addition to cell-free systems can be confounded by the enzymatic and chemical instability of the diphosphate (PP) component. Therefore, we and other workers have developed more stable, synthetic versions of PP-InsPs in which the PP groups are replaced with phosphonoacetate (PA)^{3,4} or methylenebisphosphonate (PCP)^{4–7} groups (Fig. 1).⁸ Thus, a tethered version of 5-PCP-InsP₅ was recently used to search for novel binding proteins for PP-InsPs.⁹ Metabolically-stable PP-InsP analogues can also be informative for structural analysis of enzyme/substrate crystal complexes. For example, we have used 5-PA-InsP₅ to reveal a previously unidentified ligand capture site on PPIP5K2.¹⁰

While PA analogues are relatively easy to synthesise, the carboxylic ester of the PA group could be prone to chemical hydrolysis at high pH and/or enzymatic cleavage by cellular esterases.¹¹ In both PA and PCP analogues, the bridging oxygen of the diphosphate group is replaced by a methylene (CH₂) group, making the resulting carbon–phosphorus bonds resistant to hydrolysis in comparison to the native oxygen–phosphorus bonds.¹² However, the CH₂ group in methylene phosphonates is less electronegative than the bridging oxygen atom of phosphates, causing an increase in the pK_a value of the methylene phosphonic acid in its second deprotonation. This can mean that a methylene phosphonate analogue is less strongly ionised than the phosphate equivalent at physiological pH, potentially leading to a decreased affinity of the

^a Medicinal Chemistry and Drug Discovery, Department of Pharmacology, University of Oxford, Mansfield Road, Oxford OX1 3QT, UK.
E-mail: barry.potter@pharm.ox.ac.uk; Fax: +44 (0)1865 271853;
Tel: +44 (0)1865 271945

^b Inositol Signaling Group, Laboratory of Signal Transduction, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, North Carolina, USA

† Electronic supplementary information (ESI) available: Data deposition: atomic coordinates and structure factors have been deposited in the Protein Data Bank (PDB ID code 6N5C). See DOI: 10.1039/c9md00163h







Fig. 1 A. Inositol hexakisphosphate (InsP_6) is phosphorylated on P-5 by IP6K to give 5-PP- InsP_5 , also known as “5- InsP_7 ”. Further phosphorylation at P-1 by PPIP5K gives 1,5-[[PP] $_2$ - InsP_4 , (“ InsP_8 ”). B. Stabilised analogues of 5-PP- InsP_5 containing methylenebisphosphonate (PCP), phosphonoacetate (PA) and phosphonodifluoroacetamide (PCF_2Am) mimics of the natural PP group. IP6K, inositol hexakisphosphate kinase; PPIP5K, diphosphoinositol pentakisphosphate kinase.

Scheme 1 Synthesis of 5-deoxy-5-amino-inositol **7**. Reagents and conditions: a. see ref. 22; b. i. Tf_2O , CH_2Cl_2 , pyridine, -78°C to r.t.; ii. NaN_3 , DMF, r.t., 86%; c. i. $\text{LiOH}\cdot\text{H}_2\text{O}$, MeOH, THF, H_2O ; ii. NaH, BnBr, DMF, 93%; d. i. CH_2Cl_2 , 95% aqueous TFA; ii. NaH, BnBr, DMF, 91%; e. LiAlH_4 , THF, 100%. Bn, benzyl.

analogue for protein binding sites.¹² A well-established approach to this problem, originally developed by Blackburn and co-workers,¹³ involves replacing the phosphonate CH_2 group with difluoromethylene (CF_2). The more electronegative CF_2 group increases the acidity of the phosphonic acid and, in addition, the CF_2 group itself has greater electronic and steric similarity to the bridging oxygen atom of a phosphate than does CH_2 .¹⁴ The difluoromethylenephosphonate (PCF_2) group has been particularly useful as a phosphate mimic in the development of protein tyrosine phosphatase inhibitors.^{15,16}

In nucleotide chemistry, the difluoromethylenebisphosphonate (PCF_2P) motif has been used to mimic PP in stable analogues of nucleoside diphosphates and triphosphates.¹⁷ Although the PCF_2P group has been proposed as a potential PP mimic in stabilised analogues of PP- InsP_5 , this was anticipated to present a considerable synthetic challenge,^{5,18} and while several PCP- InsP_5 have been synthesised,^{4–7} no fluorinated equivalents have yet been disclosed.

In the current report, we explore an alternative approach to a fluorinated isopolar analogue of a PP- InsP_5 , building on our earlier work with phosphonoacetic acid (PA) esters. Although replacing the CH_2 group in the β -ketophosphonate fragment of a PA ester with CF_2 is possible, the resulting difluoroacetate ester would be very labile to hydrolysis. Therefore, we chose to replace the ester of PA with a stable amide linkage, allowing the inclusion of the electron-withdrawing CF_2 group in place of CH_2 . In the resulting analogue, 5- $\text{PCF}_2\text{Am-InsP}_5$ (**1**, Fig. 1), the terminal phosphonate group should more closely resemble the corresponding β -phosphate

group of 5-PP- InsP_5 in its electronic properties than in either the PA or the PCP equivalents. The α -phosphono- α,α -difluoroacetamide (PCF_2Am) unit lacks a close equivalent to the α -phosphate of PP, but the amide carbonyl retains the potential to accept H-bonds and the rigidity of the amide structure itself may confer advantages at some binding sites. To the best of our knowledge, the PCF_2Am motif¹⁹ has not previously been explored as a diphosphate isostere, although it has been used successfully in analogues of 1,3-bisphosphoglyceric acid as inhibitors of phosphoglycerate kinase²⁰ and in the design of inhibitors of aspartate carbamoyltransferase²¹ and protein tyrosine phosphatases.¹⁶


Results and discussion

Synthesis of 5- $\text{PCF}_2\text{Am-InsP}_5$ (**1**)

To construct compound **1**, we needed to synthesise an appropriately protected 5-deoxy-5-amino-*myo*-inositol intermediate (Scheme 1). We have previously shown²² that regioselective sulfonylation of butanediactal (BDA) protected *myo*-inositol **2** (ref. 3, 22 and 23) using triflic anhydride, followed by solvolysis using wet dimethylacetamide, gives inversion of configuration at C-5, to produce the *neo*-inositol acetate derivative **3** *via* an iminium ion intermediate.²² In the present work, we used a second triflate esterification of the free OH group in **3** followed by treatment with sodium azide in DMF to give a second configurational inversion, returning us to the *myo*-inositol configuration in the 5-deoxy-5-azido-*myo*-inositol derivative **4**.[‡]

[‡] A recent study²⁶ has systematically investigated the solvolysis and azidolysis of triflates derived from diol **2**.





Scheme 2 a. $(\text{EtO})_2\text{P}(\text{O})\text{CF}_2\text{COOH}$, EDAC, CH_2Cl_2 , 87%; b. H_2 , 50 p.s.i., $\text{Pd}(\text{OH})_2/\text{C}$, MeOH, THF, AcOH, 100%; c. i. $(\text{BnO})_2\text{PN}^i\text{Pr}_2$, 5-phenyl-1*H*-tetrazole, CH_2Cl_2 ; ii. *m*CPBA, 66%; d. i. TMSBr, CH_2Cl_2 ; ii. MeOH; iii. aqueous triethylammonium bicarbonate, 85%. Bn, benzyl.

Compound 4, with its combination of acid- and base-labile protecting groups, is a potentially versatile intermediate in itself. However, to simplify the current synthesis, we sequentially replaced these labile protecting groups with benzyl ethers. Thus, replacement of the 2-*O*-acetate ester with a 2-*O*-benzyl ether gave 5, a precursor for the synthesis of analogues of 5-PP-InsP₄. Next, the BDA groups were removed and replaced with benzyl ethers to give pentabenzyl 5-deoxy-5-azido-inositol 6. Re-

duction of the azide group in 6 using LiAlH_4 now gave the 5-deoxy-5-amino-inositol 7 in quantitative yield. The next step was to introduce the phosphonodifluoroacetamide unit (Scheme 2). Thus, reaction of 7 with diethyl phosphonodifluoroacetic acid²⁴ gave phosphonodifluoroacetamide 8. The benzyl protecting groups in 8 were removed by catalytic hydrogenolysis over $\text{Pd}(\text{OH})_2/\text{C}$ to give pentaol 9. Phosphitylation using bisbenzyloxydiisopropylaminophosphine activated with 5-phenyl-1*H*-tetrazole then gave the intermediate pentakisphosphite, which was observed by ³¹P NMR, but not isolated. Oxidation of phosphites with *m*CPBA yielded the fully-protected pentakisphosphate 10. Finally, the benzyl and ethyl protecting groups in 10 were cleanly removed using TMSBr in dichloromethane followed by MeOH.

Compound 1 was isolated as the triethylammonium salt, containing six TEA⁺ ions per molecule of 1. The non-fluorinated phosphonoacetamide (PCH₂Am) equivalent of 1 was also initially synthesised to explore synthetic methods (see ESI[†] for details). During this synthesis, we observed facile exchange of the CH₂ protons with deuterium from D₂O during NMR spectroscopy of the PCH₂Am-containing pentaol intermediate analogous to compound 9. The deuterium was retained after the subsequent phosphorylation and deprotection steps. Attempts to exchange deuterium back to hydrogen in the final product were unsuccessful, possibly because ionisation of the methylenephosphonate group disfavors the required enolisation (see ESI[†] for further details). This may suggest a strategy for developing tritiated versions of the non-fluorinated equivalent of 1 and related analogues of PP-InsPs.




Fig. 2 A. Inhibition of human PPIP5K2 by 5-PCF₂Am-InsP₅ (1) in the presence of 100 nM 1,5-IPP₂-InsP₄. B. Crystallographic analysis of PPIP5K2 in complex with 1 and AMPPNP (PDB code 6N5C). C. The 2*F*_o-*F*_c electron density map of 1, contoured at 1.7σ, is shown as light blue mesh. PPIP5K2 is shown in cartoon and 1 is shown as a stick model, with carbon atoms coloured dark grey, phosphorus atoms orange, oxygen atoms red, nitrogen atom blue and fluorine atoms cyan. D. Another view of 5-PCF₂Am-InsP₅ (1) and its overlap with 5-PP-InsP₅, which is shown as white sticks. Panels E and F show the moieties on position 5 of 1 and the natural substrate 5-PP-InsP₅ (PDB code, 3T9D).



The ^{31}P NMR spectrum of **1** in D_2O showed the signal corresponding to the PCF_2Am phosphorus atom at δ 0.64, shifted approximately 13.5 ppm up-field relative to the equivalent signal in the PCH_2Am equivalent (ESI† Fig. S1). This reflects the profound effect of the CF_2 group on the electronic properties of the phosphonate group in **1**. Compound **1** was very stable in aqueous solution; the NMR sample showed no sign of decomposition after >1 year in solution in D_2O .

Interaction of 5- $\text{PCF}_2\text{Am-InsP}_5$ with PPIP5K2

To determine whether **1** could act as a mimic of 5-PP- InsP_5 , we examined its interaction with the kinase domain of human PPIP5K2. We used a “reverse-kinase” assay that records ATP generated from 0.1 mM ADP during the dephosphorylation of 100 nM 1,5-[PP] $_2$ - InsP_4 . This approach avoids the need to use radiolabelled material and slow-throughput HPLC analysis. Furthermore, the assay of ATP production is inherently more sensitive compared to measuring ATP consumption. Compound **1** inhibited 1,5-[PP] $_2$ - InsP_4 metabolism by PPIP5K2 with an IC_{50} of 375 nM (Fig. 2A). We then obtained an X-ray structure of 5- $\text{PCF}_2\text{Am-InsP}_5$ (**1**) in complex with PPIP5K2 kinase domain and the stable ATP analogue AMPNP. The structure shows that 5- $\text{PCF}_2\text{Am-InsP}_5$ binds to the catalytic site of PPIP5K2 in a similar orientation to 5-PP- InsP_5 (Fig. 2B–D). While crystallography previously showed that 5-PA- InsP_5 binds to two sites in PPIP5K2 (the catalytic site and a surface-mounted ligand capture site),^{4,10} electron density for **1** was seen only in the catalytic site (Fig. 2B). This is similar to the complex structures of PPIP5K2 with 5-PCP- InsP_5 (ref. 7) or with 5-PP- InsP_5 itself.²⁵

The electron density for **1** (Fig. 2C) clearly shows that the planar amide unit adopts the *Z* conformation, and with the amide NH antiperiplanar to H-5 of *myo*-inositol. The carbonyl oxygen of the amide is orientated similarly to one non-bridging oxygen atom of the α -phosphate of bound 5-PP- InsP_5 (Fig. 2E and F). Additionally, one of the fluorine atoms in the CF_2 group stretches its position to that of the other non-bridging O-atom of an α -phosphate, which is expected to bring the negative charge distribution in this region closer to that of 5-PP- InsP_5 than occurs with the PA analogue. The terminal phosphonate group in bound **1** is held in an extended conformation (Fig. 2E) although its orientation is different to that of the corresponding β -phosphate group in 5-PP- InsP_5 bound to PPIP5K2 (Fig. 2F).

Conclusions

We have described the design and synthesis of a novel type of stable PP- InsP analogue containing a phosphonodifluoroacetamide (PCF_2Am) structure, intended to mimic the 5-diphosphate (PP) group of 5-PP- InsP_5 . As proof of principle, we examined the interaction of the new analogue, 5- $\text{PCF}_2\text{Am-InsP}_5$ (**1**), with the catalytic domain of the diphosphoinositol pentakisphosphate kinase PPIP5K2. A crystallographic study showed that **1** binds to the PP- InsP -binding site of the PPIP5K2 kinase domain in a similar orien-

tation to that of the natural substrate 5-PP- InsP_5 and that the PCF_2Am structure can mimic several functionally important aspects of the diphosphate group (PP) in 5-PP- InsP_5 .

Recent years have seen a rapid increase in reported functions of PP- InsPs , the identification of their target proteins, and the characterisation of metabolising enzymes. Stabilised analogues of PP- InsPs have often played valuable roles in this work. No single analogue of a PP- InsP can perfectly mimic its steric and electronic features in all environments, and a range of compounds with complementary properties therefore offers a more versatile analytical approach. It is in this context that we propose that 5- $\text{PCF}_2\text{Am-InsP}_5$ (**1**) brings new opportunities. For example, the presence of the CF_2 group in **1** means that its terminal phosphonate group should be more acidic than the equivalent group in 5-PA- InsP_5 or 5-PCP- InsP_5 , making it more similar in this respect to the β -phosphate of 5-PP- InsP_5 and potentially allowing stronger interactions with protein binding sites. In addition, the rigidity of the amide unit in **1** means that the PCF_2Am unit contains fewer rotatable bonds than the equivalent PP, PA or PCP structures. A more rigid ligand is less likely to lose entropy on binding than a more flexible one, and thus may bind more tightly to some sites. Note also that the 5-diphosphate in 5-PP- InsP_5 participates in complexing a hydrated Mg^{2+} ion, a property that is mimicked in the complex of 5-PCP- InsP_5 with PPIP5K2.⁷ This Mg^{2+} ion is not present in the complex with **1** (Fig. 2E). Thus, 5- $\text{PCF}_2\text{Am-InsP}_5$ may be particularly useful for studying ligand/protein interactions that do not involve Mg^{2+} .⁹ To the best of our knowledge, the PCF_2Am group has not been previously employed as a diphosphate mimic; given the large number of biomolecules that contain diphosphate or polyphosphate motifs, it may find wider applications than those outlined here.

Experimental

General chemistry methods

All reagents and solvents were of commercial quality and were used without further purification. Petroleum ether used for chromatography and crystallisations was of fractions 40–60 °C. Alcohol **3** was synthesised as previously reported.²² Diethyl phosphonodifluoroacetic acid was synthesised according to a literature procedure.²⁴ Thin-layer chromatography (TLC) was performed on pre-coated aluminium plates (Merck, silica gel 60, F254). Chromatograms were visualised under UV light and by dipping plates into either phosphomolybdic acid in EtOH, vanillin in acidic EtOH, or alkaline aqueous KMnO_4 , followed by heating. Flash column chromatography was performed on an ISCO CombiFlash Rf automated flash chromatography system using RediSep Rf disposable flash columns. NMR spectra were recorded on Bruker Avance III 400 and 500 MHz NMR spectrometers. Proton chemical shifts are reported in ppm (δ) relative to internal tetramethylsilane (TMS, 0.0 ppm) or with the solvent resonance relative to TMS employed as the internal standard (D_2O , 4.79 ppm). The following abbreviations are used to



describe resonances: br, broad; s, singlet; d, doublet; dd, double doublet; q, quartet; m, multiplet; t, triplet. ^{13}C chemical shifts are reported relative to internal TMS (TMS, 0.0 ppm) or with the solvent resonance relative to TMS employed as the internal standard. The assignments of the proton and carbon atoms are based on 2D-NMR experiments (^1H - ^1H -COSY, HSQC). ^{31}P chemical shifts are reported in ppm (δ) relative to an 85% H_3PO_4 external standard (H_3PO_4 , 0.0 ppm). ^{19}F chemical shifts are reported in ppm (δ) relative to a CFCl_3 external standard (CFCl_3 , 0.0 ppm). Melting points were determined using a Stanford Research Systems Optimelt MPA100 automated melting point system and are uncorrected. High resolution time-of-flight mass spectra were obtained on a Bruker Daltonics micrOTOF mass spectrometer using electrospray ionisation (ESI $^+$).

2-O-Acetyl-5-deoxy-5-azido-1,6:3,4-bis-[O-(2,3-dimethoxybutane-2,3-diyl)]-myo-inositol (4)

To a stirred solution of alcohol 3 (1.15 g, 2.55 mmole) in dry dichloromethane (10 mL) and dry pyridine (1 mL) under N_2 at -78°C was added trifluoromethanesulfonic anhydride (4.0 mL of a 1.0 mol dm^{-3} solution in dichloromethane, 4.0 mmol) dropwise over 15 min. The cooling bath was removed and the solution was allowed to warm to room temperature. Stirring was continued overnight (16 h) after which time TLC (ethyl acetate:petroleum ether 2:1) showed complete conversion of alcohol (R_f 0.32) into a less polar product (R_f 0.70). Excess triflic anhydride was destroyed by careful addition of deionised water, and the mixture was diluted with dichloromethane (30 mL). The organic layer was separated and washed with 1.0 mol dm^{-3} HCl, saturated NaHCO_3 and brine (40 mL of each), then dried over MgSO_4 and concentrated under reduced pressure (no heat) to give crude triflate (1.45 g 2.49 mmole, 98%) as a reddish solid. To a solution of crude triflate (1.24 g, 2.13 mmol) in dry DMF (10 mL) at room temperature was added sodium azide (415 mg, 6.38 mmol). The mixture was stirred under N_2 for 16 h, after which time TLC (ethyl acetate:petroleum ether 1:3) showed complete conversion of triflate (R_f 0.24, stains purple with vanillin) into a slightly less polar product (R_f 0.30, stains orange with vanillin). The solution was concentrated by evaporation under reduced pressure (care: explosion risk with azides!). The residue was taken up in ethyl acetate (50 mL) and the solution was washed with deionised water and brine (50 mL each), then concentrated. The residue was purified by flash chromatography on silica, eluting with ethyl acetate to give 4 as a white solid (892 mg, 1.88 mmol, 86% from 3); crystals from boiling petroleum ether, m.p. 198–199 $^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 1.23 (6H, s, $2 \times \text{CH}_3$), 1.30 (6H, s, $2 \times \text{CH}_3$), 2.12 (3H, s, CH_3CO), 3.24 (6H, s, OCH_3), 3.30 (6H, s, OCH_3), 3.52 (1H, t, J 10.0 Hz, H-5), 3.66 (2H, dd, J 10.0, 2.6 Hz, H-1 and H-3), 3.85 (2H, t, J 10.0 Hz, H-4 and H-6), 5.41 (1H, t, J 2.7 Hz, H-2); ^{13}C NMR (100 MHz, CDCl_3) δ 17.41 (CH_3), 17.62 (CH_3), 21.10 (CH_3CO), 48.00 (OCH_3), 48.16 (OCH_3), 61.04 (C-5), 67.53 (C-1 and C-3), 68.51 (C-4 and C-6), 68.94 (C-2), 99.72

(BDA quaternary C), 99.94 (BDA quaternary C), 170.23 (C=O); HRMS (m/z) [$\text{M} + \text{Na}$] $^+$ calcd. for $\text{C}_{20}\text{H}_{33}\text{N}_3\text{O}_{10}$, 498.2058; found 498.2035.

2-O-Benzyl-5-deoxy-5-azido-1,6:3,4-bis-[O-(2,3-dimethoxybutane-2,3-diyl)]-myo-inositol (5)

To a solution of 4 (530 mg, 1.11 mmol) in THF (12 mL) was added a solution of $\text{LiOH}\cdot\text{H}_2\text{O}$ (420 mg, 10 mmol) in deionised water (3 mL) and methanol (12 mL). The mixture was stirred vigorously at room temperature for 2 h, after which time TLC (ethyl acetate:petroleum ether 1:1) showed complete conversion of 4 (R_f 0.54) into a more polar product (R_f 0.28). The solution was concentrated and the residue was taken up in ethyl acetate (50 mL). The solution was washed with water (2×50 mL), dried (MgSO_4) and concentrated to give the alcohol as a white solid (484 mg). This crude alcohol was taken up in DMF (10 mL). The solution was stirred at 0°C and sodium hydride (60% suspension in mineral oil, 67 mg, 1.67 mmole) was added. After 30 min, benzyl bromide (0.16 mL, 1.3 mmol) was added and the mixture was stirred overnight (16 h) at room temperature. Excess NaH was destroyed by careful addition of water and the solution was concentrated. The residue was taken up in ethyl acetate (50 mL) and washed with water and brine (50 mL each), dried (MgSO_4) and concentrated to give a solid residue. Purification by flash chromatography on silica (ethyl acetate in petroleum ether, 0 to 50%) gave 5 as a white solid (541 mg, 1.03 mmol, 93% over two steps); R_f 0.34 (ethyl acetate:petroleum ether 1:3); crystals from petroleum ether, m.p. 175–176.5 $^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 1.31 (6H, s, $2 \times \text{CH}_3$), 1.32 (6H, s, $2 \times \text{CH}_3$), 3.23 (6H, s, OCH_3), 3.29 (6H, s, OCH_3), 3.49 (1H, t, J 10.0 Hz, H-5), 3.57 (2H, dd, J 10.0, 2.5 Hz, H-1 and H-3), 3.80 (1H, t, J 2.7 Hz, H-2), 4.03 (2H, t, J 10.1 Hz, H-4 and H-6), 4.84 (2H, s, OCH_2Ph), 7.22–7.27 (1H, m, *para*-H of Ph), 7.29–7.34 (2H, m, *meta*-H of Ph), 7.48–7.51 (2H, m, *ortho*-H of Ph); ^{13}C NMR (100 MHz, CDCl_3) δ 17.61 (CH_3), 17.70 (CH_3), 47.92 (OCH_3), 48.00 (OCH_3), 61.52 (C-5), 68.46 (C-4 and C-6), 69.70 (C-1 and C-3), 73.94 (OCH_2Ph), 76.02 (C-2), 99.51 (BDA quaternary C), 99.71 (BDA quaternary C), 127.02 (*para*-C of Ph), 127.62 (CH of Ph), 127.90 (CH of Ph), 139.49 (*ipso*-C of Ph); HRMS (m/z) [$\text{M} + \text{Na}$] $^+$ calcd. for $\text{C}_{25}\text{H}_{37}\text{N}_3\text{O}_9$, 546.2422; found 546.2406.

1,2,3,4,6-Penta-O-benzyl-5-deoxy-5-azido-myoinositol (6)

To a stirred solution of 5 (507 mg, 0.968 mmol) in dichloromethane (5 mL) was added 95% aqueous TFA (5 mL). A yellow colour (butanedione) appeared within 30 s. After 30 min, the solution was concentrated to leave the crude tetraol as a white solid (325 mg). This solid was taken up in dry DMF (10 mL) and the solution was cooled to 0°C , before sodium hydride (60% suspension in mineral oil, 240 mg, 6.0 mmole) was added. The stirred suspension was allowed to warm to room temperature and then cooled again to 0°C before benzyl bromide (0.60 mL, 5.1 mmol) was added, dropwise. The cooling bath was removed and the mixture



was stirred overnight (16 h) at room temperature. Excess NaH was destroyed by careful addition of water and the solution was concentrated. The residue was taken up in dichloromethane (50 mL) and washed with water and brine (50 mL each), dried (MgSO₄) and concentrated to give an oily residue. Purification by flash chromatography on silica (ethyl acetate in petroleum ether, 0 to 50%) gave **6** as a colourless oil, which slowly crystallised (576 mg, 0.878 mmol, 91% over two steps); *R*_f 0.40 (ethyl acetate:petroleum ether 1:4); crystals from boiling ethanol, m.p. 89.0–90.5 °C; ¹H NMR (400 MHz, CDCl₃) δ 3.33 (2H, dd, *J* 9.7, 2.2 Hz, H-1 and H-3), 3.35 (1H, t, *J* 9.9 Hz, H-5), 3.85 (2H, t, *J* 9.8 Hz, H-4 and H-6), 4.00 (1H, t, *J* 2.4 Hz, H-2), 4.57, 4.62 (4H, AB quartet, *J*_{AB} 11.7 Hz, 2 × OCH₂Ph), 4.81, 4.87 (4H, AB quartet, *J*_{AB} 10.5 Hz, 2 × OCH₂Ph), 4.85 (2H, s, OCH₂Ph), 7.25–7.41 (25H, m, Ph); ¹³C NMR (100 MHz, CDCl₃) δ 67.68 (C-5), 72.77 (2 × OCH₂Ph), 74.25 (OCH₂Ph), 74.31 (C-2), 75.73 (2 × OCH₂Ph), 79.77 (C-4 and C-6), 81.20 (C-1 and C-3), 127.46, 127.57, 127.70, 127.72, 127.78, 128.20, 128.30, 128.34 and 128.41 (CH of Ph), 138.12 (2 × *ipso*-C of Ph), 138.24 (2 × *ipso*-C of Ph), 138.79 (*ipso*-C of Ph); HRMS (*m/z*) [*M* + Na]⁺ calcd. for C₄₁H₄₁N₃O₅, 678.2938; found 678.2966.

1,2,3,4,6-Penta-*O*-benzyl-5-deoxy-5-amino-*myo*-inositol (7)

To dry THF (3 mL) under argon was added LiAlH₄ (1.0 mL of a 1.0 mol dm⁻³ solution in THF, 1.0 mmol). The solution was stirred at 0 °C under argon and a solution of the azide **6** (420 mg, 0.640 mmol) in dry THF (4 mL) was injected over 10 min. After 30 min, the cooling bath was removed and the solution was stirred at room temperature for a further 1 h. The solution was then cooled to 0 °C again and excess LiAlH₄ was destroyed by addition of ethyl acetate (2 mL). The solution was allowed to warm to room temperature and then diluted with water (20 mL). The suspension was extracted with diethyl ether (3 × 20 mL). The aqueous layer was re-extracted with dichloromethane (20 mL) and the combined organic extracts were then washed with brine (50 mL), dried (MgSO₄) and concentrated to give amine **7** as a white solid (402 mg, 0.638 mmol, 100%); ¹H NMR (400 MHz, CDCl₃) δ 1.73 (2H, broad s, NH₂), 2.81 (1H, t, *J* 9.6 Hz, H-5), 3.40 (2H, dd, *J* 9.6, 2.2 Hz, H-1 and H-3), 3.85 (2H, t, *J* 9.6 Hz, H-4 and H-6), 4.08 (1H, t, *J* 2.4 Hz, H-2), 4.61 (4H, s, OCH₂Ph), 4.69, 5.01 (4H, AB quartet, *J*_{AB} 11.0 Hz, 2 × OCH₂Ph), 4.88 (2H, s, OCH₂Ph), 7.25–7.43 (25H, m, Ph); ¹³C NMR (100 MHz, CDCl₃) δ 56.17 (C-5), 72.41 (2 × OCH₂Ph), 74.16 (OCH₂Ph), 74.26 (C-2), 75.61 (2 × OCH₂Ph), 81.66 (C-4 and C-6), 82.23 (C-1 and C-3), 127.27, 127.57, 127.63, 127.65, 127.99, 128.08, 128.12, 128.40 and 128.41 (CH of Ph), 138.33 (2 × *ipso*-C of Ph), 138.85 (2 × *ipso*-C of Ph), 139.09 (*ipso*-C of Ph). HRMS (*m/z*) [*M* + H]⁺ calcd. for C₄₁H₄₃NO₅, 630.3214; found 630.3210.

1,2,3,4,6-Penta-*O*-benzyl-5-deoxy-5-(diethylphosphonodifluoroacetamido)-*myo*-inositol (8)

To a solution of amine **7** (141 mg, 0.224 mmol) and EDAC (86 mg, 0.448 mmol) in dry dichloromethane (3 mL) under

N₂ was added a solution of diethyl phosphonodifluoroacetic acid²⁴ (104 mg, 0.448 mmol) in dry dichloromethane (2 mL). The solution was stirred at room temperature for 1 h, after which time TLC (ethyl acetate:petroleum ether 1:1) showed total conversion of amine (streak, ~*R*_f 0.20) into a less polar product (spot, *R*_f 0.46). Ethyl acetate (30 mL) was added and the solution was washed with saturated NaHCO₃, 1.0 mol dm⁻³ HCl and brine (20 mL each), then dried (MgSO₄) and concentrated. The residue was purified by flash chromatography on silica (ethyl acetate in petroleum ether, 0 to 100%) to give **8** as a colourless oil, which slowly crystallised (164 mg, 0.194 mmol, 87%); *R*_f 0.40 (ethyl acetate:petroleum ether 1:4); crystals from boiling diisopropyl ether, m.p. 132.0–133.5 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.24 (6H, td, *J* 7.1, 0.6 Hz, POCH₂CH₃), 3.46 (2H, dd, *J* 9.0, 2.2 Hz, H-1 and H-3), 3.98 (2H, t, *J* 9.1 Hz, H-4 and H-6), 4.03 (1H, t, *J* 2.2 Hz, H-2), 4.10 (1H, q, *J* 9.2 Hz, H-5), 4.13–4.21 (4H, m, POCH₂CH₃), 4.56, 4.60 (4H, AB quartet, *J*_{AB} 11.8 Hz, 2 × OCH₂Ph), 4.73, 4.80 (4H, AB quartet, *J*_{AB} 11.8 Hz, 2 × OCH₂Ph), 4.84 (2H, s, OCH₂-Ph), 6.66 (1H, d, *J* 9.2 Hz, NH), 7.20–7.33 (23H, m, Ph), 7.39–7.42 (2H, m, Ph); ¹³C NMR (100 MHz, CDCl₃) δ 16.22 (³*J*_{CP} 5.6 Hz, POCH₂CH₃), 54.68 (C-5), 65.65 (²*J*_{CP} 6.6 Hz, POCH₂CH₃), 72.77 (2 × OCH₂Ph), 74.12 (OCH₂Ph), 74.55 (C-2), 74.76 (2 × OCH₂Ph), 78.31 (C-4 and C-6), 81.34 (C-1 and C-3), 111.92 (td, ¹*J*_{CF} 271.6 Hz, ¹*J*_{CP} 197.9 Hz, CF₂), 127.40, 127.49, 127.66, 127.69, 127.83, 127.92, 128.19, 128.23 and 128.36 (CH of Ph), 138.11 (2 × *ipso*-C of Ph), 138.35 (2 × *ipso*-C of Ph), 138.77 (*ipso*-C of Ph), 161.40 (td, ²*J*_{CF} 24.6 Hz, ²*J*_{CP} 17.3 Hz, C=O), ³¹P NMR (CDCl₃, 162 MHz, ¹H-decoupled) δ 3.47 (1 P, t, ²*J*_{FP} 96.1 Hz); ¹⁹F NMR (CDCl₃, 471 MHz) δ -116.11 (2 F, d, ²*J*_{FP} 96.1 Hz); HRMS (*m/z*) [*M*]⁻ calcd. for C₄₇H₅₂F₂NO₉P, 842.3275; found 842.3264.

5-Deoxy-5-(diethylphosphonodifluoroacetamido)-*myo*-inositol (9)

To a solution of **8** (130 mg, 0.154 mmol) in methanol (8 mL), THF (2 mL), deionised water (2 mL) and acetic acid (1 mL) was added palladium hydroxide on activated charcoal (20%, 50% water, 50 mg). The suspension was shaken in a Parr hydrogenator under H₂ (50 p.s.i.) for 24 h. The catalyst was removed by filtration through a PTFE syringe filter and the resulting colourless solution was concentrated, then dried under vacuum to give pentaol **9** as a white solid (61 mg, 0.154 mmol, 100%); TLC (dichloromethane:methanol 2:1): *R*_f 0.40; ¹H NMR (400 MHz, D₂O) δ 1.38 (6H, broad t, *J* ~7 Hz, POCH₂CH₃), 3.61 (2H, broad d, *J* ~9 Hz, H-1 and H-3), 3.71–3.81 (3H, m, H-4, H-5 and H-6), 4.10 (1H, broad s, H-2), 4.35–4.42 (4H, m, POCH₂CH₃); ¹³C NMR (100 MHz, D₂O) δ 15.64 (³*J*_{CP} 5.2 Hz, POCH₂CH₃), 56.28 (C-5), 67.37 (²*J*_{CP} 7.1 Hz, POCH₂CH₃), 70.05 (C-4 and C-6), 71.91 (C-2), 72.04 (C-1 and C-3), 111.71 (td, ¹*J*_{CF} 271.8 Hz, ¹*J*_{CP} 208.7 Hz, CF₂), 163.33 (td, ²*J*_{CF} 24.5 Hz, ²*J*_{CP} 16.2 Hz, C=O); ³¹P NMR (162 MHz, D₂O, ¹H-decoupled) δ 4.39 (t, ²*J*_{FP} 101.5 Hz); ¹⁹F NMR (471 MHz, D₂O) δ -117.62 (d, ²*J*_{FP} 101.4 Hz); HRMS (*m/z*) [*M* + Na]⁺ calcd. for C₁₂H₂₂F₂NO₉P, 416.0892; found 416.0876.



5-Deoxy-5-(diethylphosphonodifluoroacetamido)-*myo*-inositol 1,2,3,4,6-*O*-pentakis(dibenzylphosphate) (10)

To a stirred suspension of pentaol **9** (30 mg, 0.076 mmol) and 5-phenyl-1*H*-tetrazole (83 mg, 0.57 mmol) in dry dichloromethane (3 mL) under N₂ at room temperature was added bis(benzyloxy)diisopropylaminophosphine (0.20 mL, 0.60 mmol). The mixture was stirred under N₂ at room temperature for 3 h and then cooled to -78 °C, before *m*CPBA (70%, 187 mg, 0.760 mmol) was added. The mixture was allowed to warm to room temperature and then diluted with EtOAc (30 mL). The clear, colourless solution was washed with 10% aq. Na₂SO₃ solution (3 × 25 mL), dried over MgSO₄ and concentrated. The residue was purified by flash chromatography (EtOAc in petroleum ether, 0 to 100%) to give **10** as a colourless oil (85 mg, 0.050 mmole, 66%); TLC (EtOAc:petroleum ether, 4:1): *R*_f 0.56; ¹H NMR (CDCl₃, 400 MHz) δ 1.27 (6H, td, *J* 7.1, 0.5 Hz, POCH₂CH₃), 4.20–4.33 (5H, m, 2 × POCH₂CH₃ and H-5), 4.39 (2H, broad t, *J* ~9.5 Hz, H-1 and H-3), 4.82 (2H, q, *J* ~10 Hz, H-4 and H-6), 4.89–5.11 (18H, POCH₂Ph), 5.15–5.19 (2H, POCH₂Ph), 5.63 (1H, broad d, *J* ~9 Hz, H-2), 7.11–7.28 (50H, m, Ph), 7.52 (1H, broad d, *J* ~9 Hz, NH); ¹³C NMR (101 MHz, CDCl₃) δ 16.31 (d, ³*J*_{CP} 5.6 Hz, POCH₂CH₃), 52.70 (broad, C-5), 65.35 (d, ²*J*_{CP} 6.4 Hz, POCH₂CH₃), 69.75–70.06 (overlapping signals with *J*_{CP} couplings, POCH₂Ph), 73.72 (with *J*_{CP} couplings, C-4 and C-6), 74.28 (with *J*_{CP} couplings, C-1 and C-3), 75.93 (with *J*_{CP} couplings, C-2), 112.1 (¹*J*_{CP} 205 Hz, ¹*J*_{CF} unreadable through noise, CF₂), 127.83, 127.95, 128.12, 128.16, 128.21, 128.26, 128.32, 128.36, 128.39, 128.45, 128.49 (CH of Ph), 135.54–135.74 (overlapping signals with *J*_{CP} couplings, *ipso*-C of Ph), 162.85 (²*J*_{CP} ~18 Hz, ²*J*_{CF} unreadable through noise, C=O); ³¹P NMR (162 MHz, CDCl₃, ¹H-decoupled) δ -2.30 (1 P, P-2), -1.64 (2 P), -0.11 (2 P), 3.20 (1 P, t, ²*J*_{PF} 93.6 Hz); ¹⁹F NMR (471 MHz, CDCl₃) δ -113.48 (d, ²*J*_{FP} 93.4 Hz); HRMS (*m/z*) [M + Na]⁺ calcd. for C₈₂H₈₇F₂NO₂₄P₆, 1716.3904; found 1716.3965; [M + H]⁺ calcd. for C₈₂H₈₇F₂NO₂₄P₆, 1694.4085; found 1694.4130.

5-Deoxy-5-(phosphonodifluoroacetamido)-*myo*-inositol 1,2,3,4,6-pentakisphosphate (1)

A stirred solution of **10** (68 mg, 40 μmole) in dry dichloromethane (2 mL) was cooled to 0 °C under N₂ and trimethylsilyl bromide (1 mL) was added dropwise over 5 min. The solution was allowed to warm gradually to room temperature, and stirring was continued for 48 h. The solution was concentrated and methanol (5 mL) was added to the residue. The resulting colourless solution was stirred at room temperature for a further 1 h, and then concentrated to give a white gum. The gum was washed with diethyl ether (3 × 2 mL), then taken up in aqueous triethylammonium bicarbonate (1.0 mol dm⁻³, pH 7.6, 5 mL). This solution was then washed with diethyl ether (3 × 5 mL) and concentrated. The residue was re-dissolved in MilliQ water and lyophilised to give the triethylammonium salt of the title compound **1** as a colourless solid (47 mg, 34 μmole, 85%); ¹H NMR (500 MHz,

D₂O) δ 1.28 (approx. 57 H, t, *J* 7.3 Hz, CH₃ of TEA⁺), 3.20 (approx. 38 H, q, *J* 7.3 Hz, CH₂ of TEA⁺), 4.16 (1H, broad t, *J* ~10 Hz, H-5), 4.30 (2H, tt, *J* 9.6, 2.0 Hz, H-1 and H-3), 4.55 (2H, q, *J* 9.7 Hz, H-4 and H-6), 4.87 (1H, dt, *J* 9.8, 2.4 Hz, H-2); ¹³C NMR (126 MHz, D₂O) δ 8.19 (CH₃ of TEA⁺), 46.54 (CH₂ of TEA⁺), 53.91 (C-5), 74.44 (C-1, C-3, C-4 and C-6), 75.86 (C-2), 114.65 (dt, ¹*J*_{CP} 176.6 Hz, ¹*J*_{CF} 268.7 Hz, CF₂), 166.30 (td, ²*J*_{CF} 25.3 Hz, ²*J*_{CP} 15.3 Hz, C=O); ³¹P NMR (162 MHz, D₂O, ¹H-decoupled) δ -0.75 (1 P, P-2), -0.32 (2 P), 0.15 (2 P), 0.64 (1 P, t, ²*J*_{PF} 87.9 Hz, P-5); ¹⁹F NMR (471 MHz, D₂O) δ -118.86 (d, ²*J*_{FP} 85.5 Hz, CF₂); HRMS (*m/z*) [M - H]⁻ calcd. for C₈H₁₉F₂NO₂₄P₆, 735.8618; found 735.8654.

Protein expression, purification, crystallisation and structure determination

The kinase domain of human PPIP5K2 (residues 41–366) was sub-cloned, expressed and purified as before.²⁵ PPIP5K2 kinase domain was crystallised by hanging drop vapour diffusion against a well buffer of 12% (w/v) PEG 3350, 20 mM MgCl₂, 0.1 M HEPES, pH 7.0, 2 mM CdCl₂, 1 mM AMPPNP at 4 °C. The crystals were transferred to a stabilising buffer containing 22% (w/v) PEG 3350, 10 mM MgCl₂, 0.1 M sodium acetate, pH 5.2 at 4 °C and the crystals were then soaked under the above stabilising buffer for three days with 2 mM compound **1**. Cryosolvent was prepared by adding 33% ethylene glycol into the soaking buffer. Diffraction data were collected using APS beamlines 22-ID. All data were processed with the program HKL2000. The structure was determined using rigid body and direct Fourier synthesis, and refined with the equivalent and expanded test sets. The structure was further manually rebuilt with COOT and refined with REFMAC from the CCP4 package. Ligand topology and parameter files were prepared using the PRODRG server. The molecular graphics representations were prepared with the program PyMol (Schrödinger, LLC). Atomic coordinates and structure factors have been deposited with the Protein Data Bank with accession code 6N5C.

PIPIP5K2 kinase assay

Human PPIP5K2 kinase domain (residues 1–366, 2.5 μg mL⁻¹) was incubated at 24 °C for 30 min with buffer containing 20 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 0.1 mM ADP, 100 nM 1,5-[PP]₂-InsP₄ and various concentration of compound **1** in a 20 μL assay. The generated ATP was measured using a Molecular Probes ATP Determination kit (Thermo Fisher Scientific catalog number A22066). The IC₅₀ value for **1** was calculated using GraphPad Prism.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

BVLP is a Wellcome Trust Senior Investigator (grant 101010). SBS and HW are supported by the Intramural Research



Program of the NIH, National Institutes of Environmental Health Sciences.

Notes and references

- 1 S. B. Shears, *J. Cell. Physiol.*, 2018, **233**, 1897–1912.
- 2 M. P. Thomas and B. V. L. Potter, *FEBS J.*, 2014, **281**, 14–33.
- 3 A. M. Riley, H. Wang, J. D. Weaver, S. B. Shears and B. V. L. Potter, *Chem. Commun.*, 2012, **48**, 11292–11294.
- 4 A. M. Riley, H. Wang, S. B. Shears and B. V. L. Potter, *Chem. Commun.*, 2015, **51**, 12605–12608.
- 5 M. Wu, B. E. Dul, A. J. Trevisan and D. Fiedler, *Chem. Sci.*, 2013, **4**, 405–410.
- 6 M. Wu, L. S. Chong, S. Capolicchio, H. J. Jessen, A. C. Resnick and D. Fiedler, *Angew. Chem., Int. Ed.*, 2014, **53**, 7192–7197.
- 7 A. Hager, M. Wu, H. Wang, N. W. Brown, S. B. Shears, N. Veiga and D. Fiedler, *Chem. – Eur. J.*, 2016, **22**, 12406–12414.
- 8 N. W. Brown, A. M. Marmelstein and D. Fiedler, *Chem. Soc. Rev.*, 2016, **45**, 6311–6326.
- 9 M. Wu, L. S. Chong, D. H. Perlman, A. C. Resnick and D. Fiedler, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, E6757–E6765.
- 10 H. Wang, H. Y. Godage, A. M. Riley, J. D. Weaver, S. B. Shears and B. V. L. Potter, *Chem. Biol.*, 2014, **21**, 689–699.
- 11 S. Vertuani, A. Baldisserotto, K. Varani, P. A. Borea, B. De Marcos Maria Cruz, L. Ferraro, S. Manfredini and A. Dalpiaz, *Eur. J. Med. Chem.*, 2012, **54**, 202–209.
- 12 T. S. Elliott, A. Slowey, Y. Ye and S. J. Conway, *Med. Chem. Commun.*, 2012, **3**, 735–751.
- 13 G. M. Blackburn, D. A. England and F. Kolkman, *J. Chem. Soc., Chem. Commun.*, 1981, 930–932.
- 14 D. O'Hagan and H. S. Rzepa, *Chem. Commun.*, 1997, 645–652; V. D. Romanenko and V. P. Kukhar, *Chem. Rev.*, 2006, **106**, 3868–3935; K. Panigrahi, D. L. Nelson and D. B. Berkowitz, *Chem. Biol.*, 2012, **19**, 666–667; M. V. Ivanova, A. Bayle, T. Besset, X. Pannecoucke and T. Poisson, *Chem. – Eur. J.*, 2016, **22**, 10284–10293.
- 15 A. P. Combs, *J. Med. Chem.*, 2010, **53**, 2333–2344; M. Bahta, G. T. Lountos, B. Dyas, S. E. Kim, R. G. Ulrich, D. S. Waugh and T. R. Burke, *J. Med. Chem.*, 2011, **54**, 2933–2943.
- 16 O. L. Kobzar, M. V. Shevchuk, A. N. Lyashenko, V. Y. Tanchuk, V. D. Romanenko, S. M. Kobelev, A. D. Averin, I. P. Beletskaya, A. I. Vovk and V. P. Kukhar, *Org. Biomol. Chem.*, 2015, **13**, 7437–7444.
- 17 G. M. Blackburn, D. E. Kent and F. Kolkman, *J. Chem. Soc., Perkin Trans. 1*, 1984, 1119–1125; O. Baszczyński and Z. Janeba, *Med. Res. Rev.*, 2013, **33**, 1304–1344; V. D. Romanenko and V. P. Kukhar, *ARKIVOC*, 2018(i), 1–49.
- 18 S. B. Shears, *Biochem. Soc. Trans.*, 2016, **44**, 191–196.
- 19 R. Pajkert, M. Milewska, G. V. Rösenthaller and H. Koroniak, *J. Fluorine Chem.*, 2009, **130**, 695–701.
- 20 G. M. Blackburn, D. L. Jakeman, A. J. Ivory and M. P. Williamson, *Bioorg. Med. Chem. Lett.*, 1994, **4**, 2573–2578; D. L. Jakeman, A. J. Ivory, M. P. Williamson and G. M. Blackburn, *J. Med. Chem.*, 1998, **41**, 4439–4452; E. Kotsikorou, G. Sahota and E. Oldfield, *J. Med. Chem.*, 2006, **49**, 6692–6703.
- 21 C. Grison, P. Coutrot, C. Comoy, L. Balas, S. Joliez, G. Lavecchia, P. Oligier, B. Penverne, V. Serre and G. Hervé, *Eur. J. Med. Chem.*, 2004, **39**, 333–344.
- 22 A. M. Riley, D. J. Jenkins and B. V. L. Potter, *Carbohydr. Res.*, 1998, **314**, 277–281.
- 23 J. L. Montchamp, F. Tian, M. E. Hart and J. W. Frost, *J. Org. Chem.*, 1996, **61**, 3897–3899; D. K. Baeschlin, A. R. Chaperon, L. G. Green, M. G. Hahn, S. J. Ince and S. V. Ley, *Chem. – Eur. J.*, 2000, **6**, 172–186.
- 24 C. J. Hamilton and S. M. Roberts, *J. Chem. Soc., Perkin Trans. 1*, 1999, 1051–1056.
- 25 H. Wang, J. R. Falck, T. M. Hall and S. B. Shears, *Nat. Chem. Biol.*, 2012, **8**, 111–116.
- 26 A. Ravi, S. Z. Hassan, A. N. Vanikrishna and K. M. Sureshan, *Chem. Commun.*, 2017, **53**, 3971–3973.

