






RESEARCH ARTICLE

REVISED Anesthesia modality does not affect clinical outcomes of intra-arterial vasodilator treatment in patients with symptomatic cerebral vasospasms [version 2; peer review: 2 approved]

Corinne Fischer ¹, Sonja Vulcu¹, Johannes Goldberg¹, Franca Wagner², Belén Rodríguez¹, Nicole Söll¹, Pasquale Mordasini², Matthias Haenggi ³, Joerg C. Schefold ³, Andreas Raabe¹, Werner J. Z'Graggen¹

¹Department of Neurosurgery, Inselspital, University Hospital Bern, Bern, 3010, Switzerland

²University Institute for Diagnostic and Interventional Neuroradiology, Inselspital, University Hospital Bern, Bern, 3010, Switzerland

³Department of Intensive Care Medicine, Inselspital, University Hospital Bern, Bern, 3010, Switzerland

V2 First published: 25 May 2021, 10:417
<https://doi.org/10.12688/f1000research.52324.1>
 Latest published: 02 Aug 2021, 10:417
<https://doi.org/10.12688/f1000research.52324.2>

Abstract

Background: Delayed cerebral ischemia and cerebral vasospasm remain the leading causes of poor outcome in survivors of aneurysmal subarachnoid hemorrhage. Refractory cerebral vasospasms can be treated with endovascular vasodilator therapy, which can either be performed in conscious sedation or general anesthesia. The aim of this study is to compare the effect of the anesthesia modality on long-term clinical outcomes in patients undergoing endovascular vasodilator therapy due to cerebral vasospasm and hypoperfusion.




Methods: Modified Rankin Scale (mRS) scores were retrospectively analyzed at time of discharge from the hospital and six months after aneurysmal subarachnoid hemorrhage. Additionally, National Institutes of Health Stroke Scale (NIHSS) was assessed 24 hours before, immediately before, immediately after, and 24 hours after endovascular vasodilator therapy, and at discharge and six months. Interventional parameters such as duration of intervention, choice and dosage of vasodilator and number of arteries treated were also recorded.


Results: A total of 98 patients were included in this analysis and separated into patients who had interventions in conscious sedation, general anesthesia and a mix of both. Neither mRS at discharge nor at six months showed a significant difference for functionally independent outcomes (mRS 0-2) between groups. NIHSS before endovascular vasodilator therapy was significantly higher in patients

Open Peer Review

Reviewer Status  

Invited Reviewers

	1	2
version 2 (revision) 02 Aug 2021		 report
version 1 25 May 2021	 report	 report

1. **Roland Roelz**, University of Freiburg, Freiburg, Germany
2. **Giovanna Brandi** , University of Zurich, Zurich, Switzerland
Stefan Yu Bögli, University of Zurich, Zurich, Switzerland

Any reports and responses or comments on the article can be found at the end of the article.

receiving interventions in general anesthesia but did not differ anymore between groups six months after the initial bleed.

Conclusion: This study did not observe a difference in outcome whether patients underwent endovascular vasodilator therapy in general anesthesia or conscious sedation for refractory cerebral vasospasms. Hence, the choice should be made for each patient individually.

Keywords

aneurysmal subarachnoid hemorrhage, Nimodipine, Papaverine, delayed cerebral ischemia, general anesthesia, conscious sedation, functional outcome, hypoperfusion

Corresponding author: Werner J. Z'Graggen (werner.zgraggen@insel.ch)

Author roles: **Fischer C:** Formal Analysis, Investigation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Vulcu S:** Data Curation, Writing – Review & Editing; **Goldberg J:** Data Curation, Writing – Review & Editing; **Wagner F:** Data Curation, Writing – Review & Editing; **Rodriguez B:** Formal Analysis, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Söll N:** Data Curation, Writing – Review & Editing; **Mordasini P:** Data Curation, Writing – Review & Editing; **Haenggi M:** Conceptualization, Methodology, Writing – Review & Editing; **Schefold JC:** Conceptualization, Methodology, Writing – Review & Editing; **Raabe A:** Conceptualization, Methodology, Writing – Review & Editing; **Z'Graggen WJ:** Conceptualization, Formal Analysis, Methodology, Project Administration, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Fischer C *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Fischer C, Vulcu S, Goldberg J *et al.* **Anesthesia modality does not affect clinical outcomes of intra-arterial vasodilator treatment in patients with symptomatic cerebral vasospasms [version 2; peer review: 2 approved]** F1000Research 2021, 10:417 <https://doi.org/10.12688/f1000research.52324.2>

First published: 25 May 2021, 10:417 <https://doi.org/10.12688/f1000research.52324.1>

REVISED Amendments from Version 1

To address key issues in the reviewer comments, text was added to the methods section of the paper clarifying how the choice of anesthesia was made in each patient. Additionally, WFNS, Hunt & Hess, BNI and Fisher Scores were statistically analyzed. The corresponding p-values were added to Table 1.

Any further responses from the reviewers can be found at the end of the article

Introduction

Cerebral vasospasms (CVS) and delayed cerebral ischemia still remain among the leading causes of morbidity and mortality in survivors of aneurysmal subarachnoid hemorrhage (aSAH). Up to 40% of aSAH patients experience symptomatic CVS, resulting in disability in up to 50% thereof.¹ CVS, a narrowing of cerebral arteries thought to be caused by blood breakdown products, mostly develop between 5 to 14 days after aSAH.² So far, there is no therapy known, which was shown in randomized trials to improve cerebral perfusion and thus to avoid brain ischemia and infarction in symptomatic patients. Commonly used rescue treatments for symptomatic CVS include induced hypertension, and in refractory CVS angioplasty or intra-arterial application of vasodilators, e.g. nimodipine or papaverine.^{1,3,4} Both have been shown in case series to improve neurological outcome in said patients.⁵⁻⁷

In recent years, several studies investigated the best method of anesthesia for endovascular treatment in acute ischemic stroke.⁸⁻²⁰ While initially in mostly retrospective studies, data showed conscious sedation (CS) to be superior,^{8,9} a recent meta-analysis showed no significant difference in outcomes for CS and general anesthesia (GA)¹⁷ if only randomized controlled trials were considered.^{15,16,18} To the best of our knowledge, no studies comparing CS and GA in endovascular treatments for refractory CVS after aSAH have been performed.

The aim of this study is to compare six-month outcomes for choice of sedation in patients treated with endovascular vasodilators for CVS after aSAH.

Methods

This is a single-center retrospective case-control study analyzing clinical outcomes in patients with symptomatic CVS after aSAH treated with endovascular vasodilators at the University Hospital Bern, Bern, Switzerland.

Study design

The University Hospital Bern conducts a prospective database for patients treated with aSAH. This database was retrospectively searched for patients hospitalized between September 2011 and October 2019. Only patients aged >18 and <85 years were included. Inclusion criteria were: 1) aSAH of all severities (World Federation of Neurosurgeons (WFNS) score I–V), 2) secured aneurysm either by endovascular or surgical treatment, 3) refractory CVS treated by intra-arterial admission of either nimodipine and/or papaverine. Exclusion criteria were: 1) incomplete data, 2) loss of follow up, 3) continuous intra-arterial nimodipine treatment, 4) re-rupture of aneurysm during the hospital stay.

Patients were divided into three treatment groups: patients who underwent treatment with endovascular vasodilators in CS only (“CS”), in GA only (“GA”), or patients who received intra-arterial treatments in CS and GA (“both”). Choice of anesthesia modality was made by the treating physician on an individual basis. However, according to institutional guidelines, GA was preferred in patients with impaired consciousness (GCS ≤ 8) or insufficient swallowing.

Data collection

All data was acquired from patient records and the institutional electronic Patient Data Management System (Centricity™ Critical Care, General Electric Company, GE Healthcare, United States of America). Vital signs are automatically recorded and the bedside team additionally enters clinical scores and administered drugs into the system.

The primary endpoint of this study was functional outcome at six months, analyzed by the modified Rankin Scale (mRS). Secondary outcome parameters included mRS at discharge and National Institutes of Health Stroke Scale (NIHSS) assessed 24 hours before the (first) intra-arterial vasodilator treatment (t_1), directly before (t_2), directly after (last) treatment (t_3), 24 hours after (last) treatment (t_4) and consecutively at discharge from the hospital (t_5) and after six months (t_6). Further parameters consisted of interventional parameters such as duration of intervention, choice of vasodilator (nimodipine or papaverine), number of treated arteries and vasodilator dosage.

Patient characteristics such as age, sex, aneurysm location and treatment and Barrow Neurological Institute (BNI), Fisher, Hunt & Hess and WFNS scores were obtained from institutional patient records.

Statistical analysis

The statistical analysis was performed using SPSS Statistics 21.0 (IBM, Armonk, NY, USA). The Shapiro-Wilk normality test was used to test for normal distribution.

Univariate Analysis of Variance (ANOVA) test was used to compare “CS”, “GA” and “both” groups for differences in age as well as BNI, Fisher, Hunt & Hess and WFNS scores. Differences in sex and aneurysm treatment were explored with Pearson Chi Squared analysis.

For mRS at discharge and six months, outcomes were divided in functionally independent (mRS 0-2) and functionally dependent (mRS 3-6). A Chi Squared test was used to test for significant group differences between CS, GA and both. An additional subgroup analysis was performed using a Chi Squared test with “CS” and “GA” groups divided into single versus multiple interventions, resulting in five groups (“single CS”, “single GA”, “multiple CS”, “multiple GA”, “both”).

For the analysis of NIHSS, a 3×6 analysis of variance (ANOVA) for repeated measures with post hoc Bonferroni correction for multiple comparisons was conducted. The factors were (i) treatment (“CS”, “GA” and “both”) and (ii) time ($t_1 - t_6$).

Interventional parameters were analyzed for each intervention separately and therefore compared between those performed in CS and GA. For the duration of the intervention, a Welch's two sample t-test was performed. The choice of vasodilator was analyzed by Pearson Chi Squared test. Vasodilator dosage as well as number of treated arteries were analyzed with an independent samples t-test.

Data are presented as mean with standard deviation (SD) in brackets and in figures as mean with +1 SD as error bars. A p-value of $p < 0.05$ was considered statistically significant.

Ethics statement

This study was carried out in accordance with the recommendations of the local ethics committee (Kantonale Ethikkommission Bern, Switzerland). All subjects gave written general consent in accordance with the Declaration of Helsinki. The protocol was approved by the local ethics committee (Kantonale Ethikkommission Bern, Switzerland).

Results

In total, 109 patients with refractory CVS treated by intra-arterial admission of either nimodipine and/or papaverine between September 2011 and October 2019 at the University Hospital Bern, Bern, Switzerland were included. Of those, 11 patients had to be excluded. Reasons for exclusion were incomplete data ($n = 2$), re-rupture of aneurysm during the hospital stay ($n = 2$), continuous intra-arterial nimodipine treatment ($n = 1$) and loss of follow-up at six months ($n = 6$). The final study population consisted of 98 patients, 23 patients in the “CS” group, 53 patients in the “GA” group and 22 patients in the “both” group. In the “CS” group, 16 patients received a single intervention (“single CS”) and seven patients received up to five interventions (“multiple CS”). In the GA group, 26 patients received one intervention (“single GA”) and 27 patients received 2-10 treatments (“multiple GA”). As per definition, all patients in the “both” group received more than one and up to 10 interventions.

Patient characteristics

Table 1 shows patient characteristics of the three anesthesia groups. Overall mean age was 54.7 years (range 24-81). All groups showed higher percentages of female patients. Age, sex, aneurysm treatment modality did not significantly differ between the three groups.

Primary outcome

mRS at six months is displayed in **Figure 1a**. There was a tendency for a slightly higher percentage of functionally independent patients (mRS 0-2) in the “CS” group (78.3%) at six months. However, this difference did not prove to be statistically significant ($p = 0.109$). The subgroup analysis comparing single and multiple intra-arterial interventions separately for each anesthesia modality is displayed in **Table 2**. This analysis also revealed no significant difference in functional outcome at six months between “single CS”, “single GA”, “multiple CS”, “multiple GA” and “both” groups ($p = 0.089$).

Secondary outcomes

Figure 1b shows mRS at discharge from hospital. This analysis displays no significant difference between “CS”, “GA” and “both” groups ($p = 0.056$). The subgroup analysis for single and multiple interventions separately also revealed no statistical significance ($p = 0.156$), as listed in **Table 2**.

The NIHSS time course analysis is presented in **Figure 2**. ANOVA for repeated measures displayed a significant interaction of “time*treatment” ($p = 0.008$) and of “time” ($p < 0.001$). Post-hoc analysis revealed that significant group differences only occur when comparing “GA” to the two other groups. All of these significant differences were between t_1

Table 1. Patient characteristics.

	CS	GA	Both	Total	p-value
Number of patients	23	53	22	98	
Mean age (years, range)	57.0 (32 – 74)	54.0 (24 – 81)	53.7 (36 – 72)	54.7 (24 – 81)	0.550 [#]
Sex					0.219 ⁺
Female	20 (87%)	40 (75%)	21 (91%)	80 (82%)	
Male	3 (13%)	13 (25%)	2 (9%)	18 (18%)	
Admission WFNS Score					0.023 ^{**#}
I	12 (52%)	15 (28%)	8 (36%)	35 (36%)	
II	1 (4%)	7 (13%)	9 (41%)	17 (17%)	
III	3 (13%)	5 (9%)	1 (5%)	9 (9%)	
IV	4 (17%)	14 (26%)	4 (18%)	22 (22%)	
V	3 (13%)	12 (23%)	0 (0%)	15 (15%)	
Hunt & Hess Score					0.023 ^{**#}
1	3 (13%)	6 (11%)	4 (18%)	13 (13%)	
2	11 (48%)	16 (30%)	13 (59%)	40 (40%)	
3	3 (13%)	7 (13%)	2 (9%)	12 (12%)	
4	0 (0%)	7 (13%)	1 (5%)	8 (8%)	
5	6 (26%)	17 (32%)	2 (9%)	25 (26%)	
BNI Score					0.999 [#]
1	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
2	7 (30%)	15 (28%)	8 (36%)	30 (31%)	
3	8 (35%)	22 (42%)	7 (32%)	37 (38%)	
4	5 (22%)	8 (15%)	2 (9%)	15 (15%)	
5	3 (13%)	8 (15%)	5 (23%)	16 (16%)	
Fisher score					0.675 [#]
1	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
2	2 (9%)	0 (0%)	0 (0%)	2 (2%)	
3	14 (61%)	38 (72%)	18 (82%)	70 (71%)	
4	7 (30%)	15 (28%)	4 (18%)	26 (27%)	
Aneurysm location					
Choroideal artery	0 (0%)	1 (2%)	0 (0%)	1 (1%)	
ACA	1 (4%)	0 (0%)	0 (0%)	1 (1%)	
ACOM	9 (39%)	18 (34%)	5 (23%)	32 (33%)	
Basilar	2 (9%)	4 (7%)	2 (9%)	8 (8%)	
ICA	1 (4%)	6 (11%)	1 (5%)	8 (8%)	
MCA	5 (21%)	9 (17%)	6 (27%)	20 (20%)	
PCA	1 (4%)	0 (0%)	0 (0%)	1 (1%)	
PCOM	1 (4%)	10 (19%)	7 (32%)	18 (18%)	
A.pericallosa (A2)	2 (9%)	0 (0%)	0 (0%)	2 (2%)	
PICA	0 (0%)	3 (6%)	0 (0%)	3 (3%)	
Superior cerebellar artery	1 (4%)	1 (2%)	0 (0%)	2 (2%)	
Vertebral	0 (0%)	1 (2%)	1 (5%)	2 (2%)	

Table 1. Continued

	CS	GA	Both	Total	p-value
Aneurysm treatment					0.916 [†]
Clipping	5 (21%)	10 (19%)	4 (18%)	19 (19%)	
Coiling	18 (78%)	42 (79%)	18 (82%)	78 (80%)	
Flow diverter	0 (0%)	1 (2%)	0 (0%)	1 (1%)	


Where not stated otherwise, values represent the number of patients with their respective percentages in brackets. WFNS: World Federation of Neurological Surgeons Score; CS: conscious sedation; GA: general anesthesia.

^{*}univariate ANOVA.

[†]Chi-squared test.

^{*}p < 0.05.

a) Modified Rankin Scale at 6 months



b) Modified Rankin Scale at discharge




Figure 1. Distribution of modified Rankin Scale (mRS) categories according to anesthesia modality (Conscious Sedation = “CS”, General Anesthesia = “GA” and Conscious Sedation as well as General Anesthesia = “both”). Numbers represent the percentages for each mRS category per group. a) “CS”, “GA” and “both” at six-month follow up. b) “CS”, “GA” and “both” at discharge from hospital.

Table 2. Subgroup analysis of modified Rankin Scale (mRS) at 6 months and at discharge.

		single CS	multiple CS	single GA	multiple GA	both	p-value
mRS 6 months	mRS 0-2	11 (68.8%)	7 (100%)	16 (61.5%)	12 (44.4%)	14 (63.6%)	0.089 [†]
	mRS 3-6	5 (31.3%)	0 (0%)	10 (38.5%)	15 (55.6%)	8 (36.4%)	
mRS discharge	mRS 0-2	9 (56.3%)	4 (57.1%)	9 (34.6%)	6 (22.2%)	7 (31.8%)	0.156 [†]
	mRS 3-6	7 (43.8%)	3 (42.9%)	17 (65.4%)	21 (77.8%)	15 (68.2%)	

Values represent the number of patients with their respective percentages in brackets. mRS: modified Rankin Scale; CS: conscious sedation; GA: general anesthesia.

[†]Chi-squared test.




Figure 2. Bar graph depicting National Institutes of Health Stroke Scale (NIHSS) scores of the three anesthesia groups. Patients were treated in conscious sedation (“CS”), general anesthesia (“GA”) or a combination of both (“both”). Each bar represents a different group. Each cluster represents a different time point. Time points are 24 hours before (first) intra-arterial vasodilator intervention (t₁), immediately before (t₂), immediately after (t₃), 24 hours after (last) intervention (t₄), at discharge from the hospital (t₅) and at six-month follow up (t₆). Data are presented as mean + 1 Standard Deviation (SD) as error bars. Significant inter-group differences are highlighted with asterisks. One asterisk represents p < 0.05, two asterisks represent p < 0.01 and three asterisks represent p < 0.001.

Table 3. Interventional parameters for intra-arterial vasodilator treatments.

	CS	GA	p-value	Total
Number of interventions	65	172		237
Duration of intervention (min.)	80 (31)	96 (46)	0.002 ^{**+}	92 (44)
Medication (number of interventions)			0.517 ^x	
Nimodipine	59	147		206
Papaverine	3	15		18
Nimodipine and papaverine	2	7		9
Medication dosage (mg)				
Nimodipine	4.8 (1.7)	4.8 (1.7)	0.928 [#]	4.8 (1.7)
Papaverine	200.0 (69)	224.2 (100)	0.696 [#]	212.1 (94)
Nimodipine and papaverine	2.5 (0.07) and 120.0 (0)	3.5 (2) and 123.9 (85)	0.507 [#] and 0.953 [#]	3.0 (1.8) and 121.9 (74)
Number of treated arteries	1.8 (0.6)	2 (0.8)	0.057 [#]	1.9 (0.8)

Where not stated otherwise, values represent the means with the standard deviation in brackets. CS: conscious sedation; GA: general anesthesia.

^{*}Welch’s two sample t-test.

^xChi-squared test.

[#]Independent samples t-test.

and t₅, meaning between 24 hours before (first) intervention and discharge from the hospital. At the six-month follow up appointment, “CS”, “GA” and “both” did not differ significantly regarding NIHSS.

Interventional parameters

Interventional parameters are displayed in Table 3. Overall, a total of 237 intra-arterial vasodilator treatments were performed, 65 of which were performed in CS and 172 in GA. Mean duration of intervention was significantly longer if performed in GA (p = 0.002). A total of four interventions had to be excluded from further analysis because of missing

data about medication (n = 3) and number of treated arteries (n = 1). Neither choice and dosage of intra-arterial vasodilator, nor number of treated arteries showed significant differences between groups.

Discussion

This retrospective study found no significant differences in functionally independent outcomes (mRS 0-2) six months after aSAH in patients who were treated with intra-arterial vasodilators in CS, GA or a combination of both. While NIHSS was significantly higher in patients undergoing endovascular therapy in GA compared to patients of the “CS” or “both” group in the time window 24 hours before intervention up to discharge from the hospital, this difference was no longer found at six months.

To the best of our knowledge, the effect of anesthesia modality (CS versus GA) on functional outcome has not yet been studied for intra-arterial admission of either nimodipine and/or papaverine in patients with refractory CVS after aSAH. Albeit, there have been several research papers published regarding anesthesia in aneurysm treatment.^{21–23} Most articles describe both CS and GA to be generally safe for treatment of unruptured aneurysms or aSAH with no clear recommendation for either one.

Additionally, multiple studies have had similar research questions in relation to the choice of anesthesia during endovascular therapy of acute ischemic stroke. Abou-Chebl *et al.* (2010) showed in their analysis of the “North American SOLITAIRE Stent-Retriever Acute Stroke” (NASA) registry, that patients treated in GA experienced poorer neurologic outcome at 90 days and higher mortality rates than patients treated in CS.⁸ Berkhemer *et al.* (2016) reported similar results in a post-hoc analysis of a prospective trial.¹⁵ Correspondingly, a recent analysis of the “Endovascular Therapy Following Imaging Evaluation for Ischemic Stroke 3” (DEFUSE 3) trial by Powers *et al.* (2019) showed higher rates of functional independence (mRS 0-2) and a lower NIHSS score at 24 hours for patients treated in CS. At discharge they did not find a statistically significant difference in NIHSS scores anymore.¹¹

In contrast, most recent randomized controlled trials found a non-inferiority of GA when compared to CS. Hendén *et al.* (2017) reported no difference in outcome at three months or NIHSS after 24 hours in their Anesthesia During Stroke (AnStroke) Trial.¹⁶ Schönerberger *et al.* (2016) and Simonsen *et al.* (2018) report, that GA produced better 3-month outcomes, with the former even finding this result to be significant.^{18,19} Finally, a recent meta-analysis determined no significant difference between GA and CS if only randomized controlled trials were considered.¹⁷

Our results are in line with these recent randomized controlled trials published for endovascular treatment in acute ischemic stroke. Similar to Schönerberger *et al.* (2016), Hendén *et al.* (2017), Simonsen *et al.* (2018) and Kim *et al.* (2019), we also found no significant difference in functional independency (mRS 0-2) at discharge or six months.^{16–19}

Similar considerations regarding choice of anesthesia hold true in intra-arterial vasodilator therapy after aSAH and treatment for acute ischemic stroke alike. Disadvantages of GA may be a delay in treatment, hemodynamic changes and complications associated with intubation such as an increased risk for pneumonia.⁸ Disadvantages of CS may be procedural discomfort for patients, more difficult interventions because of patients’ movements, emergency conversion to GA or increased risk for aspiration.⁸ Many of the conceived disadvantages of GA have also been analyzed in prospective studies concerning ischemic stroke treatment. For example, Berkhemer *et al.* (2016) as well as Hendén *et al.* (2017) found no treatment delay in the GA group.^{15,16} Schönerberger *et al.* (2016) reported no difference in feasibility, safety and intra-interventional complication between the groups.¹⁸ They did however discover more postprocedural complications after GA such as delayed extubation and pneumonia. Different authors mention the possibility of blood pressure drops and decreased cerebral blood flow as possible additional complications of GA, which could potentially worsen outcomes because of an increase of the ischemic area.¹⁵ Others argue that GA on its own has a neuroprotective effect by lowering the neuronal oxygen need.^{24,25} Overall, the similar functional outcome in our study as well as in the above cited studies in ischemic stroke suggest that these factors are of minor relevance.

The major limitations of this study are its retrospective design and the single-center approach. A potential bias could lie in the choice of sedation for treatment. Even before intra-arterial vasodilator treatment, patients who would go on to receive treatments in GA showed significantly higher NIHSS scores when compared to the “CS” and “both” groups. This indicates that patients in clinically and neurologically worse conditions were more likely to be treated in GA. However, this bias was not reflected in our results, as we did not find a significant difference in mRS scores between the anesthesia groups at discharge or at six months and also no significant difference in NIHSS scores at six months. The “GA” group were therefore in an initially worse state but still managed to reach similar outcomes in the long-term clinical course. This suggests to an even greater degree, that GA will not negatively affect long term outcome in these patients.

Furthermore, some subgroups consisted of a small number of patients. The results of this study will have to be replicated by a larger prospective trial in the future.

Conclusion

Our preliminary results indicate that choice of anesthesia method does not negatively affect six-month outcome in aSAH patients who undergo intra-arterial vasodilator treatment for CVS. Treating physicians should therefore decide between CS and GA individually based on patient characteristics and circumstances.

Data availability

Underlying data

Dryad: Functional Outcome after intraarterial vasodilator therapy in CS vs GA, <https://doi.org/10.5061/dryad.g4f4qrfq5>.²⁶

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

References

1. Baggott CD, Aagaard-Kienitz B: **Cerebral vasospasm**. *Neurosurg Clin N Am*. 2014; **25**: 497–528.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. van Gijn J, Kerr RS, Rinkel GJ: **Subarachnoid haemorrhage**. *Lancet*. 2007; **369**: 306–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Cho WS, Kang HS, Kim JE, et al.: **Intra-arterial nimodipine infusion for cerebral vasospasm in patients with aneurysmal subarachnoid hemorrhage**. *Interv Neuroradiol*. 2011; **17**: 169–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Labeyrie MA, Gaugain S, Boulouis G, et al.: **Distal balloon angioplasty of cerebral vasospasm decreases the risk of delayed cerebral infarction**. *Am J Neuroradiol*. 2019; **40**: 1342–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Dabus G, Nogueira RG: **Current Options for the Management of Aneurysmal Subarachnoid Hemorrhage-Induced Cerebral Vasospasm: A Comprehensive Review of the Literature**. *Interv Neurol*. 2014; **2**: 30–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Andereggen L, Beck J, Z'Graggen WJ, et al.: **Feasibility and safety of repeat instant endovascular interventions in patients with refractory cerebral vasospasms**. *Am J Neuroradiol*. 2017; **38**: 561–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Hänggi D, Turowski B, Beseoglu K, et al.: **Intra-arterial nimodipine for severe cerebral vasospasm after aneurysmal subarachnoid hemorrhage: Influence on clinical course and cerebral perfusion**. *Am J Neuroradiol*. 2008; **29**: 1053–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Abou-Chebl A, Lin R, Hussain MS, et al.: **Conscious Sedation Versus General Anesthesia During Endovascular Therapy for Acute Anterior Circulation Stroke**. *Stroke*. 2010; **41**: 1175–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Jumaa MA, Zhang F, Ruiz-Ares G, et al.: **Comparison of safety and clinical and radiographic outcomes in endovascular acute stroke therapy for proximal middle cerebral artery occlusion with intubation and general anesthesia versus the nonintubated state**. *Stroke*. 2010; **41**: 1180–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Nichols C, Carrozella J, Yeatts S, et al.: **Is periprocedural sedation during acute stroke therapy associated with poorer functional outcomes?** *J Neurointerv Surg*. 2010; **2**: 67–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Powers CJ, Dornbos D, Mlynash M, et al.: **Thrombectomy with conscious sedation compared with general anesthesia: A DEFUSE 3 analysis**. *Am J Neuroradiol*. 2019; **40**: 1001–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Takahashi C, Liang CW, Liebeskind DS, et al.: **To tube or not to tube? The role of intubation during stroke thrombectomy**. *Front Neurol*. 2014; **5** AUG.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Talke PO, Sharma D, Heyer EJ, et al.: **Republished: Society for neuroscience in anesthesiology and critical care expert consensus statement: Anesthetic management of endovascular treatment for acute ischemic Stroke***. *Stroke*. 2014; **45**.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Molina CA, Selim MH: **General or local anesthesia during endovascular procedures: Sailing quiet in the darkness or fast under a daylight storm**. *Stroke*. 2010; **41**: 2720–1.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Berkhemer OA, Van Den Berg LA, Franssen PSS, et al.: **The effect of anesthetic management during intra-Arterial therapy for acute stroke in MR CLEAN**. *Neurology*. 2016; **87**: 656–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Hendén PL, Rentzos A, Karlsson JE, et al.: **General Anesthesia Versus Conscious Sedation for Endovascular Treatment of Acute Ischemic Stroke: The AnStroke Trial (Anesthesia during Stroke)**. *Stroke*. 2017; **48**: 1601–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Kim C, Kim S-E, Jeon JP: **Influence of Anesthesia Type on Outcomes after Endovascular Treatment in Acute Ischemic Stroke: Meta-Analysis**. *Neurointervention*. 2019; **14**: 17–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Schönenberger S, Uhlmann L, Hacke W, et al.: **Effect of conscious sedation vs general anesthesia on early neurological improvement among patients with ischemic stroke undergoing endovascular thrombectomy: A randomized clinical trial**. *JAMA - J Am Med Assoc*. 2016; **316**: 1986–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Simonsen CZ, Yoo AJ, Sørensen LH, et al.: **Effect of general anesthesia and conscious sedation during endovascular therapy on infarct growth and clinical outcomes in acute ischemic stroke a randomized clinical trial**. *JAMA Neurol*. 2018; **75**: 470–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Froehler MT, Fifi JT, Majid A, et al.: **Anesthesia for endovascular treatment of acute ischemic stroke**. *Neurology*. 2012; **79**: S167 LP–S173.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Kan P, Jahshan S, Yashar P, et al.: **Feasibility, safety, and periprocedural complications associated with endovascular treatment of selected ruptured aneurysms under conscious sedation and local anesthesia**. *Neurosurgery*. 2013; **72**: 216–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Lee CZ, Young WL: **Anesthesia for Endovascular Neurosurgery and Interventional Neuroradiology**. *Anesthesiol. Clin*. 2012; **30**: 127–147.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Ogilvy CS, Yang X, Jamil OA, et al.: **Neurointerventional procedures for unruptured intracranial aneurysms under procedural sedation and local anesthesia: A large-volume, single-center experience - Clinical article**. *J Neurosurg*. 2011; **114**: 120–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Hoffmann U, Sheng H, Ayata C, et al.: **Anesthesia in Experimental Stroke Research**. *Transl Stroke Res*. 2016; **7**: 358–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Sivasankar C, Stiefel M, Miano TA, et al.: **Anesthetic variation and potential impact of anesthetics used during endovascular management of acute ischemic stroke**. *J Neurointerv Surg*. 2016; **8**: 1101–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Fischer, et al.: **Functional Outcome after intraarterial vasodilator therapy in CS vs GA**. *Dryad [dataset]*. 2021.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 09 August 2021

<https://doi.org/10.5256/f1000research.59014.r91020>

© 2021 Brandi G et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Giovanna Brandi 

Institute of Intensive Medicine, Zurich University Hospital, University of Zurich, Zurich, Switzerland

Stefan Yu Bögli

Institute of Intensive Medicine, Zurich University Hospital, University of Zurich, Zurich, Switzerland

I thank the authors for the changes they made in the revised manuscript and for the answers to our comments.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Neurocritical care

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 13 July 2021

<https://doi.org/10.5256/f1000research.55588.r88709>

© 2021 Brandi G et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Giovanna Brandi 

Institute of Intensive Medicine, Zurich University Hospital, University of Zurich, Zurich, Switzerland

Stefan Yu Bögli

Institute of Intensive Medicine, Zurich University Hospital, University of Zurich, Zurich, Switzerland

This is a retrospective study.

How were the NIHSS extracted from the patients files? Is NIHSS performed routinely? If the patient was already intubated, how was NIHSS evaluated?

While aneurysma location as well as Fisher score seem to be similar, WFNS/ H&H seem to differ between CS + GA (e.g. 52% vs 28% WFNS 1). Is there a statistical difference? If so, how do you explain the lack of association thereof with the outcome?

Patients with high WFNS + Fisher scores commonly remain intubated in case of unsuccessful sedation-reduction trial. How many patients were already under GA before intra-arterial vasodilatation. Is there difference in delay between non-intubated patients to CS or GA intra-arterial vasodilatation?

When mRS at 6 months is compared between CS and GA only (excluding both) irrespective of if they received a singular or multiple CS/GA statistical significance is reached. How do you explain this finding?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Neurocritical care

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Author Response 24 Jul 2021

Corinne Fischer, Inselspital, University Hospital Bern, Bern, Switzerland

In our institution, NIHSS is routinely assessed in stroke patients at least once a day as well as before and after interventions. In intubated patients, the NIHSS is determined after depth of sedation is decreased as much as clinically possible. The NIHSS assessment is performed according to the recommendation of the National Institute of Neurological Disorders and Stroke (https://www.ninds.nih.gov/sites/default/files/NIH_Stroke_Scale_Booklet.pdf). Most notably, intubated patients get a default score of 1 for verbally assessed categories such as orientation if they are unable to speak or write. Additionally, aphasia is assessed through writing. Dysarthria is not assessable in the intubated patient.

The p-values for WFNS, Hunt&Hess, BNI and Fisher scores were added to Table 1. Only the WFNS and Hunt&Hess scores differed significantly between the groups. However, as it has been shown in two earlier publications cited below, WFNS 5 scores at admission are often over-estimated due to accompanying factors e.g. early seizures. In addition, if factors such as intubation, ventilation, sedation, muscle relaxation or insufficient pain stimuli inhibit said motor response, the WFNS grading can be over-estimated while the outcome is not necessarily impacted. Similar effects can alter Hunt&Hess gradings. In contrast the radiologically determined scores (BNI and Fisher) are not influenced by the above factors, which can at least partially explain that these two scores are not significantly different while the clinical ones are. (Fung *et al.*, 2016; Fung *et al.*, 2015).

In addition, more severely ill patients corresponding to patients with higher WFNS and Hunt&Hess scores are more probable to undergo intra-arterial vasodilator therapy in general anesthesia (see also below.)

Twenty-two patients were already ventilated before the first intra-arterial vasodilator treatment (this corresponds to 37% of all patients who were intubated for the first intra-arterial vasodilator treatment). Unfortunately, due to the retrospective study design, the additional delay of intra-arterial vasodilatation therapy due to intubation was not assessable within the obtained documentation.

Considering only mRS at 6 months and reducing the analysis to patients who were only treated with one modality (CS or GA), the outcome differed significantly between the two groups (Fisher exact $p = 0.044$).

In our opinion this analysis has to be interpreted with caution as the number of patients undergoing intra-arterial vasodilator treatment in CS who showed an unfavorable outcome is very small ($n = 5$). Furthermore, if the above mentioned patients who remained intubated after the initial bleeding are excluded from this analysis, the result is no longer statistically significant ($p = 1.000$). This indicates that the difference in outcome of this additional analysis is probably mainly due to the initial disease severity and not due to the anesthesia modality chosen for intra-arterial vasodilator treatment.

Competing Interests: No competing interests were disclosed.

Reviewer Report 13 July 2021

<https://doi.org/10.5256/f1000research.55588.r88125>

© 2021 Roelz R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Roland Roelz

Department of Neurosurgery, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany

This is an interesting preliminary study on the anesthesia modality for endovascular rescue therapies in patients with aneurysmal subarachnoid hemorrhage (aSAH). This single-center retrospective study included 98 aSAH patients (admitted in an 8-year period) selected for endovascular vasospasm therapy. Circa half of these patients were treated in general anesthesia, 25% were treated in conscious sedation and 25% had multiple interventions and both anesthesia modalities were applied. No differences in outcome (6-months mRS) were observed between these groups.

Carefully interpreted and fully aware of the limitations associated with the study design and data collection, this study may indicate towards safety and non-inferiority of both general anesthesia and conscious sedation for endovascular interventions after aSAH. These findings are in keeping with the pertinent literature on anesthesiologic management of endovascular interventions for stroke.

I thank the authors for sharing their first experience with conscious sedation as an alternative to general anesthesia for endovascular vasospasm therapy. I encourage investigating the clinical algorithm that triggers the choice for either method. 50% of patients treated in general anesthesia had admission WFNS grades 4 or 5 (i.e. poor grade aSAH) compared to 30% of patients selected for conscious sedation. A fact that may very well explain the presumed tendency towards better outcomes of the latter group. The rate of delayed infarction should have been reported. Given that safety of conscious sedation seems to apply to endovascular vasospasm therapy, future studies are warranted. They should follow a prospective and randomized design. Potential benefits of conscious sedation (i.e. more rapid return to neurological assessability after intervention, less complications of mechanical ventilation etc.) might be of high clinical relevance and could improve outcome of patients who develop clinical vasospasm.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Neurosurgery

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 24 Jul 2021

Corinne Fischer, Inselspital, University Hospital Bern, Bern, Switzerland

The following text was added to the methods section of the manuscript: "Choice of anesthesia modality was made by the treating physician on an individual basis. However, according to institutional guidelines, GA was preferred in patients with impaired consciousness (GCS \leq 8) or insufficient swallowing."

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research