

The plasmamembrane calmodulin-dependent calcium pump: a major regulator of nitric oxide synthase I

Kai Schuh,¹ Stjepan Uldrijan,² Myriam Telkamp,¹ Nicola Röhlein,¹ and Ludwig Neyses³

¹Department of Medicine, University of Wuerzburg, D-97080 Wuerzburg, Germany

²Masaryk Memorial Cancer Institute, 656 53 Brno, Czech Republic

³Department of Medicine, University of Manchester, M13 9WL Manchester, United Kingdom

The plasma membrane calcium/calmodulin-dependent calcium ATPase (PMCA) (Shull, G.E., and J. Greeb. 1988. *J. Biol. Chem.* 263:8646–8657; Verma, A.K., A.G. Filoteo, D.R. Stanford, E.D. Wieben, J.T. Penniston, E.E. Strehler, R. Fischer, R. Heim, G. Vogel, S. Mathews, et al. 1988. *J. Biol. Chem.* 263:14152–14159; Carafoli, E. 1997. *Basic Res. Cardiol.* 92:59–61) has been proposed to be a regulator of calcium homeostasis and signal transduction networks of the cell. However, little is known about its precise mechanisms of action. Knock-out of (mainly neuronal) isoform 2 of the enzyme resulted in hearing loss and balance deficits due to severe inner ear defects, affecting formation and maintenance of otoconia (Kozel, P.J., R.A. Friedman, L.C. Erway, E.N. Yamoah, L.H. Liu, T. Riddle, J.J. Duffy, T. Doetschman, M.L. Miller, E.L. Cardell, and G.E. Shull. 1998. *J. Biol. Chem.* 273:18693–18696). Here we

demonstrate that PMCA 4b is a negative regulator of nitric oxide synthase I (NOS-I, nNOS) in HEK293 embryonic kidney and neuro-2a neuroblastoma cell models. Binding of PMCA 4b to NOS-I was mediated by interaction of the COOH-terminal amino acids of PMCA 4b and the PDZ domain of NOS-I (PDZ: PSD 95/Dlg/ZO-1 protein domain). Increasing expression of wild-type PMCA 4b (but not PMCA mutants unable to bind PDZ domains or devoid of Ca²⁺-transporting activity) dramatically downregulated NO synthesis from wild-type NOS-I. A NOS-I mutant lacking the PDZ domain was not regulated by PMCA, demonstrating the specific nature of the PMCA–NOS-I interaction. Elucidation of PMCA as an interaction partner and major regulator of NOS-I provides evidence for a new dimension of integration between calcium and NO signaling pathways.

Introduction

In initial attempts to gain insight into the unknown regulatory function of the plasma membrane calcium/calmodulin-dependent calcium ATPase (PMCA)* (Carafoli, 1997; Shull and Greeb, 1988; Verma et al., 1988), we have shown previously that the PMCA is controlled by β -adrenergic stimulation (Neyses et al., 1985), that it regulated differentiation in a skeletal muscle cell line (Hammes et al., 1994, 1996), modulated hypertrophic stimuli in the myocardium of transgenic animals expressing PMCA under the control of the ventricle-specific myosin light chain-2 promoter, and was not involved in beat-to-beat regulation of heart contraction (Hammes et al., 1998). Furthermore, we and others have shown that PMCA 4 is local-

ized to caveolae (Fujimoto, 1993; Hammes et al., 1998), membrane structures (sometimes also called “signalosomes” or microdomains) containing a wide variety of effectors of cellular signal transduction (Sargiacomo et al., 1993; Chang et al., 1994; Lisanti et al., 1994; Robbins et al., 1995; Schnitzer et al., 1995; Song et al., 1996a) including nitric oxide synthase (NOS)-I (Venema et al., 1997; Segal et al., 1999) and the cytoskeletal protein dystrophin (Song et al., 1996b). Because PMCA 4b has recently been shown to bind to members of the membrane-associated guanylate kinase protein family via their PDZ domains (Kim et al., 1998), we tested whether it interacts with NOS-I, which has been demonstrated recently to be part of a multiprotein complex largely organized by PDZ domains (through direct interaction or via adapter proteins; Brenman et al., 1995, 1996; Hillier et al., 1999) and whether this interaction was physiologically relevant.

Address correspondence to Ludwig Neyses, University Department of Medicine, Manchester Royal Infirmary, Oxford Road, M13 9WL Manchester, UK. Tel.: (44) 161-276-6631. Fax: (44) 161-276-5152. E-mail: ludwig.neyses@mhc.cmht.nwest.nhs.uk

*Abbreviations used in this paper: GST, glutathione *S*-transferase; NO, nitric oxide; NOS, NO synthase; PMCA, plasma membrane calcium/calmodulin-dependent calcium ATPase.


Key words: PMCA; NOS; calcium; nitric oxide; PDZ domain

Results and discussion

Physical interaction of the PMCA and NOS-I was demonstrated by conventional immunoprecipitation (Fig. 1, D and

Figure 1. Schematic overview of constructs used, expression of NOS-I constructs in HEK293 cells, and physical interaction of PMCA with NOS-I. (A)

Plasmid constructs used in transfection assays. pCMV-PMCA 4b contains the wild-type human PMCA 4b cDNA under the control of the CMV promoter. pCMV-PMCA 4bmut carries a mutation Asp⁶⁷²→Glu, thereby reducing the ATPase activity to ~10% (Adamo et al., 1995). In pCMV-hPMCA 4b cap the stop codon is mutated and the myc/6×His tag is added in frame, resulting in capping of the PDZ-binding COOH-terminal amino acids. NOS-I and NOS-III are also under the control of the CMV promoter, in pCMV-ΔNOS-I the interacting PDZ domain is deleted. (B) NOS-I was expressed in pCMV-NOS-I-transfected HEK293 cells, rat brain homogenate served as control. ΔNOS-I was expressed at comparable levels, the size difference is due to deletion of the PDZ domain. In untransfected HEK cells, NOS-I was undetectable. (C) GST pull-down assay (Brennan et al., 1995) demonstrating specific interaction of PMCA 4b with PDZ domain of NOS-I. Lysate of pCMV-hPMCA4b-transfected HEK293 cells was used as expression control (first lane), GST alone did not bind PMCA (second lane), whereas GST-PDZ fusion protein was able to bind PMCA 4b (last lane). (D) NOS-I-PMCA 4b interaction in cotransfected HEK293 cells. In the first lane NOS-I coprecipitation by a hPMCA 4b-specific antibody (JA3) from lysates of cotransfected cells is shown. In the third lane, lysates of NOS-I/PMCA-transfected cells were used as expression controls. Irrelevant antibody (anti-FLAG) or protein G beads alone showed no binding of NOS-I (second and fourth lane). Cotransfection with pCMV-hPMCA 4b cap, containing an additional COOH-terminal Myc/6×His epitope, prevented interaction of PMCA and NOS-I (penultimate lane), although NOS-I was present in the cells (last lane). (E) Coimmunoprecipitation using NOS-I-specific antibodies. The NOS-I-specific polyclonal antibody (ABR) coprecipitated the PMCA from lysates of cotransfected HEK293 cells (lane 4). Negative controls: unlabeled beads (lane 1), beads plus irrelevant antibodies (lane 2). Positive controls: calmodulin-coated beads (CaM beads, lane 3), anti-PMCA-specific coated beads (with polyclonal antibody "2A," lane 5, and lysates of hPMCA 4b-transfected HEK293 cells (lane 6). (F) Immunoprecipitation using rat brain extracts and different PMCA-specific antibodies demonstrating interaction of PMCA and NOS-I in brain. Antibodies specific for NOS-I (lane 3), PMCA (monoclonal, clone 5F10, lane 4, rabbit polyclonal "2A", lane 5), as well as calmodulin beads (CaM beads, lane 2) were capable of precipitating NOS-I from rat brain extracts. First lane, control using irrelevant antibodies; last lane, rat brain extract (input).



E) as well as glutathione *S*-transferase (GST) pull-down assays, using the PDZ domain of NOS-I as a bait (Fig. 1 C). "Capping" the PMCA 4b COOH terminus (carrying the PDZ domain binding sequence) with an myc/6×His tag resulted in a loss of physical interaction (Fig. 1 D). Expression of wild-type NOS-I and deletion constructs (overview in Fig. 1 A) was confirmed by Western blotting (Fig. 1 B). This established that PMCA 4b and NOS-I interact physically, either directly or through unknown adapters. Full-length PMCA 4b interacted with purified GST-PDZ fusion protein, directly demonstrating specificity of interaction with the PDZ domain (Fig. 1 C). Similar interaction of PMCA and NOS-I was also observed in extracts of rat brain. Using PMCA-specific antibodies 5F10 and 2A, NOS-I was also coprecipitated from rat brain extracts (Fig. 1 F).

In a simple working model, a PMCA molecule interacting with NOS-I might reduce the local calcium concentration, leading to reduced activity of NOS-I, a calcium/calmodulin-dependent isoform of NOS. This hypothesis was tested us-

ing HEK293 cells transfected with expression constructs shown schematically in Fig. 1 A. Increasing PMCA 4b expression reduced NOS-I-dependent cGMP dramatically in a dose-dependent manner (Fig. 2 A, black columns). The model also predicts that it is the calcium transport function of the PMCA rather than some other yet unknown module of the molecule that regulates NOS-I activity: Asp⁶⁷²→Glu point mutation has been shown previously to render PMCA almost (but not totally) inactive with respect to its ATPase and Ca²⁺-transporting activity (reduction of calcium transport activity by ~90%; Adamo et al., 1995). As predicted, this mutation resulted in a strong reduction of inhibition of NOS-I (Fig. 2 A, grey columns) compared with wild-type PMCA cotransfections (Fig. 2 A, black columns).

To test whether binding of PMCA 4b to the complex via PDZ domains was a prerequisite for its regulatory action, a constitutively active mutant of the pump ("ct120") with a deletion of both the autoinhibitory and the COOH-terminal PDZ domain binding motif (Enyedi et al., 1993) was




Figure 2. Functional interaction of PMCA 4b and NOS-I.

(A) Dose-dependent inhibition of NOS activity by PMCA 4b. Background cGMP production in untransfected HEK293 cells was set as one (first column), induced cGMP production is shown as fold increase of cGMP levels in relation to protein content of cell lysates. PMCA overexpression showed no effect on cGMP production (second column), whereas NOS-I expression resulted in an ~60-fold increase in cGMP levels (third column). Increasing amounts of PMCA led to a dose-dependent reduction of NOS-I activity, insertion of the point mutation Asp⁶⁷²→Glu in the PMCA reduced inhibition dramatically (PMCAmut, grey columns). The ~10-fold higher amounts of mutant PMCA necessary to inhibit NOS-I are in line with the fact that the mutation abolishes ~90% of Ca²⁺ transport activity (Adamo et al., 1995). Transfection with pPMCAmut alone had no effect on cGMP production (column 12), treatment of NOS-I transfected cells with L-NAME (L-ω-nitro-L-arginine methyl ester; 1 mM) resulted in the expected complete inhibition of cGMP production (penultimate column). Coexpression of the constitutively active mutant of PMCA 4b ("ct120", last column) did not result in inhibition of NOS-I ($n = 16$, mean \pm SEM, asterisk indicates $P < 0.05$ PMCA vs. PMCAmut). Representative Western blots demonstrated expression of relevant proteins: antibody JA3 showed expression of hPMCA4b and hPMCA4bmut, antibody 5F10, specific for a more NH₂-terminal epitope of PMCA, demonstrated expression of PMCA4b(ct120), parallel to constant NOS-I expression in cotransfected cells. (B) Deletion of PDZ domain of NOS-I (Δ NOS-I) results in comparable NOS-I activity and complete loss of regulation by increasing amounts of wild-type hPMCA 4b ($n = 16$, mean \pm SEM, changes in fold induction not significant). (C) NOS-III expression results in a highly increased production of cGMP, too, but this NO-dependent cGMP production was not inhibited by wild-type PMCA. The NOS inhibitor L-NAME (L-N) abolished cGMP production, proving the NOS-III-dependent cGMP production ($n = 2 \times 8$, mean \pm SEM, changes in fold induction not significant).




Figure 3. (A) Dose-dependent inhibition of NOS-I activity in neuro-2a cells. Coexpression of increasing amounts of PMCA 4b and constant levels of NOS-I resulted in a dose-dependent inhibition of NOS-I activity, comparable to the effects observed in HEK293 cells. In this cellular system smaller amounts of PMCA (~0.5 μ g transfected plasmid) were sufficient to obtain maximum inhibition of NOS-I, suggesting that an upper limit of PMCA expression is reached earlier in this cellular system ($n = 10$, mean \pm SEM, asterisk indicates $P < 0.01$). (B) Representative Western blot demonstrating constant NOS-I expression despite dose-dependent expression of PMCA 4b in transfected neuro-2a cells.

cotransfected with NOS-I. No regulation of NOS activity by this construct was observed (Fig. 2 A, last column). An NOS-I mutant carrying a deletion of the PDZ domain showed no regulation by PMCA 4b (Fig. 2 B). Endothelial NOS (eNOS or NOS-III) is also regulated by calcium/calmodulin, but carries no PDZ domain, and we have been unable to coprecipitate NOS-III with PMCA (not shown). In keeping with the absence of a physical interaction between PMCA 4b and NOS-III, the activity of this enzyme was not regulated by PMCA 4b (Fig. 2 C).

The potential physiological relevance of NOS-I regulation by PMCA 4b was tested in neuro-2a neuroblastoma cells, a commonly used model system in neuronal biology (Olmsted et al., 1970). Similarly to HEK293 cells, tight functional coupling of PMCA4b and NOS-I was observed: NOS-I activity was strongly downregulated by the PMCA4b (Fig. 3 A). This effect was reversed by the NO donor NOC-18 (2,2'-[hydroxynitrosohydrazino]bis-ethanamine), suggesting

that interaction of these proteins not only occurs in HEK293 cells, but also in a neuroblastoma-derived cell line.

These results show that the plasma membrane calmodulin-dependent calcium pump 4b is an interaction partner and a major regulator of neuronal NOS-I and also that this regulation very likely is of physiological relevance. The strategic localization to caveolae (Fujimoto, 1993; Hammes et al., 1998) also suggests that local control of calcium and/or NO might have further regulatory effects in caveolae-mediated signal transduction.

The pivotal role of NOS-I in neuronal tissue is well established, exemplified by the observation that NOS-I deficiency leads to reduced susceptibility to cerebral ischemic damage (Huang et al., 1994). Its function in other excitable cells has been extensively characterized in recent years (Christopherson and Brecht, 1997). In contrast, the specific role of the PMCA, beyond the general concept of a calcium transporter, has been elusive. Our present results assign a function to isoform PMCA 4b, i.e., regulation of NOS-I activity. In a simple model (see Fig. 4), NOS-I, a Ca^{2+} /calmodulin-dependent isoform of NOS, is regulated by the PMCA through adjustment of local calcium concentration. Both interact either directly or through an adapter protein via the PDZ domain of NOS-I and the COOH-terminal amino acids of PMCA 4b. Our results are in line with two previous reports showing the interaction of COOH-termini of different PMCA "b" splice variants (PMCA 2b and 4b) with PDZ domains and PDZ domain containing proteins of the MAGUK (membrane-associated guanylate kinase) protein family, whereas the splice variants 2a and 4a failed to bind to PDZ domains (Kim et al., 1998; DeMarco and Strehler, 2001). Indeed, it is conceivable that the predominantly neuronal isoform 2b regulates NOS-I in the context of (at least some specialized) neurones in vivo. This important issue will need to be addressed in further studies. These results and the reduced inhibition of NOS-I activity by the mutated PMCA (PMCAmut, Fig. 2, in which the Ca^{2+} -transporting and ATPase activity are reduced) strongly suggest that it is the calcium transport function of the PMCA which affects regulation of NOS-I. However, it cannot be entirely excluded that the COOH-terminal peptide of the PMCA 4b, interacting with the PDZ domain of NOS-I, mediates inhibition directly; in this case one would have to postulate that the PMCAmut (Asp⁶⁷² mutation) undergoes a conformational change conducive to reduced interaction with NOS-I. A definitive answer to this and other unlikely possibilities may have to await direct measurements of calcium concentrations in the vicinity of the pump which is currently not possible.

The coupling of interaction and function of the two proteins is in close analogy to the observation that loss of the PDZ-binding COOH-termini of NMDA receptors results not only in loss of interaction, but also in complete loss of receptor function (Sprengel et al., 1998). Different PMCA isoforms and their splice variants (overview in Strehler and Zacharias, 2001) are expressed ubiquitously and NOS-I is a widespread isoform of NOS in neurons and in excitable tissues (Christopherson and Brecht, 1997). Therefore, the physical and functional interaction of PMCA isoforms and NOS-I or other PDZ domain-containing proteins is most likely relevant in a large variety of cells. These enzymes may define a modular signal trans-




Figure 4. **Model for the regulation of NOS-I by PMCA 4b.** NO production by NOS-I is reduced through Ca^{2+} extrusion via the PMCA, resulting in alteration of downstream signaling.

duction system which subserves different functions in different cellular contexts.

Materials and methods

Immunoprecipitations and GST pull-downs

Samples were lysed in RIPA buffer (1× PBS containing 1% Igepal CA-630, 0.5% sodium deoxycholate, 0.1% SDS and Complete™ protease inhibitor (Roche Diagnostics). Debris was removed by centrifugation (10,000 g, 5 min, 4°C). After a preclearing step with 0.5 μg irrelevant antibody (anti-FLAG epitope, Neomarkers) and 25 μl of protein G-Sepharose™ (~50% slurry; Amersham Pharmacia Biotech) 500 μg remaining protein lysate was coincubated together with 3 μg of precipitating antibodies (polyclonal rabbit anti-NOS-I, Affinity Bioreagents catalog no. PA3-032A; monoclonal mouse anti-hPMCA4b, clone JA3; and monoclonal mouse anti-PMCA, clone 5F10, Neomarkers, catalog no. MS-1305-P1 and MS-1304-P1; polyclonal rabbit anti-hPMCA4b was generated using a peptide spanning the 15 COOH-terminal amino acids of the PMCA4b; polyclonal rabbit anti-PMCA ("2A") was a gift from D. Guerini, ETH Zürich, Switzerland) and 25 μl protein G-Sepharose™ at 4°C for 2 h. Calmodulin-Sepharose beads (CaM beads; Amersham Pharmacia Biotech) served as control. After washing four times with RIPA buffer (see above) beads were resuspended in 100 μl sample buffer and boiled. 20 μl of each sample was loaded and 5 μg (1/100 input) protein extract served as control. NOS-I was detected using a monoclonal antibody (clone "16"; Transduction Laboratories, catalog no. N31020) in a dilution of 1:1,000, and transfected hPMCA4b was detected using the PMCA 4 isoform-specific antibodies JA3 (1:1,000; Figs. 1, C and E, 2 A, and 3 B) and 5F10 (dilution 1:1,000, Neomarkers), necessary to detect appropriate expression of PMCA(ct120). Immunodetection of proteins on membrane was done using the ECL™ system according to the manufacturers protocol (Amersham Pharmacia Biotech).

GST pull-down assays were performed using purified GST-PDZ (NOS-I) fusion protein, containing the PDZ domain of NOS-I (Brenman et al., 1995, 1996) bound to glutathione-Sepharose™ 4B beads (Amersham Pharmacia Biotech). Pelleted beads were washed four times with RIPA buffer, resuspended in electrophoresis sample buffer, boiled for 3 min, and proteins were separated by 10% SDS-PAGE and blotted to Hybond™ ECL™ nitrocellulose membrane (Amersham Pharmacia Biotech). Immunodetection was performed as described above.

Construction of DNA vectors

Expression vector pCMV-hPMCA 4b has been described previously (pCMV-hPMCA4CI; Hammes et al., 1996). The point mutation Asp⁶⁷²→Glu was inserted with the QuikChange™ site-directed mutagenesis kit (Stratagene) using the hPMCA 4b-modifying primers GTG GGC ATT GAG GAG CCT GTG CGC CCA GAG GTG and CAC CTC TGG GCG CAC AGG CTC CTC AAT GCC CAC. The pCMV-hPMCA4b cap vector was constructed by insertion of hPMCA 4b cDNA in BamHI-KpnI sites of pcDNA3.1(-)/Myc-His vector (Invitrogen) and subsequent mutation of

the hPMCA 4b stop codon to obtain a c-myc/6×His tag at the PMCA COOH terminus. The constitutively active form of the PMCA 4b (pHPMCA4b[ct120]; Adamo et al., 1995), was a gift of E. Strehler (Mayo Clinic, Rochester, NY). NOS-I expression plasmid (pCMV-NOS-I; Brenman et al., 1996) and pGST-PDZ (of NOS-I; Brenman et al., 1995) were a gift of David S. Bredt (University of California, San Francisco, CA). The NOS-III cDNA (Lamas et al., 1992; EMBL/GenBank/DBJ accession no. M89952) was cloned in the EcoRI site of pcDNA3 (Invitrogen), similar to the NOS-I expression construct (see above). The ΔNOS-I vector, coding for NOS-I and lacking the PDZ domain, was constructed by excision of nucleotides 1–975 of the rat cDNA (numbering according to EMBL/GenBank/DBJ accession no. X59949), resulting in a truncated NOS-I, which is expressed in transfected cells and exhibits NO synthase activity.

Transfections and cGMP measurement

HEK293 and Neuro-2a cells (DSMZ) were grown in 90% DME/10% fetal bovine serum/1× nonessential amino acids (Sigma-Aldrich) to a density of 50–70% before transfection. HEK293 and neuro-2a cells were transfected using the LipofectaminPLUS™ system according to the manufacturer's protocol (Life Technologies). After medium replacement cells were incubated overnight. cGMP was measured using cGMP (low pH) colorimetric competitive ELISA (R&D Systems). Cells were saturated with L-arginine (30 min, 1 mM) and, 5 min before lysis, nascent NO was stabilized by addition of superoxide dismutase (100 U/ml culture medium; Roche Diagnostics). In parallel, 3-isobutyl-1-methylxanthine (IBMX, 1 mM; Sigma-Aldrich) was added to the culture 5 min before lysis (to inactivate phosphodiesterase activity) and NO synthesis was induced by addition of 0.5 μM ionomycin 3 min before lysis. Subsequent determination of NO-dependent cGMP production was estimated in relation to untransfected control cells.

We thank D.S. Bredt, E. Strehler, J.T. Penniston, and A. Filoteo for kindly providing reagents, G. Ertl and K. Kochsiek for continuous support, and S. Lohmann and U. Walter for critically reading the manuscript.

This work was supported by the German Research Foundation (Deutsche Forschungsgemeinschaft; SFB 355, TP B6) and by an International Appointment Grant from the Medical Research Council (UK) to L. Neyes.

Submitted: 30 April 2001

Accepted: 7 September 2001

References

- Adamo, H.P., A.G. Filoteo, A. Enyedi, and J.T. Penniston. 1995. Mutants in the putative nucleotide-binding region of the plasma membrane Ca²⁺-pump. A reduction in activity due to slow dephosphorylation. *J. Biol. Chem.* 270: 30111–30114.
- Brenman, J.E., D.S. Chao, H. Xia, K. Aldape, and D.S. Bredt. 1995. Nitric oxide synthase complexed with dystrophin and absent from skeletal muscle sarcolemma in Duchenne muscular dystrophy. *Cell.* 82:743–752.
- Brenman, J.E., D.S. Chao, S.H. Gee, A.W. McGee, S.E. Craven, D.R. Santillano, Z. Wu, F. Huang, H. Xia, M.F. Peters, S.C. Froehner, and D.S. Bredt. 1996. Interaction of nitric oxide synthase with the postsynaptic density protein PSD-95 and α1-syntrophin mediated by PDZ domains. *Cell.* 84:757–767.
- Carafoli, E. 1997. Plasma membrane calcium pump: structure, function and relationships. *Basic Res. Cardiol.* 92:59–61.
- Chang, W.J., Y.S. Ying, K.G. Rothberg, N.M. Hooper, A.J. Turner, H.A. Gambliel, J. De Gunzburg, S.M. Mumby, A.G. Gilman, and R.G. Anderson. 1994. Purification and characterization of smooth muscle cell caveolae. *J. Cell Biol.* 126:127–138.
- Christopherson, K.S., and D.S. Bredt. 1997. Nitric oxide in excitable tissues: physiological roles and disease. *J. Clin. Invest.* 100:2424–2429.
- DeMarco, S.J., and E.E. Strehler. 2001. Plasma membrane Ca²⁺-ATPase isoforms 2b and 4b interact promiscuously and selectively with members of the membrane-associated guanylate kinase family of PDZ (PSD95/Dlg/ZO-1) domain-containing proteins. *J. Biol. Chem.* 276:21594–21600.
- Enyedi, A., A.K. Verma, A.G. Filoteo, and J.T. Penniston. 1993. A highly active 120-kDa truncated mutant of the plasma membrane Ca²⁺ pump. *J. Biol. Chem.* 268:10621–10626.
- Fujimoto, T. 1993. Calcium pump of the plasma membrane is localized in caveolae. *J. Cell Biol.* 120:1147–1157.
- Hammes, A., S. Oberdorf, E.E. Strehler, T. Stauffer, E. Carafoli, H. Vetter, and L. Neyes. 1994. Differentiation-specific isoform mRNA expression of the calmodulin-dependent plasma membrane Ca²⁺-ATPase. *FASEB J.* 8:428–435.
- Hammes, A., S. Oberdorf-Maass, S. Jenatschke, T. Pelzer, A. Maass, F. Gollnick, R. Meyer, J. Afflerbach, and L. Neyes. 1996. Expression of the plasma membrane Ca²⁺-ATPase in myogenic cells. *J. Biol. Chem.* 271:30816–30822.
- Hammes, A., S. Oberdorf-Maass, T. Rother, K. Nething, F. Gollnick, K.W. Linz, R. Meyer, K. Hu, H. Han, P. Gaudron, et al. 1998. Overexpression of the sarcolemmal calcium pump in the myocardium of transgenic rats. *Circ. Res.* 83:877–888.
- Hillier, B.J., K.S. Christopherson, K.E. Prehoda, D.S. Bredt, and W.A. Lim. 1999. Unexpected modes of PDZ domain scaffolding revealed by structure of nNOS-syntrophin complex. *Science.* 284:812–815.
- Huang, Z., P.L. Huang, N. Panahian, T. Dalkara, M.C. Fishman, and M.A. Moskowitz. 1994. Effects of cerebral ischemia in mice deficient in neuronal nitric oxide synthase. *Science.* 265:1883–1885.
- Kim, E., S.J. DeMarco, S.M. Marfatia, A.H. Chishti, M. Sheng, and E.E. Strehler. 1998. Plasma membrane Ca²⁺ ATPase isoform 4b binds to membrane-associated guanylate kinase (MAGUK) proteins via their PDZ (PSD-95/Dlg/ZO-1) domains. *J. Biol. Chem.* 273:1591–1595.
- Lamas, S., P.A. Marsden, G.K. Li, P. Tempst, and T. Michel. 1992. Endothelial nitric oxide synthase: molecular cloning and characterization of a distinct constitutive enzyme isoform. *Proc. Natl. Acad. Sci. USA.* 89:6348–6352.
- Lisanti, M.P., P.E. Scherer, J. Vidugiriene, Z. Tang, A. Hermanowski-Vosatka, Y.H. Tu, R.F. Cook, and M. Sargiacomo. 1994. Characterization of caveolin-rich membrane domains isolated from an endothelial-rich source: implications for human disease. *J. Cell Biol.* 126:111–126.
- Neyes, L., L. Reinlib, and E. Carafoli. 1985. Phosphorylation of the Ca²⁺-pumping ATPase of heart sarcolemma and erythrocyte plasma membrane by the cAMP-dependent protein kinase. *J. Biol. Chem.* 260:10283–10287.
- Olmsted, J.B., K. Carlson, R. Klebe, F. Ruddle, and J. Rosenbaum. 1970. Isolation of microtubule protein from cultured mouse neuroblastoma cells. *Proc. Natl. Acad. Sci. USA.* 65:129–136.
- Robbins, S.M., N.A. Quintrell, and J.M. Bishop. 1995. Myristoylation and differential palmitoylation of the HCK protein- tyrosine kinases govern their attachment to membranes and association with caveolae. *Mol. Cell. Biol.* 15: 3507–3515.
- Sargiacomo, M., M. Sudol, Z. Tang, and M.P. Lisanti. 1993. Signal transducing molecules and glycosyl-phosphatidylinositol-linked proteins form a caveolin-rich insoluble complex in MDCK cells. *J. Cell Biol.* 122:789–807.
- Schnitzer, J.E., J. Liu, and P. Oh. 1995. Endothelial caveolae have the molecular transport machinery for vesicle budding, docking, and fusion including VAMP, NSF, SNAP, annexins, and GTPases. *J. Biol. Chem.* 270:14399–14404.
- Segal, S.S., S.E. Brett, and W.C. Sessa. 1999. Codistribution of NOS and caveolin throughout peripheral vasculature and skeletal muscle of hamsters. *Am. J. Physiol.* 277:H1167–H1177.
- Shull, G.E., and J. Greeb. 1988. Molecular cloning of two isoforms of the plasma membrane Ca²⁺- transporting ATPase from rat brain. Structural and functional domains exhibit similarity to Na⁺/K⁺- and other cation transport ATPases. *J. Biol. Chem.* 263:8646–8657.
- Song, K.S., S. Li, T. Okamoto, L.A. Quilliam, M. Sargiacomo, and M.P. Lisanti. 1996a. Co-purification and direct interaction of Ras with caveolin, an integral membrane protein of caveolae microdomains. Detergent-free purification of caveolae microdomains. *J. Biol. Chem.* 271:9690–9697.
- Song, K.S., P.E. Scherer, Z. Tang, T. Okamoto, S. Li, M. Chafel, C. Chu, D.S. Kohtz, and M.P. Lisanti. 1996b. Expression of caveolin-3 in skeletal, cardiac, and smooth muscle cells. Caveolin-3 is a component of the sarcolemma and co-fractionates with dystrophin and dystrophin-associated glycoproteins. *J. Biol. Chem.* 271:15160–15165.
- Sprengel, R., B. Suchanek, C. Amico, R. Brusa, N. Burnashev, A. Rozov, O. Hvalby, V. Jensen, O. Paulsen, P. Andersen, et al. 1998. Importance of the intracellular domain of NR2 subunits for NMDA receptor function in vivo. *Cell.* 92:279–289.
- Strehler, E.E., and D.A. Zacharias. 2001. Role of alternative splicing in generating isoform diversity among plasma membrane calcium pumps. *Physiol. Rev.* 81: 21–50.
- Venema, V.J., H. Ju, R. Zou, and R.C. Venema. 1997. Interaction of neuronal nitric-oxide synthase with caveolin-3 in skeletal muscle. Identification of a novel caveolin scaffolding/inhibitory domain. *J. Biol. Chem.* 272:28187–28190.
- Verma, A.K., A.G. Filoteo, D.R. Stanford, E.D. Wieben, J.T. Penniston, E.E. Strehler, R. Fischer, R. Heim, G. Vogel, S. Mathews, et al. 1988. Complete primary structure of a human plasma membrane Ca²⁺ pump. *J. Biol. Chem.* 263:14152–14159.