



# Draft Genome Sequences of 9 Actinobacteria from the Family *Microbacteriaceae* Associated with Insect- and Nematode-Damaged Plants

 Sergey V. Tarlachkov,<sup>a</sup> Yury V. Ospennikov,<sup>a</sup> Alexander V. Demidov,<sup>a</sup> Irina P. Starodumova,<sup>a</sup> Lubov V. Dorofeeva,<sup>a</sup> Natalia V. Prisyazhnaya,<sup>a</sup> Vladimir N. Chizhov,<sup>b</sup>  Sergei A. Subbotin,<sup>c</sup> Lyudmila I. Evtushenko<sup>a</sup>

<sup>a</sup>All-Russian Collection of Microorganisms (VKM), G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, Pushchino Scientific Center for Biological Research, Russian Academy of Sciences, Pushchino, Russia

<sup>b</sup>Center of Parasitology of A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia

<sup>c</sup>California Department of Food and Agriculture, Sacramento, California, USA

**ABSTRACT** Draft genome sequences of 9 strains of known and putative new species of *Microbacteriaceae* isolated from insect- and nematode-damaged plants were generated using Illumina technology. The data obtained will contribute to the development of the genome-based prokaryote taxonomy and the knowledge on the biology of the microbial group investigated.

**A**ctinobacteria of the family *Microbacteriaceae* are distributed widely in various terrestrial and aquatic environments and often inhabit plants as endophytes or pathogens (1–3). Many putative new species of several genera of this family, along with representatives of known species, were found in plants infested by different nematodes (4, 5).

Novel strains were isolated from affected and unaffected parts of leaves of different herbaceous and woody plants infected by nematodes, leaf-mining insects, and plant-parasitic mites (Table 1).

The air-dried plant samples were soaked in distilled water for 1 h, washed twice with sterile distilled water, placed in 0.85% NaCl solution, and milled. One drop of the obtained suspension was plated onto Reasoner's 2A (R2A) agar (Fluka Analytical, USA) and incubated for 1 to 3 weeks at room temperature (18 to 24°C). The universal bacterial primers 27F (5'-AGAGTTGATCTGGCTAG-3') and 1525R (5'-AAGGAGGTGATCCA GCC-3') with standard cycling conditions (95°C for 5 min; 30 cycles at 94°C for 30 s, 55°C for 30 s, and 72°C for 1 min 20 s; and 72°C for 4 min) were used for 16S rRNA gene PCR amplification followed by sequencing. Preliminary identification based on nearly entire 16S rRNA gene sequences revealed that the newly isolated strains belong to four genera of the family *Microbacteriaceae*, exhibiting high sequence identities to the known species (Table 1).

For genome sequencing, biomass was grown in liquid peptone-yeast-glucose (PYG) medium (5 g peptone, 3 g yeast extract, 5 g glucose, 0.2 g KH<sub>2</sub>PO<sub>4</sub>, and 1 L distilled water [pH 7.2]) or on R2A agar inoculated with cells from a single colony, followed by cultivation at 28°C for 2 to 3 days. Genomic DNA was extracted using a QIAamp DNA minikit (Qiagen, Germany). DNA library construction and sequencing were performed by using Novogene Co., Ltd. Libraries were generated with the NEBNext DNA library prep kit for Illumina (New England Biolabs, USA) following the manufacturer's recommendations. Pooled DNA libraries were sequenced on an Illumina NovaSeq 6000 instrument to obtain 150-bp paired-end reads.

The quality of the reads was checked with FastQC v0.11.8 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). Adapter sequences and low-quality regions in raw reads

**Editor** David A. Baltrus, University of Arizona

**Copyright** © 2022 Tarlachkov et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Sergey V. Tarlachkov, sergey@tarlachkov.ru, or Sergei A. Subbotin, sergei.a.subbotin@gmail.com.

The authors declare no conflict of interest.

**Received** 11 May 2022

**Accepted** 15 August 2022

**Published** 31 August 2022

**TABLE 1** Strain data, characteristics, and DDBJ/ENA/GenBank accession numbers of genome sequences

Organism	Plant	Plant parasite	Geography	No. of reads	No. of scaffolds	Scaffold N <sub>50</sub> (bp)	Genome size (Mbp)	G+C content (%)	No. of proteins	Closest species by ANI (identity %)	Closest species by dDDH (identity %)	16S rRNA gene GenBank accession no.	Genome GenBank accession no.	Genome SRA accession no.
<i>Clavibacter michiganensis</i> VRM Ac-2921	<i>Tilia</i> sp.	<i>Eriophyes tiliae</i>	Moscow region, Russia	14,417,302	660	8	2,063,997	3.18	73.4	<i>Clavibacter michiganensis</i> subsp. <i>phaseoli</i> (100)	<i>Clavibacter michiganensis</i> (97.30)	ON158671.1	JALGR000000000	SRR18694026
<i>Curtobacterium</i> sp. VRM Ac-2922	Fern	<i>Aphelenchoides</i> sp.	Washington state, USA	13,428,696	526	29	222,655	3.73	71.0	<i>Clavibacter michiganensis</i> flaccumfaciens (99.93)	<i>Curtobacterium flaccumfaciens</i> (25.4)	ON158672.1	JALGR000000000	SRR18694025
<i>Microbacterium</i> sp. VRM Ac-2923	<i>Tellima grandiflora</i>	<i>Aphelenchoides</i> sp.	Münster, Germany	14,406,260	536	37	362,450	3.93	69.8	<i>Microbacterium testaceum</i> (98.16)	<i>Microbacterium proteolyticum</i> (87.05)	ON158673.1	JALGR000000000	SRR18694024
<i>Rathayibacter caricis</i> VRM Ac-2924	<i>Umbelliferae<sup>a</sup></i>	Unknown insect (leaf miner)	Moscow region, Russia	12,318,900	426	23	880,020	4.21	71.5	<i>Rathayibacter caricis</i> (99.79)	<i>Rathayibacter caricis</i> (82.9)	ON158674.1	JALGR000000000	SRR18694023
<i>Rathayibacter festucae</i> VRM Ac-2925	<i>Umbelliferae<sup>a</sup></i>	Unknown insect (leaf miner)	Moscow region, Russia	14,839,660	479	7	2,978,450	4.49	72.2	<i>Rathayibacter festucae</i> (100)	<i>Rathayibacter festucae</i> (97.80)	ON158675.1	JALGR000000000	SRR18694022
<i>Rathayibacter</i> sp. VRM Ac-2926	<i>Umbelliferae<sup>a</sup></i>	Unknown insect (leaf miner)	Moscow region, Russia	17,536,744	590	30	366,154	4.32	72.3	<i>Rathayibacter festucae</i> (100)	<i>Rathayibacter festucae</i> (96.17)	ON158676.1	JALGR000000000	SRR18694021
<i>Rathayibacter</i> sp. VRM Ac-2927	<i>Tilia</i> sp.	Unknown insect (leaf miner)	Moscow region, Russia	18,677,542	598	13	749,896	4.55	72.1	<i>Rathayibacter festucae</i> (100)	<i>Rathayibacter festucae</i> (96.18)	ON158677.1	JALGR000000000	SRR18694020
<i>Rathayibacter</i> sp. VRM Ac-2928	<i>Aesculus</i> sp.	<i>Cameraria ohridella</i> (leaf miner)	Moscow region, Russia	13,232,092	446	13	849,744	4.32	72.3	<i>Rathayibacter festucae</i> (100)	<i>Rathayibacter festucae</i> (96.17)	ON158678.1	JALGR000000000	SRR18694019
<i>Rathayibacter</i> sp. VRM Ac-2929	<i>Artemisia</i> sp.	Unknown insect (leaf miner)	Moscow region, Russia	16,532,532	529	19	764,283	4.53	72.1	<i>Rathayibacter festucae</i> (100)	<i>Rathayibacter festucae</i> (96.03)	ON158679.1	JALGR000000000	SRR18694018

<sup>a</sup> Unaffected parts of leaves were used for the isolation of strains.

were cut with Trimmomatic 0.39 (6). Trimmed reads were assembled using SPAdes 3.15.4 (7). Assemblies were annotated with NCBI PGAP 6.1 (8). The pairwise similarity between the 16S rRNA gene sequences was determined using TaxonDC 1.3.1 (9). The genome relatedness indices, *viz.*, the average nucleotide identity (ANI) and digital DNA-DNA hybridization (dDDH) values, were calculated using JSpecies 1.2.1 (10) and GGDC 2.1 (11) tools, respectively. Accession numbers and characteristics of genomes are provided in Table 1.

The ANI and dDDH values obtained for strains VKM Ac-2922 and VKM Ac-2923 and their closest relatives were well below the accepted species threshold (12), indicating the strains to be members of two putative new species in the genera *Curtobacterium* and *Microbacterium*. Four strains (VKM Ac-2926, VKM Ac-2927, VKM Ac-2928, and VKM Ac-2929) may represent a new species as well. The genome relatedness indices for these strains toward their closest relative, namely, *Rathayibacter festucae*, were at the borderline for species definition or slightly below (12). From the genome relatedness indices, three remaining strains (VKM Ac-2921, VKM Ac-2924, and VKM Ac-2925) belonged to the known *Clavibacter* and *Rathayibacter* species.

**Data availability.** The whole-genome shotgun projects and the 16S rRNA gene sequences have been deposited in DDBJ/ENA/GenBank under the accession numbers listed in Table 1.

## ACKNOWLEDGMENTS

This work was sponsored by the United States Department of Agriculture Animal and Plant Health Inspection Service (research project agreement number AP21PPQS&T00C169, CDFA agreement number 21-0425-000-FR) and the Ministry of Science and Higher Education of the Russian Federation (grant agreement number 075-15-2021-1051).

## REFERENCES

- Evtushenko LI. 2015. *Microbacteriaceae*. In Whitman WB, Rainey F, Kämpfer P, Trujillo M, Chun J, DeVos P, Hedlund B, Dedysh S (ed), Bergey's manual of systematics of archaea and bacteria. Wiley, Hoboken, New Jersey. <https://doi.org/10.1002/9781118960608.fbm00035>.
- Murray TD, Schroeder BK, Schneider WL, Luster DG, Sechler A, Rogers EE, Subbotin SA. 2017. *Rathayibacter toxicus*, other *Rathayibacter* species inducing bacterial head blight of grasses, and the potential for livestock poisonings. *Phytopathology* 107:804–815. <https://doi.org/10.1094/PHYTO-02-17-0047-RWV>.
- Dorofeeva LV, Starodumova IP, Krauzova VI, Prisyazhnaya NV, Vinokurova NG, Lysanskaya VY, Tarlachkov SV, Evtushenko LI. 2018. *Rathayibacter oskolensis* sp. nov., a novel actinobacterium from *Androsace koso-poljan-skii* Ovcz. (Primulaceae) endemic to Central Russian Upland. *Int J Syst Evol Microbiol* 68:1442–1447. <https://doi.org/10.1099/ijsem.0.002681>.
- Tarlachkov SV, Starodumova IP, Dorofeeva LV, Prisyazhnaya NV, Leyn SA, Zlamal JE, Elane ML, Osterman AL, Nadler SA, Subbotin SA, Evtushenko LI. 2020. Complete and draft genome sequences of 12 plant-associated *Rathayibacter* strains of known and putative new species. *Microbiol Resour Announc* 9:e00316-20. <https://doi.org/10.1128/MRA.00316-20>.
- Tarlachkov SV, Starodumova IP, Dorofeeva LV, Prisyazhnaya NV, Roubtsova TV, Chizhov VN, Nadler SA, Subbotin SA, Evtushenko LI. 2021. Draft genome sequences of 28 actinobacteria of the family *Microbacteriaceae* associated with nematode-infected plants. *Microbiol Resour Announc* 10:e01400-20. <https://doi.org/10.1128/MRA.01400-20>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prijibelski AD, Pyshkin AV, Sirotnik AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Tarlachkov SV, Starodumova IP. 2017. TaxonDC: calculating the similarity value of the 16S rRNA gene sequences of prokaryotes or ITS regions of fungi. *J Bioinform Genom* 3:1–4. <https://doi.org/10.18454/jbg.2017.3.5.1>.
- Richter M, Rosselló-Móra R. 2009. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* 106:19126–19131. <https://doi.org/10.1073/pnas.0906412106>.
- Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <https://doi.org/10.1186/1471-2105-14-60>.
- Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, da Costa MS, Rooney AP, Yi H, Xu XW, De Meyer S, Trujillo ME. 2018. Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 68:461–466. <https://doi.org/10.1099/ijsem.0.002516>.