

High-quality human preimplantation embryos stimulate endometrial stromal cell migration via secretion of microRNA hsa-miR-320a

Robbert P. Berkhout^{1,2}, Remco Keijser³, Sjoerd Repping¹,
Cornelis B. Lambalk², Gijs B. Afink³, Sebastiaan Mastenbroek^{1,*†},
and Geert Hamer^{1,*†}

¹Reproductive Biology Laboratory, Center for Reproductive Medicine, Amsterdam Reproduction & Development Research Institute, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands ²Department of Obstetrics and Gynaecology, Amsterdam Reproduction & Development Research Institute, Amsterdam UMC, VU University, Amsterdam, The Netherlands ³Reproductive Biology Laboratory, Amsterdam Reproduction & Development Research Institute, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands

*Correspondence address. Sebastiaan Mastenbroek, Reproductive Biology Laboratory, Center for Reproductive Medicine, Amsterdam Reproduction & Development Research Institute, Amsterdam UMC, University of Amsterdam, Meibergdreef 9, Amsterdam, The Netherlands. E-mail: s.mastenbroek@amsterdamumc.nl (S.M.) or Geert Hamer, Reproductive Biology Laboratory, Center for Reproductive Medicine, Amsterdam Reproduction & Development Research Institute, Amsterdam UMC, University of Amsterdam, Meibergdreef 9, Amsterdam, The Netherlands. E-mail: g.hamer@amsterdamumc.nl (G.H.)

Submitted on February 13, 2020; resubmitted on April 22, 2020; editorial decision on May 26, 2020

STUDY QUESTION: How do high-quality human preimplantation embryos influence the endometrium to promote their own implantation?

SUMMARY ANSWER: High-quality human preimplantation embryos secrete a specific microRNA (miRNA), hsa-miR-320a, which promotes migration of human endometrial stromal cells (hESCs).

WHAT IS KNOWN ALREADY: We have previously shown that high-quality human preimplantation embryos excrete unknown factors that influence migration of hESCs.

STUDY DESIGN, SIZE, DURATION: Embryo excreted miRNAs, specifically those excreted by high-quality embryos, were identified and their effect on hESCs was determined by measuring the migration capacity and gene expression patterns of primary isolated hESCs.

PARTICIPANTS/MATERIALS, SETTING, METHODS: Embryo conditioned medium (ECM) from routine ICSI procedures was used to identify embryo excreted miRNAs. miRNome analyses were performed on ECM from individually cultured embryos with high morphological quality, with low morphological quality or empty control medium. MiRNA mimics and inhibitors were then used to further study the effect of miRNAs of interest on migration and gene expression of hESCs. Migration assays were performed using hESCs that were obtained from endometrial biopsies performed on hysterectomy specimens from women that received surgery for spotting due to a niche in a cesarean section scar.

MAIN RESULTS AND THE ROLE OF CHANCE: By using miRNA mimics and inhibitors, we showed that hsa-miR-320a alone can stimulate migration of decidualized hESCs, accurately resembling the response typically triggered only by high-quality embryos. Transcriptome analysis further demonstrated that this effect is very likely mediated via altered expression of genes involved in cell adhesion and cytoskeleton organization.

LIMITATIONS, REASONS FOR CAUTION: The effect of hsa-miR-320a on hESCs was measured *in vitro*. Further studies on the *in vivo* effect of hsa-miR-320a are warranted.

WIDER IMPLICATIONS OF THE FINDINGS: Implantation failure is one of the major success limiting factors in human reproduction. By secreting hsa-miR-320a, high-quality human preimplantation embryos directly influence hESCs, most likely to prime the endometrium at

[†]These authors contributed equally to this article.

© The Author(s) 2020. Published by Oxford University Press on behalf of European Society of Human Reproduction and Embryology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

the implantation site for successful implantation. Together, our results indicate that hsa-miR-320a may be a promising target to further increase success rates in assisted reproduction.

STUDY FUNDING/COMPETING INTEREST(S): The study was funded by the Amsterdam University Medical Centers and the Amsterdam Reproduction & Development Research Institute. R.P.B., G.H. and S.M. have a patent on the use of hsa-miR-320a in assisted reproduction treatments pending.

TRIAL REGISTRATION NUMBER: N/A.

Key word: preimplantation embryos / miRNAs / hsa-miR-320a / endometrial stromal cells / cell migration / implantation

Introduction

Human reproduction is characterized by its inefficiency and relatively low success rate. In natural conception, live birth rates per cycle range from 20% to 30% with a miscarriage rate of 10%, suggesting that most embryos are lost before or right after implantation (Wilcox et al., 1988; Zinaman et al., 1996; Macklon et al., 2002). Despite advances in reproductive technologies, implantation failure remains a major limiting factor in improving pregnancy rates in IVF/ICSI.

For successful implantation to occur, properly synchronized development of both embryo and endometrium is required. Embryonic development starts when the zygote initiates consecutive cleavage divisions that ultimately result in formation of a morula at Day 4 and a blastocyst at Day 5 of development (Niakan et al., 2012). In IVF/ICSI, embryonic development is monitored daily by morphological scoring, including number, size and symmetry of blastomeres, multinucleation and the percentage of fragmentation of blastomeres (Puissant et al., 1987; Van Royen et al., 1999; Balaban et al., 2011). Each of these traits contributes to a comprehensive grade of each individual embryo that is predictive for its chances to implant successfully (Balaban et al., 2011; van Loendersloot et al., 2014). Based on these grades, embryos are selected to be used for transfer into the uterus or cryopreservation (Ebner et al., 2003).

In preparation for implantation, also the endometrium undergoes profound changes. After a surge in LH, the endometrium is exposed to rising levels of progesterone leading to decidualization of human endometrial stromal cells (hESCs) (Gellersen and Brosens, 2014). Decidualization alters the phenotypical characteristics of hESCs, such as its transcriptome and morphology, in preparation for possible implantation of an embryo (Gellersen and Brosens, 2014). It has been suggested that directed migration of decidualized hESCs is crucial for implantation to succeed, and that altered migration of hESCs may be associated with reproductive failure (Grewal et al., 2008; Weimar et al., 2012; Macklon and Brosens, 2014). Indeed, when comparing the effects of both low- and high-quality preimplantation embryos, we have recently shown that only embryos of high morphological quality, i.e. with a low percentage of fragmentation, stimulate migration of decidualized hESCs in an *in vitro* migration assay (Berkhout et al., 2018). In addition, specifically high-quality embryos, and not low-quality embryos, inhibit migration of non-decidualized hESCs (Berkhout et al., 2018).

When an embryo enters the uterine cavity, embryo and endometrium are thought to start interacting to initiate implantation (Quenby and Brosens, 2013). Before they physically connect, both have been suggested to already secrete a wide variety of signals, the so-called embryo-endometrium crosstalk, which includes hormones, growth factors and cytokines (Haouzi et al., 2011; Peter Durairaj et al., 2017).

After embedding into uterine epithelial and stromal cells, paracrine and cellular signaling has been suggested to further steer the progress of implantation (Cha et al., 2012; Fritz et al., 2014).

Besides more traditional signaling factors, research has also focused on the roles of more recently discovered factors, such as microRNAs (miRNAs) (Galliano and Pellicer, 2014). These 20–25 nucleotide small non-coding RNA molecules play an important signaling roles in various mechanisms of health and disease, such as angiogenesis and carcinogenesis (Calin and Croce, 2006; Esquela-Kerscher and Slack, 2006; Lou et al., 2017). Biogenesis of miRNAs starts in the nucleus where precursor miRNAs are transcribed. Following transfer to the cytoplasm, precursor miRNAs are spliced by Dicer into single-stranded mature miRNAs (He and Hannon, 2004; Ha and Kim, 2014). Next, these mature miRNAs are assembled into RNA-induced silencing complexes (RISC) that can bind to the 3'-UTR of an analogue target mRNA, leading to translational repression and mRNA decay (He and Hannon, 2004).

Preimplantation embryos have been shown to secrete miRNAs during multiple steps of development and implantation in human and various other mammalian species (McCallie et al., 2010; Kropp et al., 2014; Rosenbluth et al., 2014; Cuman et al., 2015; Capalbo et al., 2016; Abu-Halima et al., 2017; Bidarimath et al., 2017; Gross et al., 2017). Several studies have also reported that embryos secrete specific miRNAs that may be associated with the method of fertilization, the underlying infertility diagnosis, their chromosomal status or the pregnancy outcome (McCallie et al., 2010; Rosenbluth et al., 2014; Cuman et al., 2015; Capalbo et al., 2016; Abu-Halima et al., 2017). However, no clear link between embryo-derived miRNAs and a possible biological mechanism of action during implantation has been described. In this study, we identify a human embryo-derived miRNA that influences migration of decidualized hESCs during implantation.

Materials and methods

Ethical approval and informed consent

Regarding the use of endometrial biopsies, this study was approved by the Medical Review Ethics Committee of the Amsterdam University Medical Centers and written informed consent was obtained from all subjects who agreed to donate endometrial biopsies for research purposes. Embryo conditioned medium (ECM) was collected from patients who did not object to storage and use of medical waste tissues and was stored anonymously. According to Dutch law, the use of ECM for this research was allowed without ethical permission.

Endometrial biopsies

Biopsies were performed on hysterectomy specimens from patients that received surgery for spotting due to a niche in a cesarean section scar ($n=2$). Patients were pre-menopausal, were 35 and 39 years of age, had a history of proven fertility and at least one live birth, and received no hormonal treatment 3 months prior to surgery. Biopsies were performed randomly in the menstrual cycle and biopsied material was immediately suspended in Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12 phenol red-free medium (DMEM/F12; L-glutamine, HEPES; Thermo Fisher Scientific, Waltham, MA, USA) supplemented with 1% penicillin/streptomycin (Gibco by Life Technologies, CA, USA) at 37°C. Next, the biopsied material was cut into small pieces and further digested enzymatically in DMEM/F12 phenol red-free medium supplemented with >125 U/ml collagenase IV (Gibco) and 1% penicillin/streptomycin (Gibco). The suspended material was filtered through a 76 μm sterile filter and subsequently cultured in DMEM/F12 phenol red-free medium supplemented with 10% heat-inactivated fetal calf serum (FCS; Thermo Fisher Scientific, Waltham, MA, USA) and 1% penicillin/streptomycin (C-medium). The medium was refreshed after 3 h to discard all non-attached cells and retain the attached hESCs. Purity of hESCs cultures was assessed by immunostaining for vimentin (M0725, Agilent, CA, USA) and cytokeratin 18 (M7010, Agilent, CA, USA). Only hESCs cultures that were passaged <6 times were used for subsequent experiments.

Embryo conditioned medium

All assigned ECM was collected from regular ICSI-cycles in which routine ICSI procedures were followed and during which all embryos were handled according to local regulations and standard operating procedures. Embryos were cultured in Sage medium (Quinn's advantage cleavage medium, Cooper Surgical, USA) supplemented with 5% human serum albumin (HSA; Vitrolife, Göteborg, Sweden) from Day 1 to 3 after fertilization. Thereafter, embryos were transferred to fresh Sage medium (Quinn's advantage blastocyst medium, Cooper Surgical, USA) supplemented with 5% HSA, on Day 3 after fertilization. Embryo morphology was assessed daily by an independent laboratory technician. Droplets (20 μl) from individually cultured embryos, now referred to as ECM, were collected in sterile 1.5 ml collection tubes after all embryos had been removed from the culture dishes for cryopreservation on Day 4. All ECM samples were collected on Day 4. Empty medium droplets, in which no embryo was grown but that otherwise underwent the same procedures, from the same dishes were used as controls. Samples were snap-frozen in liquid nitrogen and stored at -80°C .

Migration assays

Experimental groups.

ECM was used from individually cultured embryos with high morphological quality (≥ 8 blastomeres and fragmentation $\leq 20\%$), low morphological quality (≤ 7 blastomeres and fragmentation $>20\%$) or empty control medium in which no embryo had been cultured, but that otherwise underwent the same laboratory procedures. Per the experimental group, ECM from five individually cultured embryos was pooled (90 μl).

Experimental procedures.

Thirty thousand thawed hESCs were cultured in single wells of a 48-well plate and decidualized by adding DMEM/F12 phenol red-free medium supplemented with 10% heat-inactivated charcoal stripped FCS (Thermo Fisher Scientific, Waltham, MA, USA), 0.5 mM 8-bromoadenosine 3',5'-cyclic monophosphate (Sigma-Aldrich, Saint Louis, MO, USA), 1 μM medroxyprogesterone acetate (Sigma-Aldrich, Saint Louis, MO, USA) and 1% penicillin/streptomycin (D-medium) for 5 consecutive days. Monolayers of hESCs were scratched with a p200 pipet tip to create a migration zone. hESCs were then cultured in 90 μL of pooled ECM derived from five individual embryos. Cultures were overlaid with 150 μL mineral oil (Irvine Scientific, Santa Ana, CA, USA) to prevent evaporation. For addition of RNase, we used 1 mg/ml of RNase in PBS (10109169001, Roche, Basel, Switzerland). At 0 and 18 h after creation of the migration zone, phase contrast images were taken using a monochrome digital camera (Leica DFC365 FX) attached to an inverted microscope (Olympus IX71) with a U Plan FL 4x/0.13 PhL objective. The migration response of hESCs was quantified by assessing the reduction of the surface area of the migration zone by using Image J software (version 1.50i).

miRNome PCR

Experimental groups.

ECM from individually cultured embryos of high-quality (fragmentation $\leq 20\%$; 48 samples), low-quality (fragmentation $>20\%$; 48 samples) or empty control medium, in which no embryo had been cultured (48 samples), was used. Per experimental group, three biological replicates were performed by pooling ECM samples from 16 individually cultured embryos (320 μl pooled ECM). To ascertain that cellular fragmentation was the only difference among these experimental groups, samples from embryos with different developmental stages (ranging from six blastomeres to the morula stage) were evenly distributed across both experimental groups.

Experimental procedures.

All procedures for miRNome analysis were performed at Qiagen, Hilden, Germany. Total RNA was extracted using miRCURY RNA isolation kit—Biofluids (300112, Exiqon, Denmark) according to manufacturer's instructions and RNA was eluted in 50 μl RNase-free H_2O . Next, RNA was polyadenylated and cDNA was synthesized according to manufacturer's instructions (miRCURY LNA RT Kit, 339340, Qiagen, Hilden, Germany). Expression of miRNAs was determined by Ready-to-Use Human panel I+II PCR, using ExiLENT SYBR[®] Green master mix (339322, Qiagen, Hilden, Germany). Amplification reactions were performed in a LightCycler 480 Real-Time PCR System (05015243001, Roche, Basel, Switzerland) in 384 well plates. Amplification curves were analyzed using LC software (04994884001, Roche, Basel, Switzerland), both for determination of cycle quantification (Cq) values and for melting curve analysis. Only samples with Cq <37 were included in the analysis.

MiRNA transfections

Hsa-miR mimics (339173, miRCURY LNA miRNA Mimic) and inhibitors (339121, miRCURY LNA miRNA Inhibitors) were ordered from

Qiagen, Hilden, Germany. Additionally, negative control mimics (YM00479902, Negative Control miRCURY LNA miRNA Mimic) and negative control inhibitors (Y100199006, Negative control A), which both have no homology to any known miRNA or mRNA in human, were ordered from Qiagen, Hilden, Germany. The following miRNAs were labeled with a fluorescent 5'-carboxyfluorescein (FAM) tag: hsa-miR-19a mimic + inhibitor, hsa-miR-19b mimic + inhibitor, negative control A mimic, negative control inhibitor. Thirty thousand decidualized or non-decidualized hESCs were cultured in single wells of a 48 wells plate and transfected with 25 nM miRNA mimic or negative control mimic. Alternatively, 30 000 hESCs were transfected with 150 nM miRNA inhibitor or negative control inhibitor. All transfections were performed in D- or C-medium by using HiPerFect Transfection Reagent according to manufacturer's instructions (301704, Qiagen, Hilden, Germany). After transfection for 24 h, hESCs were washed with PBS and transfection efficiency was determined by calculating the percentage of fluorescently labeled hESCs from the total number of hESCs per image.

RNA-seq

RNA isolation.

Per well of a 6-well plate 300 000 hESCs were cultured and decidualized for 5 consecutive days, after which hESCs were transfected for 24 h with 25 nM hsa-miR-320a mimic or 25 nM negative control mimic. Directly after, transfection medium was removed and hESCs were harvested by using a cell scraper (3010, Corning, NY, USA) and passed through a 20-gauge syringe needle. Next, total RNA was isolated using the ISOLATE II RNA Mini Kit (BIO-52072, Bioline, London, UK) according to manufacturer's instructions.

Library construction and RNA sequencing.

Total RNA was sent to BGI Tech Solutions (Hong Kong, China) for library construction and sequencing (BGISEq500, paired end 100 bp). Twelve libraries were constructed for non-stranded pair-end cDNA sequencing on the BGISEQ-500 RNA-Seq platform. Cleaned sequence reads were mapped against the reference human Hg38 build (USC Genome Browser) using HISAT2 (v2.1.0). The counts of the aligned reads were summarized using featureCounts in R (version 3.4.4) using the Rsubread Bioconductor package (v1.28.1). Multiple mapped reads were excluded from counting. Fold changes were calculated, and differential analysis was performed with the R packages edgeR-Voom (v3.20.9). Genes exhibiting differential expression between MIR_320A and MIR_CTRL transfected hESCs with $P < 0.05$ were included in further analyses.

Statistical analysis

Mean differences among experimental groups were tested using the one-way ANOVA or the Student's t-test. To correct for multiple testing, we used Dunnett and Tukey post-hoc tests when applicable. In all cases, IBM SPSS Statistics 23 software was used. P -values < 0.05 were considered statistically significant.

Results

The migratory effect of ECM from high-quality embryos can be abolished by RNase

We first confirmed that, in correspondence with our previous study (Berkhout et al., 2018), decidualized hESCs migration was stimulated by ECM from high-quality embryos compared with empty control medium ($P = 0.006$, Fig. 1A and B). To determine whether this stimulatory signal was caused by secreted RNA molecules, we repeated these migration assays but with the addition of RNase. We found that the addition of RNase completely abolished the pro-migratory effect of ECM from high-quality embryos on decidualized hESCs ($P = 0.017$, Fig. 1A and B). ECM from individually cultured embryos in these experiments was pooled based on morphological quality of the embryos, and morphological quality was indeed not different between experimental groups (comparison of number of blastomeres, $P = 0.167$; and percentage of fragmentation, $P = 0.207$).

High- and low-quality human preimplantation embryos secrete specific subsets of miRNAs

To identify the nature of the soluble RNA molecules that stimulated migration of decidualized hESCs, we performed a PCR-array analyzing the presence of 752 human miRNAs in ECM. ECM was collected and pooled based on the morphological quality of the embryos: high-quality embryos (ECM droplets from 48 individually cultured embryos; fragmentation $\leq 20\%$), low-quality embryos (ECM droplets from 48 individually cultured embryos; fragmentation $> 20\%$) or empty control medium (48 droplets that were prepared and cultured similar to the other groups, but without an embryo being present) (Fig. 2A). From




Figure 1. High-quality human preimplantation embryos stimulated decidualized hESCs migration through paracrine RNA signaling. (A and B) Decidualized hESCs migration was stimulated by culture in high-quality ECM compared with control. (B) Stimulatory effect of high-quality ECM on migration of decidualized hESCs was antagonized by simultaneously adding RNase. Data are expressed as mean \pm SEM. Values that are significantly different are labeled by asterisks ($*P < 0.05$). ECM, embryo conditioned medium; hESCs, human endometrial stromal cells; w/o, without.




Figure 2. High- and low-quality human preimplantation embryos secrete specific subsets of miRNAs. (A) For both high- and low-quality embryos, 48 individually cultured samples were collected that were pooled per 16 to create three biological replicates. (B) High-quality embryos secreted 20 unique miRNAs, low-quality embryos secreted 25 unique miRNAs and eight miRNAs were overlapping. (C) The most frequently expressed miRNAs were present in both high- and low-quality ECM, except for hsa-miR-320a which is high-quality ECM specific, and hsa-miR-19a-3p and hsa-miR-19b-3p which are low-quality ECM specific. miRNAs, microRNAs.

each set of 48 ECM samples of either high- or low-quality, 16 samples each were pooled resulting in three biological replicates for both high- and low-quality ECM (Fig. 2A).

In total, 53 unique human miRNAs were identified in at least one biological replicate, that were not present in control medium: 20 miRNAs unique to high-quality embryos, 25 miRNAs unique to low-quality embryos, and eight miRNAs that were overlapping between high- and low-quality embryos (Fig. 2B). The miRNAs that were most frequently detected in ECM, regardless of the experimental group, were hsa-miR-16-5p (5/6 experiments), hsa-miR-372-3p (5/6 experiments), hsa-miR-345-5p (4/6 experiments) and hsa-miR-21-5p (3/6 experiments) (Fig. 2C). MiRNAs that were detected in 2/6 experiments were hsa-miR-320a, hsa-miR-19b-3p, hsa-miR-19a-3p, hsa-let-7a-5p, hsa-miR-1913, hsa-miR-877-3p and hsa-miR-671-5p (Fig. 2C). We then identified which miRNAs were exclusively detectable in ECM from either high- or low-quality embryos, and found that hsa-miR-320a

was specific to high-quality ECM (2/3 experiments), while hsa-miR-19b-3p and hsa-miR-19a-3p were specific to low-quality ECM (2/3 experiments) (Fig. 2C). All other miRNAs that were specific for either high- or low-quality embryos were only expressed in one biological replicate. Additional family members of the aforementioned miRNAs were not detected in any of the samples.

Endometrial stromal cell migration is regulated by miRNA hsa-miR-320a secretion from high-quality embryos

To study the effects of miRNAs that were exclusively secreted by either high- or low-quality embryos on decidualized hESCs migration, we transfected hESCs with mimics or inhibitors for either hsa-miR-320a (high-quality ECM) or hsa-miR-19b-3p and hsa-miR-19a-3p (low-quality ECM) (Fig. 3A). Both mimics and inhibitors were successfully transfected into decidualized hESCs, as visualized by a green fluorescent signal for those miRNAs that could be designed with a fluorescently labeled 5'-FAM tag (Fig. 3B). No significant difference in transfection efficiency, determined by the percentage of fluorescently labeled hESCs, was detected between negative controls and miRNA-specific mimics (68.2% vs 57.6%, $P=0.422$) or inhibitors (64.7% vs 53.1%, $P=0.100$) (Fig. 3C).

Next, we identified which of the detected miRNAs was able to mimic ECM induced changes in decidualized hESCs migration, in the absence of ECM. To do so, we first transfected miRNAs into monolayers of decidualized hESCs in the presence of regular culture medium alone. We found that hsa-miR-320a stimulated migration of decidualized hESCs in the absence of ECM ($P=0.002$), while there was no effect from hsa-miR-19b-3p ($P=0.995$) or hsa-miR-19a-3p ($P=0.837$), compared to transfection with negative control mimics (Fig. 3D). These negative controls were designed to have no homology to any known miRNA or mRNA in human and are, therefore, reflective for baseline conditions.

We then investigated whether transfection with hsa-miR-320a mimics would increase the migration response of decidualized hESCs that were subsequently cultured in ECM, derived from low-quality embryos (<8 blastomeres and $>20\%$ fragmentation) or empty control medium. To do so, decidualized hESCs were first transfected with hsa-miR-320a mimics, after which migration assays were performed with ECM (Fig. 4A). Indeed, decidualized hESCs that were transfected with hsa-miR-320a mimics and that were subsequently exposed to control ECM or low-quality ECM, showed an increased migration response, compared to transfection with negative control mimics (control ECM, $P=0.028$; low-quality ECM, $P=0.040$) (Fig. 4B). Included samples of pooled ECM were not significantly different among experimental groups (number of blastomeres, $P=0.900$; fragmentation, $P=0.812$).

Next, we examined whether transfection of decidualized hESCs with hsa-miR-320a inhibitors would interfere with signals present in high-quality ECM (≥ 8 blastomeres and $\leq 20\%$ fragmentation). First, we confirmed that high-quality ECM stimulated migration of decidualized hESCs after transfection with negative control inhibitors ($P=0.034$) (Fig. 4C). Again, these negative controls were designed to have no homology to any known miRNA or mRNA in human, and are therefore reflective for baseline conditions. Then, after transfection with hsa-miR-320a inhibitors, we demonstrated that decidualized hESCs migration was no longer stimulated by high-quality ECM ($P=0.003$;




Figure 3. Hsa-miR-320a mimics high-quality ECM-induced migration of decidualized hESCs. (A) Endogenous miRNA is assembled into RISCs and suppress expression of target genes; miRNA mimics are likewise assembled into RISCs and suppress expression of target genes; miRNA inhibitors bind to complementary mature miRNA, thereby inhibit miRNA function. (B) Transfection of miRNA mimics and inhibitors into decidualized hESCs visualized by presence of fluorescent tags (5'-carboxyfluorescein), scale bar is 15 μ m. (C) Transfection percentages of miRNA mimics and inhibitors into decidualized hESCs is not significantly different compared to negative control mimics or inhibitors. (D) Mimics for hsa-miR-320a stimulated migration of decidualized hESCs whereas all other mimics had no effect. Data are expressed as mean \pm SEM. Values that are significantly different are labeled by asterisks ($*P < 0.05$). INH, inhibitor; ctrl, control; RISCs, RNA-induced silencing complexes.

compared to control $P = 0.732$) (Fig. 4C). In other words, inhibition of hsa-miR-320a completely antagonized any pro-migratory effects of high-quality ECM. Included samples of pooled ECM were not significantly different among experimental groups (number of blastomeres, $P = 0.911$; fragmentation, $P = 1.000$).

We then questioned whether decidualization was required for hESCs to respond to hsa-miR-320a. To do so, we transfected non-decidualized hESCs with mimics and inhibitors for hsa-miR-320a and repeated the migration assays during co-culture with high-quality ECM (≥ 8 blastomeres and $\leq 20\%$ fragmentation) or empty control




Figure 4. Hsa-miR-320a mimics and inhibitors interfere with ECM to regulate migration of decidualized hESCs. (A) Decidualized hESCs were first transfected with miRNA mimics or inhibitors for 24 h, after which migration assays (18 h) were performed during co-culture with ECM. (B) Mimics for hsa-miR-320a stimulated decidualized hESCs migration when co-cultured with control ECM or low-quality ECM. (C) Inhibitors for hsa-miR-320a neutralized the pro-migratory effect of high-quality ECM. INH, inhibitor; ctrl, control. Data are expressed as mean \pm SEM. Values that are significantly different are labeled by asterisks (* P < 0.05).

medium. We found that mimicking or inhibiting hsa-miR-320a function in non-decidualized hESCs did not affect the response to high-quality ECM or empty control medium (Supplementary Fig. S1).

Hsa-miR-320a may influence decidualized hESCs migration by targeting cell adhesion and cytoskeleton organization

Finally, we investigated which specific genes were targeted by embryo-derived hsa-miR-320a in hESCs. To do so, we performed RNA-sequencing on decidualized hESCs derived from two patients that

were treated with either hsa-miR-320a mimics ($n = 3$ per patient) or negative controls ($n = 3$ per patient). Principle component analysis demonstrated a clear effect of hsa-miR-320a mimics on the transcriptome landscape of decidualized hESCs (Fig. 5A). In total, hsa-miR-320a mimics induced differential expression of 223 transcripts. Of these transcripts, 140 were downregulated and 83 were upregulated in decidualized hESCs treated with hsa-miR-320a mimics compared to negative controls (Fig. 5B and Supplementary Data S1A). Next, we used gene ontology analysis to estimate which biological processes in hESCs could be affected by hsa-miR-320a (Huang *et al.*, 2009a,b). By doing so, we identified various processes that were significantly affected by hsa-miR-320a, of which, e.g. cell adhesion, regulation of




Figure 5. MiRNA hsa-miR-320a causes transcriptional alterations in decidualized hESCs. (A) Transcriptional changes in decidualized hESCs from 2 separate patients (2846 versus 2850) are clustered by patient type and transfection type in a multidimensional scaling plot. (B) Upon transfection of decidualized hESCs with hsa-miR-320a mimics 83 genes were upregulated and 140 genes were downregulated compared to cells transfected with negative control mimics.

cytoskeleton organization and cell migration, could be directly related to the observed migration response of decidualized hESCs in response to high-quality ECM (false discovery rate of < 0.05) (Table 1 and Supplementary Data S1B).

Discussion

Our work provides evidence that high-quality human preimplantation embryos secrete a specific paracrine signal, miRNA hsa-miR-320a, that drives the endometrial response during implantation. Among 752 miRNAs investigated, hsa-miR-320a was the only miRNA that is specifically secreted by high-quality embryos. Our results further suggest

that hsa-miR-320a stimulates decidualized hESCs migration by regulating cytoskeleton organization, a crucial biological process involved in cellular motility (Fletcher and Mullins, 2010).

Migration of various cancerous cells has been previously reported to be affected by miRNAs, including hsa-miR-320a. Interestingly, hsa-miR-320a was most often identified as a tumor suppressor that reduced cellular migration (Zhang *et al.*, 2012; Guo *et al.*, 2014; Qi *et al.*, 2014; Zhao *et al.*, 2014, 2018; Xishan *et al.*, 2015; Yu *et al.*, 2016; Li *et al.*, 2017; Liu *et al.*, 2017; Lv *et al.*, 2017a,b; Zhang *et al.*, 2017). However, also cases in which hsa-miR-320a stimulated or had no effect on cellular migration have been described (Yao *et al.*, 2012; Gao *et al.*, 2018). This discrepancy in reported literature may be attributed to the wide variety of different cell types that have been investigated previously, and potential side effects of hsa-miR-320a on other cellular mechanisms such as proliferation or apoptosis. In addition, the exact molecular mechanism by which hsa-miR-320a acts to regulate cellular motility remains to be clarified. Certainly, our results indicate that organization of the cellular cytoskeleton may be targeted, possibly involving microfilaments, intermediate filaments or microtubules (Fletcher and Mullins, 2010).

It was reported previously that hsa-miR-320a, which we identified as being secreted specifically by high-quality embryos, as well as miR-19a-3p and miR-19b-3p, which we identified as being specific to low-quality embryos, are being secreted by human blastocysts (Capalbo *et al.*, 2016; Abu-Halima *et al.*, 2017). Here, we demonstrate that these miRNAs are already secreted by embryos from as early as the cleavage stage. It was recently shown that hsa-miR-320a is already secreted by oocytes, particularly in those that developed into embryos with high morphological quality (Feng *et al.*, 2015). Moreover, it was shown that hsa-miR-320a functions in an autocrine manner to improve morphological development of preimplantation embryos (Feng *et al.*, 2015). However, the observed effects were attributed to hsa-miR-320 in general, without specification of the exact family member involved. Of note, many miRNAs exist as members of a larger family of miRNAs that have highly similar nucleotide sequences. Although speculative, research has suggested that miRNAs with great sequential similarity, for example, members of the same family, may be functionally redundant (Marco *et al.*, 2012; Rhee *et al.*, 2013). Therefore, considering hsa-miR-320 in general, these studies may together point toward two important aspects of embryo-derived hsa-miR-320. First, presence of hsa-miR-320 distinguishes high-quality oocytes and embryos from their low-quality counterparts. And second, hsa-miR-320 seems to play an important role in autocrine and paracrine signaling by first stimulating proper morphological development and later, by promoting directed migration of decidualized hESCs, successful implantation. Whether these aspects of human embryo implantation function in a similar way *in vivo*, and whether they contribute to improved pregnancy rates, still needs further research.

Any biomarker that reflects embryo quality in IVF/ICSI may hold great promise for future clinical applications. In this context, miRNA hsa-miR-320a, whose secretion we demonstrate to be specific to high-quality preimplantation embryos, may be suitable for further validation of its predictive value in selecting embryos with the highest potential to implant for transfer. Additionally, artificial supplementation of hsa-miR-320a during embryo culture, to improve morphological development, or during embryo transfer, to improve implantation rates, may be an interesting avenue for future research. In this study, we focused

Table 1 Hsa-miR-320a may influence decidualized hESCs migration by targeting cytoskeleton organization.

Representative GO-term	Description	Enrichment
GO:0043229	Intracellular organelle	4.1
GO:0097190	Apoptotic signaling pathway	3.5
GO:0044446	Intracellular organelle part	3.2
GO:0050839	Cell adhesion	3.1
GO:0071840	Cellular component biogenesis	2.8
GO:0048514	Vascular development	2.6
GO:0009653	Cell development	2.5
GO:0009966	Signal transduction	2.3
GO:2001233	Extrinsic apoptosis	2.2
GO:0005783	Endoplasmic reticulum	2.2
GO:0001525	Angiogenesis	2.1
GO:1902911	Kinase activity	2.1
GO:0005178	Integrin binding	2.0
GO:0005178	Integrin signaling	1.9
GO:0070273	Phospholipid binding	1.8
GO:0007049	Cell cycle regulation	1.8
GO:0048523	Cell signaling	1.7
GO:0000902	Cell morphology	1.6
GO:0051493	Cytoskeleton organization	1.5
GO:0010812	Cell-substrate adhesion	1.5
GO:0016032	Viral interaction	1.5
GO:0048013	Receptor signaling	1.4
GO:0005794	Golgi apparatus	1.4
GO:1902235	Endoplasmic reticulum stress signaling	1.4
GO:0016020	Membrane components	1.4
GO:0098552	Cell surface	1.4
GO:0031410	Intracellular vesicles	1.3
GO:0098805	Membranes	1.3
GO:0051270	Cell migration	1.3

Gene ontology (GO) analysis, in which enrichment equals $-\log_{10}(p)$, where 1.3 is equivalent to $P=0.05$ and P represents the geometric mean of P -values in an annotation cluster. Only a description of the first term of each statistically significant (enrichment > 1.3) annotation cluster is shown. Full results are shown in corresponding Supplementary Data S1B. hESCs, human endometrial stromal cells.

on pools of ECM from embryos that had been removed from the culture dishes for cryopreservation and focused solely on hESCs migration as a measure for implantation. Obviously, the perspective of future clinical application requires further research into hsa-miR-320a and its downstream effects on embryo development and implantation.

Conclusion

In conclusion, we identified a soluble miRNA, hsa-miR-320a, that not only marks high-quality human preimplantation embryos but also directly stimulates decidualized hESCs migration *in vitro*. This makes hsa-miR-320a a unique and promising target for future research to ultimately improve pregnancy rates in IVF/ICSI.

Data availability

All sequence data have been submitted to the Sequence Read Archive (NCBI) and are available under the accession number PRJNA601880.

Supplementary data

Supplementary data are available at *Human Reproduction* online.

Acknowledgments

We thank Judith Huirne for providing endometrial biopsies.

Authors' roles

S.M., G.H. and R.P.B. conceived the study and designed the experiments. S.M. and G.H. supervised the project. R.P.B. performed the experiments and wrote the first draft of the manuscript. All authors reviewed the manuscript. G.B.A. and R.K. provided technical support and performed bioinformatics analysis.

Funding

The study was funded by the Amsterdam UMC and the Amsterdam Reproduction & Development Research Institute.

Conflict of interest

R.P.B., G.H. and S.M. have a patent on the use of hsa-miR-320a in assisted reproduction treatments pending.

References

- Abu-Halima M, Häusler S, Backes C, Fehlmann T, Staib C, Nestel S, Nazarenko I, Meese E, Keller A. Micro-ribonucleic acids and extracellular vesicles repertoire in the spent culture media is altered in women undergoing *in vitro* fertilization. *Sci Rep* 2017;**7**:13525.
- Balaban B, Brison D, Calderón G, Catt J, Conaghan J, Cowan L, Ebner T, Gardner D, Hardarson T, Lundin K et al. The Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Hum Reprod* 2011;**26**:1270–1283.
- Berkhout RP, Lambalk CB, Huirne J, Mijatovic V, Repping S, Hamer G, Mastenbroek S. High-quality human preimplantation embryos actively influence endometrial stromal cell migration. *J Assist Reprod Genet* 2018;**35**:659–667.
- Bidarimath M, Khalaj K, Kridli RT, Kan FW, Koti M, Tayade C. Extracellular vesicle mediated intercellular communication at the porcine maternal-fetal interface: a new paradigm for conceptus-endometrial cross-talk. *Sci Rep* 2017;**7**:40476.
- Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nat Rev Cancer* 2006;**6**:857–866.
- Capalbo A, Ubaldi FM, Cimadomo D, Noli L, Khalaf Y, Farcomeni A, Ilic D, Rienzi L. MicroRNAs in spent blastocyst culture medium are derived from trophectoderm cells and can be explored for human embryo reproductive competence assessment. *Fertil Steril* 2016;**105**:225–235.e221–e223.
- Cha J, Sun X, Dey SK. Mechanisms of implantation: strategies for successful pregnancy. *Nat Med* 2012;**18**:1754–1767.
- Cuman C, Van Sinderen M, Gantier MP, Rainczuk K, Sorby K, Rombauts L, Osianlis T, Dimitriadis E. Human blastocyst secreted microRNA regulate endometrial epithelial cell adhesion. *EBioMedicine* 2015;**2**:1528–1535.
- Ebner T, Moser M, Sommergruber M, Tews G. Selection based on morphological assessment of oocytes and embryos at different stages of preimplantation development: a review. *Hum Reprod Update* 2003;**9**:251–262.
- Esquela-Kerscher A, Slack FJ. Oncomirs—microRNAs with a role in cancer. *Nat Rev Cancer* 2006;**6**:259–269.
- Feng R, Sang Q, Zhu Y, Fu W, Liu M, Xu Y, Shi H, Xu Y, Qu R, Chai R et al. MiRNA-320 in the human follicular fluid is associated with embryo quality *in vivo* and affects mouse embryonic development *in vitro*. *Sci Rep* 2015;**5**:8689.
- Fletcher DA, Mullins RD. Cell mechanics and the cytoskeleton. *Nature* 2010;**463**:485–492.
- Fritz R, Jain C, Armant DR. Cell signaling in trophoblast-uterine communication. *Int J Dev Biol* 2014;**58**:261–271.
- Galliano D, Pellicer A. MicroRNA and implantation. *Fertil Steril* 2014;**101**:1531–1544.
- Gao T, Deng M, Wang Q. MiRNA-320a inhibits trophoblast cell invasion by targeting estrogen-related receptor-gamma. *J Obstet Gynaecol Res* 2018;**44**:756–763.
- Gellersen B, Brosens JJ. Cyclic decidualization of the human endometrium in reproductive health and failure. *Endocr Rev* 2014;**35**:851–905.
- Grewal S, Carver JG, Ridley AJ, Mardon HJ. Implantation of the human embryo requires Rac1-dependent endometrial stromal cell migration. *Proc Natl Acad Sci U S A* 2008;**105**:16189–16194.
- Gross N, Kropp J, Khatib H. Sexual dimorphism of miRNAs secreted by Bovine *In vitro*-produced embryos. *Frontiers in Genetics* 2017;**8**:doi:10.3389/fgene.2017.00039.
- Guo T, Feng Y, Liu Q, Yang X, Jiang T, Chen Y, Zhang Q. MicroRNA-320a suppresses in GBM patients and modulates glioma cell functions by targeting IGF-1R. *Tumour Biol* 2014;**35**:11269–11275.
- Ha M, Kim VN. Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol* 2014;**15**:509–524.

- Haouzi D, Dechaud H, Assou S, Monzo C, de Vos J, Hamamah S. Transcriptome analysis reveals dialogues between human trophectoderm and endometrial cells during the implantation period. *Hum Reprod* 2011;**26**:1440–1449.
- He L, Hannon GJ. MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet* 2004;**5**:522–531.
- Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 2009a;**37**:1–13.
- Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 2009b;**4**:44–57.
- Kropp J, Salih SM, Khatib H. Expression of microRNAs in bovine and human pre-implantation embryo culture media. *Front Genet* 2014;**5**:91.
- Li H, Yu L, Liu J, Bian X, Shi C, Sun C, Zhou X, Wen Y, Hua D, Zhao S et al. miR-320a functions as a suppressor for gliomas by targeting SND1 and β -catenin, and predicts the prognosis of patients. *Oncotarget* 2017;**8**:19723–19737.
- Liu J, Song Z, Feng C, Lu Y, Zhou Y, Lin Y, Dong C. The long non-coding RNA SUMO1P3 facilitates breast cancer progression by negatively regulating miR-320a. *Am J Transl Res* 2017;**9**:5594–5602.
- Lou W, Liu J, Gao Y, Zhong G, Chen D, Shen J, Bao C, Xu L, Pan J, Cheng J et al. MicroRNAs in cancer metastasis and angiogenesis. *Oncotarget* 2017;**8**:115787–115802.
- Lv G, Wu M, Wang M, Jiang X, Du J, Zhang K, Li D, Ma N, Peng Y, Wang L et al. miR-320a regulates high mobility group box 1 expression and inhibits invasion and metastasis in hepatocellular carcinoma. *Liver Int* 2017a;**37**:1354–1364.
- Lv Q, Hu JX, Li YJ, Xie N, Song DD, Zhao W, Yan YF, Li BS, Wang PY, Xie SY. MiR-320a effectively suppresses lung adenocarcinoma cell proliferation and metastasis by regulating STAT3 signals. *Cancer Biol Ther* 2017b;**18**:142–151.
- Macklon NS, Brosens JJ. The human endometrium as a sensor of embryo quality. *Biol Reprod* 2014;**91**:98.
- Macklon NS, Geraedts JPM, Fauser BCJM. Conception to ongoing pregnancy: the 'black box' of early pregnancy loss. *Hum Reprod Update* 2002;**8**:333–343.
- Marco A, Hooks K, Griffiths-Jones S. Evolution and function of the extended miR-2 microRNA family. *RNA Biol* 2012;**9**:242–248.
- McCallie B, Schoolcraft WB, Katz-Jaffe MG. Aberration of blastocyst microRNA expression is associated with human infertility. *Fertil Steril* 2010;**93**:2374–2382.
- Niakan KK, Han J, Pedersen RA, Simon C, Pera RA. Human pre-implantation embryo development. *Development* 2012;**139**:829–841.
- Peter Durairaj RR, Aberkane A, Polanski L, Maruyama Y, Baumgarten M, Lucas ES, Quenby S, Chan JK, Raine-Fenning N, Brosens JJ et al. Deregulation of the endometrial stromal cell secretome precedes embryo implantation failure. *Mol Hum Reprod* 2017;**23**:478–487.
- Puissant F, Van Rysselberge M, Barlow P, Deweze J, Leroy F. Embryo scoring as a prognostic tool in IVF treatment. *Hum Reprod* 1987;**2**:705–708.
- Qi X, Li J, Zhou C, Lv C, Tian M. MicroRNA-320a inhibits cell proliferation, migration and invasion by targeting BMI-1 in nasopharyngeal carcinoma. *FEBS Lett* 2014;**588**:3732–3738.
- Quenby S, Brosens JJ. Human implantation: a tale of mutual maternal and fetal attraction. *Biol Reprod* 2013;**88**:81.
- Rhee JK, Shin SY, Zhang BT. Construction of microRNA functional families by a mixture model of position weight matrices. *PeerJ* 2013;**1**:e199.
- Rosenbluth EM, Shelton DN, Wells LM, Sparks AE, Van Voorhis BJ. Human embryos secrete microRNAs into culture media—a potential biomarker for implantation. *Fertil Steril* 2014;**101**:1493–1500.
- van Loendersloot L, van Wely M, van der Veen F, Bossuyt P, Repping S. Selection of embryos for transfer in IVF: ranking embryos based on their implantation potential using morphological scoring. *Reprod Biomed Online* 2014;**29**:222–230.
- Van Royen E, Mangelschots K, De Neubourg D, Valkenburg M, Van de Meerssche M, Ryckaert G, Eestermans W, Gerris J. Characterization of a top quality embryo, a step towards single-embryo transfer. *Hum Reprod* 1999;**14**:2345–2349.
- Weimar CH, Kavelaars A, Brosens JJ, Gellersen B, de Vreeden-Elbertse JM, Heijnen CJ, Macklon NS. Endometrial stromal cells of women with recurrent miscarriage fail to discriminate between high- and low-quality human embryos. *PLoS One* 2012;**7**:e41424.
- Wilcox AJ, Weinberg CR, O'Connor JF, Baird DD, Schlatterer JP, Canfield RE, Armstrong EG, Nisula BC. Incidence of early loss of pregnancy. *N Engl J Med* 1988;**319**:189–194.
- Xishan Z, Ziyang L, Jing D, Gang L. MicroRNA-320a acts as a tumor suppressor by targeting BCR/ABL oncogene in chronic myeloid leukemia. *Sci Rep* 2015;**5**:12460.
- Yao J, Liang LH, Zhang Y, Ding J, Tian Q, Li JJ, He XH. GNAI1 suppresses tumor cell migration and invasion and is post-transcriptionally regulated by Mir-320a/c/d in hepatocellular carcinoma. *Cancer Biol Med* 2012;**9**:234–241.
- Yu J, Wang JG, Zhang L, Yang HP, Wang L, Ding D, Chen Q, Yang WL, Ren KH, Zhou DM et al. MicroRNA-320a inhibits breast cancer metastasis by targeting metadherin. *Oncotarget* 2016;**7**:38612–38625.
- Zhang Y, He X, Liu Y, Ye Y, Zhang H, He P, Zhang Q, Dong L, Dong J. microRNA-320a inhibits tumor invasion by targeting neuropilin 1 and is associated with liver metastasis in colorectal cancer. *Oncol Rep* 2012;**27**:685–694.
- Zhang Z, Li X, Sun W, Yue S, Yang J, Li J, Ma B, Wang J, Yang X, Pu M et al. Loss of exosomal miR-320a from cancer-associated fibroblasts contributes to HCC proliferation and metastasis. *Cancer Lett* 2017;**397**:33–42.
- Zhao H, Dong T, Zhou H, Wang L, Huang A, Feng B, Quan Y, Jin R, Zhang W, Sun J et al. miR-320a suppresses colorectal cancer progression by targeting Rac1. *Carcinogenesis* 2014;**35**:886–895.
- Zhao W, Sun Q, Yu Z, Mao S, Jin Y, Li J, Jiang Z, Zhang Y, Chen M, Chen P et al. MiR-320a-3p/ELF3 axis regulates cell metastasis and invasion in non-small cell lung cancer via PI3K/Akt pathway. *Gene* 2018;**670**:31–37.
- Zinaman MJ, Clegg ED, Brown CC, O'Connor J, Selevan SG. Estimates of human fertility and pregnancy loss. *Fertil Steril* 1996;**65**:503–509.